MARROW ED8

Medicine

Comprehensive Question Bank

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Contents

Chapter	Title	Page
1	Clinical examination	7
2	Acid-Base Disorders	27
3	General Principles of Endocrinology	50
4	Disorders of Anterior Pituitary	57
5	Pituitary Tumours and Sheehan's Syndrome	69
6	Posterior Pituitary - ADH, Oxytocin & Diabetes ...	80
7	Syndrome of Inappropriate Anti-Diuretic Hormone...	92
8	Thyroid Disorders - Clinical Features	98
9	Thyroid Disorders - Management	114
10	Multiple Endocrine Neoplasia	125
11	Pheochromocytoma	132
12	Cushing's Syndrome	141
13	Adrenal Insufficiency and Hyperaldosteronism	154
14	Diabetes Mellitus: Types, Clinical Features and...	168
15	Diabetes Mellitus: Complications and their Mana...	187
16	Reproductive Endocrinology	209
17	Disorders of Parathyroid and Calcium Homeostasis	219
18	Obesity	238
19	Hyperbilirubinemias and Tests of Liver Function	247
20	Alcoholic Liver Diseases and Non-Alcoholic Fatt...	263

Sold by @itachibot

Chapter	Title	Page
21	Viral Hepatitis	272
22	Autoimmune Disorders of Hepatobiliary System	292
23	Acute Liver Failure and Liver Transplantation	302
24	Hemochromatosis and Wilson's disease	315
25	Cirrhosis and Complications	330
26	Peptic Ulcer Disease and Related Disorders	350
27	Evaluation of Diarrhea	364
28	Irritable Bowel Syndrome	374
29	Inflammatory Bowel Disease - Clinical Features ...	383
30	Inflammatory Bowel Disease - Complications and ...	401
31	Malabsorption Syndromes	417
32	Other Gastrointestinal Conditions	437
33	Large and medium vessel vasculitis	449
34	Small vessel vasculitis	461
35	Crystal Arthropathies	473
36	Sjogren's Syndrome and Scleroderma	484
37	Sarcoidosis	500
38	Dermatomyositis and Related Disorders	511
39	Seronegative Spondyloarthritides	526
40	SLE and APS	540
41	Rheumatoid Arthritis	558
42	Asthma & COPD	573

Sold by @itachibot

Chapter	Title	Page
43	Pneumonia	595
44	Interstitial Lung Diseases and pneumoconiosis	615
45	Bronchiectasis and Lung Abscess	632
46	Pulmonary Function Tests	649
47	Respiratory Failure and ARDS	660
48	Neoplasms of the Lung	667
49	Sleep apnea	677
50	Urine Analysis	685
51	Chronic Kidney Disease	695
52	Acute Kidney Injury	714
53	Renal Replacement Therapy	731
54	Cystic and Inherited Disorders of the Kidney	748
55	Renal Tubular Disorders	759
56	Vascular Diseases of Kidney	770
57	Diagnosis of cardiovascular disorders	779
58	ECG Basics	803
59	Supraventricular Arrhythmias	829
60	Ventricular Arrhythmias and Heart Blocks	860
61	Valvular Heart Diseases	888
62	Infective Endocarditis and Acute rheumatic fever	907
63	Hypertensive Vascular Disease	928
64	Ischemic Heart Disease - Presentation and Diagn...	940

Sold by @itachibot

Chapter	Title	Page
65	Ischemic Heart Disease - Complications and Mana...	958
66	Cardiomyopathy and Myocarditis	976
67	Heart Failure	991
68	Pericardial Diseases	1006
69	Pulmonary Hypertension	1020
70	Diseases of Aorta and Peripheral Arteries	1025
71	DVT and Pulmonary Embolism	1036
72	General Neurology, Brain Death, and GCS	1052
73	Seizures and Epilepsy	1065
74	Ischemic Stroke	1087
75	Brainstem Syndromes and Hemorrhagic Stroke	1110
76	Peripheral Neuropathy	1132
77	Guillain-Barre Syndrome and Other Immune-mediat...	1142
78	Diseases of the Spinal Cord	1156
79	Myasthenia gravis and Other Neuromuscular Junct...	1170
80	Multiple Sclerosis & Other Demyelinating Disorders	1183
81	Headache Disorders	1194
82	Movement Disorders	1209
83	ALS and Other Motor Neuron Diseases	1234
84	Ataxic Disorders	1242
85	Cranial Nerve Disorders	1251
86	Infections of the Nervous System	1265

Sold by @itachibot

Chapter	Title	Page
87	Hypoproliferative Anemia	1276
88	Macrocytic Anemia	1295
89	Hemolytic Anemia	1308
90	Myeloproliferative Disorders and Aplastic Anemia	1328
91	Platelet Disorders	1343
92	Acute Leukemia	1367
93	Plasma Cell Disorders	1387
94	Chronic Myeloid Leukemia and Chronic Lymphoid L...	1403
95	Lymphomas	1415
96	Coagulation Disorders	1438
97	Haemoglobinopathies	1458
98	Blood Bank & Transfusion Medicine	1465
99	HIV / AIDS - Epidemiology and Clinical Features	1475
100	HIV / AIDS - Management and Prophylaxis	1493
101	COVID 19 - Epidemiology and Clinical Features	1508
102	COVID 19 - Investigations	1521
103	COVID 19 - Treatment	1535
104	COVID 19 - Prophylaxis and Complications	1547

Sold by @itachibot

Clinical examination

Question 1:

A soldier posted in Siachen is brought unconscious to the emergency ward. In order to rule out hypothermia, how will you best measure his temperature?

a) Oral

b) Rectal

c) Infrared tympanic thermometer

d) Axillary

Question 2:

A patient in the emergency ward stays still while lying supine and hesitates to move. He also has very shallow breathing. This attitude of the patient is most likely due to which of the following conditions?

a) Renal colic

b) Peritonitis

c) Acute pancreatitis

d) Meningitis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

An alcohol addict presented with easy fatiguability. On examination, he is found to have pallor. Peripheral smear showed macrocytes and anisocytosis. Which of the following findings is most likely to be seen in this patient?

a) Atrophic glossitis

b) Migratory glossitis

c) Strawberry tongue

d) Macroglossia

Question 4:

Sold by @itachibot

A woman presents with bluish discoloration of the skin and mucous membranes. All of the following may be responsible for the given symptoms except:

a) Methemoglobin exceeds 1.5 gm%

b) Sulfhemoglobin exceeds 0.5 gm%

c) Reduced hemoglobin exceeds 2 gm%

d) Oxygen saturation below 85%

Question 5:

Which of the following skin findings is not associated with an endocrine abnormality?

a) Acanthosis nigricans

b) Diffuse hyperpigmentation

c) Casal's necklace

d) Purple striae

Question 6:

What are petechiae?

a) Large extravasation of blood in the skin

b) Skin hemorrhages measuring 2-3 mm

c) Skin hemorrhages measuring 1-2 mm

d) Skin hemorrhages measuring 3-4 mm

Question 7:

A 28-year-old woman presented to the clinic with the lesions shown below. Which of the following conditions is/are associated with this lesion?

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) 1 and 5

b) 2, 3 and 5

c) 1, 3 and 4

d) 1, 2 and 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

You observed the following finding in a patient in the OPD. Which among the following conditions is least likely to cause this finding?

a) Endocarditis

Sold by @itachibot

b) Bronchiectasis

c) Tricuspid atresia

d) Aortic dissection

Question 9:

A 40-year-old patient presents with loss of appetite, jaundice, and easy bruisability. He is a chronic alcohol user, and the following abnormality is noted on examination. What is the most likely cause of this nail change?

a) Scurvy

b) Hypoalbuminemia

c) Chronic iron deficiency

d) Folate deficiency

Question 10:

A resident is examining a patient with bilateral leg swelling below the knees. He notes that on applying pressure, the indentation does not persist on release. What is the likely cause of the edema?

a) Myxedema

b) Nephrotic syndrome

c) Chronic lymphangitis

d) Acute lymphatic filariasis

Sold by @itachibot

Question 11:

While examining a patient's pulse, you note a tapping impulse when you raise his arm above his head. Which of the following conditions would you not suspect in the patient?

a) Left-sided heart failure

b) Anemia

c) Thiamine deficiency

d) Aortic regurgitation

Question 12:

True about the definition of Postural Hypotension:

a) Decrease in Systolic blood pressure 20 mmHg after 6 mins of standing

b) Decrease in Systolic blood pressure 20 mmHg after 3 min of standing

c) Decrease in Diastolic blood pressure 20 mmHg after 6 mins of standing

d) Decrease in diastolic blood pressure 20 mmHg after 3 mins of standing

Question 13:

A 30-year-old African farmer presents with Winterbottom's sign. This sign is due to the involvement of:

a) Deep cervical nodes

b) Supraclavicular nodes

c) Posterior cervical nodes

d) Occipital nodes

Question 14:

Relative bradycardia occurs in:

a) Typhoid fever

b) Q fever

c) Leptospirosis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) All of the above

Question 15:

A patient with rheumatic heart disease now presents with fever and the following nail finding. The nail lesions most likely are?



a) Roth spots

b) Janeway lesions

c) Osler nodes

d) Splinter hemorrhages

If you purchase
you may have
else,
d.

Question 16:

A 25-year-old man presented to the OPD with a fever for the past 5 days. On examination, tender lesions were found on his extremities as marked below. What is the most probable diagnosis?

Sold by @itachibot

a) Hand-foot-mouth disease

b) Infective endocarditis

c) Cat scratch disease

d) Rocky Mountain Spotted fever

Question 17:

On examination of the chest wall, the presence of thrills indicate:

a) Palpable S3

b) Palpable S4

c) Palpable murmur

d) Palpable right ventricular systole

Question 18:

As an intern in the emergency ward, which of the following method would you consider as correct regarding nasogastric tube length estimation?

a) Tip of nose to ear to xiphisternum

b) Tip of nose to angle of ear to umbilicus

c) Mouth to ear to umbilicus

d) Mouth to ear to midway between xiphisternum and umbilicus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 19:

A patient presents with excessive thirst, unexplained weight loss, visual blurring, genital thrush, and lethargy. What is the most likely diagnosis?

a) Hyperglycemia

b) Hyperthyroidism

c) Hypoglycemia

d) Diabetic neuropathy

Question 20:

The recommended temperature of water bath for a patient with frost bite is

a) 37 degree C

b) 42 degree C

c) 32 Degree C

d) 30 degree C

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	c
5	c
6	c
7	d
8	d
9	b
10	a
11	a
12	b
13	c
14	d

Sold by @itachibot

15	d
16	b
17	c
18	a
19	a
20	a

Detailed Explanations

Solution to Question 1:

In the given clinical scenario, the rectal temperature would be the best site to measure the core body temperature.

At extremes of high or low temperatures, like in Siachen, the axillary and the oral temperatures are highly misleading and should not be used to measure the core body temperature.

The infrared tympanic thermometer measures the temperature of the eardrum which is representative of the core body temperature. However, it often ends up measuring the temperature of the ear canal. This is because of inadequate insertion into the ear canal due to poor manufacturing design or the presence of cerumen. Hence, it is not a reliable method of measuring the core body temperature.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 2:

The given clinical scenario is suggestive of peritonitis. A patient with peritonitis lies supine without movement and has shallow breathing as any movement exacerbates the pain.

Option A: In renal colic, patients have a restless attitude, often rolling around because of pain.

Option C: In acute pancreatitis, the patient usually bends forward and assumes a knee-chest position because of abdominal pain.

Option D: Patients with meningitis present with neck stiffness.

Solution to Question 3:

The given clinical scenario is suggestive of megaloblastic anemia likely due to folate deficiency caused by alcohol addiction. Folate deficiency leads to atrophic glossitis because of damage to the epithelial surfaces.

Sold by @itachibot



Option B: Migratory glossitis / geographic tongue is due to rapid loss and regrowth of filiform papillae of the tongue. It is a benign inflammatory condition of the tongue associated with psoriasis and reactive arthritis.

Option C: Strawberry tongue is due to the hypertrophy of fungiform papillae. It is seen in scarlet fever and Kawasaki disease.

Sold by @itachibot

Option D: Macroglossia is an abnormal enlargement of the tongue and can be congenital or acquired. New-onset macroglossia in adults is highly suggestive of amyloidosis.

Solution to Question 4:

Bluish discoloration of the skin and mucous membranes is suggestive of cyanosis. Cyanosis is seen when the concentration of deoxygenated or reduced hemoglobin exceeds 4 gm/dL.

Cyanosis is detectable with methemoglobin ≥1.5 gm% and/or sulfhemoglobin ≥0.5 gm%.

Central cyanosis is detectable when the oxygen saturation <85%. In dark-skinned individuals, it may be detectable at an oxygen saturation of 75%.

The types of cyanosis are:

Sold by @itachibot

Central cyanosis	Peripheral cyanosis
Due to diminished arterial oxygen saturation/presence of abnormal hemoglobin derivative.	Due to diminished peripheral blood flow.
Mucous membranes and skin are both affected	Mucous membranes of the oral cavity or those beneath the tongue are spared.
Decreases after O2supplementation	Does not decrease after O2supplementation

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

Casal's necklace is not associated with an endocrine abnormality. It is a feature of pellagra, which occurs due to the deficiency of niacin.

Option A: Acanthosis nigricans is seen in multiple endocrinopathies like acromegaly, Cushing's syndrome, polycystic ovary syndrome, and insulin-resistant diabetes mellitus.

Option B: Diffuse hyperpigmentation is seen in Addison's disease and ectopic ACTH syndrome.

Option D: Purple striae are seen in Cushing's disease.

Solution to Question 6:

Petechiae are defined as skin hemorrhages appearing as minute non-blanching spots that measure less than 2 mm.

A few conditions causing petechiae have been stated below:

• Infectious diseases: Cytomegalovirus (CMV) infection, infective endocarditis, meningococcemia, and viral hemorrhagic fevers.

• Non-infectious conditions: Vasculitis, thrombocytopenia, leukemia, scurvy, and vitamin K deficiency

• Drugs: Anticoagulants (warfarin, heparin), quinine, etc.

Large extravasation of blood in the skin is known as ecchymosis.

Purpura are also non-blanching spots but unlike petechiae, they measure ≥ 3 mm in size.

Solution to Question 7:

The lesions shown in the image are spider naevi (nevus araneus). They are seen in:

• Estrogen excess states such as pregnancy and OCP use

Sold by @itachibot

• Chronic liver disease - especially alcoholic liver disease

• Thyrotoxicosis

They are vascular lesions characterized by the tortuous dilatation of arterioles just underneath the skin. On applying and releasing pressure over the lesion they typically fill from the centre outward.

Solution to Question 8:

The given image shows digital clubbing. Among the given options, aortic dissection is least likely to cause digital clubbing.

Clubbing is a selective bulbous enlargement of the distal segments of the fingers and toes due to the proliferation of connective tissue, particularly on the dorsal surface.

Causes of clubbing are:

• Pulmonary

• Bronchogenic carcinoma

• Lung abscess, empyema, bronchiectasis

• Interstitial lung diseases

• Cystic fibrosis

• Sarcoidosis

• Asbestosis

• GI

• Inflammatory bowel disease (Ulcerative colitis, Crohn's disease)

• Sprue, celiac disease

• Neoplasms - esophagus, liver, bowel

• Cirrhosis - juvenile and biliary

• Chronic active hepatitis

• Cardiovascular

• Cyanotic congenital heart disease

• Subacute bacterial endocarditis

• Aortic aneurysm

• Arteriovenous fistula

• Patent ductus arteriosus with right to left shunt

• Atrial myxoma

• Endocrine

• Grave's disease (thyroid acropachy)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• CNS

• Hemiplegia (unilateral clubbing)

• Skin disease

• Palmoplantar keratoderma

• Fischer-Volavsek syndrome.

• Malignancies

• Thyroid cancer

• Thymus cancer

• Hodgkin disease

• Disseminated chronic myeloid leukemia.

Note: Chronic obstructive pulmonary disease (COPD) does not cause clubbing. If clubbing is present in a patient with COPD, underlying lung cancer and bronchiectasis must be ruled out.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

The nail change seen in the given image is leukonychia totalis. It is a prominent feature of hypoalbuminemic states. The given clinical scenario is suggestive of liver cirrhosis, which leads to hypoalbuminemia.

Leukonychia is an opaque whitish discoloration of the nail and is seen in the following conditions:

• Dietary deficiency: Protein-energy malnutrition, pellagra, zinc deficiency

• Drugs: Cytotoxic agents, Steroids, Retinoids, Pilocarpine

• Dermatological pathology: Alopecia areata, erythema multiforme, exfoliative dermatitis, and SLE

• Systemic diseases: Liver cirrhosis, renal failure, and cardiac failure

• Miscellaneous: Trauma, malignancy, and toxins (arsenic, fluoride)

• Idiopathic

Note: The leukonychia secondary to arsenic poisoning is termed leukonychia striata or Aldrich–Mees lines.

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Bilateral non-pitting edema in the lower limbs is suggestive of myxedema.

While chronic lymphangitis and acute lymphatic filariasis can also cause non-pitting edema, they usually have a unilateral/localized presentation.

Nephrotic syndrome and heart failure cause bilateral pitting edema.

Causes of generalized edema include:

- Heart failure
- Chronic renal failure
- Nephrotic syndrome
- Liver cirrhosis

Causes of localized edema include:

- Thrombophlebitis
- Chronic lymphangitis
- Resection of regional lymph nodes
- Filariasis

Solution to Question 11:

The given clinical scenario is suggestive of water hammer pulse, which is not seen in left-sided heart failure.

Sold by @itachibot

Collapsing pulse/Watson's water hammer pulse is a rapidly rising pulse that collapses suddenly. This is because arterial pressure falls rapidly during late systole and diastole. It is also called Corrigan's pulse.

The various causes of the collapsing pulse include

• Physiologically, it can be present in fever and pregnancy. This is attributed to the combination of increased cardiac output and the reduction in systemic vascular resistance.

• Cardiac lesions such as aortic regurgitation, patent ductus arteriosus, systolic hypertension, marked bradycardia, aneurysm of sinus of Valsalva

• High-output states such as anemia, cor pulmonale, thyrotoxicosis, arteriovenous fistula, Paget's disease, beriberi, and liver cirrhosis (secondary to chronic alcoholism)

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 12:

Orthostatic hypotension or postural hypotension is defined as a sustained drop in systolic BP of ≥20 mmHg or diastolic BP of ≥10 mmHg after 3 min of standing.

It can manifest as lightheadedness/dizziness, weakness or tiredness, difficulty thinking/concentrating, blurred vision, and anxiety.

In non-neurogenic causes of orthostatic hypotension such as hypovolemia, the drop in the blood pressure is accompanied by a compensatory increase in heart rate of >15 beats/min. In neurogenic causes, there is aggravation or precipitation of orthostatic hypotension by autonomic stressors like a meal, hot bath, or exercise.

Non-neurogenic causes of orthostatic hypotension include

• Cardiac pump failure - myocardial infarction, myocarditis, aortic stenosis

• Hypovolemia - dehydration, diarrhea, hemorrhage

• Metabolic - adrenocortical insufficiency, pheochromocytoma

• Venous pooling - sepsis, prolonged standing, or recumbency

• Medications - alpha-blockers, diuretics, phenothiazines, tricyclic antidepressants

Treatment includes removing reversible causes like drugs causing hypotension.
Non-pharmacological interventions include patient education regarding staged moves from supine to upright, raising the head of the bed, and increasing dietary fluid and salt. If these fail, drugs like fludrocortisone and midodrine can be given.

Solution to Question 13:

Winterbottom's sign is due to the involvement of posterior cervical nodes. It is seen in African trypanosomiasis.

African trypanosomiasis is a vector born disease caused by Trypanosoma brucei gambiense/rhodesiense. It is transmitted through the bite of a Tsetse fly.

Sold by @itachibot

The clinical syndrome includes hemolymphatic disease with associated Winterbottom's sign and a maculopapular rash primarily involving the trunk. At a later stage, the patient can present with features suggestive of meningoencephalitis.

Solution to Question 14:

In the given options typhoid, Q fever, and leptospirosis are infectious causes of relative bradycardia.

Relative bradycardia, also known as Faget's sign, is observed when patients exhibit a lower heart rate than anticipated in relation to an increase in body temperature. Body temperature is commonly regarded as a factor that independently influences heart rate. It is estimated that for each 1°C (1.8°F) rise in temperature, the heart rate typically increases by approximately 15-20 beats per minute.

While this pulse-temperature dissociation is not highly sensitive or specific for establishing a diagnosis, it can still prove beneficial in resource-limited settings due to its easy accessibility and simplicity.

Infectious causes of relative bradycardia include:

- Salmonella typhi
- Francisella tularensis
- Coxiella burnetii (Q fever)
- Leptospira interrogans
- Legionella pneumophila
- Rickettsia spp.
- Orientia tsutsugamushi (scrub typhus)
- Corynebacterium diphtheriae
- Plasmodium spp. (malaria)

Noninfectious causes include:

- Drug fever
- Beta-blocker use
- Central nervous system lesions
- Malignant lymphoma
- Factitious fever

Solution to Question 15:

The given clinical scenario is suggestive of infective endocarditis (IE). Splinter hemorrhages are seen in the nails of these patients.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Subungual splinter hemorrhages are classically identified as dark-red, linear petechiae in the nail bed.

Other peripheral manifestations of infective endocarditis include the following:

• Petechiae: Common but nonspecific finding

• Purulent purpura

• Osler nodes: Tender subcutaneous nodules usually found on the distal pads of the digits

• Janeway lesions: Nontender maculae on the palms and soles

• Roth spots: Relatively rare retinal hemorrhages with small, clear centers

Solution to Question 16:

The most probable diagnosis for this patient is infective endocarditis.

Tender violaceous subcutaneous nodules in the fingers shown in the image are Osler nodes. They usually occur in the pads of fingers and toes but can also occur on the thenar and hypothenar eminences.

Other lesions that are highly suggestive of infective endocarditis include:

• Roth spots: These are exudative, edematous hemorrhagic lesions of the retina with pale centres.

• Janeway lesions: They are nontender, slightly raised haemorrhages on the palms and soles. The image is shown below.

Solution to Question 17:

On examination of the chest wall, the presence of thrills indicates a palpable murmur. Thrills are the palpable vibrations produced by turbulent blood flow causing murmurs.

Sold by @itachibot
If you paid this from someone else, you have been scammed

Sold by @itachibot

Palpable S3 and S4 cause a double thrust in the apical impulse.

Palpable right ventricular systole is known as heave which is felt in the left parasternal area.

Solution to Question 18:

The commonly used method to determine the length of a nasogastric tube is the NEX method: the tip of the nose to the ear to the xiphisternum.

The approximate length is 120-125 cm. The markings indicate the traversing of the tube through various parts of the GIT:

• 40 cm: Cardia of the stomach

• 50 cm: Body of the stomach

• 60 cm: Pylorus

• 70 cm: Duodenum

A nasogastric tube or Ryle's tube is inserted in the sitting position with the neck slightly flexed. The tube is then gently inserted through the nostril and advanced until it reaches 10 cm. The patient is encouraged to swallow, allowing the tube to move down the esophagus with successive swallows until it reaches the measured distance.

Subjective confirmation involves insufflating air through the proximal opening of the NG tube and listening for the sound of air in the left upper quadrant where the stomach is located. Objective confirmation can be obtained through pH measurement, chest or abdominal radiographs, ultrasound, or calorimetric capnometry.

While widely practiced, blind NG tube insertion can lead to complications such as coiling of the tube in the nasopharynx, incorrect placement in the bronchial passage, or even esophageal perforation.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 19:

The most likely diagnosis in the given scenario is hyperglycemia.

The symptoms of hyperglycemia include:

• Polyuria

• Polydipsia

• Weight loss

• Fatigue

• Blurry vision

• Frequent superficial infections (vaginitis, fungal skin infections)

• Slow healing of skin lesions after minor trauma.

Sold by @itachibot

Note: The metabolic derangements occurring in these patients are due to osmotic diuresis and a catabolic state caused by urinary loss of glucose and increased muscle breakdown. Blurred vision results from changes in the water content of the lens.

Solution to Question 20:

The temperature of the water bath for a patient with frostbite should be maintained between 37-40°C (Option A). Circulating water is preferred for treatment.

In frostbite, tissue damage results from severe environmental cold exposure or from direct contact with a very cold object. Tissue injury results from both freezing and vasoconstriction. It usually affects the distal aspects of the extremities or exposed parts of the face, such as the ears, nose, chin, and cheeks.

The treatment includes - extraction from cold environment, rapid rewarming, analgesics for pain during thawing, antithrombotics or thrombolytics, cleaning of the injured area with soap or antiseptics, tetanus prophylaxis, monitoring for compartment syndrome and finally, elective late amputation after imaging in severe cases.

@itachibot

Sold by @itachibot

Acid-Base Disorders

Question 1:

Which of the following is the most effective extracellular buffer system at physiological pH?

a) Bicarbonate buffer

b) Phosphate buffer

c) Protein buffer

d) Ammonia buffer

Question 2:

Which of the following is not done before drawing blood for arterial blood gas analysis?

a) Allen's test

b) Rinse syringe with heparin

c) Flexion of wrist

d) Placing needle at 45 degree angle

Question 3:

Which of the following is not a cause of metabolic alkalosis?

a) Mineralocorticoid deficiency

b) Hypokalemia

c) Thiazide diuretic therapy

d) Recurrent vomiting

Question 4:

Which metabolic abnormality are patients with bulimia nervosa least likely to develop?

a) Hyponatremia

b) Hypochloremia

c) Metabolic alkalosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) Metabolic acidosis

Question 5:

Which of the following is false about metabolic alkalosis?

a) It is associated with hyperkalemia

b) It is associated with decreased ionic calcium concentration

c) Can be caused due to Conn's syndrome

d) Can be caused due to a renin secreting tumor

Question 6:

Metabolic alkalosis is suspected in a patient presenting with multiple episodes of vomiting. Which of the following laboratory results would be seen in his arterial blood gas analysis?

a) Option 1

b) Option 3

c) Option 2

d) Option 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

The arterial blood gas analysis of a patient reveals a pH of 7.55, pCO2- 54 mmHg, and HCO3- 40 mEq/L. What is the underlying disorder?

a) Mixed acid-base disorder

b) Metabolic alkalosis with compensatory respiratory acidosis

c) Respiratory acidosis with compensatory metabolic alkalosis

d) Respiratory alkalosis with compensatory metabolic acidosis

Question 8:

The arterial blood gas analysis of a 50-year-old diabetic patient revealed the following values: pH- 7.2 and HCO3- 15 mEq/L. Which of the following statements is false regarding renal compensation in this patient?

a) Hydrogen ion secretion is increased

Sold by @itachibot

b) Bicarbonate reabsorption is decreased

c) Urinary acidity is increased

d) Urinary ammonia is increased

Question 9:

Calculate the anion gap from the following values: Na+ = 137 mmol/L K+ = 4 mmol/L Cl- = 100 mmol/L HCO3- = 15 mmol/L

a) 22 mmol/L

b) 16 mmol/L

c) 10 mmol/L

d) 12 mmol/L

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

In which of the following patients would you expect to find an increased anion gap?

a) Renal tubular acidosis

b) Addison's disease

c) Diarrhoea

d) Diabetic ketoacidosis

Question 11:

High anion gap acidosis with low bicarbonate levels is seen in all the following conditions except:

a) Renal tubular acidosis

b) Diabetic ketoacidosis

c) Lactic acidosis

d) Salicylate poisoning

Question 12:

A 40-year-old man presented with nausea, dizziness, and hyperventilation. His arterial blood gas analysis and serum electrolyte levels revealed metabolic acidosis with a normal anion gap.

Sold by @itachibot

Which of the following is the possible reason for this condition?

a) Lactic acidosis

b) Chronic renal failure

c) Renal tubular acidosis

d) Methanol toxicity

Question 13:

Which of the following is not associated with type-A lactic acidosis?

a) Acute pyelonephritis with hypotension

b) Acute MI with free wall rupture

c) Cyanide toxicity

d) Metformin toxicity

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

A patient with chronic kidney disease came with complaints of vomiting and diarrhea. His blood gas reports show pH = 7.40, HCO_3^- = 23 mEq/L, Na^+ = 145 mEq/L, Cl^- = 100 mEq/L. What is your probable diagnosis?

a) No acid base abnormality

b) Respiratory acidosis

c) Metabolic alkalosis

d) High anion gap metabolic acidosis with metabolic alkalosis

Question 15:

A 5-year-old boy presents with generalized edema. Urine analysis showed protein 3+ and fat bodies. Which of the following would you expect to see in this condition?

a) Decrease in unmeasured anions

b) Decrease in measured anions

c) Normal anion gap

d) High anion gap

Sold by @itachibot

Question 16:

A 54-year-old male patient with renal failure has missed a round of dialysis. He complains of lethargy, anorexia, and vomiting. What is the probable acid-base disorder that we can expect in him?

a) High anion gap metabolic acidosis

b) Metabolic alkalosis with compensatory respiratory acidosis

c) Normal anion gap metabolic acidosis

d) Mixed acid-base disorder

Question 17:

ABG analysis of a patient is shown below. What is the probable underlying acid-base disorder in this patient?

a) Metabolic acidosis with compensatory respiratory alkalosis

b) Metabolic alkalosis with compensatory respiratory acidosis

c) Respiratory acidosis with compensatory metabolic alkalosis

d) Mixed acid–base disorder

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Question 18:

A 30-year-old man presents to the emergency department with dyspnea, and altered sensorium. These symptoms were preceded by an asthma attack. What is the probable reason for the patient's presenting symptoms?

a) Metabolic acidosis

b) Metabolic alkalosis

c) Respiratory acidosis

d) Respiratory alkalosis

Question 19:

A 60-year-old man with a history of chronic obstructive pulmonary disease presents with increased shortness of breath. His ABG analysis is as shown. What is the most probable underlying acid-base disorder?

a) Metabolic acidosis

Sold by @itachibot

b) Metabolic alkalosis

c) Respiratory acidosis

d) Respiratory alkalosis

Question 20:

A young asthmatic boy was rushed to ED with confusion and gasping for breath. His ABG analysis is as shown. What is the acid-base disorder?

a) Respiratory alkalosis, compensated

b) Respiratory acidosis, partially compensated

c) Respiratory acidosis, fully compensated

d) Metabolic alkalosis, uncompensated

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 21:

The arterial blood gas findings of a patient are pH = 7.12, pCO_2 = 50, HCO_3 = 28. What is the diagnosis?

a) Respiratory acidosis with metabolic compensation

b) Metabolic acidosis with respiratory compensation

c) Respiratory alkalosis with metabolic compensation

d) Metabolic alkalosis with respiratory compensation

Question 22:

A 40-year-old woman was rushed to the ER with chest pain, palpitations, and sweating. On examination, her respiratory rate was 32/min, and carpopedal spasm was noted while measuring BP. Her ECG was normal. Which of the following acid-base disorders is she likely to have?

a) Metabolic acidosis

b) Metabolic alkalosis

c) Respiratory acidosis

d) Respiratory alkalosis

Question 23:

Sold by @itachibot

In the given acid-base nomogram diagram, areas marked A and D indicate which of the following conditions?



a) Chronic respiratory acidosis, acute metabolic alkalosis

b) Acute Metabolic alkalosis, acute respiratory acidosis

c) Acute respiratory acidosis, acute respiratory alkalosis

d) Acute respiratory acidosis, acute metabolic alkalosis

Question 24:

The arterial blood gas analysis findings of a patient are as follows: pH= 7.30, HCO3= 10 mEq/L, paCO2= 30 mmHg, which is suggestive of a partially compensated metabolic acidosis. What is the mechanism for the secondary disorder?

a) Hypoventilation causing decreased CO2 washout

b) Hyperventilation causing increased CO2 washout

c) Increased tubular reabsorption of HCO3-

d) Increased excretion of H+

Question 25:

The arterial blood gas analysis of an unconscious man showed pH- 6.9, HCO3- 13 mEq/L, paCO2- 68 mmHg. The acid-base status of this patient is best described as:

a) Uncompensated metabolic acidosis

Sold by @itachibot
If you purchased this from elsewhere, you may have been scammed.

Sold by @itachibot

b) Uncompensated respiratory acidosis

c) Simultaneous respiratory and metabolic acidosis

d) Respiratory acidosis with partial renal compensation

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	d
5	a
6	d
7	b
8	b
9	a
10	d
11	a
12	c
13	d
14	d
15	a
16	a
17	a
18	c
19	c
20	b
21	a
22	d
23	c
24	a
25	c

Detailed Explanations

Sold by @itachibot

Solution to Question 1:

The most effective extracellular buffer system at physiological pH is the bicarbonate buffer system.

A molecule with a pKa close to the physiological pH (7.4) is considered an effective buffer. Although the pKa of HCO_3^- is relatively low (6.1), it makes up 92% of the plasma buffering.

This is often attributed to the enormous quantity of HCO_3^- present in the body and the ability of this system to accept protons and form carbonic acid (H_2CO_3). This further in the presence of carbonic anhydrase is converted to readily excretable H■O and CO_2. It is therefore considered the most efficient extracellular buffering system in the body.

$HCO_3^- + H^+ \rightleftharpoons H_2CO_3 \rightleftharpoons$ H■O + CO_2

Sold by @itachibot
If you found this from someone else, it has been stolen.

Solution to Question 2:

Flexion of the wrist is not done just before drawing blood for ABG analysis.

During arterial blood gas analysis, the ideal position for drawing arterial blood from the radial artery is to maintain the wrist in extension. This provides stability to the artery and prevents it from slipping away during the puncture. To draw blood for ABG analysis, the skin is pierced at a 45° angle.



Allen's test is done prior to radial cannulation to identify the adequacy of radial and ulnar collateral blood flow to the hand.

Sold by @itachibot



The syringe used to draw blood for ABG analysis is rinsed with heparin prior to the procedure. Prior to collecting blood, 0.1 ml heparin is drawn into the syringe and then flushed out. The residue that lubricates the inner wall of the syringe is sufficient to prevent the blood from clotting.

Sold by @itachi... if you purchased this from some... you may have been scammed. else,

Solution to Question 3:

Mineralocorticoid deficiency does not cause metabolic alkalosis. Its excess causes metabolic alkalosis.

Causes of metabolic alkalosis:

Loss of hydrogen:

- Gastrointestinal loss
- Removal of gastric secretions - vomiting or nasogastric suction
- Chloride-losing diarrhea
- Gastrocolic fistula
- Villous adenoma
- Antacid therapy, particularly if combined with cation exchange resin
- Renal loss
- Loop or thiazide diuretics
- Mineralocorticoid excess (primary aldosteronism, Cushing's, steroids, licorice)
- Posthypercapnic states
- Hypercalcemia, including milk-alkali syndrome
- H+ movement into cells

Sold by @itachibot

• Hypokalemia

Exogenous Alkali:

• Administration of $NaHCO_3$, sodium citrate/gluconate/acetate/antacids

• Massive blood transfusion

• Antacids- Milk alkali syndrome

Contraction alkalosis (loss of ECF volume):

• Loop or thiazide-type diuretics

• Sweat losses in cystic fibrosis

• Gastric losses in achlorhydria

Miscellaneous:

• Bartter's syndrome

• Gitelman's syndrome

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

The metabolic abnormality patients with bulimia nervosa are least likely to develop is metabolic acidosis. Chronic vomiting, as seen in bulimia nervosa leads to metabolic alkalosis, not acidosis.

Conditions associated with chronic vomiting lead to hypochloremia, hyponatremia, hypokalemia, and metabolic alkalosis.

Solution to Question 5:

Metabolic alkalosis is associated with potassium depletion, not hyperkalemia.

While hypercalcemia can be a cause of metabolic alkalosis, alkalosis itself can cause a decrease in ionized calcium. The patients can present with tetany and show signs of hypocalcemia, i.e., Trousseau sign and Chvostek's sign.

Conn's syndrome is primary hyperaldosteronism that leads to sodium reabsorption in exchange for hydrogen in the collecting ducts. This causes hypernatremia, hypokalemia, and hydrogen loss in urine and may lead to metabolic alkalosis.

Renin-secreting tumors also lead to an increase in aldosterone levels leading to alkalosis.

Causes of metabolic alkalosis:

Loss of hydrogen:

• Gastrointestinal loss

• Removal of gastric secretions- vomiting or nasogastric suction

• Chloride-losing diarrhea

• Gastrocolic fistula

Sold by @itachibot

• Villous adenoma

• Antacid therapy, particularly if combined with cation exchange resin

• Renal loss

• Loop or thiazide diuretics

• Mineralocorticoid excess (primary aldosteronism, Cushing's, steroids, licorice)

• Posthypercapnic states

• Hypercalcemia, including the milk-alkali syndrome

•

• H+ movement into cells

• Hypokalemia

Exogenous Alkali:

• Administration of $NaHCO_3$, sodium citrate/gluconate/acetate/antacids

• Massive blood transfusion

• Antacids- Milk alkali syndrome

Contraction alkalosis (loss of ECF volume):

• Loop or thiazide-type diuretics

• Sweat losses in cystic fibrosis

• Gastric losses in achlorhydria

Miscellaneous:

• Bartter's syndrome

• Gitelman's syndrome

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 6:

Among the given options, high pH (7.55), high HCO_3^- (30 mEq/L), and high pCO_2 (48 mmHg) is suggestive of metabolic alkalosis with respiratory acidosis, as a result of compensatory hypoventilation.

Multiple episodes of vomiting result in loss of acids from the body and thus increases the pH.

	pH	pCO_2(mmHg)	HCO_3^-(mEq/L)	Interpretation
1	7.48(high)	32(low)	21(low)	Respiratory alkalosis with compensatory metabolic acidosis
2	7.32(low)	30(low)	22(low)	Metabolic acidosis with compensatory respiratory alkalosis

Sold by @itachibot

	pH	pCO_2(mmHg)	HCO_3^-(mEq/L)	Interpretation
3	7.28(low)	50(high)	32(high)	Respiratory acidosis with compensatory metabolic alkalosis
4	7.55(high)	48(high)	30(high)	Metabolic alkalosis with compensatory respiratory acidosis

Solution to Question 7:

The underlying disorder in this patient is metabolic alkalosis with compensatory respiratory acidosis.

Interpretation of the ABG:

Step 1: A pH of 7.55 indicates alkalosis.

Step 2: Plasma HCO_3- is 40 mEq/L, which in the presence of a high pH indicates metabolic alkalosis.

Step 3: pCO_2 is 54 mmHg indicating respiratory acidosis. The increase in pCO_2 along with the increase in pH suggests that this is a compensatory response.

Step 4: Assessing the compensatory response:

On average, the pCO_2 rises 0.75 mmHg for every 1.0 mEq/L increment in the plasma HCO_3-

In the above scenario, the plasma HCO_3- concentration is increased to 40 mEq/L (The average normal value is considered as 24 mEq/L)

Thus, the HCO_3- elevation is 40 – 24 = 16 mEq/L

Therefore, pCO_2 elevation should be 0.75 × 16 = 12 mmHg

Now, considering that the average value of normal pCO_2 is 40 mmHg, the estimated pCO_2 measured should be 40 + 12= 52 mmHg.

As pCO_2 is higher than the expected estimate, it is a case of metabolic alkalosis with compensatory respiratory acidosis.

Solution to Question 8:

The patient has low pH and decreased levels of bicarbonate which is indicative of metabolic acidosis. Bicarbonate reabsorption is increased in renal handling of metabolic acidosis.

In response to metabolic acidosis, acidification of urine occurs by the following compensatory mechanisms:

• The proximal tubule reabsorbs all the filtered HCO_3- (as any loss of HCO_3- is equal to the addition of an equimolar amount of H^+).

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• The collecting duct excretes the daily H+ load (loss of H+ is equal to the addition of an equimolar amount of HCO3-) primarily in the form of urinary ammonia.

Solution to Question 9:

The anion gap is the difference between measured cations and measured anions and is estimated as:

Anion gap = [Na+] – ([Cl-] + [HCO3-])

Substituting the given values, AG is 22 mmol/L. Normal AG is 6 to 12 mmol/L with an average of 10 mmol/L.

The concentrations of anions and cations in plasma must be equal to maintain electrical neutrality. Therefore, there is no real “anion gap” in the plasma.

However, only certain cations and anions are routinely measured in the clinical laboratory. The cation normally measured is Na+, and the anions are usually Cl– and HCO3–. The value of plasma potassium is omitted from the calculation of the anion gap.

Usually, the unmeasured anions exceed the unmeasured cations.

The anion gap will increase if unmeasured anions rise or if unmeasured cations fall.

The unmeasured anions include albumin, phosphate, sulfate, and other organic anions. The unmeasured cations include calcium, magnesium, and potassium.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 10:

The anion gap is increased in ketoacidosis. The ketone bodies act as unmeasured anions and increase the anion gap.

The anion gap is calculated as:

AG = (Na+) – [(Cl-) + (HCO3-)]

The normal anion gap is 6 - 12 mEq/L (optimally <10 mEq/L).

Solution to Question 11:

Renal tubular acidosis causes normal anion gap metabolic acidosis, not high anion gap acidosis.

Normally, in the serum, the total number of cations is equal to the total number of anions. But, a few of the anionic proteins (e.g., albumin), phosphates, sulfates, and organic anions are not measured and this is called an anion gap. It is the difference between measured cations and measured anions and is estimated as:

Plasma anion gap = [Na+] –([Cl-]+[HCO3-])

The normal anion gap ranges between 6-12 mEq/L.

Sold by @itachibot

Solution to Question 12:

Metabolic acidosis with a normal anion gap is a feature of renal tubular acidosis.

Renal tubular acidosis is a form of hyperchloraemic metabolic acidosis which occurs when the renal damage primarily affects tubular function without much effect on glomerular function. The bicarbonate loss leading to acidosis is mirrored by an increase in chloride concentration. Hence the unmeasured anions are unaffected and thus, the anion gap remains within normal limits.

The result is a decrease in H+ excretion which is greater than that can be explained by any change in the glomerular filtration rate (GFR). However, if the GFR is significantly depressed eg., renal failure, the retention of fixed acids leads to a high anion gap metabolic acidosis.

If you purchased this from someone else, Sold by @itachibot may have been scammed.

Solution to Question 13:

Metformin toxicity is associated with type B lactic acidosis.

Lactic acidosis is defined as an increase in plasma lactate levels. The various types of lactic acidosis are:

D-lactic acidosis is an unusual form of metabolic acidosis. It occurs due to the formation of D-lactate by gut bacteria. It occurs in patients with:

- Short bowel syndrome
- Post-jejunoileal bypass
- Intestinal obstruction

Type A	Type B
Occurs secondary to circulatory insufficiency or tissue hypoxia.	Occurs in the absence of systemic hypoperfusion.
Associated with:ShockCardiac failureSevere anemiaCarbon monoxide toxicityCyanide toxicity	Associated with:Malignancies Renal failureHepatic failureDrugs like metformin, ethanol, ethylene glycol, and methanol

Solution to Question 14:

The probable diagnosis is a high anion gap metabolic acidosis (HAGMA) with metabolic alkalosis.

Interpretation of the ABG:

Step 1: The pH being 7.4 is within normal limits.

Sold by @itachibot

Step 2: Plasma HCO3- is 23 mEq/L and again is within normal limits.

At this juncture, we have ruled out options B and C as this particular clinical scenario is not suggestive of a simple acid-base disorder.

Step 3: Calculating the Anion gap:

Anion gap = [Na+] – ([Cl-] + [HCO3-])

= 145-(100+23)

= 22 mEq/L

The presence of a high anion gap indicates that there's indeed an acid-base disorder, thus ruling out option A.

A high anion gap indicates metabolic acidosis. However, HCO3- levels within normal limits suggest that there is also coexisting metabolic alkalosis in this patient.

This can be confirmed by calculating the delta gap.

Delta gap = ΔAG - ΔHCO3-

In this scenario: Delta gap = (22-10) - (24-23) = 11 mEq/L

Delta gap is > +6 suggesting the presence of metabolic alkalosis along with HAGMA.

If the delta gap is significantly positive (>+6), metabolic alkalosis is present because the rise in the anion gap is more than the fall in bicarbonate. If the delta gap is significantly negative (<6) then hyperchloremic acidosis is usually present because a rise in the anion gap is less than a fall in bicarbonate.

Alternatively, this can be confirmed by calculating the delta ratio.

Delta ratio = the increase in Anion Gap / the decrease in HCO3-

It is used in the presence of a high anion gap metabolic acidosis (HAGMA), to check it is a 'pure' HAGMA or if there is coexistant normal anion gap metabolic acidosis (NAGMA) or metabolic alkalosis.

Delta ratio values:

• 1-2 indicates HAGMA

• < 1 indicates NAGMA + HAGMA

• > 2 indicates HAGMA + Metabolic alkalosis

In this scenario,

Delta ratio = 12/1 = 12

Thus, the acid-base disorder for the above scenario is HAGMA with metabolic alkalosis.

Solution to Question 15:

The given clinical scenario is suggestive of nephrotic syndrome. This leads to a decrease in serum albumin which is an unmeasured anion.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Albumin at physiological pH is the major contributor to the unmeasured anions of the body. Unmeasured anions consume bicarbonate in the body and reduce its levels. This leads to an increase in the anion gap. Therefore, when these unmeasured anions are reduced the anion gap also reduces due to decreased consumption of the bicarbonate.

Nephrotic syndrome causes hypoalbuminemia and thus patients have a reduced anion gap. A fall in serum albumin by 1 g/dL from the normal value (4.5 g/dL) decreases the anion gap by 2.5 mEq/L.

Decreased anion gap (<6 mEq/L) is seen in:

• Increase in unmeasured cations

• An addition of abnormal cations to the blood, such as:

• Lithium (lithium intoxication)

• Cationic immunoglobulins (plasma cell dyscrasias)

• Reduction in the plasma anion albumin concentration (nephrotic syndrome, liver disease, or malabsorption)

• Bromide intoxication

• Hyperviscosity and severe hyperlipidemia, which can lead to an underestimation of sodium and chloride concentrations.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

In renal failure, there is a loss of kidney function due to the damage affecting glomeruli and tubules. This leads to a high anion gap metabolic acidosis.

Solution to Question 17:

The underlying acid-base disorder in this patient is metabolic acidosis with compensatory respiratory alkalosis.

Interpretation of the ABG:

Step 1: A pH of 7.1 signifies acidosis.

Step 2: Plasma HCO_3^- is 6 mEq/L, along with a low pH it signifies metabolic acidosis.

Step 3: pCO_2 is 16 mmHg indicating respiratory alkalosis. The decrease in pCO_2 along with the decrease in pH suggests that this is a compensatory response.

Step 4: The expected compensatory decrease in pCO_2 can be calculated using Winter's formula (Expected PCO_2= {[1.5 x HCO_3^-] +8} ± 2).

Expected PCO_2 = {[1.5 x 6] +8} ± 2 = 15 to 19 mmHg

A pCO_2 level of 16 mmHg falls within the range given above. Hence, metabolic acidosis is associated with compensatory respiratory alkalosis.

Sold by @itachibot

(pCO2 values above or below 15 - 19 mmHg would indicate mixed acid-base disorders)

Alternatively:

The other method to calculate the expected compensation is a 1.2 mmHg decrease in pCO2 for every 1 mmol/L decrease in bicarbonate.

The decrease in bicarbonate is 24 - 6 = 18 mEq/L

Decrease in pCO2 is 18 x 1.2 = 22

Expected pCO2 = 40 - 22 = 18 mmHg (roughly = 16 mmHg since the standard values are approximations)

This method also indicates primary metabolic acidosis with compensatory respiratory alkalosis. However, Winter's formula is more accurate.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

The probable reason for the patient's presenting symptoms is respiratory acidosis.

Asthma attack leads to the narrowing of respiratory passages. This leads to reduced O2 inhalation and reduced CO2 exhalation contributing to the rise of pCO2 in the blood, thereby causing respiratory acidosis.

Causes of respiratory acidosis:

- Central:
- Drugs (anesthetics, morphine, sedatives)
- Stroke
- Infection
- Airway:
- Obstruction
- Asthma
- Lung parenchyma:
- Emphysema
- Pneumoconiosis
- Bronchitis
- Adult respiratory distress syndrome
- Barotrauma
- Neuromuscular:
- Poliomyelitis
- Kyphoscoliosis
- Myasthenia

Sold by @itachibot

- Muscular dystrophies
- Miscellaneous:
- Obesity
- Hypoventilation
- Permissive hypercapnia.

Solution to Question 19:

The underlying acid-base disorder is respiratory acidosis.

Interpretation of the ABG:

Step 1: A pH of 7.24 signifies acidosis.

Step 2: A plasma HCO_3^- is 27 mEq/L is within normal limits. This is very likely to be due to compensation as it does not reflect the direction of the pH change.

Step 3: A pCO_2 of 60 mmHg along with an acidotic pH signifies respiratory acidosis.

Step 4: The expected compensatory increase in HCO_3^- can be calculated using the following formula:

Increase in HCO_3^- = 0.4 X Δ $PaCO_2$ (as COPD is a chronic condition and the presentation of the patient is not acute)

Thus, the increase in HCO_3^- = 0.4 x (60-40) = 8 mEq/L

Therefore expected HCO_3^- is 24 + 8 = 32 mEq/L.

The actual HCO_3^- measured is 27 mEq/L (which is less than the expected value) indicating that there could be a component of metabolic acidosis also.

But, the predominant disorder is respiratory acidosis and thus it is the best answer.

Solution to Question 20:

The given clinical scenario is suggestive of acute severe asthma. The acid-base disorder is respiratory acidosis, partially compensated.

Interpretation of the ABG:

Step 1: pH is decreased. This signifies acidosis. In acidosis, either pCO_2 is increased or plasma HCO_3^- is decreased.

Step 2: In this case, pCO_2 is increased by 30 mmHg. This signifies respiratory acidosis.

Step 3: Plasma HCO_3^- is increased by 26-24 = 2 mEq/L. This signifies compensation is taking place.

In acute conditions, for every 10mmHg rise in pCO_2 expected rise in plasma HCO_3^- is 1mEq/L. In this case, pCO_2 has risen by 30 mmHg, so the expected rise in HCO_3^- is 3 mEq/L.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

However, plasma HCO_3^- has risen only by 2 mEq/L, which implies compensation is only partial.

Solution to Question 21:

The above findings give the diagnosis of respiratory acidosis with metabolic compensation.

Interpretation of the ABG:

Step 1: pH is reduced (7.12), hence acidosis.

Step 2: Since pCO_2 is raised (50 mmHg), it is respiratory acidosis.

Step 3: HCO_3 is raised (28 mEq/L) suggesting there is metabolic compensation.

Solution to Question 22:

The clinical scenario is suggestive of a panic attack. The patient is likely to have respiratory alkalosis.

Panic attacks are characterized by hyperventilation which leads to a washout of CO_2 (reduced $PaCO_2$). This further increases the ratio of bicarbonate concentration to $PaCO_2$ and, thereby, increases the pH level causing alkalosis.

Alkalosis promotes the binding of calcium to albumin and reduces the fraction of ionized calcium in the blood. This leads to hypocalcemia without reducing the total calcium levels. Thus, alkalosis can precipitate tetany and laryngospasm, especially in calcium-deficient patients.

Induction of carpopedal spasm when the BP cuff is inflated 20 mm Hg over the systolic BP for three minutes or more is called Trousseau sign. It is a clinical manifestation of hypocalcemia.

Causes of respiratory alkalosis:

- Central nervous system stimulation:
- Pain
- Anxiety, psychosis
- Fever
- Cerebrovascular accident
- meningitis, encephalitis
- Tumor
- Trauma
- Hypoxemia or tissue hypoxia:
- High altitude
- Pneumonia, pulmonary edema
- Aspiration
- Severe anemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Drugs or hormones:

• Pregnancy

• Salicylates

• Cardiac failure

• Stimulation of chest receptors:

• Hemothorax

• Flail chest

• Cardiac failure

• Pulmonary embolism

• Miscellaneous:

• Septicemia

• Hepatic failure

• Mechanical hyperventilation

• Heat exposure

• Recovery from metabolic acidosis

Note: While metabolic alkalosis can also cause hypocalcemia, the above scenario of hyperventilation is more suggestive of respiratory alkalosis. Furthermore, hypocalcemic symptoms are more common with respiratory alkalosis.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 23:

In the given illustrated acid-base nomogram, the letters A and D indicate acute respiratory acidosis and acute respiratory alkalosis.

An acid-base nomogram, also known as a Davenport diagram, is a graphical representation of acid-base balance to interpret arterial blood gas (ABG) results. It helps determine the primary acid-base disorder and identify compensatory mechanisms.

Sold by @itachibot



The y-axis of the acid-base nomogram represents the arterial bicarbonate (HCO3-) levels, and the x-axis represents the arterial pH values. The shaded areas are the range of values of the normal respiratory and metabolic compensations for primary acid-base disturbances.

Sold by @itachi... if you purchased this from some... you may have been scammed else,

Solution to Question 24:

The mechanism for the secondary disorder is hypoventilation causing decreased CO2 washout.

The given arterial blood gas findings are suggestive of a partially compensated metabolic acidosis. The compensation is partial because the fall in paCO2 much lower than what is expected for the given HCO3 levels. The fall in paCO2 is inadequate due to hypoventilation causing decreased CO2 washout. In this case, the secondary disorder is respiratory acidosis.

Hyperventilation causing increased CO2 washout (Option B) would allow for better compensation, with lower paCO2 levels than observed.

The expected paCO2 can be calculated using Winter's formula:

For metabolic acidosis, the expected paCO2 = [1 .5 X HCO3-] + 8 ± 2.

Substituting the values, expected PaCO2 = (1.5 x 10) +8 ± 2

= 15+8 ±2

= 21 to 25 mmHg

However, the actual PaCO2 is 30 mmHg, which is higher than the expected paCO2 indicating a secondary respiratory acidosis. A possible explanation for this could only be hypoventilation causing decreased CO2 washout.

Thus, the given ABG findings are suggestive of a mixed acid-base disorder characterized by metabolic acidosis with additional respiratory acidosis.

For example, in a patient with diabetic ketoacidosis and pneumonia, the lungs would fail to compensate, and the patient could develop hypoventilation leading to secondary respiratory

Sold by @itachibot

acidosis.

Solution to Question 25:

According to the ABG analysis provided, the patient has a mixed acid-base disorder with simultaneous respiratory and metabolic acidosis.

Interpretation of the ABG:

Step 1: A pH of 6.9 signifies acidosis.

Step 2: Plasma HCO_3^- is 13 mEq/L, along with a low pH it signifies metabolic acidosis.

Step 3: pCO_2 is 68 mmHg indicating co-existing respiratory acidosis. The direction of pCO_2 change is the same as that of pH change suggesting that this is a mixed acid-base disorder.

Step 4: The expected compensatory change in pCO_2 in the metabolic acidosis calculated using Winter's formula would be:

$pCO_2 = [1.5 \times HCO_3^-] + 8 \pm 2 = 26 - 30$ mmHg

The actual pCO_2 measure is very high (68 mmHg) compared to the estimated range of 26 - 30 mmHg indicating the presence of simultaneous metabolic and respiratory acidosis.

Note- If this was a case of uncompensated acidosis, either the bicarbonate or the pCO_2 values would have been within the normal limits.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

General Principles of Endocrinology

Question 1:

Vasopressin acting on its V2 receptor uses _____ as its second messenger.

a) Calcium

b) cGMP

c) cAMP

d) Phospholipase C

Question 2:

Which of the following act as the second messenger for secretin?

a) cAMP

b) cGMP

c) Serine kinase

d) Tyrosine kinase

Question 3:

Which hormone mediates its action via cyclic GMP?

a) Oxytocin

b) Vasopressin

c) Aldosterone

d) Nitric oxide

Question 4:

What is the most abundant hormone produced by the pituitary gland?

a) Vasopressin

b) GH

c) ACTH

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) TSH

Question 5:

Insulin acts on which of the following?

a) Nuclear receptor

b) Cytoplasmic receptor

c) Tyrosine kinase receptor

d) STK receptor

Question 6:

Which of the following is considered the twin hormone of prolactin?

a) ACTH

b) GH

c) Oxytocin

d) Dopamine

Question 7:

Which of the following hormones is an amino acid derivative?

a) Thyroid hormone

b) Insulin

c) Activin

d) Parathyroid hormone

Question 8:

Which of the following chemical molecules is not a steroid hormones?

a) Cortisol

b) Estrogen

c) DHEA

d) Inhibin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 9:

Which class of hormones does insulin belong to?

a) Amino acid derivatives

b) Small neuropeptides

c) Large protein hormones

d) Medium-sized peptide

Question 10:

The action of cAMP is mediated by _____

a) Adenylyl cyclase

b) Guanylyl cyclase

c) Protein kinase

d) IP3-DAG

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 11:

The second messenger, cAMP is synthesized by the enzyme _____

a) Adenylyl cyclase

b) cAMP phosphodiesterase

c) ATPase

d) Phosphorylase

Question 12:

Which of the following is not activated by the binding of epinephrine to its receptor?

a) Gi

b) Gq

c) Gp

d) Gs

Sold by @itachibot

Question 13:

The enzyme kinase adds _____ to proteins.

a) Acetates

b) Sulphates

c) Carbonates

d) Phosphates

If you bought this from anyone else,

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	b
5	c
6	b
7	a
8	d
9	c
10	c
11	a
12	c
13	d

Detailed Explanations

Solution to Question 1:

Vasopressin acts on the V2 receptor using cAMP as its second messenger.

Vasopressin acts on GPCR using different second messengers to accomplish its various actions as follows:

• V1 receptor - vasoconstriction using phospholipase C as its second messenger

• V2 receptor - reabsorption of water through aquaporin 2, using cAMP as its second messenger

Sold by @itachibot

Vasopressin has a maximum effect on the collecting ducts. It acts by inserting aquaporin 2 present in cytoplasmic vesicles of principal cells on to the apical membrane.

Solution to Question 2:

cAMP is the second messenger for the hormone secretin. It regulates fluid and electrolyte secretion and also mediates water and nutrient reabsorption.

Solution to Question 3:

Nitric oxide mediates its action via cyclic GMP or cGMP.

cGMP is the second messenger associated with smooth muscle relaxation and vasodilation. Atrial natriuretic factor (ANF) and NO (nitric oxide) are the only hormones that mediate their action via cGMP.

Note: Recent studies show that NO is a hormone. NO, which functions as a chemical messenger, has similar properties to hormones (which are known as the body's chemical messengers).

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

Growth hormone (GH) is the most abundant hormone produced by the pituitary gland.

It is produced by the acidophil cells or somatotropes of the anterior pituitary, which account for 50% of the anterior pituitary cell population.

Option A: Vasopressin is produced in the hypothalamus and only stored in the posterior pituitary.

Option C: Corticotrophs account for 10% of the anterior pituitary cell population.

Option D: Thyrotrophs account for 5% of the anterior pituitary cell population.

Solution to Question 5:

Insulin acts on the tyrosine kinase receptor, a cell-surface receptor.

Solution to Question 6:

GH and prolactin are called twin hormones as both are secreted by the acidophil cells of the anterior pituitary. They act on cytokine receptors and show JAK-STAT activity.

Sold by @itachibot

Solution to Question 7:

The thyroid hormone is synthesized from tyrosine residues and thus is an amino acid derivative.

Insulin, activin and parathyroid hormone are large protein hormones.

Solution to Question 8:

Inhibin is a large protein hormone, not a steroid hormone.

All adrenocortical and reproductive hormones are steroid hormones synthesized from cholesterol precursors except activin and inhibin.

Solution to Question 9:

Insulin is classified as a large protein hormone (>50 amino acid residues) as it is composed of 51 amino acids.

Protein and polypeptide hormones are classified as small peptides (< 50 amino acid residues) and large proteins.

Solution to Question 10:

The action of cAMP is mediated by protein kinases.

cAMP is a second messenger molecule associated with electrolyte secretion and water and nutrient reabsorption. It is synthesized by adenylyl cyclase but it mediates its actions via protein kinases.

Solution to Question 11:

cAMP is synthesized by adenylyl cyclase, but it mediates its actions via protein kinases.

Solution to Question 12:

The binding of epinephrine to its various receptors results in the activation of different types of G-protein subunits, namely Gq, Gi, and Gs.

All catecholamines act via G protein-coupled receptors. This stimulates cAMP production and signaling via protein kinases.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 13:

Kinases are a subclass of transferases that catalyze the transfer of a phosphate group from a donor to an acceptor protein. They are associated with the regulation of many essential cellular processes.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Disorders of Anterior Pituitary

Question 1:

Which of the following is not a characteristic of growth hormone (GH) deficiency?

a) Hyperglycemia

b) Micropenis

c) Delayed bone age

d) High-pitched voice

Question 2:

A 3-year-old girl is referred to your centre for evaluation of short stature. The child's parents are found to be first cousins. Further evaluation led to the diagnosis of Laron-type dwarfism. What causes this condition?

a) GH deficiency

b) GH receptor anomaly

c) GHRH deficiency

d) IGF-1 deficiency

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 9-year-old girl who presented with short stature is diagnosed with Turner's syndrome. The physician decides to initiate treatment with recombinant human growth hormone for short stature. Which of the following side effects is she not likely to develop?

a) Tinnitus

b) Hypoglycemia

c) Pseudotumor cerebri

d) Slipped capital femoral epiphysis

Question 4:

Sold by @itachibot

Paradoxical response of GH release to thyrotropin-releasing hormone (TRH) is seen in all of the following conditions except_____

a) Prolactinoma

b) Acromegaly

c) Malnutrition

d) Anorexia nervosa

Question 5:

A patient presented to the OPD with complaints of progressive enlargement of hands and feet and was diagnosed with acromegaly. What is the most common cause for this condition?

a) GHRH secretion from hypothalamic hamartoma

b) GH secretion from somatotrope adenoma

c) GH and PRL secretion from mixed somatotroph-lactotroph adenoma

d) GH and PRL secretion from mixed somatotroph-thyrotroph adenoma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which of the following features will you expect to see in a patient with acromegaly?

a) i,ii,iii

b) i ii,iv

c) ii,iii,v

d) i,ii,v

Question 7:

A middle-aged man presented to the OPD with complaints of difficulty in chewing and a gradual change in voice over the last year. Examination revealed coarse facial features. The image of his hand (right side) compared to the physician's hand is shown below. Which of the following long-term complications would you not commonly see in this patient?

Sold by @itachibot

a) Diabetes mellitus

b) Diabetes insipidus

c) Coronary heart disease

d) Cardiomyopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

The middle-aged man shown in the image, noticed that his shoes no longer fit him and his finger ring is getting tightened. Insulin-like growth factor-1 (IGF-1) levels are elevated. Which of the following tests will help to confirm the diagnosis?

a) Insulin-like Growth Factor -1 Binding Protein - 3 (IGFBP-3) level

Sold by @itachibot

b) Random serum GH

c) Oral glucose tolerance test

d) MRI brain

Question 9:

A 44-year-old man is referred to your center for management of severe hyperglycemia. During examination, you note that the patient has enlarged feet, coarse facial features and prognathism. You suspect acromegaly. Which of the following tests can you use as a screening method for this condition?

a) Oral glucose challenge test with GH level

b) Insulin-like growth factor-1 binding protein-3 (IGFBP-3) level

c) Insulin like growth factor-1 levels

d) Random growth hormone levels

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 10:

The medical management of acromegaly includes all the following therapeutic strategies except _____

a) Lanreotide

b) Pegvisomant

c) Cabergoline

d) Levodopa

Question 11:

A 27-year-old woman presents to the OPD with infertility and amenorrhea. She also complains of milky discharge from the nipple. Her urine pregnancy test was negative and her prolactin level was 45 ng/ml. Which of the following drugs cannot lead to this condition?

a) Methyldopa

b) Bromocriptine

c) Phenothiazine

d) Metoclopramide

Sold by @itachibot

Question 12:

Systemic causes of hyperprolactinemia include all of the following clinical conditions except _____

a) Chronic renal failure

b) Epileptic seizures

c) Cirrhosis

d) Hyperthyroidism

Question 13:

Which of the following is not a hallmark of hyperprolactinemia in women?

a) Amenorrhea

b) Galactorrhea

c) Infertility

d) Diminished libido

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 14:

A patient presents to your OPD with galactorrhea and amenorrhea. Apart from the determination of serum prolactin levels, which other investigation would you like to do?

a) LH

b) TSH

c) FSH

d) Urinary ketosteroids

Question 15:

A 30-year-old nullipara with a desire to conceive presents with amenorrhea and galactorrhea. UPT is negative. Investigations indicate hyperprolactinemia. What is the drug of choice for clinical management of this condition?

a) Cabergoline

b) Bromocriptine

c) Sandostatin

Sold by @itachibot

d) Pegvisomant

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	c
5	b
6	a
7	b
8	c
9	c
10	d
11	b
12	d
13	d
14	b
15	b

Detailed Explanations

Solution to Question 1:

Hyperglycemia is not a characteristic of GH deficiency. Isolated GH deficiency, especially in children, can lead to a higher propensity to develop hypoglycemia. This is attributed to the relatively unopposed action of insulin.

Isolated GH deficiency in children is associated with the following features:

- Short stature
- Micropenis
- Increased fat
- High-pitched voice
- High propensity to develop hypoglycemia due to relatively unopposed insulin action.

Sold by @itachibot

Solution to Question 2:

Laron-type dwarfism, also known as GH-insensitivity syndrome, is characterized by GH-receptor anomalies.

Laron-type dwarfism is an autosomal recessive disorder. The primary clinical symptom associated with this condition is short stature. Treatment is by subcutaneous injections of insulin-like growth factor 1 (a growth-promoting hormone), often called IGF-1.

Solution to Question 3:

GH is a potent counter-regulatory hormone to insulin and leads to hyperglycemia, thus contraindicated in patients with uncontrolled diabetes.

Side effects of GH administration include:

- Dose-related fluid retention
- Hyperglycemia
- Joint pain and carpal tunnel syndrome
- Headache, increased intracranial pressure, hypertension
- Tinnitus
- Pituitary tumor regrowth
- Progression of skin lesions
- Acromegaly
- Slipped capital femoral epiphysis
- Gynecomastia

The actions of GH are:

- Induces protein synthesis and nitrogen retention
- Impairs glucose tolerance by antagonizing insulin action
- Stimulates lipolysis
- Promotes sodium, potassium, and water retention
- Elevates serum levels of inorganic phosphate
- Linear bone growth occurs
- Stimulates epiphyseal prechondrocyte differentiation

Solution to Question 4:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

The paradoxical response of GH release to TRH is not seen in malnutrition patients.

In normal individuals, GH doesn't respond to TRH stimulation, but paradoxically, the GH levels may increase in response to TRH stimulation in the following conditions:

• Acromegaly

• Prolactinoma

• Anorexia nervosa

The exact mechanism underlying this paradoxical GH response is poorly understood.

Solution to Question 5:

The most common cause of acromegaly is GH secretion from somatotrope adenoma.

Causes of acromegaly:

• Pituitary - 98%

• Somatotrope adenoma - 60% - most common.

• Mixed somatotroph-lactotroph adenoma

• Other plurihormonal tumors.

• Extrapituitary tumor-producing GH: Pancreatic Islet cell tumor

• Excess GHRH secretion

• Central - hypothalamic hamartoma

• Peripheral - bronchial carcinoid

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

Impaired glucose tolerance, galactorrhea, and hypertension can be seen in patients with acromegaly.

GH secretion is a part of the counter-regulatory defense mechanism against hypoglycemia. Physiological GH secretion is inhibited by hyperglycemia. In acromegaly or gigantism, GH secretion is autonomous and not suppressed by glucose. Thus, its diagnosis can be confirmed by demonstrating the failure of GH suppression with an oral glucose load (impaired oral glucose tolerance test).

Galactorrhea is due to hyperprolactinemia and hypertension is secondary to increased aldosterone levels in acromegaly. GH leads to increased production of vitamin D3 which results in increased phosphate absorption from the intestine and decreased renal excretion. Hence there is increased serum phosphate level.

Solution to Question 7:

Sold by @itachibot

The given scenario and the enlarged hands in the patient, as shown in the image, point towards a diagnosis of acromegaly. Diabetes insipidus is not a feature of acromegaly, but diabetes mellitus can develop in patients with acromegaly.

Clinical features of acromegaly are:

Note: Diabetes insipidus can be seen as a complication post Transsphenoidal Surgery for pituitary adenomas and other sellar tumours not as complication of Acromegaly per se.

System	Features
Musculoskeletal	GigantismProgнathism (protrusion of jaw)Carpal tunnel syndromeParesthesia (due to nerve entrapment)Proximal myopathyHypertrophy of frontal bones
Skin and soft tissues	Acral enlargement in hands and feet (due to thickening of soft tissues)HyperhidrosisAcanthosis nigricans and skin tags
Visceralomegaly	Enlargement of most organs especially the tongue, thyroid, salivary gland, liver, spleen, kidney, and prostate
Ophthalmic	Visual field defects (macro-pituitary adenoma)Ophthalmoplegia (cranial nerve palsy)Headache
Metabolic	Low renin with increased aldosteroneDiabetes mellitus(due to GH-induced insulin resistance)Hypercalcemia, hyperphosphatemia, and hypercalciuriaHypertriglyceridemia
Reproductive	Loss of libido and impotence in malesMenstrual abnormalities and galactorrhea
Cardiopulmonary	Left ventricular hypertrophy/septal hypertrophyHypertensionObstructive sleep apnea

Sold by @itachibot



MARROW
Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

The given scenario and the presence of coarse facial features and prognathism, as shown in the image, points towards a diagnosis of acromegaly. An oral glucose tolerance test with GH levels is used to confirm acromegaly.

Solution to Question 9:

The best screening test for acromegaly is the measurement of serum insulin-like growth factor-1 (IGF-1) level.

Other options:

Options A and B: Oral glucose challenge test with growth hormone (GH) levels is used for further confirmation of the diagnosis. Increased insulin-like growth factor-1 binding protein-3 (IGFBP-3) levels can also be seen in patients with acromegaly. But they are not typically used as initial screening methods.

Option D: IGF-1 levels are more reliable than random GH levels because GH secretion is pulsatile. IGF-1 levels reflect long-term exposure to GH and provide a more stable indicator of excessive GH production.

Acromegaly is characterized by excessive GH secretion, usually caused by a somatotrope adenoma in the pituitary gland. It can also be due to ectopic GH secretion or hypothalamic tumors.

The clinical presentation of acromegaly includes frontal bossing, increased hand and foot size, coarse facial features, mandibular prognathism, widened space between lower incisors and other manifestations such as soft tissue swelling, deepening voice, carpal tunnel syndrome, arthropathy, and acanthosis nigricans.

Sold by @itachibot

It can also lead to cardiovascular complications (cardiomyopathy, left ventricular hypertrophy, and hypertension), obstructive sleep apnea, diabetes mellitus, an increased risk of colon polyps, and colonic malignancy.

The primary treatment for acromegaly is surgical resection of GH-secreting adenomas. Somatostatin receptor ligands like octreotide and lanreotide, which suppress GH and IGF-1 levels, are used as adjuvant treatment or when surgery is not possible or unsuccessful. Radiation therapy may be considered for patients who cannot tolerate or do not respond to medical therapy. Other treatment options include GH receptor antagonists like pegvisomant and dopamine agonists like cabergoline.

Solution to Question 10:

Levodopa is not used in the treatment of acromegaly. Studies have proven that it is not as effective as agents like cabergoline.

Medical management of acromegaly includes:

- Somatostatin analogs:
- Octreotide
- Lanreotide
- Pasireotide
- GH-receptor antagonists:
- Pegvisomant
- Dopamine agonists:
- Bromocriptine
- Cabergoline

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 11:

This given scenario of infertility, amenorrhea and galactorrhea with elevated prolactin level (normal level is 0-25 ng/ml in a non-pregnant woman) is suggestive of hyperprolactinemia, and bromocriptine does not lead to hyperprolactinemia. It is a dopamine-agonist and hence it inhibits prolactin synthesis.

Dopamine agonists like bromocriptine and cabergoline are used in the treatment of hyperprolactinemia.

Solution to Question 12:

Hypothyroidism predisposes to the development of hyperprolactinemia.

Sold by @itachibot

Both enhanced hypothalamic synthesis of TRH and increased pituitary-responsiveness to TRH have been described as the mechanism behind this phenomenon.

Other systemic disorders that predispose to the development of hyperprolactinemia :

• Chronic renal failure

• Cirrhosis

• Pseudocyesis

• Epileptic seizures

Solution to Question 13:

The hallmarks of hyperprolactinemia in women are:

• Amenorrhea – (Secondary > Primary)

• Galactorrhea

• Infertility

In men with hyperprolactinemia, diminished libido, infertility, and visual loss (from optic nerve compression) are the usual presenting symptoms.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

TSH should be measured along with the prolactin level to rule out hypothyroidism which can cause hyperprolactinemia.

Note: However, pregnancy (using qualitative HCG) should be ruled out in all women of the reproductive age group with amenorrhea.

Solution to Question 15:

The drug of choice for clinical management of this patient would be bromocriptine.

Bromocriptine is an ergot alkaloid and a short-acting dopamine receptor agonist. Due to its short-term action, it is preferred when pregnancy is desired.

Cabergoline is a long-acting dopamine receptor agonist with high D2 receptor affinity. It effectively suppresses PRL secretion for >14 days after a single oral dose.

Sold by @itachibot

Pituitary Tumours and Sheehan's Syndrome

Question 1:

What is the most common functioning tumor of the pituitary?

a) GH-secreting adenoma

b) ACTH-secreting adenoma

c) Gonadotrope adenoma

d) PRL-secreting adenoma

Question 2:

What are the most common cells involved in non-functioning pituitary adenoma?

a) Prolactin-producing cells

b) ACTH-producing cells

c) Gonadotropin-producing cells

d) TSH-producing cells

Question 3:

A 30-year-old woman with a suspected prolactinoma is planned for a brain MRI. What size would the tumor have to be to call it a microadenoma?

a) < 10 mm

b) < 20 mm

c) < 5 mm

d) < 15 mm

Question 4:

Which of the following would not occur in a patient with a pituitary macroadenoma?

a) Increased insulin sensitivity

b) Pigmentation of mucous membranes

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

c) Low serum thyroxine and TSH levels

d) Loss of secondary sexual characteristics

Question 5:

Which of the following can occur in a patient with prolactinoma?

a) Superior quadrantanopia

b) Inferior quadrantanopia

c) Priapism

d) Failure of lactation

Question 6:

What levels of prolactin are definitely suggestive of prolactinoma?

a) >50µg/L

b) >100µg/L

c) >150µg/L

d) >200µg/L

Question 7:

Patients with which of the following conditions do not need to be screened for pituitary tumors?

a) MEN1

b) MEN2

c) MEN4

d) Carney Complex

Question 8:

A 56-year-old woman presented with a gradually diminishing visual field. She was found to have a pituitary mass and underwent trans-sphenoidal hypophysectomy. Now she has low levels of ACTH, TSH, FSH, and LH. Which of the following hormone supplements is not needed in the patient postoperatively?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Mineralocorticoids

b) Thyroid hormones

c) Glucocorticoids

d) Estradiol

Question 9:

Which of the following is the most common clinical presentation in patients with craniopharyngiomas?

a) Growth retardation

b) Bitemporal hemianopia

c) Headache

d) Endocrinal disturbance

If you purchased this from someone else, Sold by @Itachibot have been scammed.

Question 10:

A 10-year-old boy is referred for evaluation of progressively worsening headaches and visual field defects. An MRI brain is done and shows the following findings. Which of the following statements is incorrect about this condition?

a) Composed of epithelial cells

b) Surgery is curative in most patients

c) Contains cholesterol crystals

Sold by @itachibot

d) Calcifications are seen in adamantinomatous type

Question 11:

Patients with which of the following conditions are not at risk for pituitary apoplexy?

a) Hyperthyroidism

b) Diabetes mellitus

c) Sickle cell anemia

d) Hypertension

Question 12:

Which of the following is seen in pituitary apoplexy?

a) Hypertension

b) Shock

c) Unconsciousness

d) Fatigue

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

Which hormone is the first to fall in Sheehan syndrome?

a) Prolactin

b) GH

c) FSH

d) ACTH

Question 14:

You are a first-year resident in the postpartum ward. During rounds, you see a 28-year-old woman who has been unable to breastfeed since delivery. Her baby was born by normal vaginal delivery that was complicated by severe postpartum hemorrhage. Her blood pressure is 90/50 mm Hg and her pulse is 96/min. What is the next best step in management?

a) Estrogen

b) Growth hormone

Sold by @itachibot

c) Thyroxine

d) Cortisone

Answer Key

Question No.	Correct Option
1	d
2	c
3	a
4	b
5	a
6	d
7	b
8	a
9	c
10	b
11	a
12	b
13	b
14	d

Detailed Explanations

Solution to Question 1:

Prolactin-secreting adenoma or prolactinoma is the most common type of functioning pituitary adenoma. Pituitary adenomas are the most common type of pituitary tumor, accounting for one-third of all tumors.

Solution to Question 2:

Non-functioning pituitary adenomas most commonly arise from gonadotropic cells (80%).

Pituitary adenomas are classified as functioning or non-functioning based on the secretion of hormones. Non-functioning adenomas are usually asymptomatic in the initial stages but as the tumor grows larger, mass effect produces clinical manifestations such as:

Sold by @itachibot

- Headache
- Bitemporal hemianopia
- Diplopia, ptosis
- Compression of pituitary stalk
- Hyperprolactinemia initially - reduced dopamine release leads to loss of inhibition of lactotrophs
- Hypopituitarism in later stages

Solution to Question 3:

Pituitary adenomas < 10 mm are called microadenomas.

Asymptomatic pituitary tumors can be discovered incidentally on imaging. They must also be suspected in patients with features of hormone hypersecretion or mass effect. In such cases, MRI with contrast is the best investigation to detect the tumor, as shown below.



Levels of pituitary hormones must also be measured to rule out both hypersecretion and hyposecretion.

Solution to Question 4:

Pituitary macroadenomas do not cause pigmentation of mucous membranes as ACTH levels are low.

In this condition, there is destruction of the pituitary gland by mass effect. This leads to panhypopituitarism or reduced levels of all hormones, including ACTH. Excessive pigmentation is only seen in cases of excess ACTH.

The following hormones are deficient in panhypopituitarism:

Sold by @itachibot

• ACTH - Secondary adrenal insufficiency

• TSH - Secondary hypothyroidism

• GH - Growth failure and adult hyposomatotropism

• FSH, LH - Hypogonadic hypogonadism

• Prolactin - Lactation failure

Adiponectin, secreted by fat cells normally increases insulin sensitivity. It is usually suppressed by cortisol. In panhypopituitarism, decreased cortisol levels lead to increased insulin sensitivity.

Solution to Question 5:

Superior quadrantanopia may be seen in patients with prolactinoma > 10 mm in size (macroadenoma).

In this condition, visual field defects occur due to compression of the optic chiasm. In the initial stages, the pituitary tumor tends to compress the ventral portion of the optic chiasm. This causes a bitemporal (heteronymous) superior field defect. As the tumor grows larger, the entire optic chiasm gets compressed, leading to complete bitemporal hemianopia, as shown below.

Damage to the optic radiation in the temporal lobe would result in homonymous superior quadrantanopia or pie-in-the-sky appearance.

Solution to Question 6:

The diagnosis of a prolactinoma is highly likely when prolactin levels are > 200 µg/L.

The normal range of serum prolactin (PRL) is 5-20 µg/L. Prolactin levels < 100 µg/L can be due to non-functioning pituitary adenomas or non-neoplastic causes of hyperprolactinemia. MRI brain must be done in all patients with hyperprolactinemia.

Prolactinoma are tumors arising from the lactotroph cells of the pituitary gland, they can be classified as microadenoma(<1cm in diameter, usually non invasive) or macroadenoma(>1cm, locally invasive to adjacent structures). Clinical features include amenorrhea, infertility, and galactorrhea in women, impotence, loss of libido, infertility in men and if tumor extends beyond sella, there can be visual field defects and other mass effects.

Treatment - In asymptomatic patients and those where fertility is not desired, they do not require treatment and can be monitored using regular PRL measurements and MRI. In symptomatic microadenoma cases, oral dopamine agonists (cabergoline and bromocriptine) form the main treatment for hyperprolactinemia and reduction of tumor volume. For macroadenoma, before starting dopamine agonists visual field testing is to be done, and reassess it with MRI changes every 6-12 month intervals until maximum reduction is achieved, in unresponsive cases consider surgery.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Sold by @itachibot

Solution to Question 7:

MEN2 syndrome is not associated with pituitary tumors. It is associated with medullary thyroid cancer and pheochromocytoma.

The genetic syndromes associated with pituitary tumors include:

• MEN1

• MEN4

• Carney complex tumors

• McCune-Albright Syndrome

Solution to Question 8:

The given clinical scenario describes a woman presenting with hypopituitarism following trans-sphenoidal resection. Mineralocorticoid supplementation is not needed postoperatively as it is regulated by the renin-angiotensin-aldosterone system and not by ACTH. Therefore, mineralocorticoid levels would remain intact despite ACTH deficiency.

Thyroid hormone, estradiol, and glucocorticoid supplementation are needed in this patient as their secretion is regulated by anterior pituitary hormones.

Features of hypopituitarism include growth abnormalities in children with a growth hormone deficiency, insufficient levels of gonadotropins leading to menstrual irregularities, infertility in women and reduced sexual function, infertility and loss of secondary sexual characteristics in men. TSH deficiency leads to stunted growth in children and features of hypothyroidism in both children and adults. ACTH deficiency leads to secondary adrenal insufficiency.

Polyuria and polydipsia may be seen and indicate a loss of vasopressin secretion in posterior pituitary lesions.

The following hormones must be replaced post-hypophysectomy:

Trophic hormone deficit	Hormone Replacement
ACTH	Hydrocortisone, cortisone acetate, prednisone
TSH	L-thyroxine (75–150 µg daily)
FSH/LH	Males: TestosteroneFemales: For fertility: menopausal gonadotropins, hCGPost-menopausal women: Conjugated estrogen, progesterone
GH	Somatotropin
Vasopressin	Intranasal desmopressin

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 9:

Headache is the most common presenting symptom of craniopharyngioma.

Craniopharyngiomas are benign, suprasellar cystic masses. They have a bimodal distribution, occurring predominantly in children but also between the ages of 55 and 65 years. It can present with the following clinical symptoms:

- Headache
- Hypopituitarism
- Hypogonadotropic hypogonadism
- Central diabetes insipidus
- Bitemporal hemianopia
- Raised ICT - vomiting, papilledema, hydrocephalus

Solution to Question 10:

The given MRI showing a suprasellar mass with calcifications is suggestive of a craniopharyngioma. Surgery is not curative in the majority of patients as the tumor is adherent to surrounding structures.

They are benign epithelial cell-containing tumors and are of two types:

- Adamantinomatous - calcifications and keratin nodules are seen
- Papillary - calcifications are rare

Biopsy reveals machinery or motor-oil-like fluid within the cyst that contains cholesterol crystals. The treatment of choice is transcranial or transsphenoidal resection. It must be followed by postoperative radiation to prevent recurrence.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 11:

Pituitary apoplexy does not occur in hyperthyroidism. It is a rare endocrine emergency which can occur due to haemorrhage or infarction of the pituitary gland. It may occur as a result of the following conditions:

- Pituitary adenoma (spontaneous)
- Sheehan syndrome
- Acute shock
- Diabetes mellitus
- Hypertension
- Sickle cell anemia

Sold by @itachibot

Solution to Question 12:

Shock is seen in pituitary apoplexy.

Loss of consciousness (Option C) can also be seen but it occurs secondary to adrenal insufficiency and shock. Hence, shock is the most appropriate answer.

Pituitary apoplexy refers to acute hemorrhagic infarction of the pituitary gland. This results in the compression of surrounding structures and hypopituitarism leading to acute adrenal insufficiency. It may be seen spontaneously in a pre-existing pituitary adenoma, post-partum (Sheehan's syndrome), or be associated with hypertension, diabetes mellitus or sickle cell anemia or shock.

Pituitary apoplexy is an endocrine emergency that can lead to severe hypoglycemia, hypotension, shock, central nervous system (CNS) hemorrhage, and death. Acute clinical manifestations commonly encompass intense headaches with signs of meningeal irritation, bilateral visual changes, ophthalmoplegia, and even cardiovascular collapse.

The investigation of choice is MRI.

Patients who do not exhibit overt visual impairment or altered mental status can be managed conservatively with high-dose glucocorticoids. Progressive visual loss, cranial nerve palsy, or altered mental status necessitate expeditious surgical decompression.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 13:

Growth hormone (GH) is the first to fall in Sheehan syndrome. The hormones are affected in the order of GH → Prolactin → LH and FSH → ACTH → TSH

Sheehan syndrome refers to postpartum pituitary necrosis following severe postpartum hemorrhage (PPH). During pregnancy, the pituitary gland undergoes hypertrophy with increased blood supply. This makes it susceptible to ischemia and infarction when there is severe hypovolemia.

The clinical presentation depends on the amount of damage to the pituitary.

- Mild - can go unnoticed for many years
- Moderate - develops over months
- Lactation failure
- Failure to resume menses
- Mild fatigue and anorexia
- Severe - endocrine emergency that develops within few days
- Hypotension and shock
- Severe hypoglycemia
- Lactation failure

GH deficiency is very common in Sheehan's syndrome because somatotrophs are located in the lower and lateral regions of the pituitary gland, and are most likely to be damaged by ischemic necrosis of the pituitary.

Sold by @itachibot

Solution to Question 14:

This clinical scenario is suggestive of Sheehan syndrome. Corticosteroids must be supplemented first, as life-threatening manifestations may occur due to the sudden onset of adrenal insufficiency. This is followed by the replacement of other deficient pituitary hormones including thyroid hormone and estrogen administered with or without progesterone.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Posterior Pituitary - ADH, Oxytocin & Diabetes Insipidus

Question 1:

Which of the following are functions of oxytocin?

a) 1 only

b) 1 and 4 only

c) 1, 2 and 4 only

d) 3, 4 and 5 only

Question 2:

Where are arginine vasopressin (AVP) and oxytocin stored?

a) Anterior pituitary

b) Supraoptic nucleus

c) Paraventricular nucleus

d) Herring bodies

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following statements is true?

a) AVP binds to V1 receptors on the serosal surface of renal tubular cells

b) AVP binds to V1 receptors on the luminal surface of renal tubular cells

c) AVP binds to V2 receptors on the serosal surface of renal tubular cells

d) AVP binds to V2 receptors on the luminal surface of renal tubular cells

Question 4:

Which of the following statements about ADH is incorrect?

a) It increases aquaporin expression on apical membrane

b) It increases urine concentration and decreases urine flow

Sold by @itachibot

c) Post-operative secretion is high

d) Its secretion occurs when plasma osmolality is low

Question 5:

A mutation in the gene for aquaporin channel leads to which of the following conditions?

a) Liddle's syndrome

b) Nephrogenic diabetes insipidus

c) Cystic fibrosis

d) Bartter syndrome

Question 6:

Which of the following conditions would you not expect to see in a patient with Wolfram syndrome?

a) Diabetes mellitus

b) Sensorineural deafness

c) Optic atrophy

d) SIADH

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

All of the following statements about diabetes insipidus are true, except:

a) Low urine osmolality

b) Dilutional hyponatremia

c) High plasma osmolality

d) Diagnosed by water deprivation test

Question 8:

A man presented with frequent urination and increased thirst. Investigations revealed low urine osmolality and high plasma osmolality. MRI brain was done, which revealed the absence of posterior pituitary bright spot. Which of the following conditions is least likely to lead to this presentation?

Sold by @itachibot

a) Multiple sclerosis

b) Head injury

c) Histiocytosis

d) Viral encephalitis

Question 9:

A patient presented with frequent urination, nocturia, and enuresis. Investigations revealed 24-hour urine output of 7 litres and low urinary osmolality. ADH assay was performed and the recorded values were reported as 1.5 pg/ml. Which of the following drugs would not cause this condition?

a) Lithium

b) Cisplatin

c) Isoniazid

d) Rifampin

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 10:

A patient presents with polyuria and increased thirst. Investigations reveal hypercalcemia. What could be the cause of polyuria in this patient?

a) Neurogenic DI

b) Nephrogenic DI

c) Iatrogenic polydipsia

d) Dipsogenic polydipsia

Question 11:

Which of the following findings do you expect to see in a patient with central diabetes insipidus?

a) Low plasma and low urine osmolality

b) High plasma and high urine osmolality

c) Low plasma and high urine osmolality

d) Low urine and high plasma osmolality

Sold by @itachibot

Question 12:

A 29-year-old lady presented with polydipsia and polyuria. Her symptoms started soon after a road traffic accident 6 months ago. Her blood pressure was recorded at 120/80 mmHg with no postural drop. The daily urinary output was reported to be between 6-8 liters. Clinical investigation results are given below. What is the most likely diagnosis?

a) Central diabetes insipidus

b) Cerebral salt wasting

c) Psychogenic polydipsia

d) Nephrogenic diabetes insipidus

Question 13:

A patient presented with polyuria and polydipsia. She has been on lithium therapy for 6 years for bipolar disorder. Investigations revealed low urine osmolality and high plasma osmolality. Which of the following drugs can be used for the treatment of this condition?

a) Thiazide diuretics

b) Salt supplementation

c) Desmopressin

d) Vaptans

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

Polyuria characterized by low fixed specific gravity of urine is seen in which of the following conditions?

a) Diabetes mellitus

b) Diabetes insipidus

c) Chronic glomerulonephritis

d) Potomania

Question 15:

A patient presents with frequent urination, nocturia, and enuresis. 24-hour urine volume was measured and recorded to be 7 liters. The urine osmolarity was 260 mOsm/L. ADH assay was performed and the recorded values were reported as 0.8 pg/ml. An MRI of the brain was

Sold by @itachibot

performed and T1 weighted indicated the absence of the bright spot. What is the most likely diagnosis?

a) Nephrogenic DI

b) Primary polydipsia

c) Pituitary DI

d) Mannitol infusion

Answer Key

Question No.	Correct Option
1	b
2	d
3	c
4	d
5	b
6	d
7	b
8	a
9	c
10	b
11	d
12	c
13	a
14	c
15	c

Detailed Explanations

Solution to Question 1:

Oxytocin plays a role in milk ejection. It also helps in initiation or facilitation of labour by stimulating the contraction of uterine smooth muscles.

It stimulates postpartum milk let-down in response to suckling.

Prolactin is the hormone responsible for milk production.

Sold by @itachibot

Osmoregulation is mediated by hormones such as ADH, aldosterone, etc.

Solution to Question 2:

AVP/Antidiuretic hormone (ADH) and oxytocin are stored in Herring bodies at the axon terminals of the posterior pituitary. They are synthesized by two types of magnocellular neurons in the supraoptic and paraventricular nuclei of the hypothalamus.

Solution to Question 3:

AVP binds to V2 receptors on the serosal surface (also called basolateral surface) of renal tubular cells.

This interaction activates adenylyl cyclase and leads to the insertion of water channels called aquaporin 2 on the luminal surface of the cell.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

ADH secretion reduces when the plasma osmolality becomes low. ADH causes water reabsorption and dilutes the plasma, thereby causing a reduction in plasma osmolality.

The antidiuretic effect of ADH is mediated via a G protein-coupled V2 receptor that increases intracellular cyclic AMP, thereby inducing translocation of aquaporin 2 (AQP 2) water channels into the apical membrane.

During and after surgery, surgical stress results in the release of ADH.

Solution to Question 5:

A mutation in the gene for aquaporin 2 channel leads to nephrogenic diabetes insipidus.

The antidiuretic hormone (ADH) binds to V2 receptors in the late distal tubules, collecting tubules, and collecting ducts, and stimulates the movement of an intracellular protein, called aquaporin-2 (AQP-2), to the luminal side of the cell membranes.

Diabetes insipidus (DI) is a condition in which there is the absence of or poor response to anti-diuretic hormone (ADH). This results in reduced permeability of the distal tubules and collecting ducts to water. This furthermore leads the kidneys to excrete large amounts of dilute urine that clinically manifests as polyuria and polydipsia.

DI can be of four types :

Note: Lithium - used to treat bipolar disorder, can cause acquired diabetes insipidus by decreasing the expression of the AQP2 gene.

Sold by @itachibot

Option A : Liddle's syndrome is due to an inherited gain of function mutation of genes regulating the Na channel in collecting duct ENaC, characterized by hypertension, hypokalemia, and metabolic alkalosis.

Option C : Cystic fibrosis is an autosomal recessive disorder due to mutation in CFTR gene that serves as a channel in the luminal plasma membranes of epithelial cells and regulates volume and composition of secretions in exocrine glands.

Option D : Bartter syndrome is due to loss-of-function mutations of transporters and ion channels in thick ascending loop of henle, most commonly, Na-K-2Cl transporter.

Types	Primary pathology	Caused by
Pituitary/ Central DI	Primary deficiency of ADH se cretion	Mc defect: AVP -Neurophysin II gene
Primary polydipsia	Suppression of ADH secretio n by excessive fluid intake	1. Dipsogenic DI : Following head trauma, TB meningitis, Multiple sclerosis.2. Psychog enic DI : Due to psychosis or OCD3. Iatrogenic polydipsia : Due to motivated health ben efits
Gestational DI	Deficiency due to Increased d egradation of ADH	N terminal aminopeptidase p roduced by placenta
Nephrogenic DI	Renal insensitivity to ADH or drugs or genetic mutation	Drugs like lithium - gene enc oding V2 receptor on chXq28

If you purchase... you may have...

Solution to Question 6:

Diabetes insipidus is a clinical component of Wolfram syndrome, and not SIADH.

Characteristics of Wolfram syndrome include:

• Diabetes insipidus

• Diabetes mellitus

• Optic atrophy

• Sensorineural deafness

(Mnemonic DIDMOAD)

Wolfram syndrome is caused by mutations of the WFS1 gene.

Solution to Question 7:

In diabetes insipidus, hypernatremia is observed in the affected patient.

Sold by @itachibot

Low levels of vasopressin lead to the excretion of a large amount of diluted urine. This loss of water from the body leads to hypernatremia.

Solution to Question 8:

The given scenario and the absence of a posterior pituitary bright spot on MRI are suggestive of central diabetes insipidus. Multiple sclerosis causes primary polydipsia, not central DI.

Central DI is caused by a primary deficiency in AVP secretion which usually results from agenesis or irreversible destruction of the posterior pituitary. It can be differentiated from primary polydipsia with the help of an MRI brain.



The MRI given below shows the posterior pituitary bright spot. The arrow points at the posterior pituitary and the arrowhead at the anterior pituitary.



Sold by @itachibot

Solution to Question 9:

The given scenario is suggestive of nephrogenic diabetes insipidus. Isoniazid is not a potential cause of this condition.

Nephrogenic DI develops due to resistance to the antidiuretic action of AVP. A 24-hour urine volume >3 liters is suggestive of polyuria. An ADH value >1 pg/ml is suggestive of nephrogenic DI.



Solution to Question 10:

Metabolic derangements like hypercalcemia and hypokalemia lead to the production of large amounts of dilute urine by causing nephrogenic DI.

Options C & D: Primary polydipsia is of three types:

- Dipsogenic – Due to inappropriate thirst.
- Iatrogenic – Due to an increase in fluid intake due to its presumed health benefits.
- Psychogenic – Due to psychosis or obsessive-compulsive disorder.

Solution to Question 11:

Diabetes insipidus of central or nephrogenic origin are characterized by low urine osmolality and high plasma osmolality.

Sold by @itachibot

Condition	Plasma osmolality	Urine osmolality	Sodium concentration
Central and nephrogenic DI	High	Low	Hypernatremia
Psychogenic polydipsia	Low	Low	Hyponatremia
SIADH	Low	High	Hyponatremia

Solution to Question 12:

The most likely diagnosis for the given clinical scenario would be psychogenic polydipsia.

The normal range of urine osmolality is 500-850 mOsm/L and that of plasma osmolality is 280-295 mOsm/L. Hence in the given scenario, both urine and plasma osmolality are low, along with hyponatremia suggestive of psychogenic polydipsia.

Condition	Plasma osmolality	Urine osmolality	Sodium concentration
Central and nephrogenic DI	High	Low	Hypernatremia
Psychogenic polydipsia	Low	Low	Hyponatremia
SIADH and cerebral salt wasting	Low	High	Hyponatremia

Solution to Question 13:

The given scenario is suggestive of lithium-induced nephrogenic DI. Among the given options thiazide diuretics is the best suited answer.

Thiazides reduce the kidney's ability to excrete free water by increasing proximal tubular water reabsorption (secondary to volume contraction) and block the ability of DCT to form dilute urine. This effect results in an increase in urine osmolality.

Treatment of diabetes insipidus:

• Central and gestational diabetes insipidus:

• Desmopressin an ADH analog

• Nephrogenic diabetes insipidus:

• Adequate water intake along with a low sodium diet

• Thiazide diuretics

• Potassium-sparing diuretics (triamterene and amiloride): Amiloride is the drug of choice for lithium-induced DI

• Prostaglandin synthesis inhibitor (indomethacin)

Sold by @itachibot

• Psychogenic diabetes insipidus:

• Behavioral therapy to reduce fluid consumption

Solution to Question 14:

Fixed specific gravity, in which values remain 1.010 regardless of fluid intake, occurs in chronic glomerulonephritis with severe renal damage.

In chronic kidney disease, as renal function deteriorates, polyuria can be seen due to the impairment of the kidney's ability to concentrate urine.

In diabetes insipidus, although specific gravity is low it is not fixed.

Potomania (beer potomania/beer drinker's potomania/ beer drinker's hyponatremia) is a specific hypo-osmolality syndrome related to massive consumption of beer.

Solution to Question 15:

The features of frequent urination, nocturia, with polyuria(>3L/day), low urine osmolarity(<280 mosm/L), low plasma ADH levels and MRI showing absent pituitary bright spot is suggestive of Pituitary Diabetes Insipidus.



Diabetes Insipidus(DI) :

Treatment:

• For Central DI, synthetic analog of ADH Desmopressin(DDAVP) is used.

• For Nephrogenic DI, a low sodium diet with thiazide diuretic and/or amiloride and prostaglandin synthesis inhibitor is used.

Sold by @itachibot
If this is from someone else, it has been scammed.

Sold by @itachibot

• For primary polydipsia, there are no effective drugs used but for Iatrogenic type patient education is helpful.

Other options:

Option A: In nephrogenic DI, there is polyuria, low urine osmolarity but the ADH levels will be >1pg/mL.

Option B: In primary polydipsia, there is polyuria, low urine osmolarity, ADH levels will be <1pg/mL but a pituitary bright spot present.

Option D: In solute loss (mannitol or hypertonic saline infusion), urine osmolality is >300 mOsm/L.

Types	Primary pathology	Caused by
Pituitary/ Central DI	Primary deficiency of ADH se cretion	AVP -Neurophysin II gene(m ost common gene defect)
Primary polydipsia	Suppression of ADH secretio n by excessive fluid intake	1. Dipsogenic DI : Following head trauma, TB meningitis, Multiple sclerosis.2. Psychog enic DI : Due to psychosis or OCD3. Iatrogenic polydipsia : Due to motivated health ben efits
Gestational DI	Deficiency due to Increased d egradation of ADH	N terminal aminopeptidase p roduced by placenta
Nephrogenic DI	Renal insensitivity to ADH or drugs like lithium or genetic mutation	defect in gene encoding V2 re ceptor on chXq28(most com mon gene defect)

If you pu... you may ...

Sold by @itachibot

Syndrome of Inappropriate Anti-Diuretic Hormone Secretion (SIADH)

Question 1:

Which of the following findings will you not expect to see in a patient with SIADH?

a) Normal serum urea and creatinine levels

b) Low uric acid levels

c) Hypernatremia

d) Normal blood pressure

Question 2:

A 66-year-old chronic smoker who was diagnosed with small cell carcinoma of the lung presents to the casualty with nausea, confusion, and dizziness. After the initial workup, he is diagnosed to have syndrome of inappropriate ADH secretion (SIADH). This condition is characterized by which of the following findings?

a) Hyponatremia and urine sodium excretion > 20 mEq/L

b) Hypernatremia and urine sodium excretion > 20 mEq/L

c) Hyponatremia and urine sodium excretion < 20 mEq/L

d) Hypernatremia and urine sodium excretion < 20 mEq/L

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

SIADH is not observed in which of the following conditions?

a) Multiple sclerosis

b) Oat cell carcinoma of lungs

c) Acute attack of porphyria

d) Primary pulmonary emphysema

Question 4:

Sold by @itachibot

A 47-year-old woman presented with headache and nausea. Examination did not reveal any signs of dehydration and investigations revealed the findings shown below. Which of the following drugs does not lead to this condition?

a) Demeclocycline

b) Vincristine

c) Carbamazepine

d) Nicotine

Question 5:

Which of the following tumors is not associated with syndrome of inappropriate ADH secretion (SIADH)?

a) Ewing's sarcoma

b) Thymoma

c) Chromophobe adenoma

d) Carcinoid

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A child who underwent ventriculoperitoneal shunt surgery for tubercular meningitis with hydrocephalus, develops polyuria and hyponatremia. What could be the probable cause?

a) Central diabetes insipidus

b) SIADH

c) Cerebral salt wasting

d) Shunt block

Question 7:

A patient who presented with acute confusion and vomiting was diagnosed to have SIADH. This condition can be best treated by which of the following methods?

a) Infusion of hypertonic (3%) saline

b) Infusion of hypotonic (0.45%) saline

c) Infusion of normal (0.9%) saline

d) Administration of DDAVP

Sold by @itachibot

Question 8:

What is the drug of choice for the treatment of chronic SIADH?

a) Furosemide

b) Tolvaptan

c) Demeclocycline

d) Fludrocortisone

Question 9:

What causes central pontine myelinolysis?

a) Hyponatremia

b) Hypokalemia

c) Rapid correction of hyponatremia

d) Rapid correction of hypokalemia

If you purchased this from someone else, Sold by @itachibot may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	a
5	c
6	c
7	a
8	b
9	c

Detailed Explanations

Sold by @itachibot

Solution to Question 1:

SIADH does not cause hypernatremia. SIADH is the most common cause of euvolemic hyponatremia.

The excess ADH secreted leads to the gain of water in the body. This stimulates baroreceptors to release atrial natriuretic peptide (ANP). ANP leads to the loss of water and salt in the urine, leading to urine sodium level of > 20 mEq/L in these patients. Thus, the level of water reaches a steady state (euvolemia) but the urinary loss of sodium continues to occur, giving rise to hyponatremia.

SIADH is characterized by low levels of uric acid, normal blood pressure, and normal urea and creatinine levels.

SIADH is diagnosed based on the following criteria (Bartter-Schwartz criteria):

1. Reduced serum osmolality

2. Urine osmolality > Serum osmolality

3. Clinical euvolemia

4. Increased levels of urinary sodium

5. Absence of other causes of hyponatremia like pituitary, renal, adrenal, thyroid or liver diseases.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

The syndrome of inappropriate ADH secretion (SIADH) is associated with increased urinary excretion of sodium (urine sodium excretion > 20 mEq/L), and hyponatremia.

Small cell lung cancer is commonly associated with SIADH due to the ectopic release of antidiuretic hormone.

Solution to Question 3:

Primary pulmonary emphysema is not a cause of SIADH. Respiratory causes of SIADH include asthma, pneumothorax and positive-pressure respiration.

Solution to Question 4:

The given scenario is suggestive of SIADH and demeclocycline does not cause this condition. Demeclocycline causes nephrogenic diabetes insipidus and hence is used in the treatment of chronic SIADH.

The following features in the given case point towards the diagnosis of SIADH:

• Serum osmolality less than 275 mOsm/Kg

• Urine osmolality more than 100 mOsm/Kg

Sold by @itachibot

• Urine sodium more than 20 mmol/L

• Serum sodium level less than 135 mmol/L

Solution to Question 5:

Chromophobe adenoma is a type of pituitary adenoma that causes symptoms of hypogonadism due to pressure effects on the pituitary stalk. It is not a cause of syndrome of inappropriate ADH secretion (SIADH).

Solution to Question 6:

The given scenario is suggestive of cerebral salt wasting. Both SIADH and cerebral salt wasting cause hyponatremia, but polyuria happens in cerebral salt wasting.

Cerebral salt wasting results from hypersecretion of atrial natriuretic peptide. It is seen primarily with central nervous system disorders including brain tumors, head trauma, hydrocephalus, neurosurgery, cerebrovascular accidents, and brain death. It is associated with hyponatremia, polyuria and elevated urinary sodium excretion. Treatment is by restoring intravascular volume with sodium chloride and water.

Clinical parameters to distinguish between SIADH, cerebral salt wasting, and central diabetes insipidus are:

Clinical parameter	SIADH	Cerebral salt wastin g	Central DI
Serum sodium	Low	Low	High
Urine output	Normal or lo w	High	High
Urine sodium	High	Very high	Low
Intravascular volume statu s	Normal or hi gh	Low	Low
Vasopressin level	High	Low	Low

Solution to Question 7:

The given scenario is suggestive of acute symptomatic SIADH and this condition is best treated by the infusion of hypertonic (3%) saline. This mode of treatment also helps in the correction of sodium deficiency that is partly responsible for hyponatremia.

The other form of treatment is to help reduce the concentration of body water by giving an AVP receptor-2 antagonist, conivaptan. Conivaptan blocks the antidiuretic effect of AVP and increases urine output.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Sold by @itachibot

DDAVP is a synthetic analog of AVP and should not be used as it will further increase water retention.

Solution to Question 8:

The drug of choice for the treatment of chronic SIADH is tolvaptan.

Vaptans are selective V2 receptor antagonists that act by increasing urinary water excretion by blocking the antidiuretic effect of AVP.

Treatment options for chronic SIADH include:

• Restriction of fluid intake to below the level of urine output - This is considered the mainstay of treatment. But it also has certain disadvantages as stated below, and hence adding on a drug like vaptan is necessary in most cases.

• Tolvaptan

• Demeclocycline

• Fludrocortisone

Advantages of vaptans over fluid restriction:

• Fluid restriction is usually a burden to the patient, unreliable, impractical, and slow to work.

• Increased thirst is a frequently observed characteristic of hyponatremia in SIADH, and is particularly difficult to suppress. Hence, water restriction in SIADH frequently leads to non-compliance in the long run.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

Rapid correction of hyponatremia may result in central pontine myelinolysis.

Central pontine myelinolysis is an osmotic demyelination syndrome (ODS), wherein rapid correction of hyponatremia results in water being driven out of the cell which has adapted to a chronic state of hypotonicity, leading to damage to the myelin sheath of these cells.

It is characterized by the acute onset of quadriparesis, ataxia, and abnormal extraocular movements.

Sold by @itachibot

Thyroid Disorders - Clinical Features

Question 1:

Which of the following is a specific sign for Graves' disease?

a) Goiter

b) Ophthalmopathy

c) Thyroid bruit

d) Pretibial myxedema

Question 2:

A patient with Graves' disease presents with the following finding. Which statement regarding this condition is false?



a) Acropachy is strongly associated with this condition.

b) It occurs in more than 50% of patients with Graves' disease.

c) It occurs almost always in the presence of moderate or severe form of ophthalmopathy.

d) Skin changes can occur after trauma.

Question 3:

If yo... by @itachibot ...s from someone else, ...scammed.

Sold by @itachibot

A patient from a hilly area was referred with a partial small bowel obstruction confirmed on CECT. Conservative management was tried for a week, but he continued to deteriorate and was taken up for emergency laparotomy. On post-operative day 7, he developed atrial fibrillation. Investigations revealed elevated free T3 and free T4 levels. What is the most probable cause?

a) Jod-Basedow phenomenon

b) Wolff-Chaikoff effect

c) Basedow disease

d) Plummer disease

Question 4:

What is the most common symptom of thyrotoxicosis?

a) Palpitation

b) Fatigue and weakness

c) Heat intolerance

d) Hyperactivity and irritability

Question 5:

Which of the following statements is not true about thyroid storm?

a) Symptoms are vomiting, diarrhoea, jaundice and altered mentation

b) The mortality rate is 10% with treatment

c) Can happen after non-thyroid surgeries

d) High doses of propranolol decrease T4 → T3 conversion

Question 6:

Which of the following conditions is not associated with amiodarone therapy?

a) Transient suppression of thyroid gland

b) Hypothyroidism

c) Thyrotoxicosis

d) Silent thyroiditis

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Question 7:

Which of the following statements about amiodarone-induced thyrotoxicosis is false?

a) Type 2 AIT is associated with intrinsic abnormalities of the thyroid gland

b) Type 2 AIT is due to lysosomal destruction of thyroid gland.

c) Type 1 AIT is due to Jod-Basedow phenomenon.

d) Colour flow doppler shows increased vascularity in Type 1

Question 8:

Which of the following statements is true about non-toxic multinodular goitre?

a) It is more commonly observed in women

b) It does not occur in iodine-deficient regions

c) Most patients with non-toxic MNG have hypothyroidism

d) Risk of malignancy is higher than in cases of solitary thyroid nodule

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

What is the major difference between non-toxic and toxic multinodular goitre?

a) Pathogenesis

b) Presence of functional autonomy

c) Polyclonal nodules

d) Monoclonal nodules

Question 10:

What is the most common thyroid malignancy?

a) Papillary carcinoma

b) Follicular carcinoma

c) Medullary thyroid carcinoma

d) Anaplastic carcinoma

Question 11:

Sold by @itachibot

The following patients presented with thyroid nodules. Who has the least risk of developing thyroid carcinoma?

a) A 55-year-old woman with no voice abnormalities and nodule size of 3 cms

b) A 65-year-old woman with vocal cord paralysis and hoarse voice and nodule size of 3.5 cms

c) A 60-year-old man without any voice abnormalities and nodule size of 2.5 cms

d) A 43-year-old woman with no voice abnormalities and nodule size of 4.5 cm

Question 12:

Which of the following statements is not true about thyroid hormone resistance?

a) Inherited as an autosomal recessive disorder

b) Characterized by elevated thyroid hormone levels and normal or elevated TSH

c) Classic forms are caused by mutations in TR-β gene

d) No treatment is indicated usually

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 13:

Patients with what condition present with normal thyroglobulin levels?

a) Graves' disease

b) Toxic multinodular goiter

c) Thyrotoxicosis factitia

d) Toxic adenoma

Question 14:

A patient presented with fever and painful neck swelling. She gives a history of sore throat and cough one week ago. Lab investigations revealed elevated T3 and T4 levels and elevated ESR. Which of the following statements about this condition is false?

a) Treatment of choice is supportive care and NSAIDs

b) Women are generally affected three times more often than men

c) Both thyrotoxic and hypothyroid phases can be observed

d) It is autoimmune in origin

Sold by @itachibot

Question 15:

Which of the following conditions is not associated with hypothyroidism?

a) Graves' disease

b) Hashimoto's thyroiditis

c) Hypopituitarism

d) Lithium-induced Goiter

Question 16:

A 4-month-old baby presented with a history of constipation, lethargy, and has poor activities. Clinical examination reveals jaundice, hypotonia and there was no palpable goitre. Laboratory investigations revealed low T4 and elevated TSH levels. Which of the following is not true about this condition?

a) Most common cause is inborn defects in the synthesis of thyroid hormone

b) The developmental abnormalities are more common in girls

c) Associated with macroglossia

d) High circulating T4 level is needed to normalize TSH

Question 17:

Patients with autoimmune hypothyroidism have a higher predisposition for all except:

a) Type I diabetes mellitus

b) Type II diabetes mellitus

c) Pernicious anemia

d) Vitiligo

Question 18:

Most common symptom seen in patients with hypothyroidism is?

a) Cold intolerance

b) Weight gain

c) Tiredness and weakness

d) Dry skin

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

Question 19:

An elderly patient presents to the ER with symptoms suggestive of myxedema coma. Which of the following is unlikely to be seen in this patient?

a) Hypoglycemia

b) Hyponatremia

c) Hypercapnia

d) Hyperthermia

Question 20:

A patient came to the hospital with complaints of lethargy, increased sleep, and weight gain. Investigations revealed low plasma TSH concentration. However, on the ■administration of TRH, the TSH levels increased. Which of the following conditions is likely in this patient?

a) Hyperthyroidism due to disease in the pituitary

b) Hypothyroidism due to disease in the pituitary

c) Hypothyroidism due to disease in the hypothalamus

d) Hyperthyroidism due to disease in the hypothalamus

Question 21:

A middle-aged woman with a history of constipation, dry skin and menorrhagia presents to the ER with altered sensorium, non-pitting edema, hypothermia, bradycardia, and hypotension. What is the most likely diagnosis?

a) Septic shock

b) Cardiogenic shock

c) Myxedema coma

d) Hypoglycemia

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	b
2	b
3	a
4	d
5	b
6	d
7	a
8	a
9	b
10	a
11	a
12	a
13	c
14	d
15	a
16	a
17	b
18	c
19	d
20	c
21	c

Detailed Explanations

Solution to Question 1:

Ophthalmopathy is the specific sign of Graves' disease.

Glycosaminoglycan deposition, lymphoid infiltrates and fatty infiltration are responsible for ophthalmopathy and dermopathy.

Grave's disease is an autoimmune disorder due to the production of TSH-receptor stimulating immunoglobulins (TSI) thereby causing hyperthyroidism.

It is characterized by the triad of:

• Thyrotoxicosis

• Infiltrative ophthalmopathy (exophthalmos)

• Dermopathy (pretibial myxedema)

Sold by @itachibot

Ophthalmopathy can sometimes precede or follow thyrotoxicosis, accounting for some cases of euthyroid ophthalmopathy.

Solution to Question 2:

The given scenario along with the image is suggestive of thyroid dermopathy. It occurs in <5% of patients with Graves' disease. Thyroid dermopathy is almost always associated with ophthalmopathy.

Thyroid dermopathy is most frequently observed over the anterolateral parts of the lower leg. Hence, it is termed pretibial myxedema. Skin changes can occur at other sites, particularly after trauma. The typical lesion is a noninflamed, indurated plaque with a deep pink or purple color and orange peel-textured papules.

Thyroid acropachy refers to a form of clubbing rarely observed in Graves' disease. It is strongly associated with thyroid dermopathy.

Thyroid dermopathy usually does not need any active treatment, but if necessary, treatment includes topical, high-potency glucocorticoid ointment under an occlusive dressing or octreotide can be useful in some cases.

The following image shows thyroid dermopathy and acropachy:

Solution to Question 3:

The given clinical scenario is suggestive of the Jod-Basedow phenomenon, likely due to the administration of an iodine-containing contrast agent during CECT.

This phenomenon is characterized by hyperthyroidism due to enhanced thyroid hormone production following the administration of contrast agents and other iodine-containing substances. This occurs in:

Sold by @itachibot

• Patients with autonomous thyroid nodules

• Patients with pre-existing subclinical hypothyroidism (endemic iodine deficiency in hilly areas)

Option B: Wolff-Chaikoff effect is characterized by the transient inhibition of thyroid iodide organification by excess iodine, thereby decreasing thyroid hormone synthesis.

Option C: Basedow disease is another name for Graves' disease.

Option D: Plummer disease is another name for toxic multinodular goitre.

Solution to Question 4:

Hyperactivity and irritability are considered as the most common symptoms of thyrotoxicosis.

The following table summarizes the symptoms and signs of thyrotoxicosis in descending order of frequency:

Rarely, there can be an associated mid-systolic heart murmur called the Means-Lerman scratch. This is heard in the upper part of the sternum caused by the rubbing of the pericardium and the pleura, due to the hyperdynamic circulation.

Symptoms	Signs
Hyperactivity, irritability, dysphoriaHeat intolerance and sweatingPalpitationsFatigue and weaknessWeight loss with increased appetiteDiarrhoeaPolyuriaOligomenorrhea, loss of libido	Tachycardia, atrial fibrillation in the elderlyTremorGoitreWarm, moist skinMuscle weakness, proximal myopathyLid retraction or lagGynecomastia

If you ... by @itachibot ... from someone else, you ... scammed

Solution to Question 5:

The mortality rate of thyrotoxicosis crisis is almost 30% even with treatment.

The usual causes for mortality are:

• Cardiac failure

• Arrhythmia

• Hyperthermia

Solution to Question 6:

Silent thyroiditis is not associated with amiodarone therapy.

Amiodarone has the following effects on the thyroid gland:

Sold by @itachibot

• Acute, transient suppression of thyroid function i.e., Wolff-Chaikoff effect

• Hypothyroidism in patients susceptible to the inhibitory effects of a high iodine load

• Thyrotoxicosis may be caused by Jod-Basedow phenomenon due to the iodine load, in the setting of multinodular goitre or incipient Graves' disease. It can also be caused by a thyroiditis-like condition.

Painless thyroiditis, or silent thyroiditis, occurs in patients with underlying autoimmune thyroid disease and has a clinical course similar to that of subacute thyroiditis.

The condition can occur in up to 5% of women 3–6 months after pregnancy and is then termed postpartum thyroiditis.

If yo Sold by @itachibot sed this from someone else, e been scammed.

Solution to Question 7:

Type 1 AIT is associated with intrinsic abnormalities of the thyroid gland.

Amiodarone is structurally similar to thyroid hormones and contains 39% iodine by weight. Amiodarone-induced thyrotoxicosis (AIT) is the most common manifestation of the effects of amiodarone on the thyroid.

Administration of amiodarone should be stopped if possible.

Near-total thyroidectomy rapidly decreases thyroid hormone levels and may be the most effective long-term clinical solution.

Type 1 AIT	Type 2 AIT
Anunderlying diseaseis present.E.g., Grave's disease, Multinodular goitre etc	No underlying disease
Due to theJod-Basedowphenomenon	Due todrug-inducedlysosomal destruction of thyroid tissue
Increased vascularity on colour doppler	Decreased vascularity on colour doppler

Solution to Question 8:

Multinodular goitre (MNG) is more commonly observed in women. It is common in iodine-deficient regions.

Most patients are asymptomatic and euthyroid. Sudden pain may occur due to haemorrhage. TFTs should be performed in all patients to exclude thyrotoxicosis or hypothyroidism.

The risk of malignancy due to MNG is similar to that due to solitary nodules. Hoarseness, reflecting the involvement of laryngeal nerves implicate malignancy.

Most non-toxic MNGs can be managed conservatively. Levothyroxine if used, should be started at low doses and then increased gradually. Radioiodine therapy is implemented for large goitres as it

Sold by @itachibot

decreases goitre size and can ablate regions of autonomy.

Solution to Question 9:

The major difference between toxic and non-toxic multinodular goiter (MNG) is the presence of functional autonomy in toxic MNG.

The pathogenesis of toxic MNG appears to be similar to that of non-toxic MNG. In both types, many nodules are polyclonal whereas others are monoclonal and vary in their clonal origins.

Most patients with non-toxic MNG are asymptomatic and euthyroid. However, patients with toxic MNG have subclinical or mild hyperthyroidism.

Sold by @itachibot
If you purcha
you
from someone else
scammed.

Solution to Question 10:

Papillary carcinoma is the most common thyroid malignancy.

Prevalence of different types of thyroid malignancies:

Type of malignancy	Occurence
Papillary	80-90 %
Follicular	5-10%
Medullary	<10%
Others	1-2%

Solution to Question 11:

Among the options, the female patient with age between 20 and 65 years and no voice abnormalities and nodule size < 4 cm has the least risk for development of thyroid carcinoma.

Risk factors for the development of thyroid carcinoma in patients with thyroid nodules include:

• Age less than 20 or greater than 65 years

• Male gender

• Increased nodule size > 4 cm

• Vocal cord paralysis and hoarse voice

• New or enlarged neck mass

• Family history of thyroid cancer, MEN2, or other syndromes associated with thyroid malignancy (e.g., Cowden's syndrome, familial polyposis, Carney complex)

Sold by @itachibot

• Nodule fixed to adjacent structures

• Exposure to radiation in childhood or adolescence

• Extrathyroidal extension

• Lateral cervical lymphadenopathy

Solution to Question 12:

Thyroid hormone resistance is an autosomal dominant disorder.

It is characterized by elevated thyroid hormone levels and inappropriately elevated or normal TSH levels.

Classic forms of the disease are attributed to the mutation in TR-β gene. Patients with this condition are often asymptomatic, as hormone resistance is partial and compensated by the increase in thyroid hormone levels.

In most patients, no treatment is indicated.

Solution to Question 13:

Serum thyroglobulin levels are normal in patients with thyrotoxicosis factitia, as it is caused by the self-administration of thyroid hormone. Its level is elevated in all other types of thyrotoxicosis.

Thyroglobulin (Tg) levels are particularly increased in thyroiditis, implicating destruction of thyroid tissue and release of thyroglobulin.

The primary course of action for determining the thyroglobulin levels, however, is in the clinical follow-up of thyroid cancer patients.

Following total thyroidectomy and radiofrequency ablation, thyroglobulin levels become undetectable. In the absence of anti-Tg antibodies, measurable levels of Tg are indicative of incomplete ablation or recurrent cancer.

Solution to Question 14:

The given scenario is suggestive of De Quervain's thyroiditis, and it is not autoimmune in origin. It is subacute granulomatous thyroiditis that occurs secondary to viral infections.

Many viruses such as mumps, influenza, coxsackie, adenoviruses, and echoviruses have been clinically implicated in this condition. Peak incidence is observed at 30-50 years and women are generally affected three times more often than men.

Clinically, it presents as a painfully enlarged thyroid and is sometimes accompanied by fever.

It has three distinct phases:

• Thyrotoxic phase

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Hypothyroid phase

• Recovery phase

Treatment includes large doses of aspirin, NSAIDs, steroids and/or beta-blockers to relieve the observable clinical symptoms. Antithyroid drugs do not have any role. In case the hypothyroid phase is prolonged - levothyroxine can be used.

Solution to Question 15:

Graves' disease is not associated with hypothyroidism.

Solution to Question 16:

The given scenario is suggestive of congenital hypothyroidism. The most common cause of this condition is thyroid gland dysgenesis, seen in about 80-85% of cases of neonatal hypothyroidism.

In the remaining cases of neonatal hypothyroidism, it is due to inborn defects in thyroid hormone synthesis and rarely due to TSH receptor antibodies. These developmental abnormalities are twice as common in girls.

Clinical features include:

• Prolonged jaundice

• Umbilical hernia

• Enlarged tongue

• Hypotonia

• Delayed bone maturation

Permanent neurological damage results if treatment is delayed.

T4 requirements are relatively high during the first year of life. A high circulating T4 level is needed to normalize TSH.

Early treatment with T4 results in normal IQ levels by preventing neurodevelopmental abnormalities.

Solution to Question 17:

Autoimmune hypothyroidism is usually not associated with type II diabetes mellitus.

Genetic risk factors for autoimmune hypothyroidism has been documented with HLA-DR3, HLA-DR4, HLA-DR5. A weak association also exists with CTLA4 gene polymorphism.

This may explain the relationship between autoimmune hypothyroidism and type I diabetes mellitus, Addison's disease, pernicious anemia, and vitiligo.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Sold by @itachibot

Less commonly encountered clinical associations include:

• Celiac disease

• Dermatitis herpetiformis

• Chronic active hepatitis

• Rheumatoid arthritis

• Myasthenia gravis

• Systemic lupus erythematosus

• Sjogren's syndrome.

Solution to Question 18:

The most common symptom of hypothyroidism is tiredness and weakness.

Symptoms and signs in decreasing order of frequency are given below:

Symptoms	Signs
Tiredness, weaknessDry skin Cold intoleranceHair lossDifficulty concentrating and poor memoryConstipationWeight gain with poor appetiteDyspneaHoarse voiceMenorrhagia (later oligomenorrhea or amenorrhea)ParesthesiaImpaired hearing	Dry coarse skin, cool peripheral extremitiesPuffy face, hands, and feet (myxedema)Diffuse alopeciaBradycardiaPeripheral edemaDelayed tendon reflex relaxationCarpal tunnel syndromeSerous cavity effusions

If you @itachibot you from someone else, unmed.

Solution to Question 19:

Hyperthermia is least likely in this patient. Clinical manifestations of myxedema coma include decreased consciousness, seizures and hypothermia.

Myxedema coma is a serious condition with a mortality rate of 20-40%. The clinical outcomes are independent of the detected T4 and TSH levels.

Clinical features of myxedema crisis include:

• Impaired cognition, ranging from confusion to somnolence to coma (myxedema coma)

• Convulsions and abnormal CNS signs.

• Profound hypothermia (can reach 23°C), hypoventilation, hyponatremia, hypoglycemia, hypoxemia, hypercapnia, and hypotension.

• Rhabdomyolysis and acute kidney injury.

Sold by @itachibot

Solution to Question 20:

The presenting symptoms of lethargy, increased sleep, and weight gain are suggestive of hypothyroidism. An increase in TSH on administration of TRH confirms the lack of stimulus for TSH secretion from the hypothalamus.

Although decreased TSH levels can be seen in hyperthyroidism, clinical features such as lethargy, increased sleep and weight gain are not. This pushes the diagnosis in favor of hypothyroidism.

If the disease was in the pituitary, there would be no increase in TSH secretion on administration of TRH (Option B).

Normally, the thyrotropin releasing hormone (TRH) is released by the hypothalamus. It stimulates the pituitary gland to release the thyroid-stimulating hormone (TSH), which in turn stimulates the thyroid gland to produce T3 and T4 hormones.

Low TSH levels may be seen in the following cases:

Disease	Pathogenesis	Presentation
Hyperthyroidism	Feedback inhibition of TSH r elease	Symptoms of hyperthyroidis m
Primary disease of the pituita ry	Decreased release of TSH, th ereby causing a decreased rel ease of thyroid hormones	Symptoms of hypothyroidism. Administration of TRH does not elevate the TSH levels in this case since the pituitary g land is abnormal
Primary disease of the hypot halamus	Decreased release of TRH lea ding to decreased TSH and h ypothyroidism	Administration of TRH cause s elevation of TSH levels sinc e the pituitary gland is norm al.



Sold by @itachibot

Solution to Question 21:

The given clinical history is typical of a case of hypothyroidism and the diagnosis is myxedema coma, which is a serious complication of hypothyroidism.

Myxedema coma is a part of a larger clinical picture of myxedema crisis. Myxedema crisis refers to severe, life-threatening manifestations of hypothyroidism. It is often precipitated by the administration of sedatives, antidepressants, hypnotics, anaesthetics, or opioids.

Clinical features of myxedema crisis :

• Impaired cognition, ranging from confusion to somnolence to coma (myxedema coma).

• Convulsions and abnormal CNS signs.

• Profound hypothermia (can reach 23°C), hypoventilation, hyponatremia, hypoglycemia, hypoxemia, hypercapnia, and hypotension.

• Rhabdomyolysis and acute kidney injury

• A high mortality rate

Other options:

Option A: Without a history of or focus of infection, septic shock is unlikely.

Option B: Cardiogenic shock would present with symptoms of left heart failure i.e. pitting pedal edema, abdominal distension, prominence of neck veins, and dyspnea.

Option D: Hypoglycemia usually presents with sweating, palpitations, and tachycardia

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Thyroid Disorders - Management

Question 1:

A middle-aged woman who presented with tremors, palpitations, weight loss, and menstrual irregularities is diagnosed with Graves' disease. Which of the following statements would be correct with respect to her thyroid function test?

a) Low TSH levels

b) Low free T4 levels

c) Low serum T3 levels

d) Radioactive Iodine Uptake (RAIU) at 24 hours below normal

Question 2:

What is the most common pattern of thyroid hormone levels in a patient with sick euthyroid syndrome?

a) 1

b) 3

c) 2

d) 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

On thyroid function test analysis, TSH value was much higher than normal with a low free T4. What is the diagnosis?

a) Subclinical hypothyroidism

b) Hyperthyroidism

c) Secondary hypothyroidism

d) Primary hypothyroidism

Question 4:

Sold by @itachibot

The radio-iodine uptake scan of a patient evaluated for swelling of the thyroid is shown below. Which of the following is the most likely diagnosis?



a) Hypersecreting adenoma

b) Papillary Carcinoma thyroid

c) Graves disease

d) Lateral aberrant thyroid

Sold by @ If you purchased this from you may have been scamm. else,

Question 5:

Which of the following drugs can cause the condition shown below?

Sold by @itachibot

a) Methimazole

b) Propylthiouracil

c) Radioactive iodine

d) Levothyroxine

Question 6:

Which of the following patients cannot undergo radioiodine therapy?

a) Patients with renal disease

b) Patients with cardiac disease

c) Lactating mother

d) Patients with severe ophthalmopathy

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

A 25-year-old woman presents with palpitations. She informs you that a urine pregnancy test done one week ago was positive. Her investigations reveal hyperthyroidism. Which of the following antithyroid drugs can be prescribed for her?

a) Propylthiouracil

b) Methimazole

c) Carbimazole

d) Propranolol

Question 8:

A patient undergoing radioactive iodine therapy for a non-toxic multinodular goitre later develops altered sensorium, fever, and vomiting. What is the anti-thyroid drug of choice in this patient?

a) Propranolol

b) Iodide

c) Propylthiouracil

d) Methimazole

Question 9:

Sold by @itachibot

A patient presents with a diffusely enlarged thyroid gland. However, she has no features of hyperthyroidism. Which of the following statements about her condition is false?

a) It is also referred to as simple goitre or colloid goitre

b) Most commonly caused by iodine deficiency

c) Surgery is commonly indicated

d) Thyroid function tests should be performed in all patients with goiter

Question 10:

Which of the following statements about silent thyroiditis is true?

a) Painful thyroiditis

b) Glucocorticoids are not indicated for treatment

c) It is not associated with thyrotoxicosis

d) It is not associated with hypothyroidism

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 11:

A middle-aged woman with fatigue, depression and weight gain was found to have hypothyroidism. Which of the following statements is false about the treatment of her condition?

a) The replacement dose of thyroid is 1.6 µg/kg body weight

b) Complete relief from symptoms might occur within 3-6 months after initiation of treatment

c) TSH responses should be determined 2 months following the initiation of treatment or change in dose

d) Once TSH level is stable, TSH measurement is recommended at annual intervals

Question 12:

A 24-year-old primigravida at 10 weeks of gestational age presents to your OPD. She is on oral levothyroxine supplementation for hypothyroidism. What is the target TSH level to be maintained in this patient?

a) <2.5 mIu/L

b) <3.0 mIu/L

c) <3.5 mIu/L

Sold by @itachibot

d) <4.0 mIu/L

Question 13:

An elderly woman with hypothyroidism presents to the ER with decreased consciousness, hypothermia, sinus bradycardia, and hypotension. Her family members reveal that she has not been taking her thyroid medications recently. Investigations reveal elevated TSH levels and reduced T3 and T4 levels. All of the following can be used in the treatment except:

a) IV levothyroxine

b) Liothyronine (T3)

c) Parenteral steroids

d) Hypotonic IV fluids

If you bought this from someone else, you got scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	a
5	a
6	c
7	a
8	c
9	c
10	b
11	b
12	a
13	d

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

In Graves' disease, the level of serum TSH is low, free T4 is high and serum T3 is high. In patients with thyrotoxicosis, the radioactive iodine uptake (RAIU) at 24 hours is characteristically elevated.

Grave's disease is characterized by thyrotoxicosis caused by thyroid-stimulating immunoglobulins (TSIs) synthesized by lymphocytes in the thyroid gland, bone marrow, and lymph nodes. Thyroid peroxidase (TPO) and thyroglobulin (Tg) antibodies are also seen. This results in excess production of free T3 and free T4 levels and in turn reduces serum TSH levels.

RAIU is used to differentiate Grave's from other thyroid disorders, here there is diffuse high uptake. Measurement of TRab and TPOs is also done to confirm the diagnosis if the diagnosis clinically is unclear.



Solution to Question 2:

The most common hormone pattern observed in sick euthyroid syndrome (SES) is a decrease in total and unbound T3 levels with normal levels of T4 and TSH. This pattern is called low T3 syndrome.

SES is caused by an acute severe illness in the absence of underlying thyroid disease. T4 to T3 conversion via peripheral deiodination is impaired leading to a fall in T3 corresponding to the severity of illness. There is an adaptive increase in the level of reverse T3 (rT3).

In very sick patients, there can be a dramatic fall in both total T4 and T3 levels. This pattern is called low T4 syndrome and is associated with a poor prognosis.

Solution to Question 3:

An elevated TSH value with low free T4 is suggestive of primary hypothyroidism (Option D).

Option A: In subclinical hypothyroidism, free T4 levels remain normal despite elevated TSH.

Option B: TSH is low in hyperthyroidism.

Sold by @itachibot

Option C: The diagnosis of secondary hypothyroidism is made along with a deficiency of other pituitary hormones. TSH levels may be low, normal, or slightly increased. The elevation in TSH levels can be attributed to the secretion of immunoreactive but biologically inactive forms of TSH. To confirm the diagnosis, it is essential to detect a low level of unbound T4.

The diagnostic approach to hypothyroidism is given in the flowchart below:



Solution to Question 4:

The given image showing localized overactivity (increased uptake) in the left lobe of the thyroid i.e., hot nodule, with suppression of the remainder of the gland, is suggestive of a hypersecreting adenoma.

Other options:

Option B: The likelihood of malignancies like papillary Carcinoma thyroid is low (5%) in the case of a hot nodule.

Option C: Graves disease shows diffusely increased uptake on a thyroid scan.

Option D: Lateral aberrant thyroid presents with swelling of cervical lymph nodes and not thyroid. It is due to skip metastasis from papillary thyroid carcinoma to the ipsilateral cervical lymph nodes.

The thyroid scan appearance in different conditions is shown in the image below:

Sold by @itachibot



Diagnostic isotope studies, such as 99mTc-pertechnetate and iodine-123, are becoming less common in thyroid imaging because some warm swellings (5%) which are actually malignant could be mistaken as benign. Isotope scanning is most valuable when assessing a hyperthyroid patient presenting with a nodule or nodularity in the thyroid. It helps to identify overactivity in the gland, which can differentiate between a toxic nodule that suppresses the rest of the gland and a toxic multinodular goitre with multiple areas of increased uptake.

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 5:

The given image shows the absence of scalp hair and skin, along with calvarial defects(skull bone defects), suggestive of aplasia cutis congenita. This is commonly associated with carbimazole and methimazole.

These drugs can also cause other defects such as:

- Choanal atresia
- Tracheoesophageal fistula
- Fetal goiter

Solution to Question 6:

Lactating mothers cannot undergo radioiodine therapy. Pregnancy and breastfeeding are absolute contraindications of radioiodine therapy. Patients can safely conceive 6 months after undergoing radioiodine treatment.

Radioiodine therapy in patients with severe ophthalmopathy can be given with caution along with prednisone to prevent exacerbation of ophthalmopathy.

Sold by @itachibot

Solution to Question 7:

The patient in the given clinical scenario is in the 1st trimester of pregnancy. Propylthiouracil is used in the 1st trimester of gestation because it has lower teratogenic effects than carbimazole.

Carbimazole and methimazole are associated with teratogenic effects such as

• Aplasia cutis congenita

• Choanal atresia

• Treacheosopageal fistula

• Fetal goiter

However, propylthiouracil use is limited to the first trimester due to the risk of hepatotoxicity. It is replaced with carbimazole or methimazole in the 2nd and 3rd trimesters of the pregnancy.

The lowest effective antithyroid drug dose should be used throughout gestation. This is to maintain the maternal serum-free T4 level in the upper limit of the non-pregnant normal reference range.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 8:

The given scenario is suggestive of thyroid storm and propylthiouracil is the drug of choice in this patient. This is due to its inhibitory action on peripheral T4 → T3 conversion.

PTU is administered in large doses (500–1000 mg loading dose and 250 mg every 4h) administered orally / via nasogastric tube / per rectum.

Propranolol is also given to reduce symptoms like tachycardia and other adrenergic manifestations.

Solution to Question 9:

The given scenario is suggestive of a non-toxic diffuse goitre. Surgery is rarely indicated in this condition, except for cosmetic reasons, tracheal compression or obstruction of the thoracic inlet.

It is sometimes called simple goitre - because of the absence of nodules, and colloid goitre - because of the presence of uniform follicles that are filled with colloid. It is most commonly caused by iodine deficiency and is often termed endemic goitre.

Most diffuse goitres are benign and asymptomatic. Examination reveals a symmetrically enlarged, non-tender, generally soft gland without palpable nodules. TFTs should be performed in all patients to exclude thyrotoxicosis or hypothyroidism.

If markedly enlarged, it can cause tracheal/oesophagal compression.

Substernal goitre may obstruct the thoracic inlet leading to a positive Pemberton's sign i.e., facial congestion on bilateral arm elevation due to venous occlusion.

Sold by @itachibot

Solution to Question 10:

Glucocorticoid treatment is not indicated in silent thyroiditis.

The condition is called silent thyroiditis, as it is painless. When it occurs 3-6 months after pregnancy, it is termed postpartum thyroiditis. The condition is observed in up to 5% of women and it is 3 times more common in women with Type 1 diabetes mellitus.

Typically, patients experience a brief phase of thyrotoxicosis lasting 2-4 weeks, followed by hypothyroidism for 4-12 weeks, and then clinical resolution. Often, only one phase is apparent.

In addition to painless goitre, silent thyroiditis can be distinguished from subacute thyroiditis by a normal ESR value and the presence of TPO antibodies.

Severe thyrotoxic symptoms can be managed with a brief course of propranolol, a 20–40 mg dose administered three or four times daily.

Thyroxine replacement may be needed for treating the hypothyroid phase but should be withdrawn after 6–9 months, to enable patient recovery.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

The clinical effects of levothyroxine replacement are slow to appear. Patients may not experience full relief from symptoms until 3–6 months after normal TSH levels are restored.

If there is no residual function, the daily replacement dose of levothyroxine is usually 1.6 µg/kg body weight (typically 100–150 µg), ideally taken at least 30 minutes before breakfast.

TSH responses are gradual and should be determined 2 months following the initiation of treatment or change in dose. Once the full replacement is achieved and TSH levels are stable, TSH measurement is recommended at annual intervals.

In patients who have developed hypothyroidism following treatment of Graves' disease, lower doses are needed for replacement. This is because there is often underlying autonomous function, necessitating lower replacement doses (typically 75–125 µg/d).

Solution to Question 12:

The target TSH level to be maintained in the first trimester of pregnancy is <2.5 mIU/L. For second and third trimester the target is <3.0 mIU/L.

Thyroid function must be evaluated by testing for TSH levels every 4 weeks in the first trimester and less frequently after 20 weeks. The levothyroxine dose may be increased by 50% during pregnancy. Following delivery, doses of thyroxine are usually restored to that of pre-pregnancy levels.

Solution to Question 13:

Sold by @itachibot

The given scenario is suggestive of myxedema coma. Hypotonic IV fluids are avoided in the course of treatment of this condition. However, hypertonic saline is used.

Hypotonic fluids may exacerbate water retention and dilutional hyponatremia secondary to impaired renal perfusion and inappropriate vasopressin.

Treatment of myxedema coma includes:

• Levothyroxine (T4) can be administered as a single IV bolus of 500 µg. Although further doses are strictly not required, it is continued at 50-100 µg/day.

• An alternative mode of treatment is the IV administration of liothyronine (T3) or via NG tubes with doses ranging from 10-25 µg every 8-12 hours in case T4 to T3 conversion in myxedema is impaired.

• Parenteral hydrocortisone should be administered to treat a condition of impaired adrenal reserve in profound hypothyroidism.

• Hypertonic saline or IV glucose may be recommended for treatment.

• External warming is indicated only if the temperature is less than 30oC.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Multiple Endocrine Neoplasia

Question 1:

Multiple endocrine neoplasia (MEN) type 1 is also called as _____ .

a) Wermer's syndrome

b) Sipple syndrome

c) Weber's syndrome

d) Werner syndrome

Question 2:

Which of the following conditions is not observed in MEN1 syndrome?

a) Posterior pituitary tumors

b) Foregut carcinoid

c) Parathyroid adenoma

d) Angiofibroma

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

A 35-year-old man presented with abdominal pain, nausea, and diarrhea. He had a history of hyperparathyroidism and a transsphenoidal prolactinoma resection. Which of the following is the most common functional enteropancreatic tumor seen in this condition?

a) Glucagonoma

b) Insulinoma

c) Gastrinoma

d) VIPoma

Question 4:

Which of the following is the most common site of gastrinoma in MEN 1 syndrome?

a) Jejunum

b) Ileum

Sold by @itachibot

c) Duodenum

d) Stomach

Question 5:

A 30-year-old man presents to the OPD with complaints of recurrent episodes of weakness, headache, sweating, and fainting after fasting. He tells that the symptoms improve after the consumption of food. There is a history of MEN1 syndrome running in the family. Serum C peptide levels are found to be elevated. What is the definitive treatment of choice in this patient?

a) Frequent consumption carbohydrate-rich meals

b) Octreotide

c) Diazoxide

d) Surgery

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which of the following is the most frequently observed cutaneous manifestation of MEN1 syndrome?

a) Lipoma

b) Angiofibroma

c) Collagenoma

d) Neurofibroma

Question 7:

A 22-year-old man who was diagnosed with MEN1 syndrome presents to your OPD for his annual checkup. Which of the following investigations will you not do as part of annual screening for this patient?

a) Fasting glucose levels

b) Serum calcium levels

c) Secretin study

d) Serum gastrin levels

Sold by @itachibot

Question 8:

What is the most common component of Sipple's syndrome?

a) Medullary thyroid cancer

b) Hyperparathyroidism

c) Pituitary adenoma

d) Gastrinoma

Question 9:

Which of the following features would you not expect to see in a patient with MEN2 syndrome?

a) Pituitary adenoma

b) Parathyroid adenoma

c) Medullary thyroid cancer

d) Pheochromocytoma

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Question 10:

Which of the following features would you not expect to see in a patient with MEN2B syndrome?

a) Mucosal neuromas

b) Marfanoid habitus

c) Medullated corneal nerve fibers

d) Meningioma

Question 11:

MEN4 syndrome is caused by a mutation in which of the following genes?

a) PPKAR1A

b) RET proto-oncogene

c) PRKAR1B

d) CDKN1B

Sold by @itachibot

Question 12:

Which of the following malignancies would you not expect to see in a patient with MEN4 syndrome?

a) Pituitary adenoma

b) Parathyroid adenoma

c) Medullary thyroid carcinoma

d) Renal tumors

Question 13:

A patient presented with abdominal pain and swelling in his scrotum and was diagnosed to have testicular carcinoma. There is a history of MEN syndrome running in his family. Which of the following conditions is he likely to have?

a) MEN4

b) MEN2A

c) MEN1

d) MEN2B

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	c
5	d
6	b
7	c
8	a
9	a
10	d
11	d
12	c

Sold by @itachibot

13	a

Detailed Explanations

Solution to Question 1:

MEN1 is also referred to as Wermer's syndrome.

Solution to Question 2:

Anterior pituitary tumours are associated with MEN1 syndrome, not posterior pituitary tumors.

Solution to Question 3:

The given scenario is suggestive of MEN1 syndrome and gastrinoma is the most common functional enteropancreatic tumor in MEN1 / Wermer's syndrome.

Enteropancreatic tumors in MEN1 syndrome are:

• Non-functioning pancreatic tumors (most common overall)

• Functioning pancreatic tumors :

• Gastrinoma - 50% (most common among functioning pancreatic tumours)

• Insulinoma - 30%

• Glucagonoma

• VIPoma (least common)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

Most common site of gastrinoma in MEN 1 syndrome is the first and second part of duodenum.

MEN 1 syndrome is characterized by the triad of tumors involving the parathyroids, pancreatic islets, and anterior pituitary.

Gastrinoma is the most common pancreatic islet tumor in MEN 1 syndrome and if it is associated with marked gastric acid production and recurrent peptic ulcers, the condition is called Zollinger Ellison syndrome.

Solution to Question 5:

Sold by @itachibot

The given clinical scenario along with elevated serum C peptide levels points towards a diagnosis of insulinoma in a MEN1 patient. The definitive treatment of choice in this patient is surgery.

Insulinoma is the second most common functioning enteropancreatic tumor in MEN1 syndrome. Investigation of choice is 72-hour fasting glucose and serum insulin levels. These parameters are measured when the patient develops hypoglycemic symptoms during a supervised 72-hour fasting period.

The treatment of choice is surgery, ranging from enucleation of a single tumor to pancreatectomy.

Non-surgical management of insulinoma includes frequent consumption of carbohydrate-rich meals, octreotide or diazoxide, which are not always successful.

Selpercatinib, a kinase inhibitor was recently approved by the FDA. It is used to treat tumours with a mutation in the RET gene. These are non-small cell lung cancer, medullary thyroid cancer, and other types of thyroid cancers in patients with a mutation in the RET gene.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 6:

The most frequently observed cutaneous manifestation of MEN1 syndrome is angiofibroma.

Cutaneous tumors in MEN1 Syndrome are:

- Angiofibroma – 85%
- Collagenoma – 70%
- Lipoma - 33%

Solution to Question 7:

Secretin study is not done as an annual screening test in patients with MEN1 syndrome. It is a provocative test for the diagnosis of gastrinoma.

Annual biochemical screening tests to be done in patients with MEN1 syndrome are:

Other imaging tests like MRI and CT might have to be done for screening of conditions like anterior pituitary tumors or adrenal tumors.

Tumor	Age to begin screening	Annual screening test
Parathyroid	8	Calcium, PTH
Gastrinoma	20	Gastrin(+/- gastric pH)
Insulinoma	5	Fasting glucose, insulin
Other pancreatic NETs	<10	Chromogranin A; pancreatic polypeptide, glucagon,vasoactive intestinal peptide
Anterior pituitary	5	Prolactin, IGF-I

Sold by @itachibot

Solution to Question 8:

The most common component of MEN2A syndrome, which is also called Sipple's syndrome is medullary thyroid cancer.

Solution to Question 9:

Pituitary adenoma is not a characteristic feature of the symptom complex of MEN2 and its variants, but it is observed in MEN1 and MEN4 syndrome.

Solution to Question 10:

Meningioma is not commonly seen in MEN2B Syndrome. MEN1 syndrome is associated with meningiomas.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

MEN4 is caused by genetic mutations in CDKN1B (cyclin-dependent kinase inhibitor), located on chromosome 12.

MEN1 is caused due to MEN1 gene mutation on chromosome 11.

MEN2 is associated with RET proto-oncogene mutation on chromosome 10.

Solution to Question 12:

Medullary thyroid carcinoma is not associated with MEN4 syndrome, but it is associated with MEN2 syndrome.

Solution to Question 13:

This patient is likely to have MEN4 syndrome as reproductive organ tumors are a characteristic feature of this condition.

Sold by @itachibot

Pheochromocytoma

Question 1:

The clinical triad of pheochromocytoma includes all of the following symptoms except:

a) Headache

b) Hypertension

c) Profuse sweating

d) Palpitation

Question 2:

Which of the following would you not see in a patient with pheochromocytoma?

a) Pallor

b) Postural hypotension

c) High hematocrit

d) Hypocalcemia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

In a patient with pheochromocytoma, which of the following clinical features is least likely to be present?

a) Weight loss

b) Pallor

c) Abdominal pain

d) Oliguria

Question 4:

Which of the following is an associated finding that can be seen in patients with pheochromocytoma?

a) Erythema multiforme

b) Café au lait spots

Sold by @itachibot

c) Vitiligo

d) Ash leaf macules

Question 5:

Which of the following parameters can help you diagnose a patient with a malignant pheochromocytoma?

a) Capsular invasion

b) Number of mitotic spindles

c) Lymph node involvement

d) Cellular atypia

Question 6:

Which of the following catecholamines would be elevated in a patient with extra-adrenal pheochromocytoma?

a) Norepinephrine

b) Epinephrine

c) Metanephrines

d) Vanillylmandelic acid

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

A 31-year-old woman is rushed to the ED with a severe headache and vomiting. Past history includes a total thyroidectomy for thyroid cancer 3 months ago. Her BP is 200/100 mm Hg and pulse is 110/min. On examination, a marfanoid habitus is noted along with the findings shown. Which of the following is likely to be predominantly elevated?

Sold by @itachibot

a) Norepinephrine

b) Epinephrine

c) Dopamine

d) Vanillylmandelic acid

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 8:

In a suspected case of pheochromocytoma, which of the following would you avoid while making a diagnosis?

a) MRI scan

b) MIBG scan

c) Image-guided biopsy

d) Genetic testing

Question 9:

A middle-aged male patient is being evaluated for recurrent episodes of palpitations, headaches, and profuse sweating. He was found to have elevated blood pressure readings on multiple occasions over the past 2 months. Which of the following tests has the highest sensitivity and specificity for the diagnosis of this condition?

a) MRI

b) 24-hour urinary fractionated metanephrines

c) MIBG scintigraphy

Sold by @itachibot

d) Fractionated plasma metanephrines

Question 10:

Which of the following is the best method for detecting and localization of tumors associated with pheochromocytomas?

a) Abdominal CT

b) Fluorodopa-PET

c) MIBG scintigraphy

d) Abdominal MRI

Question 11:

A 45-year-old man presented with recurrent episodes of headache, palpitations, and sweating. His blood pressure at presentation was 180/100 mm Hg. Further testing revealed elevated urinary metanephrines. CT abdomen showed a right adrenal mass. What is the drug of choice for the management of this patient?

a) Labetalol

b) Nitroprusside

c) Phenoxybenzamine

d) Phentolamine

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 12:

Which of the following should not be used for the initial management, in a patient with pheochromocytoma?

a) Prazosin

b) Atenolol

c) Nitroprusside

d) Metyrosine

Question 13:

Which of the following is not a part of Averbuch's chemotherapy protocol?

Sold by @itachibot

a) Dacarbazine

b) Vinblastine

c) Cyclophosphamide

d) Vincristine

Question 14:

A patient is diagnosed with malignant pheochromocytoma. Which of the following is used to treat this condition?

a) Strontium

b) Cobalt-60

c) Phosphorus

d) MIBG

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	b
5	c
6	a
7	b
8	c
9	b
10	b
11	c
12	b
13	b
14	d

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The clinical triad of pheochromocytoma includes the following symptoms:

• Headache

• Profuse sweating

• Palpitation (due to tachycardia)

Hypertension is the most predominant sign of pheochromocytoma. Though intermittent hypertension is a classical feature, sustained hypertension is also common.

Solution to Question 2:

Hypercalcemia is seen in patients with pheochromocytoma, not hypocalcemia. This is due to the production of parathyroid hormone-related peptide (PTHrP) by the tumor.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Polyuria, not oliguria is seen in patients with pheochromocytoma.

Excess catecholamine levels in this condition can cause polyuria and polydipsia due to hyperglycemia.

Solution to Question 4:

Café au lait spots are seen in neurofibromatosis 1, which is a pheochromocytoma-associated syndrome.

Pheochromocytoma-associated syndromes include the following:

• Neurofibromatosis type 1

• Multiple endocrine neoplasia type 2

• Von Hippel–Lindau syndrome

• Paraganglioma syndrome

Solution to Question 5:

Malignant pheochromocytoma refers to tumors that have undergone metastasis to lymph nodes or distant sites.

Sold by @itachibot

They are histologically and biochemically the same as benign ones. Hence, typical histological criteria like cellular atypia, mitotic spindles, and capsular invasion are insufficient to confirm malignancy in pheochromocytomas.

Solution to Question 6:

Norepinephrine is exclusively secreted in patients with extra-adrenal pheochromocytoma.

The PNMT (phenylethanolamine-N-methyl transferase) enzyme converts norepinephrine to epinephrine. This enzyme is absent in extra-adrenal sites. Hence, extra-adrenal pheochromocytomas secrete only norepinephrine and cannot secrete epinephrine.

On the other hand, adrenal pheochromocytomas can secrete norepinephrine, epinephrine, and dopamine in varying amounts depending on the PNMT levels in the tumor.

Extra-adrenal pheochromocytomas or catecholamine-secreting paragangliomas are most commonly found in the abdomen. They commonly arise from the organ of Zuckerkandl.

Solution to Question 7:

This clinical scenario with a history of thyroid cancer (likely medullary carcinoma) and the image showing mucosal neuromas points towards MEN2B syndrome. Her current presentation is suggestive of a pheochromocytoma. She is likely to have predominantly elevated levels of epinephrine and its metabolite, metanephrine.

This is due to the higher expression of the PNMT enzyme associated with the mutations in MEN2 syndromes (both 2A and 2B). The higher levels of epinephrine result in a more symptomatic presentation.

In contrast, low PNMT expression is seen in von-Hippel-Lindau (VHL) syndrome. This leads to a predominant elevation of norepinephrine and normetanephrine levels. It is less symptomatic compared to MEN2 syndrome.

Solution to Question 8:

Biopsy and FNAC should be avoided in patients with pheochromocytoma, as it is a highly vascular tumor.

CT and MRI have similar sensitivity but T2-weighted MRI with gadolinium contrast is the optimal test for diagnosis. MIBG (metaiodobenzylguanidine) scintigraphy can be used to localize the tumor if both CT and MRI are negative.

Genetic testing may be performed as nearly 40% of the tumors are associated with familial syndromes like VHL, neurofibromatosis, and MEN2.

Note: In cases of malignant pheochromocytoma, biopsy may be indicated to confirm adrenal gland metastasis in other sites.

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

Solution to Question 9:

The given clinical scenario is suggestive of pheochromocytoma. The measurement of 24-hour urinary fractionated metanephrines has both the highest specificity and sensitivity in the diagnosis of this condition.

Imaging modalities like CT and MRI have a sensitivity of around 98-100% and specificity of 70%.

	24-hour urinary fractionated metanephrines	Plasma fractionated metanephrines
Sensitivity	98%	96-100%
Specificity	98%	85-89%
False positives	Very rare	Can occur due to drugs like levodopa, TCAs, α-blockers, β-blockers, diuretics
Procedure	Collection of samples over 24 hours may be cumbersome	Requires only a single blood sample. May be preferred in children.
Uses	Preferred first-line test as it is highly specific	Reserved for patients with a high index of suspicion like:Family historyGenetic syndromes like MEN2Prior history of pheochromocytoma

Solution to Question 10:

Among the given options, the fluorodopa-PET scan is the best method for detecting and localization of tumors associated with pheochromocytoma.

If biochemical testing for pheochromocytoma is positive, an abdominal MRI must be performed first. If MRI is negative or if multiple tumors are suspected, the tumor can be localized by performing one of the following investigations:

- 68 Ga-DOTATATE PET scan - most sensitive, gold standard
- Fluorodopa and 18F-fluorodeoxyglucose (FDG) PET scans - 97% sensitive
- MIBG scintigraphy - least sensitive (94%)

Solution to Question 11:

This clinical scenario is suggestive of pheochromocytoma. The drug of choice to control blood pressure prior to surgery is oral phenoxybenzamine at 0.5–4 mg/kg of body weight. It is a non-specific α-blocker and is preferred due to its long action (3-4 days).

Sold by @itachibot

Other α-blockers like oral prazosin and intravenous phentolamine have a shorter duration of action and can be used to manage paroxysms. IV phentolamine is also used for intra-operative management of BP.

α-blockers are usually given for 7 days. Following this, β-blockers are started, usually 2-3 days preoperatively. β-blockers should never be started first because blockade of vasodilatory peripheral β-receptors with unopposed α-receptor stimulation can lead to a further elevation in blood pressure. (Mnemonic: A before B)

Solution to Question 12:

Beta-blockers like atenolol should never be used as the initial drug or alone in the management of pheochromocytoma.

They must be given only after adequate alpha blockade has been established. Blocking beta-receptors may cause unopposed stimulation of alpha-receptors, precipitating a hypertensive crisis.

Option A: Prazosin is an alpha-blocker.

Option C: Nitroprusside can be used in case of hypertensive emergencies.

Option D: Metyrosine inhibits catecholamine synthesis. It can be used when other agents have been ineffective.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 13:

Vinblastine is not included in Averbuch's chemotherapy protocol for the treatment of malignant pheochromocytoma.

Solution to Question 14:

131I-MIBG (metaiodobenzylguanidine) is used for the treatment of malignant pheochromocytoma.

Malignant pheochromocytomas occur in 10% of cases and are histologically and biochemically the same as benign ones. The diagnosis of malignant pheochromocytoma is based on documentation of metastatic disease. The treatment of choice is nuclear medicine therapy with any of the following:

• 131I-MIBG at 200-mCi doses at monthly intervals (3-6 cycles)

• DOTATOC (a somatostatin receptor analog) labeled with radioactive lutetium or yttrium

Chemotherapy can also be given according to Averbuch's protocol.

Note: 123I-MIBG and Gallium-DOTATATE are used for imaging.

Sold by @itachibot

Cushing's Syndrome

Question 1:

What is the most common cause of Cushing's syndrome?

a) Pituitary adenoma

b) Adrenal adenoma

c) Ectopic ACTH secretion

d) Iatrogenic

Question 2:

Which of the following causes ACTH-dependent Cushing's syndrome?

a) Cushing's disease

b) Adrenal adenoma

c) Adrenocortical carcinoma

d) McCune-Albright Syndrome

Question 3:

Which of the following statements is false regarding Cushing's syndrome?

a) It is commonly associated with MEN1 syndrome

b) Bronchial carcinoid is a possible cause

c) It is characterized by hypokalemia

d) It is associated with coronary heart disease

Question 4:

A patient with Cushing's syndrome is least likely to present with:

a) Truncal obesity

b) Osteoporosis

c) Distal myopathy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) Amenorrhea

Question 5:

Which of the following would not be seen in a patient with Cushing's disease?

a) Skin hyperpigmentation

b) hypocoagulable state

c) Frank psychosis

d) Diastolic hypertension

Question 6:

A 28-year-old woman visits a gynecologist with complaints of irregular and infrequent menstrual cycles. She is also upset that she has gained 12 kg over the past few months. Her face appears as shown in the image below. Her blood pressure at this visit is 144/98 mm Hg and she has multiple purple striae over her abdomen. Which of the following investigations should you advise this patient?

a) Cosyntropin stimulation test

b) Midnight serum cortisol measurement

c) Serum testosterone levels and abdominal USG

d) Serum LH:FSH ratio

If you bought this from someone else, it has been scammed.
Sold by @itachibot

Sold by @itachibot

Question 7:

You can advise any of the following tests for screening of a patient suspected to be having Cushing's syndrome except:

a) Midnight plasma cortisol measurement

b) Determination of plasma ACTH levels

c) 24 hour urinary free cortisol measurement

d) Overnight dexamethasone suppression test

Question 8:

In patients with which of the following conditions would you expect to see the highest basal ACTH level?

a) ACTH-secreting pituitary adenoma

b) Ectopic ACTH-secreting tumors.

c) Bilateral adrenal hyperplasia

d) Iatrogenic hypercortisolism

Question 9:

A patient with suspected Cushing's syndrome is found to have elevated ACTH levels. What is the investigation of choice?

a) MRI brain

b) CRH stimulation test

c) High-dose dexamethasone suppression test

d) Inferior petrosal sinus sampling

Question 10:

A 56-year-old man chronic smoker presented with flecks of blood in his sputum. He is currently on treatment for COPD and hypertension. His BMI is 31 kg/m² and he has central obesity. His RBS is 261 mg/dl. During further evaluation, he is found to have high ACTH levels that remained high even after administration of high-dose dexamethasone. What is the underlying cause of this patient's symptoms?

a) Exogenous steroid intake

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

b) Pituitary adenoma

c) Adrenal cortical adenoma

d) Ectopic ACTH-secreting tumor

Question 11:

Evaluation of a patient with Cushing's syndrome reveals high ACTH levels that are responsive to CRH and suppressed by high-dose dexamethasone. What is the likely diagnosis?

a) Steroid intake

b) Small cell lung cancer

c) Pituitary adenoma

d) Adrenocortical adenoma

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 12:

A 50-year-old woman was diagnosed with ACTH-dependent Cushing's syndrome. As MRI of the pituitary was normal and further testing was unequivocal, she was scheduled for inferior petrosal sinus sampling. It revealed a baseline petrosal:peripheral ACTH ratio > 2. What is the treatment of choice in this patient?

a) Bilateral adrenalectomy

b) Transsphenoidal pituitary surgery

c) Removal of ectopic tumor

d) Stereotactic radiotherapy

Question 13:

Which of the following would not be used to treat patients with Cushing's syndrome?

a) Etomidate

b) Metyrapone

c) Ketoconazole

d) Mitomycin

Question 14:

Sold by @itachibot

Which of the following is the best adrenolytic agent that can be used in the management of patients with adrenocortical carcinoma?

a) Mitotane

b) Metyrapone

c) Ketoconazole

d) Trilostane

Question 15:

Which of the following complications is associated with bilateral adrenalectomy?

a) Sheehan's syndrome

b) Addison's disease

c) Waterhouse-Friderichsen syndrome

d) Nelson's syndrome

Question 16:

Which of the following is not a cause of pseudo-Cushing's syndrome?

a) Obesity

b) Alcoholism

c) Depression

d) Malignancy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	c
5	b
6	b

Sold by @itachibot

7	b
8	b
9	a
10	d
11	c
12	b
13	d
14	a
15	d
16	d

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Detailed Explanations

Solution to Question 1:

The most common cause of Cushing's syndrome is iatrogenic due to administration of corticosteroids.

Solution to Question 2:

Cushing's disease causes ACTH-dependent Cushing's syndrome. It occurs due to an ACTH-secreting pituitary adenoma.

Solution to Question 3:

Cushing's syndrome is rarely associated with MEN1 syndrome.

Bronchial carcinoids are a source of ectopic ACTH secretion resulting in ACTH-dependent Cushing's syndrome.

Clinical features of Cushing's syndrome are attributed to the glucocorticoid, mineralocorticoid, and sex steroid actions of cortisol. They include:

System	Clinical features
Body fat	Centralortruncal obesity(most common)Buffalo hump (fat deposition behind the neck)Moon facies

Sold by @itachibot

System	Clinical features
Skin	Purple striae (over abdomen & flanks)Acne and Hirsutism Easy bruisingFacial plethora Hyperpigmentation
Musculoskeletal	Proximal myopathyOsteoporosis
Metabolic	Poorly controlled hyperglycemiaDyslipidemia
Reproductive	OligomenorrheaAmenorrhea
Renal	Hypokalemic alkalosis
CVS	AtherosclerosisDiastolic HypertensionEdema
Hematologic	Hypercoagulable stateLymphopeniaNeutrophiliaEosinopenia
CNS	DepressionPsychosisAnxiety

Sold by @ ... one else, If you purchased this from ... may have been scammed.

Solution to Question 4:

Proximal myopathy is commonly seen in Cushing's syndrome, not distal myopathy.

Clinical features of Cushing's syndrome are attributed to the glucocorticoid, mineralocorticoid, and sex steroid actions of cortisol. They include:

System	Clinical features
Body fat	Centralortruncal obesity(most common)Buffalo hump (fat deposition behind the neck) Moon facies
Skin	Purple striae (over abdomen & flanks)Acne and Hirsutism Easy bruisingFacial plethora Hyperpigmentation
Musculoskeletal	Proximal myopathyOsteoporosis
Metabolic	Poorly controlled hyperglycemiaDyslipidemia
Reproductive	OligomenorrheaAmenorrhea
Renal	Hypokalemic alkalosis
CVS	AtherosclerosisDiastolic HypertensionEdema

Sold by @itachibot

System	Clinical features
Hematologic	Hypercoagulable stateLymphopeniaNeutrophiliaEosinopenia
CNS	DepressionPsychosisAnxiety

Solution to Question 5:

Cushing's disease, and all other causes of Cushing's syndrome, can result in a hypercoagulable state. This is due to an increase in procoagulant factors and inhibition of fibrinolysis.

Cushing's disease is a form of Cushing's syndrome caused by an ACTH-secreting pituitary tumor. The high ACTH levels stimulate melanocytes and cause skin hyperpigmentation. This is especially seen over friction-prone areas like knuckles and elbows.

The majority of patients with Cushing's syndrome have psychiatric symptoms such as:

• Anxiety,

• Depression, and

• Frank psychosis

Mineralocorticoid action of cortisol may lead to diastolic hypertension, hypokalemia, and edema.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

The given clinical scenario along with the image showing moon facies with acne and hirsutism are suggestive of Cushing's syndrome. It can be diagnosed with the help of a single midnight serum cortisol measurement.

Option A: Cosyntropin (ACTH) stimulation test is used for the diagnosis of Addison's disease.

Options C and D: Serum testosterone, abdominal USG, and LH: FSH ratio, are used to confirm a diagnosis of the polycystic ovarian syndrome (PCOS). Though PCOS can present with similar features, it is not typically associated with moon facies, purple striae, or hypertension.

Solution to Question 7:

Determination of plasma ACTH levels is not advised as a screening test for a patient with Cushing's syndrome.

Plasma ACTH determination is done to distinguish between ACTH-dependent and independent causes. It is done after the screening or confirmation of Cushing's syndrome.

Sold by @itachibot

Solution to Question 8:

Basal plasma ACTH levels are highest in ectopic ACTH-secreting tumors.

Plasma ACTH > 15pg/ml is suggestive of ACTH-dependent Cushing's syndrome due to pituitary or ectopic sources. Basal ACTH in pituitary adenoma can be normal or elevated, whereas in an ectopic source the ACTH production is very high. This occurs because a pituitary adenoma retains some sensitivity to the negative feedback mechanism exerted by cortisol, but ectopic tumors remain unaffected.

Plasma ACTH < 5pg/ml is suggestive of ACTH-independent Cushing's syndrome that can be caused by an adrenal adenoma or exogenous administration of glucocorticoids.

Solution to Question 9:

An elevated ACTH level is suggestive of ACTH-dependent Cushing's syndrome. The investigation of choice for ACTH-dependent Cushing's syndrome is MRI brain, to look for the presence of a pituitary tumour.

If the MRI brain shows a mass it is suggestive of a pituitary mass of >6 mm in size. However, if the MRI brain does not show a mass it suggests a pituitary tumor of <6mm in size or an ectopic ACTH source.



Note: High-dose dexamethasone suppression test and CRH stimulation test are employed to differentiate pituitary from ectopic ACTH syndrome. A definitive cause can be established only by imaging techniques like MRI for pituitary tumors, CT scan of the chest and abdomen for ectopic ACTH sites. Hence MRI is considered the investigation of choice.

Solution to Question 10:

Sold by @itachibot

Elevated ACTH levels that do not get suppressed by a high-dose dexamethasone test are suggestive of Cushing's syndrome secondary to an ectopic ACTH-secreting tumor. Considering the patient's smoking history and presentation with hemoptysis, it is likely to be a lung tumor.

Ectopic sources of ACTH include small cell lung cancers and pulmonary carcinoid tumors. The most effective treatment strategy is the management of the underlying SCLC or carcinoid.

Dynamic testing for Cushing's syndrome		
	High dose dexamethasone suppression test	CRH stimulation test
Pituitary adenoma	Positive (suppressed cortisol)	Positive (increase in ACTH and cortisol)
Ectopic ACTH	Negative	Negative
Iatrogenic: Suppressed ACTH levels that are not affected by dynamic testing		
Adrenalmass or hyperplasia: Suppressed ACTH levels that are not affected by dynamic testing		

Sold to
if you purchased this
you may have been s

Solution to Question 11:

CRH responsiveness is seen only in ACTH-dependent hypercortisolism which is due to a pituitary cause. This is because the pituitary tumor contains corticotrophic cells that are responsive to CRH.

Dynamic testing for Cushing's syndrome		
	High dose dexamethasone suppression test	CRH stimulation test
Pituitary adenoma	Positive(suppressed cortisol)	Positive(increase in ACTH and cortisol)
Ectopic ACTH	Negative	Negative
Iatrogenic: Suppressed ACTH levels that are not affected by dynamic testing		
Adrenalmass or hyperplasia: Suppressed ACTH levels that are not affected by dynamic testing		

Sold by @itachibot

Solution to Question 12:

Cushing's syndrome with petrosal: peripheral ACTH ratio > 2 at baseline confirms the presence of an ACTH-secreting pituitary adenoma. The treatment of choice is transsphenoidal pituitary surgery. Radiotherapy is used in case of recurrence of tumors.

Inferior petrosal sinus sampling is performed to differentiate pituitary and ectopic sources of ACTH when initial imaging and testing are inconclusive.



Solution to Question 13:

Mitomycin is not used in the treatment of Cushing's syndrome. The drugs used in its management include:

- Metyrapone
- Ketoconazole
- Aminoglutethimide
- Trilostane
- Cyproheptadine
- Mitotane – Mainly in adrenal carcinoma
- Etomidate – Severe cortisol excess

Solution to Question 14:

Mitotane is used in the management of adrenocortical carcinoma due to its adrenolytic properties.

Mitotane is an insecticide derivative. It is effective in the treatment of benign Cushing's also, but is rarely used due to its side effects.

If you purchased this, you may have been else,

Sold by @itachibot

Solution to Question 15:

Nelson's syndrome is a rare complication of total bilateral adrenalectomy (TBA). TBA is a procedure used to control hypercortisolism in Cushing's disease (due to ACTH-secreting pituitary adenoma) when resection of the primary tumor is not possible.

The loss of feedback inhibition of the hypothalamic-pituitary-adrenal axis following TBA increases CRH. This leads to stimulation of ACTH-secreting corticotroph cells, causing enlargement of the original pituitary tumor or the formation of a new one.

Nelson's syndrome is associated with a clinical triad of:

• Hyperpigmentation

• Excessive ACTH secretion

• Corticotroph adenoma

Screening is done with fasting plasma ACTH levels and MRI brain is performed at regular intervals. Prophylactic pituitary irradiation may be recommended to prevent its incidence. The primary treatment is transsphenoidal surgery. Patients will also require life-long glucocorticoid replacement, as they do not produce cortisol.

The use of steroidogenic inhibitors in the treatment of Cushing's disease has decreased the need for bilateral adrenalectomy these days.

Option A: Sheehan's syndrome refers to postpartum hypopituitarism that is caused by ischemic necrosis of the anterior pituitary as a result of severe postpartum hemorrhage.

Option B: Addison's disease occurs due to autoimmune destruction of adrenal glands.

Option C: Waterhouse-Friderichsen syndrome refers to adrenal apoplexy that is caused by fulminant meningococcemia.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

Malignancy does not cause pseudo-Cushing's syndrome. It may cause ectopic ACTH production, leading to true Cushing's syndrome.

Pseudo-Cushing's syndrome refers to functional hypercortisolism caused by chronic activation of the HPA axis. It presents with clinical and biochemical features similar to Cushing's syndrome, making it difficult to differentiate between them. It may occur secondary to the following conditions:

• Severe major depressive disorder

• Polycystic ovarian syndrome

• Insulin-resistant obesity

• Obstructive sleep apnea

• Chronic alcoholism

Sold by @itachibot

• End-stage renal disease

It usually resolves when the underlying condition is treated.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Adrenal Insufficiency and Hyperaldosteronism

Question 1:

Addison's disease is commonly associated with ____

a) Autoimmune adrenalitis

b) Adrenocortical carcinoma

c) Hypernephroma

d) Medullary carcinoma thyroid

Question 2:

What is the most common cause of adrenal insufficiency in India?

a) Tuberculosis

b) Exogenous steroid use

c) Post-partum pituitary insufficiency

d) Autoimmune adrenal insufficiency

Question 3:

In a patient with reduced cortisol and elevated ACTH, which of the following would not be part of your differential diagnoses?

a) Kearns-Sayre syndrome

b) Autoimmune polyglandular syndrome type 2

c) Triple-A syndrome

d) MEN 4 syndrome

Question 4:

A 50-year-old woman was diagnosed with Hashimoto's thyroiditis and was started on levothyroxine 2 weeks ago. She now presents to your clinic with abdominal pain and dizziness. The left and right images below show her current appearance and her appearance last month, respectively. Which of the following is not a feature of this condition?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Hyperglycemia

b) Hyponatremia

c) Hyperkalemia

d) Metabolic acidosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

What is the most common biochemical abnormality seen at presentation in patients with adrenal insufficiency?

a) Hypernatremia

b) Hypokalemia

c) Hyponatremia

d) Hyperkalemia

Question 6:

A 27-year-old woman is brought to the casualty with severe abdominal pain, vomiting, and diarrhea for the past 3 days. She has been excessively tired and dizzy for the past 2 weeks. On examination, there are patches of hyperpigmented skin. Blood pressure is 70/40 mm Hg and blood glucose is 50 mg/dl. How will you confirm the diagnosis?

a) Serum Na+, K+, renin

b) Serum Na+, K+, saline suppression test

c) Determination of plasma ACTH levels

Sold by @itachibot

d) ACTH stimulation test

Question 7:

A 20-year-old woman is being evaluated for adrenal insufficiency. You decide to check her serum levels of ACTH, renin, and aldosterone. Which of the following laboratory findings would prompt you to subsequently order a pituitary MRI?

a) High renin and low aldosterone levels

b) High ACTH levels

c) Low renin and high aldosterone levels

d) Normal renin and aldosterone levels

Question 8:

A 33-year-old man collapses during a family vacation. At the hospital, blood pressure is found to be 86/52 mm Hg, and random blood sugar is 49 mg/dl. His wife says that he has been on treatment for Addison's disease for two years but had forgotten to pack his medication for the trip. What is the drug of choice for this patient?

a) Norepinephrine

b) Hydrocortisone

c) Dexamethasone

d) Fludrocortisone

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A patient who presented with an adrenal crisis is undergoing treatment with dexamethasone. Which of the following abnormalities will not improve with this treatment?

a) Hyponatremia

b) Postural hypotension

c) Hypoglycemia

d) Hyperkalemia

Question 10:

Sold by @itachibot

A 62-year-old woman who was diagnosed with a non-functioning pituitary adenoma develops features of adrenal insufficiency. Which of the following is not a feature of this condition?

a) Hyperpigmentation

b) Postural hypotension

c) Hypoglycemia

d) Lassitude

Question 11:

Which of the following is incorrect with respect to Addisonian pigmentation?

a) Presence of moles and scars

b) Represented by palmar creases

c) Does not affect the oral mucosa

d) Decreased fibrosis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 12:

What is the most common cause of mineralocorticoid excess?

a) Bilateral micronodular adrenal hyperplasia

b) Unilateral adrenal adenoma

c) Congenital adrenal hyperplasia

d) Liddle's syndrome

Question 13:

Which of the following clinical features will you not expect to see in a patient with Conn's syndrome?

a) Hypertension

b) Hypokalemia

c) Edema

d) Muscle weakness

Question 14:

Sold by @itachibot

A 39-year-old man is referred for evaluation of severe hypertension that has not improved despite being on 3 anti-hypertensive drugs. The presence of decreased plasma renin activity in this patient would point to which of the following diagnoses?

a) Conn syndrome

b) Bartter syndrome

c) Chronic kidney disease

d) Bilateral renal artery hypertension

Question 15:

Which of the following is the best initial investigation in a patient with hypokalemic hypertension?

a) ACTH stimulation test

b) CT abdomen

c) Adrenal vein sampling

d) Aldosterone-renin ratio

Question 16:

Which of the following tests is not performed in patients with an elevated aldosterone-renin ratio?

a) Saline infusion test

b) Metyrapone stimulation test

c) Oral sodium loading test

d) Fludrocortisone suppression test

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	a

Sold by @itachibot

5	c
6	d
7	d
8	b
9	d
10	a
11	c
12	a
13	c
14	a
15	d
16	b

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Primary adrenal insufficiency, also known as Addison's disease, is most commonly caused by autoimmune adrenalitis.

Tuberculous adrenalitis is still a frequent cause of primary adrenal insufficiency in developing countries like India. It is characterized by the loss of glucocorticoid, mineralocorticoid, and adrenal androgen secretion.

Glucocorticoid deficiency results in fatigue, lack of energy, weight loss, fever, myalgia, normochromic anemia, and hypoglycemia. Mineralocorticoid deficiency is characterized by nausea, vomiting, low blood pressure, postural hypotension, salt craving, hyponatremia, and hyperkalemia. Adrenal androgen deficiency manifests as a lack of energy, dry itchy skin, loss of libido, and loss of axillary and pubic hair (in women).

Hyperpigmentation is seen in primary adrenal insufficiency due to excess proopiomelanocortin-derived peptides. Hyperpigmentation is not seen in secondary adrenal insufficiency.

The diagnosis of adrenal insufficiency can be made with the short cosyntropin test (cortisol level <450-500nmol/L sampled 30 to 60 minutes after ACTH stimulation is diagnostic of adrenal insufficiency). Plasma ACTH, plasma renin, and serum aldosterone can help differentiate primary from secondary causes. Primary adrenal insufficiency will show elevated ACTH and a compensatory elevation in renin levels due to decreased aldosterone in the body. In secondary adrenal insufficiency, ACTH is low with normal renin and aldosterone.

Treatment includes glucocorticoid, mineralocorticoid, and adrenal androgen replacement.

Other options:

Sold by @itachibot

Option B: Patients with adrenocortical carcinoma present with features of glucocorticoid and adrenal androgen excess, and not adrenal insufficiency.

Options C and D: Hypernephroma and medullary carcinoma thyroid are not associated with Addison's disease.

Solution to Question 2:

Tuberculosis is the most common cause of adrenal insufficiency in India and other developing countries.

Idiopathic atrophy due to autoimmune conditions is the most common cause in developed countries.

Solution to Question 3:

Elevated ACTH and reduced cortisol levels are seen in primary adrenal insufficiency. It is not a feature of MEN 4 syndrome.

The following syndromes are associated with primary adrenal insufficiency:

- Autoimmune polyglandular syndrome type 1
- Autoimmune polyglandular syndrome type 2
- Triple-A syndrome
- Smith-Lemli-Opitz syndrome
- Kearns-Sayre syndrome
- IMAGe syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

The given image showing skin hyperpigmentation in a patient with severe dizziness and a history of autoimmune thyroiditis points to a diagnosis of Addison's disease. Hyperglycemia is not a feature of this condition. Hypocortisolism leads to hypoglycemia in these patients.

Addison's disease is primary adrenal insufficiency characterized by the disruption of glucocorticoid, mineralocorticoid, and androgen secretion. Primary adrenal insufficiency is most commonly caused by autoimmune adrenalitis and coincident autoimmune disease occurring in these patients most frequently includes thyroid autoimmune disease, vitiligo, and premature ovarian failure.

Clinical features of Addison's disease:

Sold by @itachibot

Hormone	Clinical features
Hypoaldosteronism	Abdominal pain, nausea, vo miting, salt craving
Dizziness, postural hypotensi on	
AKI due to volume depletion	
Hyponatremia	
Hyperkalemia, metabolic aci dosis	
Hypocortisolism	Anorexia, weight loss, fatigue, myalgia, arthralgia
Fever (increased IL-1)	
Depression	
Postural hypotension	
Hypoglycemia	
Hyponatremia (loss of feedba ck inhibition of ADH)	
Slightly increased TSH (loss of feedback inhibition)	
Hypoandrogenism	Loss of libido
	Loss of pubic and axillary hai r
Elevated ACTH	Hyperpigmentation

Levothyroxine can precipitate adrenal insufficiency or crisis in undiagnosed patients. In patients with concurrent adrenal insufficiency and hypothyroidism, a course of steroids must be given before giving thyroid replacement.

Solution to Question 5:

Hyponatremia is the most common biochemical abnormality seen at presentation in patients with adrenal insufficiency. It is observed in 80% of the patients at presentation.

Hyponatremia is caused by:

• Mineralocorticoid deficiency

• Diminished inhibition of ADH release by cortisol, resulting in mild SIADH during secondary adrenal insufficiency

Hyperkalemia is diagnosed in 40% of the patients during initial diagnosis.

Solution to Question 6:

Sold by @itachibot

This clinical scenario is suggestive of adrenal insufficiency. The confirmatory test for diagnosis is the ACTH stimulation test or short cosyntropin stimulation test.

Hyponatremia and hyperkalemia are seen. Plasma levels of ACTH, renin and aldosterone are used to differentiate between primary and secondary adrenal insufficiency.



Solution to Question 7:

Pituitary MRI is indicated in secondary adrenal insufficiency which presents with the following parameters:

• Low-normal ACTH level

• Normal renin level

• Normal aldosterone level

The following findings are seen in primary adrenal insufficiency:

• High ACTH levels

• High renin

• Low aldosterone levels

Pituitary MRIs are not required in primary adrenal insufficiency. The fact that a pituitary MRI is done itself gives a hint towards secondary adrenal insufficiency.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

This clinical scenario is suggestive of an adrenal crisis triggered by missed medication. In this patient who is a known case of adrenal insufficiency, IV hydrocortisone is the drug of choice.

However, in patients with newly diagnosed adrenal insufficiency, dexamethasone is used for the treatment of adrenal crisis. This is because dexamethasone is not measured in cortisol assays and would not interfere with the diagnostic tests, whereas hydrocortisone interferes with cortisol assays.

Acute adrenal insufficiency requires the immediate initiation of the following therapeutic strategies:

• Rehydration - saline infusion at an initial rate of 1 L/h with continuous cardiac monitoring.

• Glucocorticoid replacement - bolus injection of 100 mg hydrocortisone, followed by the maintenance dose of 100–200 mg over 24h.

• Mineralocorticoid replacement - initiated once the daily hydrocortisone dose has been reduced to <50 mg as higher doses of hydrocortisone have sufficient mineralocorticoid action.

Solution to Question 9:

Hyperkalemia will not improve following administration of dexamethasone. Dexamethasone is a long-acting steroid with glucocorticoid action. It has no effect on hyperkalemia, which is mediated by mineralocorticoids.

Hyponatremia is chiefly due to mineralocorticoid deficiency, however, hypocortisolism causes hyponatremia due to loss of feedback inhibition of ADH resulting in SIADH (syndrome of inappropriate antidiuretic hormone secretion).

Sold by @itachibot

Hypocortisolism can also cause postural hypotension as cortisol regulates vascular tone by potentiating the effects of vasoconstrictors. Thus on administration dexamethasone hyponatremia and postural hypotension will improve as dexamethasone has glucocorticoid action but hyperkalemia will not improve.

Dexamethasone is used in patients with newly diagnosed adrenal insufficiency as it does not interfere with serum cortisol measurement.

Hormone	Clinical features
Hypoaldosteronism	Abdominal pain, nausea, vomiting, salt craving
	Dizziness, postural hypotension
	AKI due to volume depletion
	Hyponatremia
	Hyperkalemia, metabolic acidosis
Hypocortisolism	Anorexia, weight loss, fatigue, myalgia, arthralgia
	Fever (increased IL-1)
	Depression
	Postural hypotension
	Hypoglycemia
	Hyponatremia (loss of feedback inhibition of ADH)
	Slightly increased TSH (loss of feedback inhibition)
Hypoandrogenism	Loss of libido
	Loss of pubic and axillary hair
Elevated ACTH	Hyperpigmentation

If you p one else, med.

Solution to Question 10:

A non-functioning pituitary adenoma causes hypopituitarism due to compression of the pituitary stalk. This results in secondary adrenal insufficiency. Hyperpigmentation is not a feature.

This condition is characterized by an alabaster-like paleness due to the lack of ACTH secretion. In primary adrenal insufficiency, hyperpigmentation occurs as a result of elevated ACTH.

Sold by @itachibot

Solution to Question 11:

Addison's disease causes patchy hyperpigmentation of the oral mucosa.

Hyperpigmentation is also present in areas of increased friction including:

• Palmar creases

• Dorsal aspect of the foot

• Nipples and the axillary region

Solution to Question 12:

Bilateral adrenal hyperplasia is the most common cause of mineralocorticoid excess. It causes primary hyperaldosteronism. Other causes of primary hyperaldosteronism are:

• Conn's syndrome – Unilateral adrenal adenoma

• Glucocorticoid-remediable hyperaldosteronism (< 1%)

Other causes of mineralocorticoid excess are as follows:

• Congenital adrenal hyperplasia (CAH)

• Syndrome of apparent mineralocorticoid excess (SAME)

• Adrenocortical carcinoma

• Progesterone-induced hypertension

• Cushing's syndrome

Liddle's syndrome is due to a mutation of ENaC subunits. This results in reduced degradation of ENaC channels, causing prolonged mineralocorticoid action.

Sold by @itachibot

Solution to Question 13:

In Conn's syndrome, even though there is volume expansion, an underlying salt-escape mechanism leads to the absence of pedal edema.

The volume expansion due to hypernatremia leads to right heart dilatation that stimulates the release of atrial natriuretic peptide (ANP). This leads to compensatory diuresis and lack of edema despite sodium retention.

Features of primary hyperaldosteronism:

- Hypokalemia
- Hypertension
- Metabolic alkalosis
- Polyuria (due to ANP)
- Polydipsia (consequence of polyuria)
- Muscle weakness

If you purchased this from someone else, you may have been scammed. Sold by @Itachibot

Solution to Question 14:

Conn syndrome is a cause of primary hyperaldosteronism that presents with hypertension and low levels of renin.

Elevated aldosterone levels cause negative feedback in the renin-angiotensin-aldosterone system (RAAS) and results in decreased renin production.

Solution to Question 15:

Hypokalemic hypertension is suggestive of hyperaldosteronism. The screening test of choice is a measurement of plasma renin and aldosterone levels to calculate the aldosterone-renin ratio (ARR).

ARR must be estimated in patients with hypertension and any one of the following features:

- Drug resistance
- Hypokalemia
- Adrenal mass
- Onset before 40 years of age

Prior to the determination of ARR, hypokalemia correction must be done. If the patient is on spironolactone, it must be stopped 4 weeks prior to testing. Other blood pressure medication may be continued.

ARR screen is considered positive when:

- ARR > 750 pmol/L : ng/mL per hour

Sold by @itachibot

• Aldosterone > 450 pmol/L

Option A: ACTH stimulation test is used for the diagnosis of Addison's disease. It presents with hypotension and hyperkalemia.

Options B and C: If ARR is positive, CT abdomen and adrenal vein sampling are used for confirmation of the diagnosis.

Solution to Question 16:

An elevated aldosterone-renin ratio (ARR) is indicative of primary hyperaldosteronism. Metyrapone stimulation test is used in the diagnosis of adrenal insufficiency, not hyperaldosteronism.

A failure of suppression of aldosterone levels below 140 pmol/L indicates autonomous aldosterone production, which is the hallmark of primary hyperaldosteronism. This principle is used in the following tests:

• Saline infusion test - IV administration of 2L of physiological saline over 4 hours

• Oral sodium loading test - oral sodium given over 3 days

• Fludrocortisone suppression test - given IV over 4 days

These tests may be used to confirm the diagnosis in patients with elevated ARR, but are not necessary in cases of overt hypokalemic hypertension, strongly positive ARR, and high aldosterone.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Diabetes Mellitus: Types, Clinical Features and Management

Question 1:

Which of the following is not a clinical feature of diabetes mellitus?

a) Weight loss

b) Osmotic diuresis

c) Weight gain

d) Hypovolemia

Question 2:

A 4-year-old boy is brought to the OPD by his worried mother as he urinates more than ten times a day and is always drinking water. She also reports that despite eating more than his brother did at that age, he is not gaining any weight. Random blood glucose is 252 mg/dl. Which of the following statements is true about this condition?

a) Antibodies against zinc transporters may be present

b) It is non-insulin dependent

c) It is associated with a strong family history

d) Insulin-resistance is the hallmark for pathogenesis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

Which chromosome contains the major susceptible gene for type 1 diabetes mellitus?

a) 1

b) 4

c) 6

d) 12

Question 4:

Sold by @itachibot

Which of the following mechanisms is predominantly responsible for an increase in the fasting plasma glucose level of patients with type 2 diabetes mellitus?

a) Increased hepatic gluconeogenesis

b) Decreased peripheral uptake of glucose

c) Decreased glycogenesis in muscles

d) Decreased hepatic glycogenesis

Question 5:

A 22-year-old woman presents with fatigue and weight loss for the past 6 months. She tells you that her father is also a diabetic. Further evaluation reveals the following. What is the most likely diagnosis?

a) Type 2 diabetes mellitus

b) Maturity-onset diabetes of the young

c) Type I diabetes mellitus

d) Latent autoimmune diabetes in adults

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

All of the following are true about maturity-onset diabetes of the young (MODY) except:

a) It is an autosomal-dominant trait

b) MODY 4 is the most common form

c) Diabetic ketoacidosis is uncommon

d) Insulinopenia is present

Question 7:

Which of the following would cause type 1 diabetes in an adult patient?

a) Latent diabetes

b) Maturity-onset diabetes of the young

c) Brittle diabetes

d) Latent autoimmune diabetes in adults

Sold by @itachibot

Question 8:

Patients with all of the following conditions can develop diabetes mellitus except:

a) Hemochromatosis

b) Prader-Willi syndrome

c) Fragile X syndrome

d) Friedrich's ataxia

Question 9:

Which of the following patients can be diagnosed with diabetes mellitus?

a) A patient with RBS of 210 mg/dl

b) A patient with FBS of 125 mg/dl and 2-hour PPBS of 199 mg/dl

c) A patient with symptoms of diabetes and RBS of 192 mg/dl

d) A patient with HbA1c ≥ 6.5 %

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A 46-year-old man presents to his physician for a routine health evaluation. He is concerned that he might have diabetes as his three brothers have all been diagnosed with diabetes. He does not have any complaints. Which of the following tests can be employed in this patient?

a) 1, 2, 4

b) 1, 4

c) 2, 3, 5

d) 4, 5

Question 11:

Which of the following can be termed impaired glucose tolerance?

a) Fasting blood glucose levels between 100 and 125 mg/dL

b) Random blood glucose levels between 140 and 199 mg/dL

c) Blood glucose levels between 140 and 199 mg/dL following oral glucose tolerance test

d) Fasting blood glucose levels between 100 and 199 mg/dL

Sold by @itachibot

Question 12:

You are counselling an asymptomatic middle-aged woman regarding screening for diabetes mellitus. Based on the ADA recommendations, what will you tell her?

a) All individuals > 35 years must undergo screening every 3 years

b) All individuals > 50 years must undergo screening every 5 years

c) Oral glucose tolerance test would be performed

d) Random blood glucose levels would be checked

Question 13:

A middle-aged woman has been diagnosed with type 2 diabetes mellitus. Her HbA1c level is 11.1%. What is the initial drug of choice?

a) Insulin

b) Glimepiride

c) Metformin

d) Sitagliptin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

A 45-year-old woman was brought to the emergency department by her relatives after she suffered a brief, seizure-like episode. They tell you that she is taking an oral medication for diabetes. On examination, the patient was disoriented, and CBG was 45 mg/dl. Which of the following drugs most likely played a role in the patient's current presentation?

a) Metformin

b) Pioglitazone

c) Sitagliptin

d) Repaglinide

Question 15:

All of the following drugs can be used in the treatment of patients with type 1 diabetes except:

a) Acarbose

b) Pramlintide

Sold by @itachibot

c) Metformin

d) Insulin

Question 16:

A 40-year-old male patient with type 2 diabetes mellitus is now diagnosed with hypertension. Which of the following antihypertensive drugs will you not prefer in this patient?

a) Hydrochlorthiazide

b) Losartan

c) Lisinopril

d) Furosemide

Question 17:

A type 2 diabetic patient who was recently started on treatment complains of frequent urination, fever, and burning micturition. Which of the following drugs is most likely responsible for these symptoms?

a) Dapagliflozin

b) Metformin

c) Glipizide

d) Exenatide

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A 65-year-old man is hospitalized with pneumonia. He is a known patient of type 2 diabetes. During the hospital stay, his appetite is poor and his oral intake is irregular. The physician decides to administer rapid-acting insulin based on high blood sugar values. Which of the following would he use?

a) Lente

b) Detemir

c) Glargine

d) Glulisine

Question 19:

Sold by @itachibot

Which of the following is incorrect about Afrezza?

a) It is available as a powder for inhalation

b) It can cause bronchospasm

c) It does not cause hypoglycemia

d) It is a rapid-acting form

Question 20:

A 60-year-old man presents to the casualty with confusion, vomiting, and abdominal pain. His blood sugar levels are 421 mg/dl and urine ketones are 3+. He is started on IV normal saline and insulin infusion. Which of the following should he be monitored for?

a) Hypernatremia

b) Hyperkalemia

c) Hypokalemia

d) Hyponatremia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

Dawn phenomenon is clinically suspected when a patient shows which of the following findings?

a) Early morning hyperglycemia with hypoglycemia at 3 am

b) Early morning hypoglycemia with normal blood glucose levels at 3 am

c) Early morning hyperglycemia with normal blood glucose levels at 3 am

d) Early morning hypoglycemia with hyperglycemia at 3 am

Question 22:

A patient is said to have developed insulin resistance when the required insulin dosage is which of the following?

a) > 200 units per day

b) > 100 units per day

c) > 150 units per day

d) > 250 units per day

Sold by @itachibot

Question 23:

A 12-year-old child who is known to have type 1 diabetes mellitus presents with confusion and drowsiness. Her mother says that she seems to be breathing very fast. On examination, mucous membranes are dry and blood pressure is 70/50 mm Hg. Random blood glucose is 415 mg/dl and urine ketones are 4+. What is the next best step in management?

a) 2-3 L of normal saline over 1-3 hours

b) Insulin infusion at 0.1 units/kg/hour

c) Arterial blood gas

d) Insulin bolus of 0.1 units/kg given IV

Question 24:

Which of the following is responsible for the elevated levels of TG and VLDL in diabetes mellitus?

a) Increased activity of lipoprotein lipase and decreased activity of hormone sensitive lipase

b) Increased activity of hormone sensitive lipase and decreased lipoprotein lipase activity

c) Increase in peripheral function of LDL receptors

d) Increased in activity of hepatic lipase

Question 25:

In a PHC, which of the following initial screening tests are used to diagnose diabetes?

a) 1 and 2

b) 2 and 3

c) 2 and 4

d) 1, 2 and 3

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	a

Sold by @itachibot

3	c
4	a
5	b
6	b
7	d
8	c
9	d
10	b
11	c
12	a
13	a
14	d
15	c
16	a
17	a
18	d
19	c
20	c
21	c
22	a
23	c
24	b
25	c

If yc you else, d.

Detailed Explanations

Solution to Question 1:

Weight loss is a clinical feature of diabetes mellitus, not weight gain.

Diabetes mellitus refers to a group of common metabolic disorders that result in hyperglycemia. It causes the following classical triad of symptoms:

• Polyuria - due to osmotic diuresis

• Polydipsia - triggered by the resulting hypovolemia

• Weight loss - due to catabolic state

Note: Obesity is a risk factor, not a symptom of type 2 diabetes mellitus and plays a role in its pathogenesis. Increase in the adipocyte mass leads to an increase in the level of circulating free fatty acids. Free fatty acids impair glucose utilization in skeletal muscles, promote glucose

Sold by @itachibot

production by the liver, and impair beta-cell function.

Solution to Question 2:

This clinical scenario is suggestive of type 1 diabetes mellitus. It is an autoimmune disease associated with the following immunologic markers:

- Antibodies against zinc transporter (ZnT-8)
- Islet cells antibodies (ICA)
- Anti-glutamic acid decarboxylase antibodies (GAD)
- Insulin autoantibodies (IAA)

Solution to Question 3:

The major susceptibility gene for type 1 diabetes mellitus is located in the HLA region of chromosome 6.

Type I diabetes mellitus occurs in genetically susceptible individuals and is usually triggered by infections. The HLA complex is most commonly involved. The common HLA associations include:

- Causative - HLA-DQA1*0501 and DQB*0201
- Protective - HLA-DQA1*0102, DQB1*0602, DRB1*1501

Other genomic associations of type I diabetes mellitus include the CTLA-4 gene and IL-2 receptor.

Type 2 diabetes mellitus is associated with mutations in the calpain-10 gene on chromosome 2. It results in insulin resistance.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

In type 2 diabetes mellitus increased fasting blood sugar is predominantly due to increased hepatic gluconeogenesis.

In the postprandial period, there is a rise in peripheral blood glucose levels. Due to insulin resistance in type 2 diabetes mellitus, there is reduced peripheral uptake of glucose by insulin-sensitive tissues like the liver and skeletal muscles. This results in reduced glucose utilization by skeletal muscles as well as reduced glycogenesis in the liver and skeletal muscles. This results in increased postprandial plasma glucose levels.

Increased hepatic glucose output predominantly accounts for increased FPG levels, whereas decreased peripheral glucose utilization results in postprandial hyperglycemia.

Solution to Question 5:

Sold by @itachibot

This clinical scenario is suggestive of maturity-onset diabetes of the young (MODY).

MODY is a monogenic form of diabetes mellitus that is transmitted in an autosomal dominant fashion. It affects young adults < 25 years of age. It occurs due to reduced insulin secretion. Etiological factors of type 1 and 2 diabetes such as insulin resistance, obesity, and antibodies are not associated with this condition.

Stable hyperglycemia is usually seen, hence complications like diabetic ketoacidosis are uncommon. Macrovascular complications like hyperlipidemia or hypertension are not seen. It is treated with sulfonylureas.

Latent autoimmune diabetes in adults (LADA) is a form of type 1 diabetes that occurs between 30-50 years of age.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

MODY 3 is the most common form of maturity-onset diabetes of the young (MODY), accounting for two-thirds of all the cases.

MODY is a monogenic form of diabetes mellitus that is transmitted in an autosomal dominant fashion. It affects young adults <25 years of age. There are a total of 11 types of MODY based on mutations involves. Some of the affected genes include:

• Types 1,3,5: HNF (Hepatocyte nuclear transcription factor) mutation

• Type 2: Glucokinase mutation

• Type 4: IPF 1 (Insulin Promotor Factor-1) mutation

These mutations result in impaired glucose-induced insulin secretion resulting in insulinopenia.

Solution to Question 7:

Type 1 diabetes mellitus in an adult is suggestive of latent autoimmune diabetes in adults (LADA).

LADA is common in thin-built adults between 30-50 years and is associated with autoantibodies. As the progression of the disease is less rapid than in children, clinical features may mimic type 2 diabetes. However, patients usually require insulin treatment within 5 years.

Option A: Patients with latent diabetes become diabetic under stressful conditions. However, upon removal of stress, they become non-diabetic.

Option B: MODY or maturity-onset diabetes of the young is a monogenic form of diabetes that occurs < 25 years. There is impaired insulin secretion, but no antibodies.

Option C: Brittle diabetes is seen in children with type 1 DM. It presents with wide fluctuation in blood sugar levels.

Sold by @itachibot

Solution to Question 8:

Fragile X syndrome is not associated with diabetes mellitus.

Genetic syndromes associated with diabetes include:

- Prader-Willi syndrome
- Friedreich's ataxia
- Wolfram's syndrome
- Down's syndrome
- Klinefelter's and Turner's syndrome
- Huntington's chorea
- Laurence-Moon-Biedl syndrome
- Myotonic dystrophy
- Porphyria

Diseases of the exocrine pancreas causing diabetes:

- Hemochromatosis
- Pancreatitis
- Neoplasia
- Cystic fibrosis

Endocrinopathies causing diabetes:

- Acromegaly
- Cushing's syndrome
- Glucagonoma
- Pheochromocytoma
- Hyperthyroidism

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

The patient with HbA1c ≥ 6.5 % can be diagnosed with diabetes mellitus as this is an independent criterion for diagnosis.

Solution to Question 10:

HbA1c and fasting blood glucose are used for screening diabetes mellitus in asymptomatic individuals.

Sold by @itachibot

HbA1c refers to glycated hemoglobin and it serves as a long-term indicator for blood glucose levels (8 to 12 weeks).

• Normal < 5.7 %

• Abnormal glucose homeostasis: 5.7 to 6.4 %

• Diabetes mellitus: ≥ 6.5%

HbA1c also serves as a prognostic indicator for glycemic control.

Solution to Question 11:

Impaired glucose tolerance is defined as blood glucose levels between 140-199 mg/dL following an oral glucose tolerance test (OGTT).

Impaired fasting glucose is when the fasting blood glucose is between 100 to 125 mg/dL.

Solution to Question 12:

The ADA recommends screening for type 2 diabetes mellitus in all asymptomatic individuals > 35 years every 3 years.

The tests used for screening are:

• HbA1c levels

• Fasting blood glucose levels

Individuals can be screened earlier if they are overweight with BMI >25 kg/m2 and have one of the following risk factors:

• Family history of diabetes

• History of cardiovascular diseases

• History of gestational diabetes mellitus or delivery of a baby > 4 kg

• Obesity (BMI ≥25 kg/m2)

• Physical inactivity

• Hypertension (≥140/90 mm Hg)

• HDL cholesterol < 35 mg/dL

• Triglycerides >250 mg/dL

• Polycystic ovarian syndrome or acanthosis nigricans

Note- If an individual has a test result near the margins of the diagnostic threshold, a close follow-up should be done and repeat the test in 3–6 months.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Solution to Question 13:

Insulin is the best initial drug of choice when HbA1c is > 9% at diagnosis.

Insulin is used in the treatment of type 2 diabetes in the following cases:

• Inadequate glycemic control with 2 or more oral agents

• HbA1c > 9% at diagnosis

In patients with HbA1c < 9%, metformin is the drug of choice.



Solution to Question 14:

Repaglinide is an insulin secretagogue and hence poses the risk of causing hypoglycemia.

Drugs that can cause hypoglycemia (insulin secretagogues):

• Sulfonylureas

• Meglitinides: Nateglinide, Repaglinide

The following drugs are not directly responsible for stimulating insulin secretion and hence do not cause hypoglycemia:

• Metformin

• Alpha-glucosidase inhibitor

• Pioglitazone

• DPP-4 inhibitor: Sitagliptin

Management of diabetes mell itus		
Intervention	Advantages	Disadvantages

Sold by @itachibot

Management of diabetes mellitus		
Lifestyle changes	Reduced complications and mortality	Insufficient for most patients within 1st year
Metformin	Weight neutral	GI side effectsContraindicated when GFR < 45 ml/min
Insulin	Rapidly effectiveLowers HbA1c more than other drugs (by 1.5-3%)	Weight gainHypoglycemia
GLP-1 agonists	Weight lossReduces cardiovascular complications	GI side effects
Pramlintide	Weight lossLowers postprandial glucose	GI side effectsHypoglycemia when used with insulin
Sulfonylureas	Rapidly effective	Weight gainHypoglycemia
Thiazolidinediones	Lowers insulin requirements	Weight gain, edema, CHF, fractures
Meglitinides	Rapidly effective	Weight gainHypoglycemia
SGLT-2 inhibitors	Weight lossBetter cardiovascular and renal outcomes	UTI, vulval candidiasisDehydrationEuglycemic DKA
DPP-4 inhibitors	Weight neutral	Angioedema, urticaria
Alpha-glucosidase inhibitors	Weight neutral	GI flatulenceRise in LFT

If you purchased this, you may have been

Solution to Question 15:

Metformin can only be used in the treatment of type 2 diabetes mellitus.

Drugs that can be used for treating both type 1 and type 2 diabetes mellitus include the following:

• Acarbose - reduces the breakdown of starch into sugar in the GI tract and lowers the postprandial glucose rise in both types.

• Pramlintide - an injectable drug that enhances amylin secretion leading to reduced glucagon levels

• Insulin

Solution to Question 16:

Hydrochlorthiazide belongs to the thiazide group of diuretics, that can cause dyslipidemia and hyperglycemia. Hence they are not preferred for use in diabetic hypertensives.

The antihypertensive drug of choice in diabetes mellitus include angiotensin-converting enzyme inhibitors (ACE-I) or angiotensin II receptor blockers (ARBs).

Sold by @itachibot

Solution to Question 17:

This clinical scenario is suggestive of a urinary tract infection (UTI). This adverse effect occurs as a result of dapagliflozin use.

The group of sodium-glucose co-transporter-2 or SGLT-2 inhibitors includes drugs like dapagliflozin and canagliflozin. These drugs prevent glucose reabsorption in the renal tubules and cause loss of sodium and glucose.

• UTI - due to glucosuria

• Dehydration - due to natriuresis and volume loss

• Euglycemic diabetic ketoacidosis - low blood glucose levels due to glucosuria can reduce insulin release

SGLT-2 inhibitors must be temporarily discontinued for 3-4 days before any scheduled surgery to avoid potential risk for diabetic ketoacidosis.

Management of diabetes mellitus		
Intervention	Advantages	Disadvantages
Lifestyle changes	Reduced complications and mortality	Insufficient for most patients within 1st year
Metformin	Weight neutral	GI side effectsContraindicated when GFR < 45 ml/min
Insulin	Rapidly effectiveLowers HbA1c more than other drugs (by 1.5-3%)	Weight gainHypoglycemia
GLP-1 agonists	Weight lossReduces cardiovascular complications	GI side effects
Pramlintide	Weight lossLowers postprandial glucose	GI side effectsHypoglycemia when used with insulin
Sulfonylureas	Rapidly effective	Weight gainHypoglycemia
Thiazolidinediones	Lowers insulin requirements	Weight gain, edema, CHF, fractures
Meglitinides	Rapidly effective	Weight gainHypoglycemia
SGLT-2 inhibitors	Weight lossBetter cardiovascular and renal outcomes	UTI, vulval candidiasisDehydrationEuglycemic DKA
DPP-4 inhibitors	Weight neutral	Angioedema, urticaria
Alpha-glucosidase inhibitors	Weight neutral	GI flatulenceRise in LFT

Solution to Question 18:

This clinical scenario describes the sliding scale method. Rapid-acting insulins such as glulisine are used.

Sold by @itachibot

Sliding scale refers to the administration of insulin as a bolus dose based on the blood sugar level before the meal. Using this principle, the insulin dose is titrated every time according to pre-meal glucose levels to determine insulin requirement. Hence, rapid-acting insulin is preferred.

This method requires a fixed carbohydrate intake at every meal. It is used for the short-term management of hospitalized patients, especially when the frequency or quantity of food intake is very variable.

It does not deliver insulin in a physiological manner, hence is not a preferred long-term strategy for glycemic control.

Solution to Question 19:

Afrezza is an inhalational form of insulin that can cause hypoglycemia.

It is a rapid-acting form of insulin that is administered at the beginning of each meal. Peak levels are achieved 12-15 minutes later.

Adverse effects include hypoglycemia, cough, throat pain, or irritation. It may cause acute bronchospasm in patients with asthma, COPD, as well as active smokers. Hence, it is contraindicated in these condition.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 20:

This clinical scenario is suggestive of diabetic ketoacidosis, which is treated with insulin infusion. Insulin causes an intracellular shift of K+ ions resulting in a decreased extracellular K+ ion concentration. This results in hypokalemia.

Non-diabetic clinical applications of insulin include:

• Treatment of hyperkalemia

• To determine the functions of the anterior pituitary

• To examine the completion of truncal vagotomy

Solution to Question 21:

Dawn phenomenon refers to normal blood glucose levels at 3 am followed by early morning hyperglycemia.

This occurs due to an early morning surge in plasma growth hormone and cortisol levels, which have an insulin-antagonistic action. In normal individuals, this is countered by increased early morning insulin secretion. However, in diabetic patients, insulin secretion is disrupted which results in the unopposed action of growth hormone. Hence, there will be resulting hyperglycemia.

Sold by @itachibot

Solution to Question 22:

When a patient needs > 200 units of insulin per day, they are said to have developed insulin resistance.

Some causes of insulin resistance are as follows:

• Werner syndrome

• Lipodystrophy syndrome

• Ataxia-telangiectasia

• Polycystic ovarian disease

Insulin resistance can present as non-alcoholic fatty liver disease.

Solution to Question 23:

The given clinical scenario is suggestive of diabetic ketoacidosis (DKA) and the first step is to assess the arterial blood gas (ABG) levels. This precedes correction of fluid balance, as assessment of the current status of the patient is necessary before beginning treatment. A baseline pH and electrolyte level must be obtained.

Diabetic ketoacidosis is the most serious complication of type 1 diabetes mellitus. The symptoms include dehydration, nausea, vomiting, confusion, drowsiness, and coma in severe cases.

In DKA, the insulin-to-glucagon ratio is decreased which promotes gluconeogenesis, glycogenolysis, and ketone body formation in the liver. Ketone bodies exist as ketoacids at physiological pH which are neutralized by bicarbonate. As bicarbonate stores are depleted, metabolic acidosis ensues. Increased lactic acid production also contributes to the acidosis.

The diagnostic biochemical criteria for DKA includes:

• Hyperglycemia (>250 mg/dL)

• Venous pH <7.3

• Plasma/serum bicarbonate <15-18 mmol/L

• Increased anion gap

• Ketonemia (beta-hydroxybutyrate [BOHB] >3 mmol/L) and ketonuria

Signs of ketoacidosis:

• Kussmaul breathing (deep, heavy, non-labored rapid breathing)

• Fruity breath odor (acetone)

• Prolonged Q-T interval

• Diminished neurocognitive function

• Coma

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Solution to Question 24:

The elevated levels of triglycerides (TG) and very low-density lipoproteins (VLDL) in diabetes mellitus (DM) are due to decreased activity of lipoprotein lipase (LpL) and increased activity of hormone-sensitive lipase (HSL).

Insulin normally has an inhibitory action on HSL and induces LpL on capillary walls. There is an absolute or relative deficiency of insulin in DM. In DM, the activity of LpL is decreased and HSL is increased.

The increased activity of HSL results in increased mobilisation of fat stores towards lipolysis releasing fatty acids. These fatty acids are partly cleared by the liver, resulting in increased synthesis of VLDL and TG, resulting in dyslipidemia. They can accumulate in the liver, resulting in steatosis (non-alcoholic fatty liver disease).

The reduced activity of LpL prevents the clearing of TG and VLDL in the bloodstream and their levels remain elevated.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot



Sold by @itachi... else,
If you purchased this from some...
you may have been scammed.

Solution to Question 25:

Fasting blood sugar >126 mg/dL and 2-hour postprandial blood sugar >200 mg/dl can be used as screening tests for diabetes in a primary health center (PHC).

HbA1c estimation is usually unavailable for screening at many PHCs due to low resources. In India, facilities for HbA1c estimation are usually available only at tertiary centers.

Random blood sugar >200 mg/dL can be used for diagnosis only if the patient presents with symptoms of diabetes, hence it is not a useful screening test.

Sold by @itachibot

Diabetes Mellitus: Complications and their Management

Question 1:

Which of the following is not a feature of diabetic ketoacidosis?

- **a)** Low serum sodium
- **b)** High anion gap
- **c)** Serum pH is < 7.4
- **d)** Serum bicarbonate > 15 meq/L

Question 2:

A 19-year-old woman with a history of type 1 diabetes is brought to the ER with confusion, vomiting, and abdominal pain. Investigations show blood glucose of 520 mg/dl and urine positive for ketone bodies. Which of the following is the drug of choice?

- **a)** Regular insulin
- **b)** Lente insulin
- **c)** Insulin glargine
- **d)** 70/30 insulin

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

Which of the following is the leading cause of death in children with diabetic ketoacidosis?

- **a)** Acute respiratory distress syndrome
- **b)** Cardiac arrhythmia
- **c)** Cerebral edema
- **d)** Acute kidney injury

Question 4:

A patient with diabetic ketoacidosis has severe abdominal pain. Which of the following is the recommended test to rule out acute pancreatitis?

Sold by @itachibot

a) Lipase

b) Elastase

c) Amylase

d) Trypsinogen

Question 5:

A 15-year old boy presents with persistent vomiting for one day. Laboratory investigations reveal the following. What is the best sequence of management?

a) Start IV fluids, administer IV regular insulin, and then correct hypokalemia

b) Start IV fluids, administer subcutaneous bolus of insulin and correct hypokalemia concurrently

c) Start IV fluids and correct hypokalemia first, then administer IV regular insulin

d) Administer IV regular insulin with KCl first , then start IV fluids once the K+ level is corrected

Question 6:

Distinguishing features between hyperglycemic hyperosmolar state and diabetic ketoacidosis include all of the following except:

a) Ketosis

b) Anion gap

c) Hypotension

d) Arterial pCO_2

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

An elderly man is brought to the hospital with confusion and drowsiness. His son says that he has been urinating more frequently than normal and eating very little over the past 1 week. Capillary blood glucose is 800 mg/dl. Which of the following would not be seen in this patient?

a) Serum osmolality > 350 mOsm/L

b) Normal pH

c) Kussmaul breathing

d) Prerenal azotemia

Sold by @itachibot

Question 8:

What is the most common pattern of dyslipidemia associated with diabetes mellitus?

a) Increased triglycerides and decreased HDL

b) Increased LDL and decreased HDL

c) Increased VLDL and decreased HDL

d) Increased triglycerides and increased VLDL

Question 9:

Dyslipidemia is more commonly associated with which of the following?

a) Type 2 diabetes mellitus

b) Type 1 diabetes mellitus

c) Maturity onset diabetes of the young

d) Gestational diabetes mellitus

Question 10:

You have started a 40-year-old man on treatment for type 2 diabetes mellitus and dyslipidemia. Which of the following is not an appropriate target for this patient?

a) HbA1c < 7 %

b) Blood pressure < 130 / 80 mmHg

c) Triglycerides < 150 mg/dL

d) Postprandial plasma glucose < 200 mg/dL

Question 11:

A middle-aged patient with type 2 diabetes is diagnosed with essential hypertension. What is the antihypertensive drug of choice?

a) Losartan

b) Hydrochlorothiazide

c) Atenolol

d) Amlodipine

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 12:

Which of the following is the earliest manifestation of diabetic nephropathy?

a) Microalbuminuria

b) Increased glomerular filtration rate

c) Decreased glomerular filtration rate

d) Increased urine PCR

Question 13:

A 15-year-old boy has been on treatment for type 1 diabetes mellitus since the age of 10 years. His physician decides to begin screening for diabetic nephropathy. Which of the following is not always a reliable indicator of this condition?

a) Long duration of diabetes

b) Albuminuria of 30-300 mg/day

c) Presence of diabetic retinopathy

d) Reduced glomerular filtration rate

Question 14:

Which of the following statements about diabetic nephropathy is true?

a) Pancreatic transplantation can improve proteinuria at an early stage

b) Strict glycemic control is unlikely to prevent microalbuminuria

c) Dietary protein restriction can benefit diabetic patients with albuminuria

d) Microalbuminuria is not an indicator of long-term cardiovascular morbidity

Question 15:

A 50-year-old man with type 2 diabetes mellitus is found to have a 24-hour urinary albumin of 250 mg. Which of the following drugs may be prescribed to reduce the progression of this condition?

a) Hydrochlorothiazide

b) Enalapril

c) Amiloride

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

d) Eplerenone

Question 16:

A pathologist is studying a renal biopsy sample from a patient with suspected diabetic nephropathy. Which of the following biopsy findings is pathognomonic for this condition?

a) Diffuse glomerulosclerosis

b) Nodular glomerulosclerosis

c) Armani-Ebstein reaction

d) Basement membrane thickening

Question 17:

Which of the following types of renal tubular acidosis can develop in individuals with diabetes mellitus?

a) Type I RTA

b) Type II RTA

c) Type III RTA

d) Type IV RTA

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A 45-year-old man has been diagnosed with type 2 diabetes mellitus. When should you refer him to an ophthalmologist?

a) Within one year of diagnosis

b) Immediately at the time of diagnosis

c) Within five years of diagnosis

d) Within three years of diagnosis

Question 19:

Which of the following patients would you screen for diabetic retinopathy?

a) 2, 4, 5

b) 1, 4, 5

Sold by @itachibot

c) 2, 3, 5

d) 1, 2, 4

Question 20:

What is the most common clinical presentation of diabetic neuropathy?

a) Autonomic neuropathy

b) Mononeuropathy

c) Distal polyneuropathy

d) Polyradiculopathy

Question 21:

Which of the following findings in a diabetic patient is self-limiting?

a) Macular edema

b) Peripheral neuropathy

c) Amyotrophy

d) Macroalbuminuria

Question 22:

You are evaluating a 65-year-old diabetic woman who has been admitted following a seizure. On admission, she was unconscious, and random blood glucose at admission was 37 mg/dl. The hypoglycemia was corrected and she regained consciousness. The woman says that she had felt normal and had no other symptoms prior to the onset of seizures. What is the likely cause of this presentation?

a) Medication change from metformin to insulin

b) Insulin resistance

c) Autonomic neuropathy

d) Concurrent use of thiazide diuretics

Question 23:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

You are counselling a patient with longstanding, poorly-controlled diabetes mellitus. Advising tight glycemic control in this patient would not improve which of the following?

a) Neuropathy

b) Retinopathy

c) Peripheral vascular disease

d) Nephropathy

Question 24:

Which among the following statements explains Legacy effect and metabolic memory?

a) 1 and 5

b) 1 and 2

c) 2 and 3

d) 4 and 5

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 25:

A patient with chronic diabetes presents to the OPD for his annual checkup. A small painless ulcer is noted over his right foot. His BMI is 33 kg/m2. All of the following factors can be implicated in the causation of his foot ulcer except:

a) Venous stasis

b) Neuropathy

c) Immunosuppression

d) Ischemia

Question 26:

A 25-year-old diabetic woman presents to the OPD with the following skin lesions. What is the likely diagnosis?

Sold by @itachibot

a) Necrolytic migratory erythema

b) Diabetic dermopathy

c) Necrobiosis lipoidica

d) Granuloma annulare

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 27:

Patients with diabetes mellitus are at a higher risk of developing all of the following infections except:

a) Malignant otitis externa

b) Rhinocerebral mucormycosis

c) Emphysematous pyelonephritis

d) Emphysematous appendicitis

Question 28:

A morbidly obese diabetic woman is not responsive to metformin therapy. She has a history of pancreatitis and family history of bladder cancer. She does not want to take injections. Which of the following would be suitable to reduce her glucose levels?

a) Sitagliptin

b) Pioglitazone

c) Canagliflozin

Sold by @itachibot

d) Liraglutide

Question 29:

Which of the following anti-diabetic drugs should be stopped in a patient with a creatinine of 5.6 mg/dL?

a) Metformin

b) Linagliptin

c) Metoprolol

d) Insulin

If you bought this from someone else, it was scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	a
3	c
4	a
5	c
6	c
7	c
8	a
9	a
10	d
11	a
12	b
13	b
14	a
15	b
16	b
17	d
18	b
19	a
20	c

Sold by @itachibot

21	c
22	c
23	c
24	a
25	a
26	c
27	d
28	c
29	a

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

In diabetic ketoacidosis (DKA), serum bicarbonate is usually < 15. Serum HCO3 > 15 meq/L is suggestive of alcoholic ketoacidosis.

DKA is an acute complication of type 1 diabetes mellitus. It is usually precipitated by absolute or relative insulin deficiency caused by increased metabolic demand (infections) or insulin noncompliance.

The common symptoms include nausea, vomiting, and severe abdominal pain. The common signs include Kussmaul breathing, fruity odor of the breath, dehydration, tachycardia, hypotension, and delirium.

The diagnosis of DKA is made based on the following parameters:

• Hyperglycemia > 250 mg/dl

• Ketosis or ketones in urine

• Metabolic acidosis with pH < 7.3

• HCO3 < 15-18 mEq/L

• High anion gap due to ketoacids and lactic acidosis

Hyponatremia is seen as a result of hyperglycemia (osmotic shift causes dilutional hyponatremia). Normal sodium levels on admission indicate severe dehydration. Despite a total-body potassium deficit, the serum Potassium at presentation may be mildly elevated, secondary to the acidosis and volume depletion.

Solution to Question 2:

The drug of choice for controlling hyperglycemia in a patient with diabetic ketoacidosis (DKA) is short-acting insulin, preferably regular insulin.

Sold by @itachibot

Intermediate (such as lente insulin and NPH), long-acting insulins, and oral hypoglycemic agents are never used in the initial management of DKA.

The mainstay of treatment is the administration of normal saline followed by infusion of short-acting insulin. The precipitating event must be identified and treated. Careful monitoring of K+ levels and replacement in the later stages is required, as insulin therapy causes hypokalemia.

Solution to Question 3:

Cerebral edema is the leading cause of mortality in children with diabetic ketoacidosis.

The osmolar gradient generated due to the high blood glucose level results in water shift from the intracellular fluid (ICF) to the extracellular fluid (ECF) space and leads to the contraction of cell volume. Correction with insulin and intravenous fluids can result in a rapid reduction in effective osmolarity, reversal of the fluid shift, and result in cerebral edema.

Complications of diabetic ketoacidosis include:

- Cerebral edema – especially in children
- Cardiac arrhythmia - due to electrolyte imbalance
- Acute respiratory distress syndrome (ARDS)
- Venous thrombosis
- Acute gastric dilation
- Acute kidney injury
- Intestinal necrosis
- Acute pancreatitis
- Mucormycosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

In diabetic ketoacidosis (DKA), severe abdominal pain can mimic acute pancreatitis. Serum lipase is the best test to rule out this condition. It must then be confirmed by imaging (USG or CT abdomen).

Serum amylase is an unreliable indicator as DKA is associated with hyperamylasemia of salivary origin.

Serum and urine trypsinogen may be elevated in early pancreatitis but are not used for diagnosis. Fecal elastase is elevated in chronic pancreatitis.

Solution to Question 5:

Sold by @itachibot

This clinical scenario is suggestive of diabetic ketoacidosis (DKA). On confirming the diagnosis, IV fluids must be started. If the initial serum potassium < 3.3 meq/L, insulin should be administered only after the correction of hypokalemia.

This is because treatment with insulin and fluids can worsen hypokalemia by the following mechanisms:

• Insulin causes the shift of K+ ions into cells

• Resolution of the acidosis also promotes K+ entry into cells

• Urinary loss of K+ salts of organic acids



Solution to Question 6:

Hypotension and tachycardia are common features of both hyperglycemic hyperosmolar state (HHS) and diabetic ketoacidosis (DKA).

In DKA, insulin deficiency and an increase in counter-regulatory hormones lead to shunting of free fatty acids to form ketone bodies. The ketoacids increase the anion gap and cause low pH. In HHS, there is still some insulin available. Hence, there are no ketones formed, and the patient has normal pH and pCO2.

Solution to Question 7:

This clinical scenario is suggestive of hyperglycemic hyperosmolar state (HHS). Kussmaul's breathing is not a feature of HHS.

HHS is commonly seen in older adults suffering from type 2 diabetes mellitus. Patients can present with the following features:

• Polyuria

Sold by @itachibot

• Polydipsia

• Lethargy

• Altered sensorium, coma

• Seizures

• Severe dehydration

The following lab values are seen:

• Blood glucose of 600-1200 mg/dl

• Serum osmolality > 350 mOsm/L

• Prerenal azotemia due to dehydration

• Pseudohyponatremia may be seen due to hyperglycemia

• Normal pH and ketones

The mainstay of management is fluid replacement, IV regular insulin, along with potassium replacement.

Heparin can be given prophylactically as volume contraction can increase the risk of venous thrombosis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

Dyslipidemia commonly occurs as a complication of type 2 diabetes mellitus. The most common pattern is as follows:

• Increased triglyceride levels - due to increased glucose and free fatty acids, decreased lipolysis of VLDL.

• Decreased HDL levels

Though diabetes does not increase LDL levels, there is increased atherogenicity as the existing LDL molecules are more easily glycated and susceptible to oxidation.

Solution to Question 9:

Dyslipidemia is more commonly associated with type 2 diabetes mellitus.

It is less common in type 1 and gestational diabetes. Maturity onset diabetes of the young (MODY) is not associated with dyslipidemia.

Note: Saroglitazar is a more advanced drug that reduces blood sugar and lipid levels in diabetic patients. It acts on PPAR alpha and gamma receptors.

Solution to Question 10:

Sold by @itachibot

According to the ADA guidelines, a treatment goal of postprandial plasma glucose < 180 mg/dL must be followed in adults with diabetes.

In younger individuals or those with other cardiovascular risk factors, the target blood pressure is < 130/80 mm Hg. Although not a treatment goal, the ADA also recommends triglyceride levels be maintained at < 150 mg/dl.

Note: Saroglitazar is a more advanced drug that reduces blood sugar and lipid levels in diabetic patients. It acts on PPAR alpha and gamma receptors.

Solution to Question 11:

The drug of choice in this diabetic patient with hypertension is losartan.

Angiotensin-converting enzyme inhibitors (ACE inhibitors) or angiotensin receptor blockers (ARB) are the first-line drugs in diabetic hypertensives. Among ARBs, the drug of choice is telmisartan (acts on PPAR receptor).

They reduce both macrovascular and microvascular complications and have a nephroprotective effect. However, they are contraindicated in chronic kidney disease as they can cause a further decline in GFR.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

The earliest manifestation of diabetic nephropathy is an increase in the glomerular filtration rate (GFR).

It occurs as a result of glomerular hyperperfusion and renal hypertrophy. The following sequence of changes are seen in diabetic nephropathy:

• Increased GFR

• Incipient diabetic nephropathy - microalbuminuria (30–300 mg/day albumin)

• Diagnostic marker to detect early diabetic nephropathy

• Indicator of long-term cardiovascular disease

• Overt diabetic nephropathy - macroalbuminuria (>300 mg/day albumin)

• End-stage renal disease

Sold by @itachibot



Sold by @itachi... If you purchased this from some... you may have been scammed. ...else,

Solution to Question 13:

Albumin of 30-300 mg/day refers to microalbuminuria. It can be unreliable in the diagnosis of diabetic nephropathy.

It is not required to diagnose nephropathy as it is very variable. Non-albuminuric diabetic nephropathy may occur in some patients and regression of albuminuria may be seen in type 1 diabetics.

Diabetic nephropathy can be diagnosed if the following are satisfied:

• Reduced GFR (< 60 mL/min/1.73 m2) with or without persistent albuminuria (> 30 mg/day or urine PCR > 30 mg/g)

• Long duration of diabetes or established diabetic retinopathy

• Other etiologies are ruled out

Long duration in type 1 diabetes refers to 5 years after diagnosis. On the other hand, patients with type 2 diabetes are exposed to hyperglycemia and prediabetes for a long time even before diagnosis. Hence, at the time of diagnosis, patients with type 2 diabetes are considered to have had the disorder for a duration sufficiently long to produce kidney disease.

However, a clinical diagnosis of diabetic nephropathy can be made in patients with established retinopathy, even if the duration of diabetes is short.

Solution to Question 14:

Pancreatic transplantation can help improve proteinuria at an early stage. Successful pancreatic transplantation can have the following effects:

• Prevention of diabetic nephropathy and regression of albuminuria in early stages.

Sold by @itachibot

• Stabilization and improvement of diabetic neuropathy and retinopathy

• Increase in HDL with reduction in LDL and triglycerides

Option B: Strict glycemic control can prevent or revert microalbuminuria.

Option C: The ADA does not suggest restrictions of protein intake in diabetic patients with albuminuria because studies have failed to prove its benefits.

Option D: Microalbuminuria is an indicator of long-term cardiovascular morbidity.

Solution to Question 15:

Angiotensin-converting enzyme (ACEI) inhibitors or angiotensin-receptor blockers (ARB) are recommended to reduce the progression of diabetic nephropathy. It has the following effects:

• Reduction of albuminuria

• Reduces the associated decline in GFR

The mainstay of management of diabetic nephropathy includes control of blood sugar and blood pressure, inhibitors of the renin-angiotensin-aldosterone(RAAS) system, and inhibitors of SGLT2 to halt the progression of the disease.

RAAS inhibitors include ACE inhibitors, ARBs, or direct renin inhibitors.

Solution to Question 16:

The characteristic lesion of diabetic nephropathy is nodular or intercapillary glomerulosclerosis, also called Kimmelstiel-Wilson nodules. It is pathognomonic for this condition.

The following image shows Kimmelstiel-Wilson nodules.



Other non-specific findings seen on biopsy include:

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

• Glomerular basement membrane (GBM) thickening - highly sensitive

• Diffuse glomerulosclerosis - most common pattern, due to diffuse mesangial thickening

• Fibrin cap - accumulation of hyaline material within the capillary loop

• Capsular drop - accumulation of hyaline material attached to the Bowman's capsule

• Armani-Ebstein reaction - collection of glycogen clumps within the renal tubules due to tubular injury

Renal biopsy is done to rule out other causes of renal failure. The indications for renal biopsy in diabetic nephropathy are:

• Absence of diabetic retinopathy

• Absence of edema

• Absence of frothing and proteinuria

• Absence of hypertension

• Presence of microscopic hematuria or features of nephritic syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

Type IV renal tubular acidosis (RTA) may occur in individuals with diabetes mellitus types 1 and 2.

Type IV RTA occurs due to hyporeninemic hypoaldosteronism. Low plasma renin activity is common in diabetic patients due to:

• Impaired conversion of prorenin to renin

• Volume expansion in diabetic nephropathy suppresses renin release

These individuals have a tendency to develop hyperkalemia and acidemia, which can be exacerbated by the use of ACEI and ARBs. Hence, these drugs should be avoided.

Solution to Question 18:

Patients with type 2 diabetes should undergo a comprehensive ophthalmological examination to screen for diabetic retinopathy at diagnosis, followed by an annual examination of the fundus.

In patients with type 1 diabetes mellitus, screening for retinopathy is recommended 3 to 5 years after diagnosis. Annual fundal examination is recommended thereafter.

Solution to Question 19:

Screening for diabetic retinopathy is done at 3-5 years following diagnosis in type 1 diabetes mellitus, whereas, patients with type 2 diabetes must be screened at the time of diagnosis.

Sold by @itachibot

As type 2 diabetes mellitus often has a long asymptomatic period of hyperglycemia before clinical diagnosis, these patients are more likely to have complications at the time of diagnosis itself.

Retinopathy develops in nearly 50-80% of patients by 20 years of diagnosis. The presence of diabetic retinopathy often correlates with pathological findings of diabetic nephropathy.

Solution to Question 20:

The most common form of diabetic neuropathy is distal symmetric polyneuropathy. It presents with distal sensory loss and pain in a stocking-and-glove distribution and is a risk factor for foot ulceration.

Autonomic neuropathy is a common complication and results in the following features:

- Postural hypotension
- Silent myocardial infarction
- Hypoglycemia unawareness
- Erectile dysfunction
- Overflow incontinence
- Gastroparesis
- Enteropathy with constipation or diarrhea

Mononeuropathy of peripheral (median, ulnar, radial) and cranial (oculomotor) nerves is also seen, as well as mononeuritis multiplex.

Diabetic polyradiculopathy is characterized by severe pain along the distribution of one or more nerve roots.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 21:

Amyotrophy refers to weakness of hip extensors and flexors secondary to polyradiculopathy involving the lumbar plexus and femoral nerve. It is associated with severe pain along the distribution of the nerve roots. It is self-limiting and resolves over 6-12 months.

Strict glycemic control can reduce the incidence and prevent the progression of macular edema, peripheral neuropathy, and macroalbuminuria. However, once developed, they cannot be reversed. Intermittent microalbuminuria with normal GFR can be reversed with strict glycemic control.

Solution to Question 22:

This clinical scenario is suggestive of silent hypoglycemia in a diabetic patient. The most likely cause is autonomic neuropathy.

Sold by @itachibot

Hypoglycemia presents with symptoms like palpitation, tremors, and sweating as a result of sympathetic stimulation. These symptoms are protective and provide advance warning before progression to neuroglucopenia.

However, in autonomic neuropathy, there is a reduced release of counterregulatory hormones like catecholamines. Hence, the classical symptoms of hypoglycemia do not occur and the patient directly presents with features of neuroglucopenia. This is called hypoglycemia unawareness.

The use of beta-blockers can also cause hypoglycemia unawareness.

Solution to Question 23:

In patients with longstanding diabetes mellitus, tight glycaemic control is beneficial for microvascular complications. However, it has no effect on macrovascular complications like peripheral vascular disease (PVD), myocardial infarction, and stroke.

Good glycemic control improves microvascular complications through primary and secondary prevention. It reduces the incidence of complications as well as stabilizes and prevents further progression in affected individuals. Intermittent microalbuminuria with normal GFR can even be reversed with tight glycemic control.

Complication	% risk reduction with tight glycemic control
Retinopathy	47 %
Microalbuminuria	39 %
Nephropathy	54%
Neuropathy	60 %

If you p... by @itachibot ... from someone else, you ... examined.

Solution to Question 24:

Legacy effect or metabolic memory can be observed in relation to glycemic control during early disease in patients with diabetes mellitus.

Intensive glycemic control during the early stages of the disease has a positive effect on long-term microvascular and macrovascular complications (statement 1) even if this intensive glycemic control is not maintained long term. Similarly, uncontrolled blood glucose levels in the early stages causes an increased risk of these complications (statement 5). This phenomenon is called legacy effect and is attributed to metabolic memory.

In patients with longstanding diabetes, intensive glycemic control at later stages can improve microvascular complications but has no effect on macrovascular complications.

Sold by @itachibot

Solution to Question 25:

Venous stasis is not a cause of diabetic foot ulcers and gangrene. They are usually caused by a combination of three factors:

• Ischemia - secondary to atheroma

• Peripheral neuropathy - leads to trophic skin changes

• Immunosuppression - due to excess sugar deposition in tissues, predisposing diabetic patients to infection

Common sites of ulcers include:

• Great toe

• Metatarsophalangeal areas.

Solution to Question 26:

The given image showing irregular plaques with yellowish discoloration and telangiectasias are suggestive of necrobiosis lipoidica diabeticorum.

It is an inflammatory granulomatous disorder of the skin due to collagen degeneration and lipid deposition. It is usually seen in young women with diabetes mellitus.

The lesions usually begin in the pretibial region. It begins as papules that gradually enlarge and darken to form erythematous plaques with irregular margins, atrophic centers, and central ulceration.

Option A: Necrolytic migratory erythema is a feature of glucagonoma. It is a waxing and waning rash that appears as shown below.

Option B: Diabetic dermopathy or shin spots are commonly seen in elderly men. It presents as pigmented pretibial papules as shown below.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option D: Granuloma annulare presents with small papules arranged in an annular pattern. There is no yellowish discoloration or telangiectasias, as shown below.

Solution to Question 27:

Emphysematous appendicitis is not a complication of diabetes mellitus. The potentially fatal infectious complications of diabetes include:

- Malignant otitis externa
- Rhinocerebral mucormycosis
- Emphysematous pyelonephritis
- Emphysematous cholecystitis

Sold by @itachibot

Solution to Question 28:

Among the given options, canagliflozin (Option C) is the most suitable treatment. It is an oral drug belonging to the class SGLT-2 inhibitors and causes weight loss. Although it increases the risk of urogenital infections, it has no risk of pancreatitis or bladder cancer.

Option A: Sitagliptin, a DPP-IV inhibitor, is associated with the risk of acute pancreatitis.

Option B: Pioglitazone, a thiazolidinedione, can increase the risk of bladder cancer and can cause weight gain and congestive cardiac failure.

Option D: Liraglutide is an injectable GLP-1 receptor agonist. Pancreatitis is a serious adverse effect of this drug. Hence, it is contraindicated in individuals with a history of pancreatitis and should be permanently discontinued if pancreatitis develops.

Solution to Question 29:

Metformin should be stopped in case of elevated creatinine levels (renal insufficiency) > 1.5 mg/dl in men and >1.4 mg/dl in women.

Metformin acts by reducing hepatic glucose production and improves peripheral glucose utilization. Lactic acidosis and Vitamin B12 deficiency are specific adverse effects of metformin. It is not used in patients with a glomerular filtration rate <30 mL/min (moderate renal insufficiency), acidosis, unstable congestive cardiac failure, liver disease, severe hypoxemia, and patients undergoing contrast studies.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Reproductive Endocrinology

Question 1:

Which of the following is not a feature of 11-beta-hydroxylase deficiency?

a) Metabolic acidosis

b) Hypokalemia

c) Hypertension

d) Virilization

Question 2:

A neonate with ambiguous genitalia undergoes karyotyping and is confirmed to have a 46XY genotype. His blood pressure is 140/80 mm Hg and serum potassium is 1.9 mEq/L. Which of the following is responsible for these findings?

a) 11-deoxycorticosterone

b) Cortisol

c) 17-hydroxypregnanediol

d) 17-hydroxyprogesterone

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

Which of the following abnormalities will you see in a child with salt-wasting congenital adrenal hyperplasia?

a) Hypoglycemia

b) Hypernatremia

c) Hypertension

d) Hypokalemia

Question 4:

Which of the following is the drug of choice for the prenatal treatment of congenital adrenal hyperplasia due to 21-hydroxylase deficiency?

Sold by @itachibot

a) Dexamethasone

b) Betamethasone

c) Prednisolone

d) Hydrocortisone

Question 5:

A female neonate is noted to have ambiguous genitalia at birth. Further testing reveals elevated 17-(OH)-progesterone levels and a blood glucose of 40 mg/dl. What treatment will you recommend?

a) Betamethasone

b) Beclomethasone

c) Dexamethasone

d) Hydrocortisone

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 6:

Which of the following is incorrect about galactorrhea?

a) May be bilateral

b) Associated with lactation

c) Associated with prolactinoma

d) Associated with acromegaly

Question 7:

In a patient with gynecomastia, you will order all of the following tests except:

a) Testosterone

b) Prolactin

c) Estradiol

d) Luteinizing hormone

Question 8:

Sold by @itachibot

A 32-year-old man is being evaluated for gynecomastia. Laboratory investigations reveal low levels of serum testosterone and luteinizing hormone. What is the likely diagnosis?

a) Testicular failure

b) Sertoli cell tumor

c) Gonadotropin-secreting tumor

d) Androgen-resistant state

Question 9:

A mother brings her 17-year-old daughter to the gynecologist as she has not started menstruating yet. Examination reveals well-developed breasts and the absence of pubic and axillary hair. Pelvic examination reveals a short, blind-ended vagina. What is the likely diagnosis?

a) Turner's syndrome

b) Stein-Leventhal syndrome

c) Premature ovarian failure

d) Complete androgen insensitivity syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A baby girl is referred to a pediatric surgeon for suspected bilateral inguinal hernias. Further evaluation reveals the masses to be testes in the inguinal canal. Which karyotype would you expect to find in this child?

a) 46 XX

b) 46 XY

c) 47 XXY

d) 47 XYY

Question 11:

An 18-year-old boy is brought to the casualty with minor burns following a house fire. He tells you that he did not even smell the smoke. Further probing reveals a history of bilateral orchidopexy at the age of 7. Examination findings include eunuchoid habitus, prepubertal testes, and microphallus. What is the likely diagnosis?

a) Primary hypogonadism

Sold by @itachibot

b) Klinefelter syndrome

c) Kallman syndrome

d) Hyperprolactinemia

Question 12:

What is the most common cause of hypergonadotropic hypogonadism in males?

a) Viral orchitis

b) Klinefelter's syndrome

c) Kallman syndrome

d) Noonan syndrome

Question 13:

Which of the following will you recommend in all patients with androgen insensitivity syndrome?

a) Gonadectomy

b) Testosterone replacement

c) Vaginoplasty

d) Estrogen replacement

Question 14:

A 19-year-old patient is being evaluated for primary amenorrhea. Laboratory investigations reveal decreased estrogen, FSH, and LH. What is the likely diagnosis?

a) Asherman syndrome

b) Müllerian agenesis

c) Kallman syndrome

d) Turner's syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	a
2	a
3	a
4	a
5	d
6	b
7	b
8	b
9	d
10	b
11	c
12	b
13	a
14	c

Sold by @itachibot
If you purchased this from else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

11-beta-hydroxylase deficiency is associated with metabolic alkalosis, not metabolic acidosis.

It is a rare form of congenital adrenal hyperplasia that presents with the following features:

• Glucocorticoid deficiency:

• Hypoglycemia

• Hyperpigmentation around the genitals due to excess adrenocorticotropic hormone (ACTH) secretion

• Mineralocorticoid excess:

• Hypokalemia

• Hypertension

• Metabolic alkalosis

• Adrenal androgen excess

• Virilization in females

• Precocious puberty in males

Sold by @itachibot

Solution to Question 2:

This clinical scenario with ambiguous genitalia in a male neonate is suggestive of 17 alpha-hydroxylase deficiency. It causes hypertension and hypokalemia due to the accumulation of 11-deoxycorticosterone, which is a mineralocorticoid causing sodium retention.

11-hydroxylase deficiency also leads to the accumulation of 11-deoxycorticosterone.

Solution to Question 3:

Hypoglycemia is a feature of salt-wasting congenital adrenal hyperplasia (CAH).

It is due to 21-hydroxylase deficiency. It results in hyponatremia, hyperkalemia, and hypotension. Males typically present with salt-wasting crises as they often go undiagnosed at birth due to the presence of normal genitalia.

Solution to Question 4:

The drug of choice for the prenatal treatment of congenital adrenal hyperplasia (CAH) due to 21-hydroxylase deficiency is dexamethasone, as it readily crosses the placenta.

It is an experimental approach aimed at reducing the virilization of 46XX fetuses by suppressing fetal adrenal steroid and androgen secretion. It may be considered when there is a high risk of CAH due to strong family history. It is effective only when started before 4 weeks of gestation.

Hydrocortisone is given in the children at divided doses of 1–1.5 times the normal cortisol production.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 5:

Ambiguous genitalia, elevated 17-(OH)-progesterone, and hypoglycemia are suggestive of 21-hydroxylase deficiency. The drug of choice is hydrocortisone.

Hydrocortisone acts as a potent glucocorticoid with mineralocorticoid action and thus increases salt absorption from the kidneys as well.

Betamethasone, beclomethasone, and dexamethasone have negligible effects as mineralocorticoids.

Solution to Question 6:

Galactorrhea or pathological production of milk is not associated with lactation, which is a physiological condition.

Sold by @itachibot

Galactorrhea is the inappropriate discharge of milk-containing fluid from the breast. It may be unilateral or bilateral. It is associated with acromegaly and hyperprolactinemia.

Treatment is with dopamine agonists like bromocriptine or cabergoline if it is due to hyperprolactinemia.

Solution to Question 7:

Determination of prolactin level is not useful in patients with gynecomastia.

For the diagnosis of gynecomastia, serum levels of the following hormones must be checked:

- Testosterone
- LH
- FSH
- Estradiol
- hCG

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

This clinical scenario of gynecomastia with low LH and testosterone is suggestive of a sertoli cell tumor.

Hypogonadotropic hypogonadism is also a possible diagnosis with the above clinical presentation and must be ruled out before diagnosing a Sertoli cell tumor.

The differential diagnosis of gynecomastia is as follows:

- High LH and low testosterone
- Testicular failure
- Low LH and low testosterone
- Hypogonadotropic hypogonadism
- Sertoli cell tumor - increased primary estrogen production
- High LH and high testosterone
- Androgen-resistant state
- GnRH-secreting tumor

Solution to Question 9:

The given clinical scenario is suggestive of complete androgen insensitivity syndrome.

Clinical features:

Sold by @itachibot

• Female phenotype

• Normal breast development

• Blind-ended vagina with absent uterus and ovaries

• Presence of testes in the abdomen or inguinal region

• Scanty pubic and axillary hair

• Primary amenorrhea in late adolescence

Option A: In Turner's syndrome, breasts are infantile.

Option B: Stein Leventhal syndrome or polycystic ovarian disorder presents with hirsutism and anovulatory menstrual cycles.

Option C: Premature ovarian failure presents with secondary amenorrhea.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

This clinical picture is suggestive of androgen insensitivity syndrome (AIS). Affected children have a male karyotype or 46 XY with a female phenotype.

46XY individuals with complete AIS (formerly called testicular feminization syndrome) have the following features:

• Female phenotype

• Normal breast development

• Blind-ended vagina with absent uterus and ovaries

• Scanty pubic and axillary hair

Most patients present with

• Inguinal masses in infancy - Due to undescended testes or hernias containing testes

• Primary amenorrhea in late adolescence

Gonadectomy is a recommended course of treatment due to the possible risks of malignancy.

Solution to Question 11:

Anosmia and absent secondary sexual characteristics in this patient are suggestive of Kallman syndrome.

It is an X-linked disorder that causes isolated hypogonadotrophic hypogonadism due to defective development of the olfactory bulb. This results in:

• Defective migration of GnRH-producing neurons that originate from the olfactory epithelium

• Aplasia or hypoplasia of olfactory bulb

The clinical features include:

Sold by @itachibot

• Cryptorchidism

• Absent or delayed pubertal changes due to GnRH deficiency

• Anosmia or hyposmia

Options A and B: Klinefelter syndrome is a cause of primary or hypergonadotrophic hypogonadism.

Option D: Hyperprolactinemia can cause hypogonadotrophic hypogonadism, but is also associated with gynecomastia, galactorrhea, and bitemporal hemianopia.

Solution to Question 12:

The most common cause of hypergonadotrophic hypogonadism in males is Klinefelter's syndrome

It occurs in males with a 47 XXY karyotype and is associated with testicular dysfunction and male infertility.

Causes of hypergonadotrophic hypogonadism in males and females:

• Congenital

• Klinefelter's syndrome

• Turner's syndrome

• Swyer's syndrome

• Premature ovarian failure associated with fragile X syndrome and galactosemia

• Acquired

• Mumps orchitis

• Testicular torsion, trauma

• Liver cirrhosis

• Ketoconazole

• Premature ovarian failure

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 13:

Gonadectomy is clinically recommended in all patients with androgen insensitivity syndrome (AIS) or testicular feminization syndrome due to the risk of malignancy arising from the abnormally located testes.

Options B, C, D: These treatment options may be offered to patients with AIS, but it depends on personal preference and whether they identify as males or females.

Solution to Question 14:

Sold by @itachibot

Decreased levels of estrogen, FSH, and LH are suggestive of hypogonadotrophic hypogonadism, which is caused by Kallman syndrome.

It is an X-linked disorder (KAL gene) that causes isolated hypogonadotrophic hypogonadism and anosmia.

Causes of hypogonadotrophic hypogonadism:

• Congenital

• Kallman's syndrome

• Isolated hypogonadotrophic hypogonadism (IHH)

• Prader-Willi syndrome

• CHARGE syndrome

• Acquired

• Pituitary tumors, apoplexy

• Hyperprolactinemia

• Anorexia nervosa, excess exercise

• Irradiation

• Infection

• Trauma

Option A: Asherman syndrome causes secondary amenorrhea due to intrauterine synechiae.

Option B: Müllerian agenesis presents with primary amenorrhea with normal levels of all hormones.

Option D: Turner's syndrome causes primary amenorrhea due to hypergonadotrophic hypogonadism.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Disorders of Parathyroid and Calcium Homeostasis

Question 1:

Hypercalcemia is associated with all of the following medical conditions except:

a) Multiple myeloma

b) Hyperparathyroidism

c) Sarcoidosis

d) Hypothyroidism

Question 2:

Which of the following can cause hypercalcemia in a patient?

a) Vitamin D deficiency

b) Amyloidosis

c) Vitamin A intoxication

d) Vitamin C intoxication

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

A 66-year-old woman is brought to the emergency department after being found unconscious by her neighbors. She is found to have marked dehydration and serum calcium levels of 16 mg/dl. All of the following can cause this presentation except:

a) Metastatic breast carcinoma

b) Hyperparathyroidism

c) Pancreatitis

d) Hodgkin's lymphoma

Question 4:

Which of the following will not be seen in a patient with hypercalcemia?

a) Polyuria

b) Diarrhoea

Sold by @itachibot

c) Depression

d) Vomiting

Question 5:

Which of the following is not seen in patients following excessive intake of calcium?

a) Chronic renal failure

b) Milk-alkali syndrome

c) Burnett's syndrome

d) Restrictive cardiomyopathy

Question 6:

What is the most common cause of primary hyperparathyroidism?

a) Parathyroid hyperplasia

b) Parathyroid adenoma

c) MEN 1 syndrome

d) Iatrogenic conditions

Question 7:

Which of the following sets of lab values will be seen in a patient with secondary hyperparathyroidism?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

	Calcium	PTH	Phosphate	ALP
1	Normal	Increased	Decreased	Increased
2	Increased	Increased	Decreased	Increased
3	Decreased	Increased	Increased	Increased
4	Decreased	Increased	Normal	Normal

©MARROW

MARROW

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

a) Option 3

b) Option 1

c) Option 4

d) Option 2

Question 8:

All of the following statements are true about primary hyperparathyroidism except:

a) It causes prolongation of QT interval

b) Asymptomatic disease can occur

c) Peptic ulcer disease is seen

d) Hyperuricemia can occur

Question 9:

Which of the following would you expect in a patient with primary hyperparathyroidism?

a) Gallstones

b) Recurrent abortions

c) Hypotension

d) Neuropsychiatric changes

Sold by @itachibot

Question 10:

Calcitonin levels are increased in:

a) Hyperthyroidism

b) Hypoparathyroidism

c) Hyperparathyroidism

d) Cushing's syndrome

Question 11:

The spinal x-ray of a patient with elevated ALP levels is shown below. What is the likely diagnosis?



a) Paget's disease

b) Hyperparathyroidism

c) Osteopetrosis

d) Ankylosing spondylitis

Question 12:

A middle-aged woman presents with worsening pain and an inability to bear weight on her left leg. She says that she had banged her leg against a chair 3 days ago. X-ray reveals a fracture of her left tibia. In view of this finding, a skeletal survey was performed and reveals the following additional finding. What is the likely cause?

Sold by @itachibot

a) Renal osteodystrophy

b) Osteomalacia

c) Hyperparathyroidism

d) Hypoparathyroidism

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

This x-ray finding in a patient is characteristic of which of the following conditions?

a) Hyperparathyroidism

b) Down's syndrome

Sold by @itachibot

c) Acromegaly

d) Psoriasis

Question 14:

Which of the following is not used to treat a patient with hypercalcemia due to primary hyperparathyroidism?

a) Forced diuresis with loop diuretics

b) Glucocorticoids

c) Pamidronate

d) Hydration with saline

Question 15:

Which of the following is used in the treatment of hypercalcemia due to Vitamin D toxicity?

a) Dexamethasone

b) Hydroxychloroquine

c) Chloroquine

d) Ketoconazole

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

A patient is brought to the emergency department with confusion, drowsiness, and multiple episodes of vomiting. His family members say that he was diagnosed with bipolar disorder several years ago and has been on medication ever since. Further evaluation reveals a serum calcium level of 16 mg/dl. What should be the next step in the management of this patient?

a) Saline hydration

b) Denosumab

c) IV biphosphonate

d) Subcutaneous calcitonin

Question 17:

Sold by @itachibot

In a patient with hypocalcemia and hyperphosphatemia, which of the following will you not suspect?

a) Vitamin D-dependent rickets type 1

b) Pseudohypoparathyroidism

c) Chronic kidney disease

d) DiGeorge syndrome

Question 18:

All of the following medical conditions are associated with hyperphosphatemia except:

a) Chronic renal failure

b) Prolonged phosphate intake

c) Pseudopseudohypoparathyroidism

d) Pseudohypoparathyroidism

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

A 5-year-old boy is brought by his mother with with the following abnormality. Serum levels of vitamin D are found to be low. Which of the following would not be seen in this boy?

a) Decrease in parathyroid hormone level

b) Decrease in renal calcium excretion

c) Increase in renal phosphate excretion

Sold by @itachibot

d) Increase in serum alkaline phosphatase

Question 20:

Which of the following statements is incorrect with respect to hypocalcemia?

a) Can occur due to hypermagnesemia

b) Latent tetany is observed

c) Prolonged QT interval

d) Not a feature of hyperparathyroidism

Question 21:

The following sign is elicited in a patient. What is it called?

a) Chvostek's sign

b) Trousseau's sign

c) Tinel's sign

d) Troisier's sign

Question 22:

Which of the following agents cannot be used to manage a patient with chronic hypocalcemia?

Sold by @itachibot

a) Etidronate

b) Thiazide

c) Elemental calcium

d) Vitamin D analogues

Answer Key

Question No.	Correct Option
1	d
2	c
3	c
4	b
5	d
6	b
7	a
8	a
9	d
10	c
11	b
12	c
13	c
14	b
15	a
16	a
17	a
18	c
19	a
20	d
21	b
22	a

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

Hypercalcemia is associated with all of the following conditions except hypothyroidism. Hyperthyroidism is however associated with hypercalcemia.

Other options:

1) Multiple myeloma is a hemotological malignancy associated with hypercalcemia.

2) Hyperparathyroidism is associated with sub-periosteal bone resorption resulting in hypercalcemia.

3) Sarcoidosis is associated with hypercalcemia due to increased activity of 1 alpha hydroxylase activity in the macrophages.

Solution to Question 2:

Vitamin A intoxication can cause hypercalcemia. This is due to high bone turnover. Ingestion of 50,000–100,000 units of Vitamin A daily can increase calcium levels up to 12–14 mg/dL.

Note: Vitamin D intoxication also leads to hypercalcemia.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 3:

This clinical scenario with severe hypercalcemia (> 14.8 mg/dl) is suggestive of a hypercalcemic crisis. Pancreatitis causes hypocalcemia, not hypercalcemia.

Hypercalcemic crisis is often caused by malignancies due to an acute rise in calcium levels following bone resorption due to metastases. It is a medical emergency that presents with polyuria, severe dehydration, gastrointestinal symptoms, and altered consciousness. It can lead to cardiac arrest and coma.

Solution to Question 4:

Hypercalcemia is associated with constipation and not diarrhea.

Clinical manifestations of hypercalcemia include:

The clinical signs associated with primary hyperparathyroidism can be remembered with the mnemonic 'stones, bones, abdominal groans, psychic moans'.

System	Clinical features
Renal	Renal stones (Calcium oxalat e > calcium phosphate)
Nephrocalcinosis	
Polyuriaand dehydration	
Renal failure	

Sold by @itachibot

System	Clinical features
Musculoskeletal	Bone pain, bone cysts
Myalgia and arthralgia	
Osteopenia, osteoporosis	
GIT	Constipation
Nausea,vomiting, anorexia	
Abdominal pain	
Peptic ulcer disease	
Acute pancreatitis	
Neuropsychiatric	Anxiety,depression, fatigue
Cognitive dysfunction	
Drowsiness, obtundation, co ma	
Cardiac	Short QT interval, bradycardi a
Hypertension	
Others	Ectopic calcification

Sold to you purchased this else, you may have been st d.

Solution to Question 5:

Excess calcium ingestion is not associated with restrictive cardiomyopathy. It results in milk-alkali syndrome, commonly associated with excessive intake of absorbable antacids such as milk or calcium carbonate.

In susceptible individuals, there is a loss of negative feedback, leading to excess calcium absorption when there is excess intake. It results in the following triad of clinical features:

• Hypercalcemia

• Alkalosis - due to HCO_3 reabsorption caused by PTH suppression, natriuresis, and water depletion.

• Renal failure - due to damage to renal tubules.

It is reversible once ingestion of antacid is stopped. However, in severe cases, a chronic form called Burnett's syndrome can develop. It is associated with irreversible renal damage.

Solution to Question 6:

Parathyroid adenomas are the most common cause of primary hyperparathyroidism.

Causes of primary hyperparathyroidism:

• Parathyroid adenomas: These maybe

Sold by @itachibot

- Solitary (more common)
- Multiple
- Part of MEN-1, MEN-2A or MEN-4
- Chief cell parathyroid hyperplasia
- Parathyroid gland carcinoma
- Lithium use
- Hyperparathyroidism-jaw tumor (HPT-JT) syndrome

Solution to Question 7:

Secondary hyperparathyroidism presents with decreased calcium and increased levels of PTH, phosphate, and ALP.

In this condition, calcium deficiency leads to reactive stimulation of the parathyroid. It is caused by the following:

- Chronic kidney disease
- Vitamin D deficiency
- Gastrointestinal malabsorption
- Liver disease

Solution to Question 8:

Primary hyperparathyroidism (PHPT) causes shortening of the QT interval.

The asymptomatic form is the most common presentation in the Western population. But in developing countries, the patients are usually symptomatic. The classical symptoms and signs of PHPT are due to hypercalcemia as well as increased parathyroid hormone (PTH) levels.

The features of hypercalcemia are as follows:

The features of elevated PTH levels are as follows:

- Osteitis fibrosa cystica
- Osteopenia/osteoporosis
- Hyperuricemia and gout
- Pseudogout
- Increased production of calcitriol
- Nephrolithiasis
- Hypophosphatemia
- Anemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

The clinical signs associated with primary hyperparathyroidism can be remembered with the mnemonic - “bones, stones, groans, moans with fatigue overtones”

System	Clinical features
Musculoskeletal	Bone painSkeletal deformitiesPathological fracturesMyalgia and arthralgia
Renal	Renal stones(Calcium oxalate > calcium phosphate)Nephrocalcinosis Polyuria and dehydrationRenal failure
GIT	ConstipationNausea, vomiting, anorexiaAbdominal painPeptic ulcer diseaseAcute pancreatitis
Neuropsychiatric	Anxiety, depressionCognitive dysfunctionIrritabilityHallucinations
Cardiac	Short QT intervalbradycardia Hypertension
Ectopic calcification	Cornea: Band keratopathyExtravascular tissues: calcinosis Small arteries: calciphylaxis

If you purchased this, your may have been stolen else, and is illegally being sold.

Solution to Question 9:

Neuropsychiatric changes are seen in patients with primary hyperparathyroidism (PHPT).

The clinical signs associated with primary hyperparathyroidism can be remembered with the mnemonic - “bones, stones, groans, moans with fatigue overtones”

Note: Though multiple clinical studies have found an increased incidence of gall stones in primary HPT, there is no reference to this in Harrison's and Williams's endocrinology. Hence, option D would be the most appropriate answer.

System	Clinical features
Musculoskeletal	Bone painSkeletal deformitiesPathological fracturesMyalgia and arthralgia
Renal	Renal stones(Calcium oxalate > calcium phosphate)Nephrocalcinosis Polyuria and dehydrationRenal failure

Sold by @itachibot

System	Clinical features
GIT	ConstipationNausea, vomiting, anorexiaAbdominal painPeptic ulcer diseaseAcute pancreatitis
Neuropsychiatric	Anxiety, depressionCognitive dysfunctionIrritabilityHallucinations
Cardiac	Short QT intervalbradycardia Hypertension
Ectopic calcification	Cornea: Band keratopathyExtravascular tissues: calcinosis Small arteries: calciphylaxis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

Calcitonin levels are increased in hyperparathyroidism.

Due to elevated parathyroid hormone (PTH) levels in hyperparathyroidism, there is osteoclast activation. This leads to increased calcium levels in plasma. As a response to this, there is an increase in the level of calcitonin to regulate the high calcium levels.

Calcitonin is secreted by the parafollicular C-cells of the thyroid gland. Its principal effects are to lower serum calcium and phosphate by its action on bone and kidney. In bone, calcitonin inhibits osteoclasts and reduces bone resorption. In the kidney, it inhibits the reabsorption of both calcium and phosphate.

Other options

Option A: In hyperthyroidism, the direct effect of thyroid hormones can lead to bone resorption and mild hypercalcemia. However, hyperparathyroidism is a better answer.

Options B and D: Hypoparathyroidism and Cushing's syndrome will have low serum calcium and hence, will not have increased calcitonin levels.

Solution to Question 11:

The above x-ray shows a rugger jersey spine which is a distinct feature of hyperparathyroidism. It is associated with elevated ALP.

It refers to the prominent hyperdensities at the endplates of successive vertebrae, which mimics the striped appearance of a rugby jersey.

Option A: Paget's disease causes cortical thickening, which results in increased opacity on all sides of the vertebral body. These are called picture frame vertebrae.

Option C: Osteopetrosis presents with normal ALP levels. It results in sclerotic endplates of vertebrae that are termed sandwich vertebrae. Though this finding is similar to rugger jersey

Sold by @itachibot

spine, it is denser and more sharply defined than rugger jersey spine. In addition, there is sclerosis of inner margins of vertebral bodies, resulting in a bone within a bone appearance.

Option D: Ankylosing spondylitis produces a bamboo spine appearance due to fusion of vertebral bodies.

Solution to Question 12:

This clinical scenario is suggestive of a pathological fracture of the tibia. The given x-ray shows brown tumors or osteitis fibrosa cystica, which are characteristic of hyperparathyroidism.

Solution to Question 13:

The arrows in this x-ray point to arrow-headed fingers or the spade phalanx sign. It is pathognomonic of acromegaly.

Acromegaly is the result of excessive growth hormone (GH) production, most commonly from a macroadenoma of the pituitary. There is osseous enlargement, presenting with spade-like hands and widening of the terminal phalangeal tufts, giving an arrow-head appearance.

Other features in the hand include:

• Widened joint spaces (due to cartilage hypertrophy)

• Generalized osteoporosis and cystic changes in the carpal bones

Option A: Hyperparathyroidism causes subperiosteal bone resorption, which is marked along the radial aspect of the middle phalanges of the index and middle fingers.

Option B: Down's syndrome can cause hypoplasia of the middle phalanx of the 5th finger.

Option C: Psoriasis causes a pencil-in-cup appearance of the distal phalanx.

Solution to Question 14:

Glucocorticoids have no effect on the serum calcium concentration in primary hyperparathyroidism and normal individuals.

They are effective in the treatment of hypercalcemia when associated with the following conditions:

• Malignancy

• Vitamin D intoxication

• Sarcoidosis

Initial medical management of hypercalcemia involves the following:

• Normal saline (200-500 ml/hour): It is the mainstay treatment to correct dehydration, and titrated to maintain a urine output of >100 ml/hour

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Pamidronate or zoledronate: These are bisphosphonates and are considered as the drug of choice in primary hyperparathyroidism

• Cinacalcet: This is a calcimimetic that increases the affinity of the glands to extracellular calcium, thereby decreasing parathyroid hormone secretion

• Denosumab: This inhibits RANK-ligand, which acts on osteoclasts

• Calcitonin: This is an activated vitamin D analog that may be used to reduce calcium levels. It should be used within the first 24 hours

Forced diuresis refers to aggressive fluid hydration with loop diuretic administration. Although this is an effective method to reduce calcium levels, it requires careful monitoring for adverse effects like fluid overload, hypokalemia, and hypomagnesia. Due to this reason as well as the availability of more effective treatment modalities, forced diuresis is not preferred any longer.

The definitive management of primary hyperparathyroidism is the surgical removal of the parathyroid gland.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 15:

In patients with hypercalcemia due to vitamin D toxicity, glucocorticoids like dexamethasone are the preferred therapy as they decrease 1,25 (OH2) vitamin D (calcitriol) production. This results in reduced intestinal calcium absorption and increased urinary excretion of calcium.

Hydroxychloroquine, chloroquine, and ketoconazole may also decrease calcitriol production and are used occasionally. But the preferred therapy is steroids.

Vitamin D toxicity presents acutely with features of hypercalcemia such as confusion, polyuria, polydipsia, anorexia, vomiting, and muscle weakness. Chronic intoxication may cause nephrocalcinosis, bone demineralization, and pain.

Lab features include elevated 25 (OH) vitamin D, high calcium, and suppressed PTH.

Dexamethasone can also be used in the treatment of hypercalcemia caused by the humoral hypercalcemia of malignancy (due to PTHr) and sarcoidosis.

Solution to Question 16:

This clinical scenario is suggestive of a hypercalcemic crisis. The next best step in management is saline hydration. Typically 200-500 ml/hour fluids are given to correct dehydration and titrated to maintain a urine output of >100 ml/hour.

Hypercalcemic crisis is a medical emergency associated with serum calcium >14 mg/dl. It is often caused by conditions causing an acute rise in calcium levels such as malignancies or lithium overdose.

It presents with oliguria, severe dehydration, gastrointestinal symptoms, and altered consciousness. It can lead to cardiac arrest and coma.

Initial medical management of hypercalcemia involves the following:

Sold by @itachibot

• Fluids are the mainstay treatment

• Bisphosphonates

• Cinacalcet

• Denosumab

• Calcitonin

Solution to Question 17:

Vitamin D-dependent rickets type 1 is associated with hypocalcemia and hypophosphatemia

Hypocalcemia with hyperphosphatemia is observed in the following conditions:

• True hypoparathyroidism

• Hereditary hypoparathyroidism

• Acquired hypoparathyroidism – following surgery

• Hypomagnesemia

• Pseudohypoparathyroidism

• Chronic kidney disease

• DiGeorge syndrome

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 18:

Pseudopseudohypoparathyroidism is a type of pseudohypoparathyroidism type 1A associated with normal calcium and normal phosphate. The clinical features are similar to pseudohypoparathyroidism but there is no PTH resistance.

Pseudohypoparathyroidism is an autosomal dominant condition caused by a mutation of the alpha subunit of the Gs protein. This causes PTH resistance. It is associated with Albright hereditary osteodystrophy (short stature, round facies, and short fourth metacarpal).

Types of pseudohypoparathyroidism:

PHP: pseudohypoparathyroidism, PPHP: pseudopseudohypoparathyroidism, PTH: parathyroid hormone

Type	PHP 1A	PHP 1B	PHP 1C	PHP 2	PPHP
Hypocalcemia	Yes	Yes	Yes	Yes	No
Hyperphosphatemia	Yes	Yes	Yes	Yes	No
PTH levels	↑	↑	↑	↑	Normal
GS α subunit deficiency	Yes	No	No	No	Yes

Sold by @itachibot

Type	PHP 1A	PHP 1B	PHP 1C	PHP 2	PPHP
Albright hereditary osteodystrophy	Yes	No	Yes	No	Yes
PTH resistance	Yes	Yes	Yes	Partial	No

Solution to Question 19:

The given image shows a windswept deformity of the lower limbs (combination of valgus deformity of one leg with varus deformity of the other leg). This finding along with low serum vitamin D points to a diagnosis of vitamin D deficiency rickets. It is associated with an increase in parathyroid hormone (PTH) levels.



Solution to Question 20:

Hypocalcemia is a feature of secondary hyperparathyroidism (SHPT).

Hypocalcemia refers to total serum calcium ≤8.4 mg/dl. The causes include:

- Low PTH
- Hypoparathyroidism
- High PTH (SHPT)
- Chronic kidney disease
- Vitamin D deficiency
- Pseudohypoparathyroidism

Sold by @itachibot

• Hyperphosphatemia

• Loop diuretics, calcitonin, bisphosphonates

• Massive blood transfusions

• Hyperventilation

• Hypomagnesemia - Mg is essential for PTH release

• Severe hypermagnesemia - higher Mg levels suppress PTH secretion

Hypocalcemia causes tetany or increased neuromuscular excitability. It can cause paresthesias, cramps, spasms, and seizures. Latent tetany can only be detected by performing Chvostek and Trousseau's signs. A prolonged QT interval is seen on ECG.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 21:

The above image indicates Trousseau's sign or carpopedal spasm classically seen in patients with hypocalcemia.

Trousseau's sign is elicited by making the forearm ischemic by applying a sphygmomanometer cuff to the arm and keeping the pressure above systolic blood pressure (SBP) for 5 minutes. The hand goes into a painful carpopedal spasm and assumes the accoucheur's hand position.

Hypocalcemia results in tetany or increased neuromuscular irritability. It leads to paresthesia and tingling of extremities and perioral area. Severe hypocalcemia can cause seizures.

Option A: Chvostek's sign is also seen in hypocalcemia. Tapping the facial nerve in front of the tragus will cause spasms of the facial muscles.

Option C: Tinel's sign is seen in carpal tunnel syndrome. It refers to a tingling sensation on tapping over the median nerve.

Option D: Troisier's sign refers to left supraclavicular lymphadenopathy in metastatic abdominal malignancy.

Solution to Question 22:

Etidronate is not used in the management of hypocalcemia. It is a bisphosphonate used in the management of hypercalcemia.

The goal in the management of hypoparathyroidism is to restore the calcium-phosphate balance. The mainstay is calcium and vitamin D supplements. Other underlying conditions must also be treated.

Instead of loop diuretics, thiazides can be administered on a low sodium diet as they enhance urinary calcium reabsorption.

Sold by @itachibot

Obesity

Question 1:

The hormone leptin is derived from which of the following?

a) Arcuate nucleus

b) Adipocytes

c) Chief cells

d) Paneth cells

Question 2:

A 25-year-old woman has been fasting all day as part of a new diet. Which of the following is a gut peptide that will be increased in this woman?

a) Peptide YY

b) Ghrelin

c) Cholecystokinin

d) Orexin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 16-year-old boy ate two pieces of cake, a sandwich, and two scoops of ice cream at his friend's birthday party an hour ago. Levels of which of the following are likely to be high in this boy?

a) Neuropeptide Y(NPY)

b) Melanin-concentrating hormone(MCH)

c) α- Melanocyte-stimulating hormone(α-MSH)

d) Agouti-related peptide (AgRP)

Question 4:

Which of the following is a key transcription factor involved in the differentiation of mesenchymal preadipocytes to adipose cells?

Sold by @itachibot

a) RARγ

b) Rev-erbAβ

c) RXRα

d) PPARγ

Question 5:

Which of the following statements is incorrect regarding adiponectin?

a) Its levels are increased in obesity

b) It increases insulin sensitivity

c) Promotes lipid oxidation

d) Has vascular protective effects

Question 6:

A 2-year-old girl is referred to an endocrinologist with severe obesity and insatiable hunger despite adequate nutrition. Due to the early-onset severe obesity, genetic testing was done and revealed inactivating mutations in the leptin gene. Which of the following neuroendocrine abnormalities is this child most likely to develop?

a) Central hypothyroidism

b) Growth retardation

c) Hypogonadotropic hypogonadism

d) Diabetes mellitus

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 7:

Genetic mutations causing pro-opiomelanocortin (POMC) deficiency presents with all of the following except:

a) α-MSH deficiency

b) Cachexia

c) Red hair pigmentation

d) Adrenal insufficiency

Question 8:

Sold by @itachibot

The production of which of the following hormones is increased in obesity?

a) Insulin

b) Thyroxine

c) Growth hormone

d) Adiponectin

Question 9:

In a child with obesity, hypogonadism, and mental retardation, which of the following would you not suspect ?

a) Prader-Willi syndrome

b) Laurence-Moon-Bardet-Biedl syndrome

c) Carpenter syndrome

d) Angelman syndrome

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Obesity is not seen in patients with which of the following conditions?

a) Craniopharyngioma

b) Pickwickian syndrome

c) Prader-Willi syndrome

d) Sipple syndrome

Question 11:

A 40-year-old woman is being evaluated for high blood glucose levels. On examination, acanthosis nigricans is noted, and her BMI is 34 kg/m^2. How would you classify her weight?

a) Overweight

b) Class I obesity

c) Class II obesity

d) Class III obesity

Question 12:

Sold by @itachibot

A known patient of type 2 diabetes mellitus on oral medication presents to your OPD for review. His blood pressure is 150/90 mm Hg and his BMI is 28.4 kg/m². His fasting glucose is 118 mg/dl and postprandial glucose is 154 mg/dl. He consumes a low-calorie diet and exercises regularly. How will you manage his obesity?

a) Start him on a more intense diet and exercise regimen

b) Anti-obesity drugs in combination with proper diet and exercise

c) Refer for bariatric surgery

d) Anti-obesity drugs alone as lifestyle changes have not been effective

Question 13:

Which of the following anti-obesity drugs is a selective 5-HT2C receptor agonist?

a) Orlistat

b) Phentermine

c) Lorcaserin

d) Topiramate

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	d
5	a
6	c
7	b
8	a
9	d
10	d
11	b
12	b
13	c

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Leptin is an adipocyte-derived hormone.

It acts through brain circuits (predominantly in the hypothalamus) to influence appetite, energy expenditure, and neuroendocrine function. High leptin levels reduce appetite and increase energy expenditure.

The arcuate nucleus in the hypothalamus plays an important role in the regulation of appetite.

Solution to Question 2:

In the fasting state, high levels of ghrelin are seen. It is a gut-peptide produced in the stomach and stimulates hunger and appetite.

Solution to Question 3:

This boy is likely to have high levels of α-Melanocyte-stimulating hormone (α-MSH). It is a hypothalamic peptide that decreases appetite and hence would be increased in the fed state.

Solution to Question 4:

One of the key transcription factors involved in the differentiation of mesenchymal preadipocytes to adipose cells is the peroxisome proliferator-activated receptor γ (PPARγ). It is a nuclear receptor that also binds the thiazolidinedione class of insulin-sensitizing drugs used in the treatment of type 2 diabetes mellitus.

Note: Saroglitazar has a dual action on PPARα and PPAR■ receptors. It reduces blood sugar and lipid levels in diabetic patients.

Solution to Question 5:

Adiponectin levels are reduced in obesity.

It is an abundant adipose-derived protein with the following characteristics:

• Its levels are reduced in obesity

• It enhances insulin sensitivity and lipid oxidation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• Has vascular-protective effects

On the other hand, resistin and RBP4 levels, which are also secreted by adipocytes, may induce insulin resistance.

Solution to Question 6:

The most prominent neuroendocrine abnormality seen in obese individuals with inactivating mutations in the leptin (ob gene) or leptin receptor (db gene) is hypogonadotropic hypogonadism.

Leptin mutations are responsible for a rare form of obesity that runs in families. These mutations prevent leptin from delivering satiety signals. Hence, the brain perceives starvation despite adequate nutrition. It results in morbid, early-onset obesity.

Mutations in the following genes can cause obesity in humans:

• Lep (ob)

• LepR (db)

• POMC

• PC-1

• MC4R

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 7:

Cachexia is not a feature of mutations of the pro-opiomelanocortin (POMC) gene. It is associated with the following characteristics:

• Absence of α-MSH

• Severe obesity - α-MSH inhibits appetite in the hypothalamus

• Pale skin and red hair

• Absence of ACTH

• Secondary adrenal insufficiency

Solution to Question 8:

Insulin levels are increased in obesity. It stimulates the synthesis of fatty acids and TAG and decreases the rate of lipolysis. Thyroxine, growth hormone, and adiponectin have lipolytic effects, hence their levels are reduced in obesity.

Hyperinsulinemia and insulin resistance are consistent features of obesity. Insulin resistance is more strongly linked to intra-abdominal fat than to fat in other areas.

Sold by @itachibot

Adiponectin (Option D) is an anti-inflammatory cytokine that is produced by adipocytes. It performs vital functions that are reduced in obesity - like, increasing insulin sensitivity and the inhibition of gluconeogenic enzymes. It also enhances fatty acid oxidation.

Insulin resistance in obesity occurs as a result of the following:

• Insulin receptor downregulation

• Increased FFA (free fatty acids) and capable of impairing insulin action

• Intracellular lipid accumulation

• Circulating peptides produced by adipocytes such as TNF-α, IL-6, RBP4, and resistin modify insulin action.

If yo... Sold by @itachibot ...rchased this from someone else, ...have been scammed.

Solution to Question 9:

Children with Angelman syndrome present with mental retardation but do not have obesity and hypogonadism. It occurs due to the deletion of band q12 of the maternal chromosome 15.

The following syndromes present with obesity, hypogonadism, and mental retardation

• Prader-Willi syndrome - deletion of band q12 of the paternal chromosome 15.

• Laurence-Moon-Bardet-Biedl syndrome

• Carpenter syndrome

• Cohen syndrome

The following image shows a child with Prader-Willi syndrome.

Solution to Question 10:

Sipple syndrome is characterized by pheochromocytoma which is associated with weight loss.

Sold by @itachibot

Obesity is seen in the following syndromes:

• Cushing's syndrome

• Prader-Willi syndrome

• Pickwickian syndrome

• Hypothyroidism

• Insulinoma

• Craniopharyngioma and other hypothalamic disorders

• Laurence-Moon-Bardet-Biedl syndrome

• Carpenter syndrome

• Cohen syndrome

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

According to the WHO classification, a BMI of 34 kg/m^2 would be classified as class I obesity.

The revised BMI classification for Asians/Indians is as follows:

• Underweight - BMI < 18.5 kg/m2

• Normal or lean - BMI 18.5–22.9 kg/m2

• Overweight - BMI 23.0 –24.9 kg/m2

• Obese BMI ≥25 kg/m2

Note: As the question does not specify Asian/Indian, the answer is given according to WHO classification.

Classification	BMI (kg/m^2)	Risk
Underweight	<18.5	-
Normal weight	18.5–24.9	-
Overweight	25–29.9	Increased
Obesity (Class I)	30–34.9	High
Obesity (Class II)	35–39.9	Very high
Extreme obesity(Class III)	≥ 40	Extremely high

Solution to Question 12:

Sold by @itachibot

This obese man with a BMI of 28.4 kg/m² along with comorbidities must be managed with anti-obesity drugs as an adjunct to comprehensive lifestyle modifications.

The recommendations for management of obesity are as follows:

• Diet, exercise, behavioral therapy:

• BMI - 25-29.9 with comorbidities

• BMI ≥ 30

• Adjunctive pharmacotherapy:

• BMI - 27-29.9 with comorbidities

• BMI ≥ 30

• Bariatric surgery:

• BMI - 35-39.9 with comorbidities

• BMI ≥ 40

If you purchased this from someone else, you may have been scammed. Sold by @Itachibot

Solution to Question 13:

Lorcaserin is a selective 5-HT2C receptor agonist with a functional selectivity ~15 times that of 5-HT2A receptors and 100 times that of 5-HT2B receptors.

Sold by @itachibot

Hyperbilirubinemias and Tests of Liver Function

Question 1:

A 4-month-old child was brought with a history of recurrent episodes of jaundice. Blood investigations were suggestive of hereditary spherocytosis. Which of the following liver function test findings is likely to be seen here?

a) Direct bilirubin >50% of total bilirubin

b) Direct bilirubin >85% of total bilirubin

c) Direct bilirubin >15% of total bilirubin

d) Direct bilirubin <15% of total bilirubin

Question 2:

A 2-week-old baby presented with yellowish discoloration of the skin. Investigations revealed markedly elevated direct bilirubin levels. Which of the following conditions should be suspected in the newborn?

a) Biliary atresia

b) Physiological jaundice

c) Breast milk jaundice

d) Hemolytic disease of the newborn

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

Investigations performed in a patient with recurrent jaundice gave the following results. What is the possible cause of jaundice in this patient?

a) Hemolytic anaemia

b) Gilbert syndrome

c) Criggler-Najjar syndrome

d) Dubin Johnson syndrome

Question 4:

Sold by @itachibot

Which of the following is a cause of unconjugated hyperbilirubinemia?

a) Probenecid

b) Amoxicillin-clavulanate

c) Acute hepatitis

d) Stone in the common bile duct

Question 5:

A 24-year-old woman presents with jaundice. She had been on OCPs for the past month. Her LFT and liver biopsy findings are given below. What is the probable diagnosis?



a) Gilbert syndrome

b) Crigler- Najjar syndrome

c) Rotor syndrome

d) Dubin Johnson syndrome

Question 6:

What is the defect seen in patients with Rotor syndrome?

a) Absent UDP glucuronosyl transferase

b) Mutation in organic anion transport protein

c) Mutation in multidrug resistance–associated protein 2

d) Reduced UDP glucuronosyl transferase

Sold by @itachibot

Question 7:

While evaluating a patient with jaundice, all of the following investigations would help you to differentiate between Dubin Johnson syndrome and Rotor syndrome except:

a) Oral cholecystography

b) Total urinary coproporphyrin

c) Serum bilirubin levels

d) Bromsulphalein clearance test

Question 8:

In a patient with unconjugated hyperbilirubinemia, which feature helps confirm the diagnosis of Crigler- Najjar syndrome?

a) Presence of kernicterus

b) Absence of hemolysis

c) UGT1A1 mutation

d) Autosomal recessive inheritance

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

An infant presented with yellow discolouration of the skin, poor feeding, and seizures. Investigations revealed total bilirubin- 40 mg/dL and unconjugated bilirubin- 34 mg/dL. Phenobarbitone was administered while blood was being arranged for an exchange transfusion. However, there was no improvement. What is the most probable diagnosis?

a) Gilbert syndrome

b) Crigler-Najjar syndrome type 1

c) Crigler-Najjar syndrome type 2

d) Rotor syndrome

Question 10:

An 8-year-old child presents with jaundice and easy fatiguability. He had a history of ear infection which was treated with amoxicillin-clavulanic acid. Investigations done are shown below. What is the most likely diagnosis?

Sold by @itachibot

a) Drug induced hepatitis

b) Hemolytic jaundice

c) Obstructive jaundice

d) Gilbert syndrome

Question 11:

Which of the following conditions is associated with increased activity of 5'- nucleotidase?

a) Bone diseases

b) Prostate cancer

c) Chronic renal failure

d) Cholestatic disorders

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Which among the following is the most specific indicator of liver cell injury?

a) Alkaline phosphatase

b) Gamma glutamyl transpeptidase

c) Alanine aminotransferase

d) Aspartate aminotransferase

Question 13:

In a patient with acute viral hepatitis, what level of serum transaminases will you expect?

a) 10-40 IU/L

b) <500 IU/L

c) 40-300 IU/L

d) >1000 IU/L

Question 14:

A 64-year-old man with jaundice and abdominal pain was found to have an ALP level of 200 mg/dL. His LFT reports from a month ago had shown an ALP level of 50 mg/dL. What is your inference?

Sold by @itachibot

a) Normal finding for age

b) Could be due to fatty meal intake

c) Indicative of hemolysis

d) Suggestive of cholestatic liver disease

Question 15:

Which of the following statements regarding liver enzymes is true?

a) ALT is raised to >1000 IU/L in cirrhosis

b) Absolute levels of aminotransferases correlate with the prognosis

c) AST:ALT ratio <1 is seen in nonalcoholic fatty liver disease

d) AST:ALT ratio >1 is seen in chronic viral hepatitis

Question 16:

Liver function tests of a patient show isolated elevation of gamma-glutamyltransferase with other liver enzymes within normal limits. All of the following drugs could be responsible for this finding except:

a) Barbiturates

b) Griseofulvin

c) Penicillin

d) Phenytoin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

Which of the following statements regarding prothrombin time is false?

a) Indicates prognosis in acute liver disease

b) Indicates prognosis in chronic liver disease

c) Measures factors II, V, VIII, X

d) Measures factors II, V, VII, X

Question 18:

Which among the following is a marker of chronic liver disease?

Sold by @itachibot

a) Prothrombin

b) Albumin

c) 5'- nucleosidase

d) Alkaline phosphatase (ALP)

Question 19:

Which of the following can be seen in a patient with chronic liver disease?

a) Hypergammaglobulinemia

b) Hypogammaglobulinemia

c) Rise in beta globulin levels

d) Rise in alpha globulin levels

Question 20:

Which of the following is not a marker for liver fibrosis?

a) Procollagen peptide III (PIIINP)

b) Hyaluronic acid

c) Tissue inhibitor of metalloproteinase 1(TIMP1)

d) Telopeptide

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

In which of the following patients planned for a liver biopsy, would you not be required to use the transjugular approach?

a) A patient with prolonged INR

b) A patient with significant ascites

c) A patient with splenomegaly

d) A morbidly obese patient

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	d
2	a
3	d
4	a
5	d
6	b
7	c
8	a
9	b
10	b
11	d
12	c
13	d
14	d
15	c
16	c
17	c
18	b
19	a
20	d
21	c

Detailed Explanations

Solution to Question 1:

The given clinical scenario describes recurrent hemolysis in a patient with hereditary spherocytosis leading to jaundice. In hemolytic conditions, only the unconjugated/ indirect bilirubin is raised and conjugated/direct bilirubin remains normal i.e. <15% of total bilirubin.

Bilirubin is synthesized from heme which is released primarily from the breakdown of old RBCs. This bilirubin is in its unconjugated form. It enters the circulation and reaches the liver. It is converted to conjugated bilirubin which is then secreted into the biliary tree.

Therefore, in a healthy person, most of the circulating bilirubin is unconjugated/indirect bilirubin. Hence, the normal liver function tests show indirect bilirubin >85% and direct/conjugated bilirubin <15% of the total bilirubin. The normal bilirubin values are as follows:

• Total bilirubin levels - 1 to 1.5 mg/dL

Sold by @itachibot

• Direct bilirubin - <0.3 mg/dL

Solution to Question 2:

The raised direct bilirubin levels indicate conjugated hyperbilirubinemia. Among the given options, biliary atresia causes conjugated hyperbilirubinemia.

Biliary atresia is a disorder characterised by destruction or absence of bile ducts in a newborn. It leads to obstruction to the flow of bile thereby causing leaking of conjugated bilirubin into the circulation.

Option B: Physiological jaundice of the newborn occurs due to immaturity of the newborn's liver. It is not able to conjugate the excess bilirubin produced after birth. It leads to unconjugated hyperbilirubinemia.

Option C: Breast milk may contain certain factors that inhibit the conjugation of bilirubin leading to unconjugated hyperbilirubinemia.

Option D: Hemolytic disease of the newborn causes excess production of bilirubin due to hemolysis leading to unconjugated hyperbilirubinemia.

Solution to Question 3:

The presence of urinary bilirubin and increased total bilirubin indicates conjugated hyperbilirubinemia. Among the given options, Dubin Johnson syndrome causes conjugated hyperbilirubinemia.

Only conjugated bilirubin can appear in the urine because it is water-soluble and filterable by the kidney. Unconjugated bilirubin is water-insoluble and bound to albumin, hence it cannot be filtered by the kidney.

Causes of unconjugated hyperbilirubinemia:

• Increased bilirubin production-

• Hemolytic anemias

• Hematomas

• Impaired bilirubin uptake and conjugation-

• Gilbert syndrome

• Crigler Najjar syndrome

• Drugs - Probenecid, Rifampin

Causes of conjugated hyperbilirubinemia:

• Hereditary cholestatic syndromes

• Dubin Johnson syndrome

• Rotor syndrome

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Biliary obstruction

Solution to Question 4:

Probenecid causes impaired hepatic uptake of bilirubin which leads to unconjugated hyperbilirubinemia.

Amoxicillin-clavulanate, acute hepatitis and stone in the common bile duct are causes that lead to mixed hyperbilirubinemia (both conjugated and unconjugated bilirubin elevated) or predominantly conjugated hyperbilirubinemias.

Levels of conjugated and unconjugated bilirubin are useful in determining pre-hepatic causes of jaundice but are not useful in differentiating between intrahepatic and post-hepatic causes of jaundice.

Solution to Question 5:

The given clinical scenario is suggestive of Dubin Johnson syndrome (DJS). The LFT describes conjugated hyperbilirubinemia with normal liver enzyme levels. The given image of microscopic examination of liver biopsy shows an accumulation of dark melanin-like pigment in the hepatocytes which is seen in this condition.

DJS is an autosomal recessive condition causing defective expression of MRP 2 protein which is important for the biliary excretion of conjugated bilirubin. The patient has icterus but is otherwise asymptomatic.

DJS also impairs biliary excretion of various compounds like epinephrine metabolites. The dark melanin-like pigment is derived from these metabolites causing the liver to appear black.

Rotor syndrome is also an autosomal recessive condition causing conjugated hyperbilirubinemia but it does not present with pigmentation of the liver.

Gilbert and Crigler-Najjar syndromes are characterized by unconjugated hyperbilirubinemia.

Solution to Question 6:

Rotor syndrome occurs due to mutation in OATP (organic anion transport protein) 1B1 and 1B3.

Rotor syndrome is an inherited autosomal recessive disorder with conjugated hyperbilirubinemia. It is a benign disorder with mild elevation of bilirubin levels. It is similar to Dubin Johnson syndrome but the liver does not show any black pigmentation. No treatment is required for these patients.

Solution to Question 7:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Serum bilirubin levels are not helpful in differentiating between Dubin Johnson syndrome (DJS) and Rotor's syndrome as both these conditions present with mild elevation of serum bilirubin.

These two conditions can be differentiated as follows:

Test	Rotor Syndrome	Dubin Johnson syndrome
Oral cholecystography	Gall bladder is visualized	Gall bladder not visualisedIm paired ability to concentrate t he contrast material into bile
Total urinary coproporphyrin	Elevated	NormalIncreased excretion o f coproporphyrin 1 and decre ased excretion of coproporph yrin 3
Bromsulphalein (BSP) cleara nce test	Delayed clearance of BSPNo reflux	Rise in plasma levels of BSP after 90 minutes of injection Reflux of BSP from hepatocyt es

Sold by @ita
If you purchased this from so
you may have been scammed.

Solution to Question 8:

Kernicterus is seen in Crigler-Najjar syndrome but not in Gilbert syndrome. Gilbert syndrome does not cause a significant elevation in unconjugated bilirubin to cause kernicterus.

Crigler-Najjar syndrome and Gilbert syndrome are both caused by various mutations in the UGT1A1 gene. This leads to decreased activity of bilirubin UDP-glucuronosyltransferase enzyme leading to unconjugated hyperbilirubinemia. Both are predominantly inherited in an autosomal recessive pattern.

These syndromes present with jaundice in infants. The unconjugated hyperbilirubinemia in these syndromes is not associated with any hemolysis. All the other liver function tests are usually normal.

Crigler Najjar syndrome may lead to kernicterus causing severe brain damage. Liver transplantation is required without which the infant may not survive beyond 1-2 years of age. Whereas, Gilbert syndrome is a benign asymptomatic condition that does not need any intervention.

Solution to Question 9:

The given clinical scenario with poor feeding and seizures is indicative of kernicterus. This accompanied with unconjugated hyperbilrubinemia is highly suggestive of Crigler-Najjar syndrome type 1.

No improvement in bilirubin levels following the administration of phenobarbital further supports the diagnosis.

Sold by @itachibot

Crigler-Najjar syndrome is of two types with type 1 having absent/trace enzyme activity and type 2 having <10% of the normal enzyme activity. Hence, type 1 is far more severe and leads to kernicterus and neurological damage. Type 2 may also lead to kernicterus occasionally.

Kernicterus occurs because the unconjugated bilirubin is lipid-soluble and hence crosses the blood-brain barrier.

Phenobarbital is an enzyme inducer. Its administration can be used to differentiate between type 1 and type 2 Crigler-Najjar.

• Enzyme induction by phenobarbital leads to improved conjugation of bilirubin leading to a reduction in total bilirubin level in type 2 Crigler-Najjar syndrome.

• No improvement is seen in type 1 as the enzyme activity is absent and therefore cannot be induced.

Inherited unconjugated hyperbilirubinemias (Autosomal Recessive) include:

Note: Rotor syndrome causes conjugated hyperbilirubinemia.

	Crigler-Najjar type I	Crigler-Najjar type II	Gilbert Syndrome
UDP-GT Activity	Absent	Markedly reduced	Reduced
Serum bilirubin	20 - 45 mg/dl	<20 mg/dl	Generally <3 mg/dl
Pheno-barbital effect on serum bilirubin	No effect	Reduction	Reduction
Outcome	Kernicterus	Usually benign, kernicterus is rare	Benign
Prevalence	Rare	Rare	Common

The normal serum bilirubin (between 1 and 1.5 mg/dL) present in serum represents a balance between input from the production of bilirubin and hepatic/biliary removal of the pigment.

Isolated elevation of the bilirubin:

• Conjugated hyperbilirubinemia- >15% conjugated bilirubin

• Unconjugated hyperbilirubinemia- <15% conjugated bilirubin

Solution to Question 10:

The given clinical scenario is suggestive of hemolytic jaundice. It can present with the following features:

• Low hemoglobin

• Unconjugated hyperbilirubinemia

• Pigment gall stones - excess bilirubin in bile

Sold by @itachibot

• Development of cholestasis when the gall stones obstruct the flow of bile. This would result in the elevation of conjugated bilirubin and LFTs show a cholestatic picture.

Hepatic and obstructive causes of jaundice are not likely as the other liver function tests are normal. The various causes of jaundice can be differentiated as follows:

Inherited hyperbilirubinemias like Gilbert syndrome are suspected when there is no associated hemolysis. It is unlikely in this patient because there are clear hemolytic features.

Jaundice	Pre hepatic	Hepatic	Post hepatic
Cause	Excessive amount of bilirubin is presented to the liver	Impaired cellular uptake/ defective conjugation/ defective secretion of bilirubin	Defective excretion due to mechanical obstruction to the flow of bile
LFT	Elevated unconjugated bilirubin	Elevated conjugated and unconjugated bilirubinMarked elevated levels of AST and ALT (>1000U/L)	Elevated conjugated bilirubin Elevated levels of ALP and GGT
Example	Hemolytic anaemia	Hepatitis	Choledocholithiasis

Sold by
if you purchased this fro
you may have been scam

Solution to Question 11:

5■-nucleotidase is usually elevated in cholestasis.

Enzymes elevated in cholestasis are :

• Alkaline phosphatase (ALP)

• 5■-nucleotidase

• γ-glutamyl transpeptidase (GGT)

Solution to Question 12:

Among the given options, alanine aminotransferase is the most specific indicator of liver cell injury.

Alanine aminotransferase (ALT) and aspartate aminotransferase (AST) are indicators of liver cell injury. This is because ALT is found primarily in the liver, whereas, AST is found in the liver, cardiac muscle, skeletal muscle, kidneys, brain, pancreas, etc.

GGT and ALP are enzymes that primarily reflect cholestasis.

Solution to Question 13:

Sold by @itachibot

In a patient with acute viral hepatitis, serum transaminases are elevated to >1000 IU/L

The normal levels of serum transaminases are 10-40 IU/L. Marked elevation of serum transaminases >1000 IU/L is seen in conditions with extensive hepatocellular injury. These include:

• Viral hepatitis

• Ischemic injury to the liver in conditions like acute heart failure

• Toxin/drug-induced liver injury

Minor elevations of serum transaminase levels up to 300 IU/L are non-specific and can be found in any liver disease.

Solution to Question 14:

A 4 fold rise of serum ALP (alkaline phosphatase) is suggestive of cholestatic liver disease. However, serum ALP levels cannot differentiate between extrahepatic and intrahepatic cholestasis.

The normal range for serum ALP is considered 40-125 IU/L. Serum alkaline phosphatase is found in the liver, bone, small intestines, and placenta. Hence, a mild increase in ALP, i.e., 1 - 1.5 times normal, can be seen in these conditions:

• Following a fatty meal intake - influx of intestinal alkaline phosphatase into the blood. Hence, the patient is advised to fast before the blood sample is collected for the test.

• In pregnancy - placental ALP

• In children - rapid bone growth

• In elderly people over 60 years of age

ALP can be raised even in conditions like metastatic bone disease, inflammatory bowel disease, Hodgkin's lymphoma, etc. Hence, it needs to be correlated with raised levels of gamma-glutamyltransferase (GGT) and 5' nucleotidase which are more specific for liver disease.

Solution to Question 15:

AST:ALT ratio <1 is seen in NAFLD (nonalcoholic fatty liver disease).

The AST:ALT ratio can be interpreted as follows:

Option A: Patients with cirrhosis have only mildly elevated or even normal levels of serum transaminases.

Option B: Absolute levels of aminotransferases are not used for predicting prognosis. This is because there is a poor correlation between the degree of liver cell damage and the levels of aminotransferases.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

AST: ALT	Condition
<1	Chronic viral hepatitis, Nonal coholic fatty liver disease
>1	Cirrhosis
>2:1	Suggestive of alcoholic liver d iseaseAlcohol intake is associ ated with the deficiency of py ridoxal phosphate which in t urn leads to decreased levels of ALT enzyme. So, AST>AL T.
>3:1	Highly suggestive of alcoholic liver disease.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

Among the given options, penicillin does not lead to elevated gamma-glutamyltransferase (GGT) levels.

GGT is a microsomal enzyme. Hence, drugs that are microsomal enzyme inducers can lead to an increase in GGT levels. However, other liver enzyme levels will be normal. Therefore, an isolated rise in GGT level does not indicate cholestatic liver disease.

Drugs that increase levels of GGT are:

- Barbiturates
- Carbamazepine
- Ethanol
- Griseofulvin
- Rifampicin
- Phenytoin

Solution to Question 17:

Prothrombin time (PT) can help in estimating the levels of factors II, V, VII, and X.

PT evaluates the extrinsic and the common pathway of coagulation. Factor VII is required in the extrinsic pathway and factors II, V, X are required in the common pathway.

Coagulation factors are produced by the liver (except factor VIII) and they have a short half-life. Hence, they indicate the synthetic function of the liver and are very useful in assessing acute hepatic dysfunction. Factor VIII is produced by vascular endothelial cells.

Sold by @itachibot

Prolonged PT that is not corrected by vitamin K administration indicates poor prognosis in both acute and chronic liver disease.

Solution to Question 18:

Serum albumin is a marker of chronic liver disease because it is synthesized by the hepatocytes and has a long half-life of 18–20 days. Serum albumin is not a good indicator of acute or mild hepatic dysfunction.

Hypoalbuminemia is more common in chronic liver disorders such as cirrhosis and usually reflects severe liver damage. In hepatitis, albumin levels <3 g/dL should raise the possibility of chronic liver disease.

Prothrombin time is used to assess acute parenchymal liver disease. 5' nucleosidase is elevated in cholestasis. Elevation of ALP (<3 fold) is not specific to liver diseases.

Solution to Question 19:

Chronic liver disease is associated with hypergammaglobulinemia. Chronic liver diseases with hypergammaglobulinemia include autoimmune hepatitis, primary biliary cirrhosis, and alcoholic liver disease.

The liver produces all globulins except gamma globulin which are produced by B cells. In chronic liver diseases like liver cirrhosis, there is impaired reticuloendothelial function in hepatic sinusoids. Hence the liver fails to clear the bacterial antigens. This causes an increased production of gamma globulin.

Solution to Question 20:

Telopeptide is not a marker for liver fibrosis. It is a marker for bone resorption.

Various non-invasive markers to detect hepatic fibrosis are given below:

Non-invasive test for fibrosis	Components tested
Fibrotest/FibroSure	α2-macroglobulinHaptoglobinApolipoprotein A-1GGTTotal bilirubin
Enhanced Liver Fibrosis (ELF) assay	Hyaluronic acidProcollagen peptide III(PIIINP)Tissue inhibitor of metalloproteinase(TIMP)
Transient elastography (FibroScan)	Ultrasound waves are used to measure liver stiffness

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Non-invasive test for fibrosis	Components tested
Fibrospect II	Hyaluronic acidAlpha 2 macroglobulinTissue inhibitor of metalloproteinase 1 (TIMP 1)

Solution to Question 21:

Liver biopsy can be performed in a patient with splenomegaly via the percutaneous approach.

When the percutaneous approach is not advisable, liver biopsy can be performed via the transjugular approach. In this approach, a cannula is inserted via the jugular vein which is then passed into hepatic veins through which the biopsy sample is taken. Therefore, any bleeding resulting from the procedure will drain into the hepatic veins.

Indications of transjugular liver biopsy:

- Percutaneous approach contraindicated
- Significant ascites
- Prolonged INR
- Coagulopathies
- Difficulty in accessing the liver via the percutaneous route
- Morbid obesity

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Alcoholic Liver Diseases and Non-Alcoholic Fatty Liver Disease

Question 1:

Which of the following is the most common cause of cirrhosis in India?

a) Non alcoholic fatty liver disease

b) Alcoholic liver disease

c) Hepatitis B infection

d) Hepatitis C infection

Question 2:

What is the minimum amount of ethanol a man must have consumed for him to be at increased risk for alcoholic liver disease?

a) 35 g/day

b) 20 g /day

c) 28g/day

d) 14 g/day

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 45-year-old chronic alcoholic presents with right upper quadrant abdominal pain and nausea. His ultrasound abdomen revealed grade 2 fatty liver. Which among the following findings is unlikely to be seen in his condition?

a) AST : ALT < 2

b) Prolonged prothrombin time

c) Low platelet counts

d) Elevated ALP

Question 4:

Which of the following is unlikely to be seen in patients with Zieve's syndrome?

Sold by @itachibot

a) Cholestatic jaundice

b) Hemolytic anaemia

c) Transient hyperlipidemia

d) Thrombocytosis

Question 5:

Which of the following is the most specific marker for alcoholism?

a) ALT

b) GGT

c) ALP

d) LDH

Question 6:

You are using a CAGE questionnaire to screen a patient for alcohol abuse. Which of these is not the correct expansion of this questionnaire?

a) C - drink to calm your nerves

b) A - criticism over drinking annoys you

c) G - feel guilty about drinking

d) E - had a drink as eye opener

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

What parameters are used in calculating discriminant function?

a) Prothrombin time and bilirubin

b) Bilirubin, ascites and hepatic encephalopathy

c) Prothrombin time, ascites and hepatic encephalopathy

d) Prothrombin, Bilirubin and creatinine

Question 8:

A patient with alcoholic hepatitis has just been admitted to your ward. Prior to starting treatment, the senior doctor asks you to calculate his discriminant function. What is the

Sold by @itachibot

cutoff value for determining severe disease?

a) 26

b) 36

c) 22

d) 32

Question 9:

Which among the following is true about alcoholic cirrhosis?

a) Hepatocyte hyperplasia

b) Macronodular cirrhosis

c) Mixed micro and macronodular cirrhosis

d) Absence of Mallory Denk bodies

Question 10:

Prednisolone therapy for alcoholic hepatitis can be started in a patient with all of the following except:

a) Glasgow score ≥9

b) Serum creatinine ≥1 mg/dL

c) Spontaneous hepatic encephalopathy

d) SOFA score >2

Question 11:

A 34-year-old man presents with yellowish discoloration of the skin, abdominal pain, and fever. He reports consuming one-quarter of whisky every day for the past 10 years. He is a known case of chronic kidney disease. His lab reports are given below. Which of the following is used in his treatment?

a) Nutritional support alone

b) Prednisolone

c) Betamethasone

d) Pentoxifylline

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 12:

Which of the following statements is not true about nonalcoholic fatty liver disease?

a) Results from fat accumulation in more than 5% of hepatocytes

b) Strongly associated with obesity and insulin resistance

c) Associated with reduced serum ferritin levels

d) Diagnosis of NASH requires presence of inflammation

Question 13:

A non-diabetic patient has been diagnosed with nonalcoholic steatohepatitis (NASH). Which of the following is the first-line therapy for it?

a) Vitamin A

b) Vitamin E

c) Betaine

d) Ursodeoxycholic acid

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	d
5	b
6	a
7	a
8	d
9	c
10	d
11	d
12	c
13	b

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Alcoholic liver disease is the most common cause of cirrhosis in India.

The most common cause of cirrhosis worldwide is chronic viral hepatitis - Hepatitis C and B. NASH is expected to overtake it in the near future.

Solution to Question 2:

The minimum amount of ethanol that leads to an increased risk of developing alcoholic liver disease is 28g/day. In women, the minimum is 14g/day.

The risk factors for alcoholic liver disease are:

• Quantity of alcohol consumption - The risk of developing alcoholic liver disease increases when alcohol consumption is >14 drinks/ week in men and >7drinks/week in women. A standard drink contains 14g of ethanol which is present in -

• 120 mL of wine

• 350ml of beer

• 45ml of spirits (gin, rum, whiskey, vodka, tequilla)

• Gender - Women have increased susceptibility to alcoholic liver disease

• Hepatitis C infection - Increases the severity and decreases survival in alcoholic liver disease

• Genetics - A variation in patatin-like phospholipase domain-containing protein 3 (PNPLA3) gene confers an increased risk of progressive liver disease

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Solution to Question 3:

The given clinical scenario is suggestive of alcoholic liver disease. It is usually characterized by an AST: ALT > 2. This is because there is a deficiency of pyridoxal 5'- phosphate that is a cofactor for the enzymatic activity of ALT. Therefore, ALT levels fail to rise and there is a reversal of the AST: ALT ratio.

Prothrombin time (PT) is prolonged due to the deficiency of coagulation factors and, usually, does not respond to the administration of parenteral vitamin K. Platelet counts are often reduced early in the disease, which is reflective of portal hypertension with hypersplenism.

Elevated ALP levels are seen due to enzymatic induction by alcohol and mild cholestasis.

Solution to Question 4:

Zieve's syndrome is not associated with thrombocytosis.

Sold by @itachibot

Zieve's syndrome is a rare syndrome occurring due to severe alcoholic hepatitis characterized by a triad of:

• Cholestatic jaundice

• Hemolytic anemia

• Transient hyperlipidemia

It resolves with abstinence from alcohol.

Solution to Question 5:

GGT (Gamma Glutamyl Transferase) is the most specific marker of alcoholism among the given options.

The markers with >60% sensitivity and specificity for heavy alcohol use are gamma-glutamyl transferase (>35 u) and carbohydrate-deficient transferrin (CDT), the combination of the two is likely to be more accurate than one. An isolated elevation in serum GGT or GGT elevation out of proportion to that of other enzymes (such as the ALT and ALP) may be an indicator of alcohol abuse or alcoholic liver disease.

ALT (option A) is usually elevated in acute hepatocellular conditions such as hepatitis. Alanine aminotransferase is predominantly found in the liver, hence a more specific marker of liver injury. A ratio of AST/ALT > 2:1 indicates the possibility of alcoholic liver disease and > 3:1 ratio is highly indicative of this condition. In alcoholic liver disease, ALT levels are often normal.

ALP (option C) is elevated in cholestasis, ■■is also elevated when there is rapid bone growth like in children and adolescents because of bone alkaline phosphatase and in normal pregnancies due to placental alkaline phosphatase.

LDH (option D) elevation is nonspecific.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 6:

C stands for feeling the need to cut down your drinking in the CAGE questionnaire.

The CAGE questionnaire is used for identifying alcohol abuse. It can be expanded as below:

• C - Have you ever felt you ought to cut down on your drinking?

• A - Have people annoyed you by criticizing your drinking?

• G - Have you ever felt guilty or bad about your drinking?

• E - Have you ever had a drink first thing in the morning to steady your nerves or get rid of a hangover (eye-opener)?

One 'yes' response raises a suspicion of an alcohol use problem and more than one is a strong indication of abuse or dependence.

Sold by @itachibot

Solution to Question 7:

Discriminant function (DF) is calculated using prothrombin time and bilirubin level. It is also known as the Maddrey score and it assesses the severity and mortality risk in alcoholic hepatitis.

DF = 4.6 X (the prolongation of the prothrombin time above control [seconds]) + serum bilirubin (mg/dL)

DF ≥ 32 indicates severe disease and is an indication for starting glucocorticoids.

Solution to Question 8:

Severe alcoholic hepatitis is defined as discriminant function (DF) ≥ 32.

Severe alcoholic hepatitis is defined as:

• Discriminant Function (DF) ≥ 32

• Model for End-stage Liver Disease (MELD) score ≥ 21

• Presence of hepatic encephalopathy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

Alcoholic cirrhosis is characterized by mixed micro-and macronodular. If a mixed pattern is not an option, the best choice is micronodular cirrhosis.

Cirrhosis is the late stage of alcoholic liver disease. It is characterised by extensive fibrosis. Alcoholic cirrhosis begins as micronodular cirrhosis. With the cessation of alcohol use, larger nodules may form, resulting in a mixed micronodular and macronodular pattern.

The various conditions in which cirrhosis is seen are listed below:

• Macronodular cirrhosis- Parenchymal nodules >3 mm

• Micronodular cirrhosis- Parenchymal nodules <3 mm

Macronodular cirrhosis	Micronodular cirrhosis
Chronic viral hepatitisα1 Anti trypsin deficiencyDrugs & tox insHereditary hemorrhagic te langiectasiaCryptogenic cirrh osisWilsons disease (Micro & Macro nodular >> Macronod ular)	AlcoholPrimary biliary cirrho sisLarge bile duct obstruction HemochromatosisIntestinal bypassCystic fibrosisIndian c hildhood cirrhosis

Solution to Question 10:

Sold by @itachibot

SOFA score >2 indicates sepsis and glucocorticoids are avoided in septic patients.

Glucocorticoids are indicated in severe alcoholic hepatitis which is defined as:

• Discriminant Function (DF) ≥ 32

• Model for End-stage Liver Disease (MELD) score ≥ 21

• Glasgow alcoholic hepatitis score ≥ 9

• Presence of hepatic encephalopathy

Prednisolone is given at a dose of 32 mg orally daily for 4 weeks and then slowly tapered off.

Glucocorticoid therapy is avoided in:

• Active GI bleeding requiring blood transfusions

• Active infection requiring antibiotic treatment

• Preexisting chronic renal disease with a serum creatinine level greater than 2 mg/dL

• Pancreatitis

• Obvious contraindications to glucocorticoid therapy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

The given clinical scenario is suggestive of severe alcoholic hepatitis. Since the patient is a known case of chronic kidney disease, pentoxifylline is used in his treatment.

Severe alcoholic hepatitis is defined by a discriminant function ≥ 32. In the given scenario,

DF = 4.6 X (the prolongation of the prothrombin time above control [seconds]) + serum bilirubin (mg/dL)

= 4.6 x (16-11) + 11 = 34.

Thus, the patient cannot be managed by nutritional support alone.

Pentoxifylline is a TNF alpha inhibitor used either by itself or along with glucocorticoids in the treatment of severe alcoholic hepatitis. It is preferred in patients with kidney diseases and conditions where glucocorticoids are contraindicated.

Glucocorticoid therapy is avoided in:

• Active GI bleeding requiring blood transfusions

• Active infection requiring antibiotic treatment

• Preexisting chronic renal disease with a serum creatinine level greater than 2 mg/dL

• Pancreatitis

• Obvious contraindications to glucocorticoid therapy

Solution to Question 12:

Sold by @itachibot

Nonalcoholic fatty liver disease (NAFLD) is associated with increased levels of serum ferritin.

Nonalcoholic fatty liver disease (NAFLD) is defined as hepatic steatosis in the absence of any secondary cause for fat accumulation. Histologically, this requires fat accumulation in more than 5% of hepatocytes and it can be divided into:

• Nonalcoholic fatty liver (NAFL)- steatosis without inflammation

• Nonalcoholic steatohepatitis (NASH) - steatosis with inflammation

NAFLD is strongly associated with obesity and insulin resistance. Its diagnosis requires demonstration of hepatic steatosis and exclusion of significant alcohol consumption and other causes of hepatic fat accumulation.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 13:

The first-line therapy for nondiabetic patients with nonalcoholic steatohepatitis(NASH) is vitamin E.

Management of NAFLD includes:

• Vitamin E - first-line therapy in non-diabetics

• Insulin sensitizers- first-line therapy in diabetics

• Immunizations for Hepatitis A and B

• Weight loss

• Abstinence from alcohol

Note: The FDA in 2024 has approved Resmetirom, a thyroid hormone receptor-beta (THR-beta) agonist for noncirrhotic NASH with moderate to advanced fibrosis.

Sold by @itachibot

Viral Hepatitis

Question 1:

Which among the following hepatitis viruses is a DNA virus?

a) Hepatitis A

b) Hepatitis B

c) Hepatitis C

d) Hepatitis D

Question 2:

Which of the following is the most common route of spread of hepatitis E virus?

a) Sexual transmission

b) Blood transfusion

c) Feco-oral

d) IV drug use

Question 3:

Which of the following statements is true regarding hepatitis E?

a) Can be fatal in pregnant women

b) Can cause chronic hepatitis in immunocompetent

c) Most common route of spread is vertical transmission

d) Has a carrier state

Question 4:

Which of the following is associated with the highest risk of developing fulminant hepatitis?

a) Hepatitis A

b) Hepatitis B

c) Hepatitis E

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) Hepatitis D

Question 5:

A pregnant woman comes to your clinic for her first antenatal checkup in her 3rd trimester. Laboratory workup reveals that she is HBsAg positive and HBeAg positive. What is the chance of vertical transmission of hepatitis B to the child?

a) 25-30%

b) 50%

c) 10-15%

d) >90 %

Question 6:

What is the first marker to rise in a patient with acute hepatitis B?

a) Anti-HBs

b) HBsAg

c) HBeAg

d) Anti-HBc

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which serological marker can be used to diagnose acute hepatitis B in the window period?

a) Anti HBs

b) HBsAg

c) HBeAg

d) Anti HBc

Question 8:

A 35-year-old male patient presented with right upper quadrant abdominal pain and jaundice. His lab reports are given below. What is the likely causative organism for his condition?

a) HBV surface mutant

Sold by @itachibot

b) HBV precore mutant

c) HBV escape mutatnt

d) Vaccine-derived HBV

Question 9:

The lab report of a 40-year-old man is shown below. What is the most likely cause for this serological picture?

a) Recovering from hepatitis B

b) Hepatitis B in the remote past

c) Immunization with hepatitis B vaccine

d) Window period of HBV infection

Sold by @itachibot
ed this from someone else,
heen scammed.

Question 10:

A patient with acute hepatitis B has recovered from the infection. Identify the serological marker marked in the image:



©MARROW

a) Anti-HBc Ab

b) Anti-HBs Ab

c) Anti-HBe Ab

d) HBsAg

Sold by @itachibot

Question 11:

An intern sustained a needle-prick injury while establishing an IV line on a patient in the casualty. The patient was subsequently found to be HBsAg positive. The presence of which of the following markers in the intern would reliably indicate an early infection?

a) IgM anti HBsAg

b) IgM anti HBc

c) HBsAg

d) HBeAg

Question 12:

During your casualty rotation, you accidentally injure yourself while suturing a patient. You had taken the full schedule of hepatitis B vaccination during your first year of medical school. Which of the following markers will you get measured to determine the need for HBIG administration?

a) Anti HBeAg

b) IgM Anti HBc

c) IgG Anti HBc

d) Anti HBsAg

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 13:

Which of the following markers can be used to differentiate between hepatitis D superinfection and coinfection?

a) IgM anti-HDV

b) IgM anti-HBc

c) HBV DNA

d) HBsAg

Question 14:

A male patient presents with complaints of jaundice, loss of appetite, and weight loss for the past several months. After a full laboratory workup, a liver biopsy was performed which appeared as shown in the given image. What is the most likely diagnosis?

Sold by @itachibot



a) Acute hepatitis A

b) Chronic hepatitis B

c) Chronic hepatitis C

d) Chronic hepatitis A

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 15:

Which of the following pairs of viral hepatitis and its management is incorrectly matched?

a) Acute viral hepatitis E - Supportive care

b) Acute viral hepatitis B - Antiviral therapy

c) Acute viral hepatitis A - Supportive care

d) Acute viral hepatitis C - Antiviral therapy

Question 16:

Which of the following is an indication for antiviral therapy in chronic hepatitis B infection?

a) IgG anti-HBc positive

b) HbeAg negative with HBV DNA <2000 IU/L

c) ALT > 2 times upper limit of normal

d) Normal Fibroscan report with ALT < 2 ULN

Sold by @itachibot

Question 17:

A 32-year-old man presented with complaints of fatigue for the past 1 month. His lab reports are given below. What is the most appropriate next step in the management of this patient?

a) Tenofovir

b) Lamivudine and IFN

c) Entecavir

d) Serial monitoring

Question 18:

Which of the following is recommended as first-line therapy in a patient with chronic hepatitis B?

a) Pegylated interferon

b) Telbivudine

c) Lamivudine

d) Entecavir

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

In which of the following populations should surveillance for hepatocellular carcinoma be done?

a) 1, 3, 4, 5

b) 2, 3, 4

c) 1, 2, 4, 5

d) 1, 2, 3

Question 20:

A 35-year-old male comes to the OPD with intermittent jaundice for the past 2 months. You suspect acute viral hepatitis. Which of the following will you not evaluate in this patient?

a) HBsAg

b) IgM anti HBc

c) Anti HCV

Sold by @itachibot

d) IgM anti HBe

Question 21:

A 40-year-old male chronic smoker presents to the AIIMS OPD with fever, fatigue, yellow-colored urine, and clay-colored stools. For the past few days, he has developed an aversion to cigarette smoking. On examination icterus was present. What investigations would you advise to rule out acute viral hepatitis? Liver function test results are given below:

a) HBsAg, IgM antiHBC, AntiHCV, AntiHEV

b) AntiHAV, HBsAg, IgM antiHBc, AntiHCV

c) HBsAg, IgM antiHBc, AntiHDV, AntiHCV, Anti HEV

d) AntiHAV, IgM antiHBc, AntiHCV, AntiHEV

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 22:

Which of the following groups of individuals need not be screened for Hepatitis C infection?

a) Patients with HIV infection

b) Patients with unexplained elevation of aminotransferase levels

c) Patients who have undergone long-term haemodialysis

d) Household contacts of patients with hepatitis C

Question 23:

Which of the following is unlikely to be seen as an extrahepatic manifestation in patients with hepatitis C infection?

a) Mixed cryoglobulinemia

b) Lichen planus

c) Porphyria cutanea tarda

d) Polyarteritis nodosa

Question 24:

A 35-year-old man who presented with weight loss and abdominal discomfort is diagnosed with chronic hepatitis C . All the following factors can determine the progression of liver

Sold by @itachibot

disease in this patient except:

a) Age

b) Duration of infection

c) HCV RNA level

d) Body mass index

Question 25:

A 47-year-old man presented with generalized weakness, fatigue, and fever. His lab investigations are given below. What is the preferred treatment regimen for this patient?

a) Velpatasvir and sofosbuvir

b) Ribavirin and lamivudine

c) Telaprevir and simeprevir

d) α-interferon and dasabuvir

Question 26:

Which of the following is not a non-structural protein 5A inhibitor used in the treatment of patients with hepatitis C infection?

a) Sofosbuvir

b) Daclatasvir

c) Ledipasvir

d) Velpatasvir

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	c
3	a
4	d
5	d

Sold by @itachibot

6	b
7	d
8	b
9	b
10	a
11	b
12	d
13	b
14	b
15	b
16	c
17	d
18	d
19	a
20	d
21	b
22	d
23	d
24	c
25	a
26	a

If you pu... you may ... else, ...d.

Detailed Explanations

Solution to Question 1:

Hepatitis B is a DNA virus.

The hepatotropic viruses include:

Hepatitis virus	Family	Genetic material
A	Picorna virus	RNA
B	Hepadna virus	DNA
C	Flavivirus	RNA
D	Deltavirus	RNA

Sold by @itachibot

Hepatitis virus	Family	Genetic material
E	Hepeviridae	RNA

Solution to Question 2:

The most common route of the spread of hepatitis E is feco-oral.

Hepatitis E virus is the most common cause of acute hepatitis in adults, especially in endemic countries like India. The most common cause of acute hepatitis in children is the hepatitis A virus.

Routes of transmission of hepatitis viruses:

Hepatitis	Vertical route	Needle	Blood transfusion	Sexual	Feco-oral
Hep B	HBeAg positive mother >90 %HBsAg with Anti-HBe mother 10-15%	6-30 %	1 in 2 lakh(Most common blood transfusion-related virus).	Variable	-
Hep C	5%	1.8-6 %	1 in 18 lakh	5%	-
Hep A	Nil	Nil	Rare	Rare	Most common
Hep E	Rare	Nil	Nil	Nil	Most common

Solution to Question 3:

Fatality due to Hepatitis E is highest in pregnant women. In the general population, hepatitis E can lead to fatal fulminant hepatitis in only 1–2% of cases. However, in pregnant women, this is raised to 20% of cases.

HEV does not have a carrier state and can cause chronic hepatitis only in immunocompromised individuals. Its most common route of spread is feco-oral.

Note - HEV is also known as E-NANB, i.e., Enterically transmitted Non-A Non-B hepatitis virus.It often appears in epidemics; hence, it is also called epidemic NANB.

Solution to Question 4:

Acute viral hepatitis caused by the hepatitis D virus is associated with the highest risk of developing fulminant hepatitis.

Sold by @itachibot

However, fulminant hepatitis is most commonly caused by hepatitis E, especially in endemic countries such as India. This is due to the greater prevalence of hepatitis E infections compared to hepatitis D.

Solution to Question 5:

The chance of vertical transmission of hepatitis B from a mother who is HBsAg positive and HBeAg positive is >90%. In contrast, when the mother is HBsAg positive but HBeAg negative or HBsAg positive with Anti Hbe, the risk of vertical transmission is low.

To prevent vertical transmission of HBV, HBsAg positive pregnant women with high HBV DNA titres should receive tenofovir prophylaxis from the 28th week of pregnancy until birth. Infants born to these women are given hepatitis B immunoglobulin along with the first dose of the hepatitis B vaccine.

The presence of HBeAg indicates active replication of HBV. Therefore, HBeAg is the qualitative marker of infectivity. HBV DNA titre is the quantitative marker of infectivity.

Routes of transmission of hepatitis viruses:

Hepatitis	Hep B	Hep C	Hep A	Hep E
Vertical route	HBsAg and HBeAg positive mother >90%HBsAg positive with Anti HBe mother- 10-15 %	5%	Nil	Rare
Needle	6-30%	1.8-6 %	Nil	Nil
Blood transfusion	1 in 2 lakh(Most common blood transfusion-related virus)	1 in 18 lakh	Rare	Nil
Sexual	Variable	Yes	Rare	Nil
Feco-oral	-	-	Most common	Most common

Solution to Question 6:

The first virologic marker to rise in acute hepatitis B is HBsAg i.e., Hepatitis B surface antigen. It is detectable in serum within 1–12 weeks of infection. Usually, HBsAg becomes undetectable 1–2 months after the onset of jaundice. If it persists beyond 6 months, it indicates chronic hepatitis B.

However, very rarely, patients may have an HBsAg negative status even during active infection due to infection with escape mutant HBV. These escape mutants have mutations within the S gene and they are known to be responsible for occult hepatitis B infections.

Sold by @itachibot



MARROW
Sold by @itachi... If you purchased this from some... you may have been scammed. else,

Solution to Question 7:

The diagnosis of acute hepatitis B in the window period or serological gap can be made by the detection of the anti-HBc antibody.

The window period is the period between the disappearance of HBsAg to anti-HBs seroconversion. During this window period, anti-HBc is the only detectable antibody.

Solution to Question 8:

Increased viral load and negative HBeAg suggest a pre-core mutant variety of hepatitis B virus. These strains have mutations in the pre-core region of the C gene, leading to a stop codon that prevents HBeAg synthesis. These strains can be detected in the serum by HBV DNA analysis.

Infection with hepatitis B mutants is difficult to treat and causes infections of prolonged duration. They have a higher risk of progressing to liver cirrhosis.

Types of hepatitis B mutants are:

Type	Feature
Pre-core mutants	Defect in pre-core region CNo HBeAg
Escape mutants	Mutation in S regionHBsAg resists neutralisation by anti-HBsSeen inInfants born to HBeAg positive mothersLiver transplant recipientAssociated with passive and active immunization

Sold by @itachibot

Type	Feature
YMDD mutation	Mutation in P region i.e. YMDD locus which codes forreverse transcriptaseSeen in patients on lamivudine therapy

Solution to Question 9:

The given lab findings are most likely to be in patients who had hepatitis B in the remote past i.e., patients in late convalescence period. In such conditions, the patient will have positive IgG anti-HBc and negative HBsAg.

Solution to Question 10:

The serological marker marked in the image which becomes detectable early in the infection and remains elevated for a long period after recovery from the infection is an anti-HBc antibody.

Anti-HBc is demonstrable in serum within the first 1–2 weeks after the appearance of HBsAg. It precedes detectable levels of anti-HBs by weeks to months.

IgM anti-HBcAg antibody is seen early in the infection for about 6 months and indicates a recent infection. IgG anti-HBcAg antibody is seen later, after 6 months, and indicates remote infection.

Solution to Question 11:

The most reliable indicator of recent HBV infection is IgM anti-HBc in serum. However, the first virologic marker to rise in acute hepatitis B is HBsAg i.e. surface antigen.

HBsAg is detectable in serum within 1–12 weeks of infection. Anti-HBc is demonstrable in the serum about 1-2 weeks after this, and may sometimes persist in the serum even longer than an anti-HBs antibody. IgM anti-HBcAg is seen early in the infection and persists for about 6 months. It indicates a recent infection. IgG anti-HBcAg antibody is seen later i.e. after 6 months and indicates remote infection.

HBcAg is not detectable in the serum of patients with HBV infection as it is present intracellularly.

Note: The risk of transmission of HBV from percutaneous routes (eg., needle prick) is about 6-30% and it is about 100 times more infectious than HIV.

Solution to Question 12:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

The need for administration of hepatitis B hyperimmune globulin (HBIG) in a previously vaccinated individual can be determined based on anti-HBsAg levels. It is the marker for determining the efficacy of hepatitis B vaccination.

The complete hepatitis B vaccine schedule consists of three intramuscular doses at 0, 1, and 6 months.

Post Exposure Prophylaxis:

The prophylaxis can be done using the hepatitis B vaccine or hepatitis B immunoglobulin (HBIG).

Vaccination status of the exposed person	Recommended prophylaxis
Unvaccinated	HBIG (0.06 mL/kg) and initiate hepatitis B vaccine series
Known responder to previous vaccination (anti-HBs titer>10 mIU/ml)	No action required
Non-responder to previous vaccination (anti-HBs titre <10 mIU/ml)	HBIG: 2 doses (1 month apart) OR HBIG: 1 dose + initiate hepatitis B revaccination.

Sold by
if you purchased this from
you may have been scam

Solution to Question 13:

IgM anti-HBc can be used to differentiate between hepatitis D coinfection and superinfection in patients with HBV infection.

Co-infection:

• Simultaneous infection with both HBV and HDV

• IgM anti-HBc detectable in co-infection as its presence indicates recent infection

Superinfection:

• HDV infection of an individual chronically infected with HBV

• IgM anti-HBc will not be detectable

• IgG anti-HBc will be present

Solution to Question 14:

The given histopathological image shows ground-glass hepatocytes which are a hallmark of chronic hepatitis B infection.

Sold by @itachibot



These ground-glass hepatocytes are liver parenchymal cells that have an eosinophilic, granular, and glassy appearance on light microscopy. This appearance is due to the accumulation of HBsAg (hepatitis B surface antigen) in the smooth endoplasmic reticulum.

Liver biopsy findings in viral hepatitis:

Virus	Liver Biopsy Findings
Hepatitis A	Hepatocyte swelling, monocyte infiltration, Councilman bodies
Hepatitis B	Ground glass appearance
Hepatitis C	Lymphoid aggregates with a focal area of macrovesicular steatosis
Hepatitis D	Similar to HBV
Hepatitis E	Patchy necrosis

Solution to Question 15:

Antiviral therapy is not indicated in acute hepatitis B infection. It is only indicated in chronic hepatitis B.

The majority of acute hepatitis (A, B, and E) infections are self-limiting, resulting in viral elimination and complete healing within 6 months. Hence, they need only supportive management. Only 5% of acute hepatitis infections become chronic.

Sold by @itachibot

In acute hepatitis C infection, recovery is rare, and the risk of progression to chronic hepatitis is high(~85%). Thus antiviral therapy is started irrespective of the disease stage.

Solution to Question 16:

ALT > 2 times upper limit of normal is an indication for antiviral therapy in chronic hepatitis B infection.

The decision to initiate antiviral therapy is primarily based on the presence or absence of cirrhosis, and parameters such as HBeAg, alanine aminotransferase (ALT) level, and HBV DNA level.

The following are the indications for starting antivirals in chronic hepatitis B:

- Patients without cirrhosis
- HBeAg + with HBV DNA >20,000 IU/mL and ALT level >2 times the upper limit of normal
- HBeAg -ve with HBV DNA >2000 IU/mL and ALT level >2 times the upper limit of normal
- Patients with compensated cirrhosis and an HBV DNA >2000 IU/mL
- Patients developing acute liver failure or decompensated cirrhosis

Recommended first-line antiviral drugs:

- Entecavir
- Tenofovir alafenamide
- Tenofovir disoproxil fumarate

Pegylated interferon alfa can also be used. Treatment is aimed at HBeAg seroconversion, but most patients may need to continue antiviral therapy indefinitely.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

HBeAg negative and HBV DNA <2000 IU/mL is suggestive of HBV carrier. Serial monitoring of both biochemical and virological parameters is done for such patients.

Indications for starting antiviral therapy in chronic hepatitis B are:

- Patients without cirrhosis
- HBeAg + with HBV DNA >20,000 IU/mL and ALT level >2 times the upper limit of normal
- HBeAg -ve with HBV DNA >2000 IU/mL and ALT level >2 times the upper limit of normal
- Patients with compensated cirrhosis and an HBV DNA >2000 IU/mL
- Patients developing acute liver failure or decompensated cirrhosis

Solution to Question 18:

Sold by @itachibot

First-line drugs used for the treatment of chronic hepatitis B include entecavir and tenofovir.

Second-line drugs for this condition include:

• Lamivudine

• Adefovir

• Telbivudine

Pegylated interferon can also be used in its treatment.

Solution to Question 19:

Surveillance for hepatocellular carcinoma is recommended for Asian male HBV carriers over the age of 40 years.

The most common etiological factor for hepatocellular carcinoma in the world is the hepatitis B virus (HBV) infection. Surveillance programs have been implemented to screen for HCC in high-risk individuals because it is more cost-effective than the treatment of HCC.

Surveillance for hepatocellular carcinoma (HCC) is recommended in:

• HBV carriers with cirrhosis

• Patients with high serum levels of HBV DNA and ongoing liver injury

• HBV carriers born in Africa and over the age of 20

• Asian male HBV carriers over age 40

• Asian female HBV carriers over age 50

• HBV carriers with a family history of HCC

Note:

The CDC recently updated its guidelines for Hepatitis B screening. The report includes screening for all adults >18 years old using the triple-panel test (HBsAg, Anti-HBs, total anti-HBc) and expanded risk-based testing in certain high-risk populations.

Solution to Question 20:

IgM anti-HBe is not one of the markers that are evaluated in a suspected case of acute viral hepatitis.

Investigations done in suspected acute viral hepatitis are:

• IgM anti-HAV - to detect acute hepatitis A infection which is the most common cause of acute hepatitis in children

• HBsAg and IgM anti-HBc - to detect acute hepatitis B infection, where HbsAg is the first marker to rise and IgM anti-HBcAg is the most reliable marker of acute infection

• Anti HCV - to evaluate for acute hepatitis C infection

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Sold by @itachibot

• IgM anti-HEV (rarely)- to detect acute hepatitis E infection which is one of the causes of acute viral hepatitis in adults.

Solution to Question 21:

The given clinical scenario is suggestive of jaundice. The tests that should be done in this patient to rule out acute viral hepatitis include:

• Anti-HAV

• HBsAg

• IgM anti-HBc

• Anti-HCV

We mainly test for hepatitis A, B, and C. Tests for hepatitis D and E are not used routinely.

Hepatitis D only occurs concurrently with Hepatitis B, either as a co-infection (when it is clinically indistinguishable from HBV infection) or superimposed on chronic HBV infection. In the latter scenario, it presents as a sudden worsening of the patient's condition. IgM anti-HDV, IgG anti-HDV, and HDV RNA assays are useful for diagnosis when hepatitis D is suspected.

Hepatitis E is not tested as the majority of patients who acquire hepatitis E virus (HEV) infection spontaneously clear the virus. Acute hepatic failure is more likely in those who are pregnant and in those who are malnourished or have preexisting liver disease. Chronic HEV has been described almost exclusively in immunocompromised hosts (e.g., organ transplantation). The diagnosis of acute HEV is typically based upon the detection of IgM antibodies to HEV. Both false positives and negatives are common with available assays and are thus, not very reliable. Thus, additional serologic testing or HEV RNA testing should be performed to confirm the diagnosis and not be useful as a screening test. [Please note the difference between anti-HBe and anti-HEV]

Condition Suspected	Targetted Test Performed
Acute Hepatitis A	IgM Anti-HAV
Acute Hepatitis B	HBsAg andIgM Anti-HBc*1 or both can be positive. A positive IgM AntiHBc with negative HBsAg is seen when HBsAg is below the detection threshold.
Acute Hepatitis C	Anti HCV

Solution to Question 22:

HCV screening is not indicated for household contacts of persons with hepatitis C. However, hepatitis B virus screening is recommended in household contacts of HBV-infected persons.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Indications of HCV infection screening:

• Persons with a history of using injection drugs

• Persons with HIV infection

• Long term hemodialysis

• Persons with an unexplained elevation of aminotransferases

• Recipient of blood or organs from a donor found to be positive for hepatitis C

• Children born to women with Hepatitis C

• Health care, public safety, and emergency medical personnel following needle injury or mucosal exposure to HCV contaminated blood

• Sexual partners of persons with Hepatitis C

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 23:

Polyarteritis nodosa is not characteristically associated with hepatitis C infection. However, it is a common extrahepatic manifestation of hepatitis B viral infection.

Solution to Question 24:

While HCV RNA is considered the gold standard for establishing a diagnosis of hepatitis C, it has no role in determining the progression of liver disease in patients with chronic hepatitis C.

The likelihood of progression of liver disease in patients with chronic hepatitis C is increased by:

• Longer duration of infection - most important determinant

• Older age

• Obesity - patients with a high body mass index are at an increased risk of developing fibrosis

• Diabetes mellitus and insulin resistance

• Concomitant liver disorder (eg., alcoholic liver disease, chronic hepatitis B, hemochromatosis)

• Concomitant HIV infection.

The best prognostic indicator in chronic hepatitis C is the rate of fibrosis and baseline inflammation seen in liver histology.

Solution to Question 25:

The given clinical scenario is suggestive of hepatitis C for which the preferred treatment regimen is velpatasvir 100 mg and sofosbuvir 400 mg.

Sold by @itachibot

The goal of treatment is a sustained virologic response (SVR), which is defined as the absence of a virus in the blood (undetectable HCV RNA), 12 weeks after completion of treatment. This can be achieved by the use of pan-genotypic directly acting antiviral drugs, irrespective of the disease stage.

Solution to Question 26:

Sofosbuvir is not an NS5A inhibitor but is an NS5B nucleoside/nucleotide polymerase inhibitor.

The directly acting antivirals used in the treatment of Hepatitis C infection include:

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Autoimmune Disorders of Hepatobiliary System

Question 1:

Autoimmune hepatitis is suspected in a woman presenting with fatigue, pruritus, and arthralgia. She has a past history of Hashimoto's thyroiditis. Which of the following is not part of the simplified scoring system for the given condition?

a) ANA

b) Typical histological features

c) Elevated IgG

d) Presence of viral markers

Question 2:

Which of the following drugs is associated with Autoimmune hepatitis (AIH)?

a) Minocycline

b) 5 Aminosalicylic acid

c) Piroxicam

d) Nitroglycerine

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 20-year-old lady is suspected to have autoimmune hepatitis. Which of the following antibodies is unlikely to be present in her serum?

a) Cytoplasmic antineutrophil cytoplasmic antibodies (cANCA)

b) Antinuclear antibody (ANA)

c) Smooth muscle antibody(SMA)

d) Anti-soluble liver antigen (SLA)

Question 4:

Match the following:

a) A-i , B-iii , C-ii , D-iv

Sold by @itachibot

b) A-iv , B-ii , C-iii , D-i

c) A-ii , B-iii , C-i , D-iv

d) A-i , B-iii , C-iv , D-ii

Question 5:

A 24-year-old woman presenting with abdominal pain is diagnosed with autoimmune hepatitis. The mainstay of management in this patient is:

a) Glucocorticoids

b) Azathioprine

c) Mycophenolate mofetil

d) Methotrexate

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A 23-year-old female patient with a diagnosis of autoimmune hepatitis did not respond well to glucocorticoids. Which among the following is false regarding her further management options?

a) Increase the dose of glucocorticoids

b) Glucocorticoids with high dose azathioprine

c) Switch to Mycophenolate mofetil

d) Liver transplant is contraindicated

Question 7:

A young lady has been on treatment for autoimmune hepatitis for 2 weeks, she has now come for follow-up investigations. Which of the following suggests a poor prognosis in this patient?

a) Bridging necrosis

b) Decreased globulins

c) Increased serum bilirubin

d) Increased serum albumin

Question 8:

Sold by @itachibot

The most common symptom of primary biliary cholangitis (PBC) is

a) Pruritus

b) Fatigue

c) Jaundice

d) Right upper quadrant pain

Question 9:

A 44-year-old woman with rheumatoid arthritis is now diagnosed with primary biliary cholangitis. Which of the following is false regarding her lab values?

a) Positive Anti-LKM- 2 antibodies

b) Elevated alkaline phosphatase (ALP)

c) Elevated gamma-glutamyl transferase (GGT)

d) Positive antimitochondrial antibodies

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A patient with a diagnosis of primary biliary cholangitis has been taking ursodeoxycholic acid (UDCA) twice daily. She now presents with ascites and variceal bleeding. What is the most appropriate treatment for this patient?

a) Initiating methotrexate

b) Continuing ursodeoxycholic acid

c) Add obeticholic acid to the treatment

d) Consider liver transplantation

Question 11:

A middle-aged woman presented with fatigue, pruritis, and xanthomas over the elbows. A biopsy of her bile duct revealed florid duct lesions. What is the diagnosis?

a) Klatskin tumor

b) Primary sclerosing cholangitis (PSC)

c) Primary biliary cholangitis (PBC)

d) Intrahepatic cholestasis

Sold by @itachibot

Question 12:

Which of the following statements about primary sclerosing cholangitis is false?

a) Affects both intra and extrahepatic bile ducts

b) Liver biopsy definitively establishes the diagnosis

c) more common in males

d) Leads to development of cholangiocarcinoma

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	a
5	a
6	d
7	c
8	b
9	a
10	d
11	c
12	b

Detailed Explanations

Solution to Question 1:

Presence of viral markers is not a part of the simplified scoring system for diagnosis of autoimmune hepatitis.

The simplified scoring system includes four variables. They are:

• Autoantibodies

• Elevated serum IgG level

• Typical compatible histologic features

Sold by @itachibot

• Absence of viral hepatitis markers

The clinical presentation in autoimmune hepatitis is variable and shares features with acute and chronic hepatitis of other aetiologies. They usually present with an insidious onset of non-specific symptoms such as fatigue, anorexia and eventually jaundice. It can then progress to extensive liver necrosis and liver failure.

Autoimmune hepatitis is associated with several autoimmune diseases such as:

• Rheumatoid arthritis

• Systemic lupus erythematosus

• Thyroiditis and Grave's disease

• Haemolytic anemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

Minocycline is associated with the development of drug-induced AIH.

Other drugs associated with AIH include nitrofurantoin, methyldopa, diclofenac, and possibly statins.

Solution to Question 3:

cANCA is not seen in autoimmune hepatitis. It is positive in Wegner's granulomatosis. pANCA is what is seen in autoimmune hepatitis.

There are 2 types of autoimmune hepatitis:

Type I (classic) autoimmune hepatitis is often seen in young women. It is linked to HLA-DR3 or HLA-DR4. It responds well to steroids and has a better prognosis. The autoantibodies that are seen in type I are:

• ANA - antinuclear antibodies

• ASMA - anti-smooth muscle actin

• pANCA - perinuclear anti-neutrophil cytoplasmic

• anti-SLA - anti-soluble liver antigen

Type II autoimmune hepatitis is often seen in children. It is linked to HLA-DRB1 or HLA-DQB1. It usually leads to fibrosis, cirrhosis and liver failure. The autoantibodies that are seen in type II are:

• Anti-LKM 1 - anti-liver kidney microsomes

• Anti-liver cytosol 1

Solution to Question 4:

Sold by @itachibot

• Anti LKM 1- Autoimmune hepatitis II

• Anti SLA - Autoimmune hepatitis I

• Anti LKM 2 - Drug-induced hepatitis

• Anti LKM 3 - Chronic hepatitis D

Solution to Question 5:

The mainstay of management in autoimmune hepatitis is glucocorticoid therapy.

Glucocorticoids are started at high doses and tapered down to a low maintenance dose. The dose of glucocorticoids is often reduced by combining with azathioprine.

The benefits of glucocorticoid + azathioprine combination therapy are:

• Reduction in the duration of therapy.

• Avoid life-threatening complications of steroids (e.g., cushingoid features, hypertension, diabetes, osteoporosis).

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

Liver transplant is not contraindicated in refractory autoimmune hepatitis even though the risk of recurrence is high.

The medical management of refractory autoimmune hepatitis include:

• High dose glucocorticoids

• Glucocorticoids + high dose azathioprine

• Addition of cyclosporine, tacrolimus, or mycophenolate mofetil.

Solution to Question 7:

Persistently increased serum bilirubin after 2 weeks of therapy suggests a poor prognosis.

Early signs of poor prognosis include:

• Failure of serum bilirubin to reduce after 2 weeks of therapy.

Early signs of good prognosis include:

• An increase in serum albumin.

• A decrease in serum bilirubin, globulins and aminotransferases.

• Improvement of symptoms.

Histopathologic improvement can often take months to years.

Sold by @itachibot

Solution to Question 8:

The most common symptom of primary biliary cholangitis(PBC) is fatigue.

Primary biliary cholangitis(PBC) is a granulomatous autoimmune condition characterized by the destruction of interlobular hepatic bile ducts.

Most patients with PBC are asymptomatic at the time of the diagnosis. Fatigue and pruritus are both common presenting symptoms. However, the most common symptom is fatigue.

Other symptoms and signs are -

• Right upper quadrant pain

• Hyperpigmentation

• Xanthomas and xanthelasma

• Jaundice

• Hepatosplenomegaly.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

Anti-LKM 2 antibodies are not seen in primary biliary cholangitis. They are seen in drug-induced hepatitis.

The laboratory findings of primary biliary cholangitis include :

• Liver function tests

• Elevated ALP

• Elevated GGT and 5'- nucleotidase

• Elevated ALT and AST

• Serology -

• Antimitochondrial antibodies - positive in 95% of patients. These are mostly directed against the pyruvate dehydrogenase complex.

• Antinuclear antibodies

• Elevated HDL, LDL, and VLDL

• Elevated serum IgM

Hyperbilirubinemia is seen once cirrhosis has developed.

Solution to Question 10:

Sold by @itachibot

The presence of ascites and variceal bleeding suggest decompensated liver disease and the appropriate treatment is liver transplantation.

The first-line drug in primary biliary cholangitis is ursodeoxycholic acid (UDCA). It only slows the rate of progression of the disease. It does not provide any symptomatic benefit or cure the disease.

Obeticholic acid is another drug used in PBC as monotherapy or in combination with UDCA. It is a farnesoid X receptor agonist.

Liver transplantation is considered when there are signs of decompensation or portal hypertension such as ascites, variceal bleeding, encephalopathy, coagulopathy, or malnutrition.

Solution to Question 11:

Florid duct lesion is characteristic of primary biliary cholangitis (PBC). It is the lymphocytes and plasma cells all centered around bile ducts.



Primary sclerosing cholangitis has the characteristic "onion-skin" concentric fibrosis surrounding the bile ducts.

Sold by @itachibot



Sold by @itachi... else,
If you purchased this from some... you may have been scammed.

Solution to Question 12:

Liver biopsy does not establish the diagnosis in primary sclerosing cholangitis. The diagnosis requires imaging with MRCP.

Primary sclerosing cholangitis (PSC) is a chronic progressive disorder characterized by inflammation and fibrosis of intra and extrahepatic biliary ducts resulting in strictures and cholestasis. The cause is unknown. Most patients have underlying ulcerative colitis. It has an association with HLA DRw52. Patients present with symptoms of cholestasis such as fatigue and pruritus.

The laboratory findings include:

• Liver function tests - Elevated ALP and aminotransferases.

• Increased serum IgM levels

• Positive p-ANCA

MRCP is the investigation of choice. It shows multiple strictures in intra and extrahepatic bile ducts. Liver biopsy reveals a concentric "onion skin" fibrosis. This is however not pathognomic of PSC.

There is no definitive treatment. It involves the administration of UDCA and liver transplantation.

PSC can lead to the development of cholangiocarcinoma

The image below shows onion-skin fibrosis in PSC.

Sold by @itachibot



Sold by @Itachi... else,
If you purchased this from some
you may have been scammed.

Sold by @itachibot

Acute Liver Failure and Liver Transplantation

Question 1:

In acute liver failure, encephalopathy occurs within _____ of the development of clinical symptoms.

a) 6 months

b) 6 weeks

c) 4 months

d) 4 weeks

Question 2:

What is the most common cause of acute liver failure in India?

a) Alcohol

b) Viral infection

c) Drug induced

d) Malaria

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A small child was rushed to the ED by his parents. He had accidentally consumed wild mushrooms while they were trekking. Examination showed features of encephalopathy and his liver function tests were deranged. Which of the following is least likely to be seen in this condition?

a) Hyperglycemia

b) Hypomagnesemia

c) Hypokalemia

d) Hyponatremia

Question 4:

A patient is said to have hyperacute liver failure if encephalopathy occurs within?

Sold by @itachibot

a) 3 days

b) 7 days

c) 10 days

d) 14 days

Question 5:

A young patient with Hepatitis E virus infection develops acute liver failure. Which of the following scores can be used in assessing the prognosis of this patient?

a) Pitt score

b) Child Pugh score

c) MELD score

d) MDF score

Question 6:

A young patient is admitted to the hospital after ingestion of a large number of acetaminophen tablets. To predict the prognosis using King's College hospital criteria in this patient, which of the parameters are not used?

a) Arterial pH

b) Prothrombin time

c) Serum creatinine

d) Serum bilirubin

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

For patients with acute liver failure, which of the following is not an absolute contraindication for liver transplantation?

a) AIDS

b) Active alcohol abuse

c) Metastatic hepatocellular carcinoma

d) Portal vein thrombosis

Question 8:

Sold by @itachibot

Which among the following is the most common indication of liver transplantation in children?

a) Neonatal hepatitis

b) Alpha-1 antitrypsin deficiency

c) Biliary atresia

d) Congenital hepatic fibrosis

Question 9:

When should a liver transplant assessment be considered in a patient with cirrhosis of the liver?

a) When he develops spontaneous bacterial peritonitis

b) When he has ascites which responds to diuretics

c) MELD score more than 10

d) Child Pugh class A

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A surgeon has removed a part of the liver from the live donor and has stored it in the University of Wisconsin (UW) solution. Which of the following is not a component of this solution?

a) Lactobionic acid

b) Glutathione

c) Allopurinol

d) Mannitol

Question 11:

Which of the following is a false statement regarding liver transplantation (LT)?

a) The ideal graft weight to recipient weight ratio (GRWR) in living donor transplant is >0.8 %.

b) In domino LT, liver grafts with metabolic diseases is used as they are structurally and functionally normal.

c) Extended criteria donors and paired-exchange LT decrease waitlist mortality.

Sold by @itachibot

d) Reduced size LT involves dividing the whole liver into right lobe for an adult patient and left lateral segments for pediatric patients.

Question 12:

Auxiliary liver transplantation is beneficial in which of the following conditions:

a) Acute liver failure

b) Hepatocellular carcinoma

c) Genetic Hemochromatosis

d) Chronic viral hepatitis

Question 13:

Which of the following sentences is true regarding split liver transplantation?

a) It is done in living donor

b) Left lateral lobe is donated to the paediatric recipient

c) It is the most common type of liver transplant

d) Caudate lobe is donated to the adult recipient

Question 14:

A patient who underwent liver transplantation is started on tacrolimus for immunosuppression. His family physician prescribed him fluconazole for the treatment of a fungal infection. This will lead to

a) Toxicity of fluconazole

b) Transplant rejection

c) No effect on both the drugs

d) Toxicity of tacrolimus

Question 15:

A patient presented with fever and pain abdomen. He had undergone liver transplantation a month ago and is taking immunosuppressants. On examination, abdominal guarding was present and on auscultation, no bowel sounds were heard. Doppler showed hepatic artery thrombosis. Which of the following medications is he most likely on?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Tacrolimus

b) Sirolimus

c) Cyclosporine

d) Mycophenolate mofetil

Question 16:

All of the following complications can be seen in patients with liver transplantation except

a) Arrhythmias

b) Seizures

c) Hepatic artery thrombosis

d) T cell lymphoma

Question 17:

A liver biopsy is performed in a patient who is suspected to have acute rejection following liver transplantation. Which of the following findings in the biopsy will confirm the presence of acute rejection?

a) Fibrosis

b) Mononuclear infiltration

c) Endotheliitis

d) Focal parenchymal necrosis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A 40-year-old male patient developed fever, malaise, and anorexia after a month of liver transplantation. On evaluation, he has leukopenia, thrombocytopenia, and deranged liver function tests. Which of the following organisms is most likely to have caused this?

a) Cytomegalovirus

b) Varicella zoster virus

c) Epstein-Barr virus

d) Herpes simplex virus

Question 19:

Sold by @itachibot

Which of the following primary diseases has the higher chance of recurrence, after liver transplantation?

a) Sclerosing cholangitis

b) Autoimmune hepatitis

c) Alpha-1 antitrypsin deficiency

d) Primary biliary cirrhosis

Answer Key

Question No.	Correct Option
1	d
2	b
3	a
4	b
5	c
6	d
7	d
8	c
9	a
10	d
11	d
12	a
13	b
14	d
15	b
16	d
17	c
18	a
19	b

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

In acute liver failure, encephalopathy occurs within 4 weeks of the development of the clinical symptoms.

Acute liver failure is the rapid worsening of the liver function in patients without cirrhosis or any preexisting liver disease. The patient initially suffers from an acute injury to the liver by viruses, drugs, etc. Following this, the function of the liver rapidly worsens and patient develops coagulopathy and encephalopathy within 4 weeks of the development of jaundice or other clinical symptoms.

Solution to Question 2:

The most common cause of acute liver failure (ALF) in India is a viral infection. Among the viruses, hepatitis E virus is the most common cause of acute liver failure.

In developed countries, acetaminophen overdose is the most common cause of acute liver failure.

Solution to Question 3:

The given scenario is suggestive of Amanita phalloides (wild mushrooms) toxicity causing acute liver failure. Hypoglycemia is seen in acute liver failure and not hyperglycemia. It occurs due to impaired gluconeogenesis and glycogenolysis.

Features of acute hepatic failure are:

• Hepatic encephalopathy and cerebral edema- There are increased levels of ammonia (neurotoxin) in the body as a result of decreased elimination by the liver.

• Coagulopathy- There is a decrease in the synthesis of coagulation factors leading to the elevated prothrombin time and INR.

• Increased risk of infections- It occurs as a result of decreased reticuloendothelial activity, decreased synthesis of complements, acute phase reactants, etc.

• Renal failure- This is more commonly seen in drug/toxin-induced hepatic failure.

• Metabolic complications like hypoglycemia, hyponatremia, hypokalemia, hypophosphatemia, hypomagnesemia.

Solution to Question 4:

Hyperacute liver failure is defined as the development of encephalopathy within 7 days of the appearance of symptoms.

Acute liver failure can be subcategorized into three groups depending on the time interval between the onset of jaundice and encephalopathy.

• Hyperacute- encephalopathy develops within 7 days or less than 7 days

• Acute- encephalopathy develops between 8-28 days

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• Subacute- encephalopathy develops between 28 days to 6 months

Patients with hyperacute liver failure are more likely to develop coagulopathy and cerebral edema, but most of the patients will recover without the need for liver transplantation. On the other end, patients with subacute liver failure have less severe coagulopathy and cerebral edema, but most of them will need liver transplantation.

Solution to Question 5:

In a patient with acute liver failure, MELD score and King's College Hospital criteria can be used to assess the prognosis.

Option A: Pitt score is used used to predict mortality in patients having bloodstream infections.

Option B: The child-Pugh score is used to assess the severity of cirrhosis.

Option D: Maddrey's Discriminant Function (MDF) score suggests which patients with alcoholic hepatitis may have a poor prognosis and benefit from steroid administration.

Solution to Question 6:

Serum bilirubin is not included in King's College Hospital criteria in acute liver failure due to paracetamol overdose.

The King's College Hospital Criteria is a prognostic tool for patients with acute liver failure (ALF). The criteria is divided into acetaminophen and non-acetaminophen causes of liver failure.

Liver transplantation is indicated in acetaminophen-induced ALF if there is a presence of

• Either pH <7.3

• Or all of the following

• Prothrombin time >100 sec or INR >6.5

• Serum creatinine level >3.4 mg/dL (300 µmol/L) or anuria

• Grade 3-4 encephalopathy

Liver transplantation is indicated in non-acetaminophen causes of ALF if there is a presence of

• Either prothrombin time >100 sec or INR >6.7

• Or any three of the following

• Unfavorable etiology (seronegative hepatitis or drug reaction)

• Age <10 or >40 yr

• Acute or subacute category (duration of jaundice >7 days)

• Serum bilirubin level >17.5 mg/dL (300 µmol/L)

• Prothrombin time >50 sec or INR >3.5

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 7:

Portal vein thrombosis is not an absolute contraindication to liver transplantation. This is because portal vein thrombosis can be overcome by constructing a graft from the donor liver portal vein to the recipient's superior mesenteric vein.

Absolute contraindications of liver transplantation are:

• Uncontrolled extra-hepatobiliary infection or untreated sepsis

• Uncorrectable life-limiting congenital anomalies

• Active substance or alcohol abuse

• Malignancies- Hepatocellular carcinoma with metastatic spread, incurable malignancies outside the liver, cholangiocarcinoma

• AIDS

• Advanced cardiopulmonary disease

Note: Milan criteria- Patients with primary hepatocellular malignancy can be considered for liver transplantation if there is:

• A single lesion ≤5 cm.

• Up to three separate lesions all <3 cm without any evidence of gross vascular invasion, nodal or distant metastases.

Sold by @itachibot
If you purchased this from someone else you may have been scammed

Solution to Question 8:

Biliary atresia is the most common indication of liver transplantation in children.

Indications of liver transplant in children:

• Biliary atresia

• Neonatal hepatitis

• Congenital hepatic fibrosis

• Alpha-1-antitrypsin deficiency

• Wilsons disease

• Glycogen storage disorders

• Lysosomal storage disorders

The most common indication of liver transplant in an adult is the cirrhotic liver. Cirrhosis may be the consequence of viral hepatitis, autoimmune hepatitis, chronic use of alcohol, or biliary cirrhosis.

Solution to Question 9:

Sold by @itachibot

Liver transplant assessment should be considered in a patient with cirrhosis of the liver when he develops spontaneous bacterial peritonitis.

Cirrhosis is the most common indication of liver transplantation in adults. However, not all cases of cirrhosis require liver transplantation. It is considered

1. When the patient either develops the complication of cirrhosis like:

• Persistent hepatic encephalopathy

• Spontaneous bacterial peritonitis

• Portal hypertensive bleeding

• Diuretic resistant ascites

• Hepatorenal syndrome

2. Or has a poor liver function

• MELD score > 12

• Child-Pugh grade C

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

Mannitol is not a component of the University of Wisconsin (UW) solution. It is the most commonly used preservation solution during the transplantation of the liver and kidneys. The organ can be stored in this solution for up to 20 hours.

Components of UW solution are:

• Lactobionate- Prevents cellular swelling and reperfusion injury

• Raffinose and hydroxyethyl starch- Prevents endothelial injury

• Glutathione and allopurinol- free radical scavengers

• Adenosine- required for regeneration of ATP

Solution to Question 11:

The false statement regarding liver transplantation is option D. Reduced-size liver transplantation involves the resection of a full-size liver to appropriately fit in the abdominal cavity of small adults/adolescents. It is reserved only for urgent transplantation. On the contrary, split LT involves dividing the whole donor liver into a left lateral segment (segments II and III) for a pediatric patient and a right lobe (segments I, IV-VIII) for an adult patient.

Option A: The required graft volume in a living donor transplant is measured by graft weight to recipient weight ratio (GRWR) or ratio of graft volume (GV) to recipient standard liver volume (SLV). The ideal GRWR is >0.8 % or GV/SLV is > 35%. The remnant donor liver mass is kept at >30% for safe regeneration.

Sold by @itachibot

Option B: Domino liver transplant involves transplanting a liver from a donor with metabolic disease (eg: familial amyloid polyneuropathy) to a recipient with end-stage liver disease (eg: hepatocellular carcinoma). The donor also receives an LT (healthy donor) and is prioritized over the end-stage liver disease recipient. As a liver with metabolic diseases is structurally and functionally normal, it can prolong the recipient's life by at least 15-20 years as they will not develop the metabolic disease immediately.

Option C: Extended criteria donors/ECDs (liver grafts extracted from donors after circulatory death) and paired-exchange technique are used to decrease waitlist mortality. However, ECDs have poor graft function due to prolonged hypoxia and are associated with higher infection transmission and biliary complications.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

Auxiliary liver transplantation is beneficial in acute liver failure.

Auxiliary liver transplantation involves implanting a healthy liver graft heterotopically or orthotopically while leaving the whole or part of the native liver intact. The auxiliary partial orthotopic LT (APOLT) technique is used in patients with acute liver failure and metabolic diseases (primary hypercholesterolemia, glycogen storage disorders, organic acidemia, etc). In acute liver failure, this technique helps in native liver regeneration and immunosuppression withdrawal. In children with metabolic liver diseases, this technique helps in future gene therapy.

Options B, C, and D are chronic liver diseases not suitable for auxiliary liver transplantation.

Solution to Question 13:

In a split liver transplant, the left lateral lobe is donated to the pediatric recipient.

Split liver donation: This method is done when the liver from the deceased donor is suitable for transplant in an adult as well as a pediatric recipient. The liver is split into two. The left lateral segment (2 and 3) is donated to the pediatric recipient, left medial and right segments (4, 5, 6, 7, and 8) is donated to the adult recipient.

Solution to Question 14:

In a patient who is on tacrolimus, the addition of fluconazole may lead to toxicity of the tacrolimus.

Tacrolimus and cyclosporine are calcineurin inhibitors and are the common agents used for the immunosuppression following liver transplantation. They are metabolized by the CYP3A enzyme. Fluconazole can inhibit this enzyme and lead to increased levels of tacrolimus and cyclosporine.

Solution to Question 15:

Sold by @itachibot

The given clinical scenario is suggestive of peritonitis due to hepatic artery thrombosis. Sirolimus is known to cause hepatic artery thrombosis in liver transplant recipients.

Sirolimus is an mTOR inhibitor and prevents T cell activation. It is not used commonly in liver transplant recipients because of the increased frequency of hepatic artery thrombosis in the first month posttransplantation. However, they can be used in patients with calcineurin inhibitor-related nephrotoxicity.

Solution to Question 16:

Patients with liver transplantation are more likely to develop B cell lymphomas and not T cell lymphoma.

Complications of liver transplantation

• Rejection of the transplanted liver

• Hepatic artery thrombosis

• Posttransplantation lymphoproliferative disorders- It can range from a mild polyclonal proliferation of B cells to B cell lymphomas.

• Congestive heart failure and arrhythmias

• Increased susceptibility to infections as a result of immunosuppression

• Renal dysfunction due to acute tubular necrosis or nephrotoxicity of the tacrolimus and cyclosporine

• Seizures are commonly associated with cyclosporine or tacrolimus toxicity.

Solution to Question 17:

The presence of endotheliitis confirms the presence of acute rejection of the liver transplant.

Rejection following liver transplantation can be acute or chronic.

Patients usually present with

• Malaise

• Fever

• Right upper quadrant pain

Laboratory investigations will show

• Elevated levels of serum bilirubin

• Aminotransferases

• Gamma-glutamyl transpeptidases

• Liver biopsy- for confirming the diagnosis and planning the treatment

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Liver biopsy of the acute rejection is characterized by the presence of mixed portal cellular infiltrate, bile duct injury, and endothelial inflammation (endotheliitis).

Chronic rejection is characterized by the presence of focal parenchymal necrosis, mononuclear cell infiltration, progressive cholestasis, and fibrosis. In some patients, this process will lead to the loss of intrahepatic bile ducts, which is called vanishing bile duct syndrome.

Solution to Question 18:

The given clinical scenario is suggestive of post-solid transplant infection most likely due to cytomegalovirus(CMV). It is the most common cause of post-solid organ transplants in India. Epstein-Barr virus (EBV), herpes, and varicella are also known to cause infection post-solid organ transplantation but to a lesser extent.

CMV disease in recipients can present with fever, leukopenia, thrombocytopenia, malaise, and arthralgia. They usually present within 1-4 months after transplant. The diagnosis of CMV infection is established by nucleic acid testing (NAT) using the polymerase chain reaction (PCR).

Management includes the use of anti-viral drugs like ganciclovir, valganciclovir, foscarnet, cidofovir and intravenous immunoglobulins.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 19:

Among the given options, autoimmune hepatitis has the highest chance of recurrence after liver transplantation.

Although some cases of sclerosing cholangitis and primary biliary cirrhosis recur, data for recurrence of autoimmune hepatitis is more supportive.

Alpha-1 antitrypsin deficiency does not recur after liver transplantation.

Sold by @itachibot

Hemochromatosis and Wilson's disease

Question 1:

What is the most common gene mutation associated with hemochromatosis?

a) H49D

b) C282Y

c) G97C

d) A480H

Question 2:

Patients with which of the following conditions are unlikely to develop iron overload?

a) Sideroblastic anemia

b) Non-alcoholic steatohepatitis

c) Porphyria cutanea tarda

d) Thalassemia minor

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following clinical manifestations of hemochromatosis is incorrectly matched with its mechanism of onset?

a) 1

b) 2

c) 3

d) 4

Question 4:

A 7-year-old boy undergoes genetic testing as his brother has been diagnosed with hereditary hemochromatosis. He is found to be homozygous for the C282Y mutation. Which of the following is the strongest indicator that he will develop symptoms of the disease?

a) Plasma iron >180 µg/dL

Sold by @itachibot

b) Total iron binding capacity >200 µg/dL

c) Transferrin saturation >50%

d) Serum ferritin level >1000 µg/L

Question 5:

A 47-year-old diabetic man is worried as the skin over his face, neck, and dorsum of hands has turned greyish over the past two months. He also complains of abdominal and joint pain. On examination, gynecomastia and hepatomegaly are noted. What is the treatment of choice for this condition?

a) Phlebotomy

b) Deferoxamine

c) Deferasirox

d) Liver transplantation

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 6:

A 30-year-old man with hereditary hemochromatosis has been started on weekly phlebotomy. All of the following features will improve with this treatment except:

a) Hepatomegaly

b) Skin pigmentation

c) Cardiac failure

d) Arthropathy

Question 7:

Which of the following is incorrect about Wilson's disease?

a) Gain-of-function mutation of ATP7B

b) AP1S1 gene is essential for normal functioning

c) Siblings of affected children have 1 in 4 risk of the disease

d) Mutation on chromosome 13

Question 8:

Sold by @itachibot

Which of the following are true regarding KF ring?

a) 1, 2 and 5 only

b) 1 and 5 only

c) 1, 2, 3, 4 and 5

d) 2, 3, 4 and 5 only

Question 9:

In patients with Wilson's disease, Kayser-Fleischer rings are most commonly associated with:

a) Hepatitis

b) Dystonia

c) Osteoarthritis

d) Cholelithiasis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A 22-year-old woman is being evaluated in the emergency department for a 3-day history of dark-colored urine and abdominal distension. On examination, she is conscious and oriented. Icterus is present. The following investigations are done. Which of the following would you suspect?

a) Hemochromatosis

b) Acute viral hepatitis

c) Primary biliary cirrhosis

d) Wilson's disease

Question 11:

A child presents with hepatitis and progressive neurological degeneration. A picture of his eye has been provided below. Among the following options, what would be the initial step regarding the investigations done for this child's diagnosis?

Sold by @itachibot

a) Serum ceruloplasmin levels

b) Serum copper levels

c) Enzyme assay

d) Karyotyping

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 12:

Which of the following findings would be seen in a patient with Wilson's disease?

a) 2, 3, 4

b) 1, 3, 4

c) 2, 4, 5

d) 2, 3, 5

Question 13:

A 15-year-old boy is brought with complaints of abnormal posture. His scholastic performance used to be above average but has deteriorated over the last 3 months. Dystonia and increased muscle tone are noted on clinical examination. The following picture is seen on ophthalmic examination. What is the gold standard for the diagnosis of this condition?

Sold by @itachibot



a) Serum copper

b) Serum ceruloplasmin

c) Urinary copper

d) Liver copper

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 14:

A pathologist is preparing a tissue sample from the liver biopsy of a patient with Wilson's disease. What stain will he use?

a) Perl stain

b) Rubeanic acid

c) Supravital stain

d) Schmorl's method

Question 15:

A 17-year-old girl was brought with involuntary, writhing movements of her limbs. On examination, she is anicteric with dystonia and hyperreflexia. Investigations revealed mildly elevated transaminases and low serum ceruloplasmin levels. Which of the following is the safest drug that can be initiated for this patient?

a) Zinc

b) D-Penicillamine

c) Trientine

Sold by @itachibot

d) Tetrathiomolybdate

Question 16:

For a patient with Wilson's disease, you would administer pyridoxine along with which of the following drugs?

a) Zinc

b) Penicillamine

c) Trientine

d) Tetrathiomolybdate

Question 17:

The Nazer's prognostic index for Wilson's disease includes all of the following parameters except:

a) Serum bilirubin

b) AST

c) ALT

d) Prothrombin time

Question 18:

Low serum copper levels attributed to ATP7A gene is due to:

a) Dubin-Johnson syndrome

b) Wilson's disease

c) Menke's disease

d) Gilbert's syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	b

Sold by @itachibot

2	d
3	d
4	d
5	a
6	d
7	a
8	b
9	b
10	d
11	a
12	b
13	d
14	b
15	c
16	b
17	c
18	c

If you purchased this ... you may have been ... else, ... d.

Detailed Explanations

Solution to Question 1:

The most common mutation seen in hemochromatosis is a cysteine to tyrosine substitution at position 282 (C282Y) in the HFE gene.

HFE mutations are the most common cause of hereditary hemochromatosis and are inherited as an autosomal recessive trait.

Solution to Question 2:

Thalassemia minor does not result in iron overload. Secondary hemochromatosis or iron overload is characterized by the deposition of iron in parenchymal cells.

Solution to Question 3:

Sexual dysfunction (decreased libido, impotence, amenorrhea) in hemochromatosis occurs as a result of hypogonadotropic hypogonadism. This is due to iron deposition in the pituitary gland, not testes.

Sold by @itachibot

Iron deposition occurs in the following sites:

Organ	Manifestation
Liver	Hepatomegaly
Skin	Slate-grey or bronze skin hyperpigmentation:increased melanin synthesisdue to iron deposition
Pancreas	pancreatic diabetes/type 3C diabetes
Pituitary	Hypogonadotropic hypogonadismSecondary hypothyroidism
Heart	Dilated cardiomyopathyCongestive heart failureArrhythmias
Joints	Arthropathy is due to:Iron depositionCalcium pyrophosphate deposition (pseudogout)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

Serum ferritin > 1000 µg/L is the strongest predictor of disease expression among individuals homozygous for the C282Y mutation.

Many patients with C282Y mutations are initially asymptomatic and do not show features of iron overload. In such patients, serum ferritin can indicate whether they are likely to develop the disease as ferritin levels reflect the total iron stores.

As ferritin levels may be high following inflammatory conditions, combined measurement of ferritin and transferrin saturation is done in a practical setting to avoid false positives. It is also repeated once inflammation subsides. If these tests are abnormal, genetic testing for HFE mutations is done.

Solution to Question 5:

This clinical scenario is suggestive of hemochromatosis. The treatment of choice is the removal of excess iron by phlebotomy.

Phlebotomy is an inexpensive, convenient, and safe method of treatment. In this process, 500 mL of blood is removed 1-2 times weekly which removes 200-250 mg of iron. Weekly phlebotomy should be continued until the serum ferritin level is ≤100 µg/L and every 3 months thereafter. It is contraindicated in severe anemia and hypoproteinemia.

Chelating agents bind to iron to form complexes that get excreted and are the drugs of choice. Drugs such as deferoxamine and deferasirox are commonly used. However, they are less effective

Sold by @itachibot

and more expensive than phlebotomy.

Liver transplantation is indicated either in end-stage liver disease or in refractory cases.

Supportive management includes avoiding iron-containing substances such as red meat, multi-vitamin tablets, etc. Alcohol should be completely avoided as it can aggravate the disease.

Solution to Question 6:

In hemochromatosis, removal of excess iron through phlebotomy has no effect on arthropathy and hypogonadism.

Solution to Question 7:

Wilson's disease is caused due to a loss-of-function mutation in the ATP7B gene on Chr.13q. The gene encodes a copper transporting ATPase (ATP7B) which has 2 functions:

- Maintains copper in bound form by binding copper to apoceruloplasmin
- Helps excrete free copper in bile.

Thus, an ATP7B deficiency leads to excess deposition of free copper within tissues. It is an autosomal recessive condition. So, siblings of affected children have a 1 in 4 chance.

Intracellular trafficking of ATP7B (Wilson's disease) and ATP7A (Menke's disease) genes are regulated by an adaptor protein encoded by the AP1S1 gene. Mutations in this gene result in a syndrome with features of both Wilson's and Menke's diseases. It is called MEDNIK syndrome:

- Mental retardation
- Enteropathy
- Deafness
- Neuropathy
- Ichthyosis
- Keratodermia.

Solution to Question 8:

Among the given options only statements 1 and 5 are true.

Kayser-Fleischer (KF) rings are seen in almost all patients of Wilson's disease with neurological involvement. It initially appears as a crescent in the superior pole of Descemet's membrane and gradually extends inferiorly, eventually becoming circumferential in shape.

KF rings occur due to copper deposition in the descemet's membrane. It is golden-brown to greenish-brown in color and can be best seen by slit-lamp examination.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Sold by @itachibot

The image below shows a KF ring :



While it is a characteristic sign of Wilson's disease (statement 2), it can also be seen in:

• Primary biliary cholangitis

• Primary sclerosing cholangitis

• Cholestasis

KF rings are present in almost 95% of patients of Wilson's disease with neurological/psychiatric symptoms compared to around 65% of those with hepatic presentations (statement 4).

They disappear on treatment with copper-chelating agents like d-penicillamine (while desferrioxamine is an iron chelating agent) and reappear with disease progression (statement 3).

Sold by @ita... If you purchased this from so... you may have been scammed. ...else,

Solution to Question 9:

The presence of the Kayser-Fleischer (KF) ring in Wilson's disease is most commonly associated with neuropsychiatric manifestations such as dystonia.

In majority of the cases, neuropsychiatric manifestations follow hepatic manifestations. It is due to the deposition of free copper in the basal ganglia (leading to bilateral degeneration and atrophy) and cortex. The features include:

• Dysarthria (most common): Irregular word spacing, hypophonic or spastic speech

• Dystonia: Vacuous smile i.e mask-like face with an open mouth

• Parkinsonism: Bradykinesia, postural instability, rigidity

• Subcortical dementia: Slow mental processing, memory loss

• Psychiatric symptoms: Loss of emotional control, depression, hyperactivity

• Decreased scholastic performance: Deterioration in handwriting, clumsiness

• Autonomic symptoms: Excessive sweating and salivation, sexual dysfunction

Sold by @itachibot

• Wing beating tremors (seen in advanced disease): High amplitude tremor that is prominent with arms abducted, elbows flexed and palms facing downward.

Solution to Question 10:

The given laboratory values indicate hepatic disease and hemolytic anemia (low Hb with high reticulocyte count). These lab findings along with clinical features point to a diagnosis of Wilson's disease.

The clinical features of Wilson's disease include:

Option A: Hemochromatosis presents with clinical features of iron overload. Serum ferritin and total iron are elevated.

Option B: Acute viral hepatitis presents with positive HbsAg, whereas anti-Hbs IgG indicates past infection. There is tender hepatomegaly with very high levels of transaminases (>1000 IU/L).

Option C: Primary biliary cirrhosis presents with jaundice, severe itching, and pale stools. There are high levels of direct bilirubin, ALP, and positive anti-mitochondrial antibodies.

System	Clinical features
Hepatic(most common)	JaundiceAbdominal painHepatomegalyFatigueAscitesVariceal bleedCoagulation defects High AST & ALT with low ALP
Neurological	DysarthriaDystoniaParkinsonismSubcortical dementiaDepressionWing beating tremors Decreased scholastic performance
Ophthalmic	Kayser-Fleischer (KF) ringsSunflower cataract
Others	Hemolytic anemia (Coombs negative)Acute pancreatitisFanconi's syndromeHyperparathyroidismCardiomyopathySpontaneous abortionsInfertility

Solution to Question 11:

The clinical features of hepatitis, progressive neurological degeneration, and the Kayser-Fleischer (KF) ring seen in the image are suggestive of Wilson's disease. The initial step regarding investigations done for the diagnosis is the measurement of serum ceruloplasmin levels.

Kayser-Fleischer (KF) ring is a golden to greenish-brown ring of deposition of copper in the Descemets membrane of the peripheral cornea. Patients with neurological manifestations

Sold by @itachibot

manifest with KF ring in >95% of cases.

Lab investigations in Wilson's disease would show:

The gold standard technique for diagnosis of Wilson's disease is percutaneous needle liver biopsy for measurement of hepatic copper levels >200 µg/g of dry weight.

Lab markers	Change in Wilson's disease	Reason
Serum ceruloplasmin	↓↓↓	ATP7B deficiency results in reduced bound copper
24h urinary copper	↑↑ (>100 µg/24h)	Excess free copper excreted in urine
Total serum copper	↓	Due to marked reduction in ceruloplasmin

Option B: The role of total serum copper in the diagnosis of Wilson's disease is limited. Measurement of serum ceruloplasmin and urinary copper is preferred.

Option C: ATP7B enzyme assays have shown promise in early studies as a neonatal screening test for Wilson's disease. However, it is not yet part of routine clinical practice.

Option D: Karyotyping is done for every asymptomatic 1st and 2nd degree relative of a patient with Wilson's disease. This helps differentiate the carriers (heterozygous) from the patients (homozygous) and can lead to early initiation of treatment before the onset of symptoms.

If you purchased this from someone, you may have been scammed.
Sold by @itachibot

Solution to Question 12:

The following findings will be seen in a patient with Wilson's disease:

• Decreased total serum copper

• Decreased serum ceruloplasmin

• Increased urinary copper

Normally, >90% of copper exists in bound form and <10% in free form. In Wilson's disease, despite elevated levels of free copper, the marked reduction in ceruloplasmin results in a net reduction of total serum copper i.e.,

↓total serum copper =↑free copper + ↓↓ceruloplasmin

Lab markers	Change in Wilson's disease	Reason
Serum ceruloplasmin	↓↓↓	ATP7B deficiency results in reduced bound copper
24h urinary copper	↑↑	Excess free copper excreted in urine

Sold by @itachibot

Lab markers	Change in Wilson's disease	Reason
Total serum copper	↓	Due to marked reduction in ceruloplasmin

Solution to Question 13:

This clinical scenario and image showing Kayser-Fleischer (KF) ring with sunflower cataract are suggestive of Wilson's disease. The gold standard for diagnosis is the estimation of liver copper following biopsy.

A copper level of >200 µg/g dry liver tissue is diagnostic for Wilson's disease. It is measured using neutron activation analysis or atomic absorption spectrometry.

Sunflower cataract is a rare finding of Wilson's disease. It occurs due to the deposition of copper in the center of the lens.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

In Wilson's disease, liver biopsy samples are stained using rubeanic acid to visualize the cytoplasmic copper.

Another stain used for copper is the rhodanine stain. However, copper staining is not reliable for diagnosis and a quantitative copper assay must be performed.

The given image shows the staining of copper by rubeanic acid.



Solution to Question 15:

Sold by @itachibot

This clinical scenario is suggestive of Wilson's disease with predominantly neurological features. The safest drug that can be initiated for this patient is trientine. This is because D-penicillamine can worsen neurological symptoms and would not be tolerated by this patient.

The copper chelating agents, D-penicillamine and trientine are the first-line drugs in the management of Wilson's disease. D-penicillamine is the recommended drug due to its availability and cost-effectiveness compared to trientine.

However, D-penicillamine can precipitate various adverse effects including aplastic anemia, myasthenia gravis and membranous nephropathy. It can also worsen neurological symptoms. Hence, trientine is administered in those who cannot tolerate D-penicillamine.

Elemental zinc is commonly used as a maintenance therapy along with chelating agents. It can also be used as first-line therapy in pregnancy.

Tetrathiomolybdate is currently under phase 2 clinical trials. It has been identified to be effective in patients with neuropsychiatric manifestations but is not approved for clinical use.

Liver transplantation is the definitive treatment for advanced liver disease.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

Penicillamine should always be co-administered with pyridoxine (25 mg/day) as it is an antagonist to pyridoxine.

Solution to Question 17:

ALT is not a part of Nazer's prognostic index.

Nazer's prognostic index is used to establish disease severity in Wilson's disease patients who present with hepatic decompensation.

Solution to Question 18:

Low serum copper level attributed to the ATP7A gene is due to Menke's disease (Option C).

Menke's disease or Menke-Kinky hair syndrome is an X-linked recessive condition occurring due to a mutation in the ATP7A gene on chromosome X. This causes impaired transport of dietary copper from the small intestine to the liver leading to low levels of copper-containing enzymes. It presents within 3 years of age and is characterized by:

• Depigmented kinky hair

• Failure to thrive

• Neurodegenerative symptoms like loss of developmental milestones, truncal hypotonia, epilepsyLoose, wrinkled skin

• Multiple fractures

Sold by @itachibot

• Poor visual acuity

• Recurrent infections.

Lab investigations show low serum copper (<65 µg/dL) and low ceruloplasmin(<20 mg/dl). The disease is fatal as it has no curative treatment. But, administration of L-histidine and copper chloride injections may be beneficial, if started early.

Aceruloplasminemia is an autosomal recessive disease characterized by tissue iron overload, mental deterioration, microcytic anemia, and low serum iron and copper concentrations.

Option B: Low serum copper and low ceruloplasmin is also seen in Wilson's disease, but is associated with the gene defect ATP7B on chromosome 13. It is an autosomal recessive disorder of copper metabolism with features of hepatolenticular degeneration and Kayser Fleischer ring.

Option A: Dubin Johnson Syndrome is an autosomal recessive disorder characterized by impaired canalicular excretion of bilirubin due to mutation in MRP-2 protein leading to conjugated hyperbilirubinemia.

Option D: Gilbert's Syndrome is characterized by decreased UDP glucuronyl transferase activity leading to unconjugated hyperbilirubinemia, which can be autosomal recessive or dominant.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed

Sold by @itachibot

Cirrhosis and Complications

Question 1:

Which of the following conditions is unlikely to lead to cirrhosis in a patient?

a) Alpha 1 antitrypsin deficiency

b) Cystic fibrosis

c) Nonalcoholic steatohepatitis

d) Hepatitis A

Question 2:

While examining a chronic alcoholic patient, you notice the given finding. Which of the following is false regarding it?



a) Occurs due to alterations in the metabolism of estrogen

b) Presence of even 1 lesion is abnormal

c) Commonly seen over the superior vena cava territory

d) Compression results in blanching followed by reappearance after release

Question 3:

If yo... by @itachibot ...s from someone else, ...scammed.

Sold by @itachibot

Clinically significant portal hypertension is defined as a hepatic venous pressure gradient greater than _____ mm Hg.

a) 3

b) 10

c) 5

d) 12

Question 4:

Which of the following is a posthepatic cause of portal hypertension?

a) Budd-Chiari syndrome

b) Schistosomiasis

c) Portal vein thrombosis

d) Banti's syndrome

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

Which of the following drugs is used to reduce the intrahepatic resistance in portal hypertension?

a) Propranolol

b) Carvedilol

c) Octreotide

d) Terlipressin

Question 6:

Ascitic tap done on a patient with abdominal distention revealed a SAAG of 1.4g/dL. What is the likely cause of his ascites?

a) Portal hypertension

b) Nephrotic syndrome

c) Peritoneal tuberculosis

d) Biliary leak

Sold by @itachibot

Question 7:

A patient presented with massive abdominal swelling. Paracentesis fluid was sent for lab analysis and results show serum protein to be 3g/dL and ascitic fluid protein to be 1g/dL. What is the likely diagnosis?

a) Congestive heart failure

b) Early Budd Chiari syndrome

c) Late Budd Chiari syndrome

d) Tuberculosis

Question 8:

A patient is said to have refractory ascites if the ascites does not resolve even after:

a) 1, 2 and 4

b) 4 and 5

c) 1, 3 and 5

d) 2 and 3

Sold by @itachibot
If you purchased this from someone else, you may have been scammed!

Question 9:

What is the minimum absolute neutrophilic count that should be present in the ascitic fluid to diagnose a patient with spontaneous bacterial peritonitis?

a) >250/µL

b) >150/µL

c) >300/µL

d) >200/µL

Question 10:

A cirrhotic patient admitted for ascites complained of abdominal pain on the 2nd day of his hospital stay. On examination, he was febrile. His lab reports are given below. Which of the following is false about his condition?

a) E.coli is the most commonly isolated organism

b) Prophylaxis against this condition is recommended in upper GI bleed

Sold by @itachibot

c) Third generation cephalosporins are used for treatment

d) Weekly antibiotic prophylaxis is recommended to prevent recurrence

Question 11:

Which of the following cannot be used to predict the risk of bleeding of oesophageal varices?

a) Severity of cirrhosis

b) Size of the varices

c) Location of the varices

d) Number of varices

Question 12:

In a patient with which of the following conditions would you not recommend a transjugular intrahepatic portosystemic shunt?

a) Ectopic varices

b) Refractory ascites

c) Budd Chiari syndrome

d) Hepatic encephalopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

A middle-aged man with hematemesis is rushed to the casualty. On examination, he is icteric, PR-122 /min and BP- 88/64 mmHg. The following finding is seen on his abdomen. Which of the following is false regarding his initial management?

Sold by @itachibot

a) pRBC is used to maintain Hb between 7-9g/dL

b) Prophylactic antibiotics given before endoscopy

c) IV octreotide is used to acheive hemostasis

d) Upper GI endoscopy is only performed after 12 hours

Question 14:

Which of the following is false about the given image?



a) Esophageal balloon should be deflated for about 5 mins every 6 hours

b) Gastric balloon should be inflated first

Sold by @itachi... purchased this from some... else, may have been scammed.

Sold by @itachibot

c) Pressure in the oesophageal balloon should not exceed 60 mm Hg.

d) Endotracheal intubation prior to tube insertion reduces the risk of pulmonary aspiration

Question 15:

A patient with hereditary hemochromatosis shows marked confusion and gross disorientation. He is drowsy but responds to pain and voice stimuli. What is the most likely grading of hepatic encephalopathy in this patient?

a) Grade 1

b) Grade 2

c) Grade 3

d) Grade 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

What is the earliest symptom of hepatic encephalopathy?

a) Mood changes

b) Asterixis

c) Altered sleep pattern

d) Disorientation

Question 17:

A cirrhotic patient was admitted following variceal bleeding. On the 3rd day of his hospital stay, he appears to be drowsy and confused. His wife reports that he had been awake throughout the previous night. On examination, flapping tremors were noticed. Which of the following is least likely to have precipitated his symptoms?

a) Antibiotics

b) Peritoneal tap

c) Diuretics

d) Urinary tract infection

Question 18:

Sold by @itachibot

Which of the following antibiotics could be used in the treatment of patients with hepatic encephalopathy?

a) Rifaximin

b) Azithromycin

c) Doxycycline

d) Rifampicin

Question 19:

Which of the following is not part of the diagnostic criteria of hepatorenal syndrome?

a) 50% increase in creatinine from baseline within 7 days

b) Rise in creatinine of 0.3 mg/dl in 48 hours

c) Presence of shock

d) No recent use of nephrotoxic agent

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 20:

Which of the following clinical features is most likely to be seen in patients with hepatopulmonary syndrome?

a) Trepopnea

b) Nocturnal dyspnea

c) Orthopnea

d) Platypnea

Question 21:

A 52-year-old man presents with oliguria, abdominal distension, and pedal edema. His medical records show long-standing cirrhosis with multiple hospitalisations for ascites and hepatic encephalopathy. His lab reports are given below. What is the best treatment for his condition?

a) Midodrine

b) Liver transplant

c) Intravenous albumin

d) TIPS

Sold by @itachibot

Question 22:

Which of the following conditions is not a MELD exception?

a) Hepatopulmonary syndrome

b) Portopulmonary hypertension

c) Primary hyperoxaluria

d) Chronic Hepatitis B

Question 23:

Which among the following is false?

a) Components of MELD (Model for end stage liver disease) scoring system are creatinine, bilirubin, International normalized ratio (INR), serum sodium (mEq/L).

b) Components of CTP(Child Turcotte Pugh) score are albumin, bilirubin, INR, ascites, encephalopathy

c) Components of CTP (Child Turcotte Pugh) are creatinine, bilirubin, International normalized ratio (INR)

d) MELD score is used to assess and prioritize patients awaiting liver transplantation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 24:

What is the Child-Turcotte-Pugh class for the patient who has a serum bilirubin 2.5 mg/dl, serum albumin 3 g/dl, prothrombin time increased by 5 seconds (INR = 2), no encephalopathy, and mild ascites?

a) Class A

b) Class B

c) Class C

d) Class D

Answer Key

Question No.	Correct Option

Sold by @itachibot

1	d
2	b
3	b
4	a
5	b
6	a
7	c
8	c
9	a
10	d
11	d
12	d
13	d
14	c
15	c
16	c
17	a
18	a
19	c
20	d
21	b
22	d
23	c
24	b

Detailed Explanations

Solution to Question 1:

Hepatitis A infection is an acute self-limiting condition that does not lead to cirrhosis.

Cirrhosis is a late stage of chronic liver disease pathologically characterized by hepatic fibrosis, distortion of hepatic architecture, and formation of regenerative nodules.

The causes of cirrhosis are:

• Chronic viral hepatitis - Hepatitis B and C

• Alcoholic liver disease

• Nonalcoholic fatty liver disease

Sold by @itachibot

• Autoimmune hepatitis

• Biliary cirrhosis -

• Primary biliary cholangitis

• Primary sclerosing cholangitis

• Cardiac cirrhosis

• Inherited metabolic liver diseases -

• Cystic fibrosis

• Hemochromatosis

• Wilson's disease

• α1 Antitrypsin deficiency

• Cryptogenic cirrhosis

• Drugs - methotrexate, isoniazid

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

The given image shows spider telangiectasia and the presence of >2 to 3 of these are considered abnormal.

Spider telangiectasias are dilated arterioles with a prominent central arteriole with radiating vessels appearing in the shape of a spider. They are generally seen in the superior vena cava territory. Compression results in blanching followed by reappearance after the release of pressure. They occur due to a hyperestrogenic state which is caused by defective estrogen metabolism in liver cirrhosis.

Clinical features seen in cirrhosis include:

• Spider telangiectasias

• Palmar erythema

• Terry's nails - pallor of the proximal nail bed

• Clubbing

• Gynaecomastia

• Caput medusae - dilated abdominal veins.

• Parotid enlargement

The image below shows Terry's nails.

Sold by @itachibot

Sold by @itachi... else, If you purchased this from some... you have been scammed.

Solution to Question 3:

Clinically-significant portal hypertension is defined as hepatic venous pressure gradient (HVPG) ≥10 mm Hg.

HVPG is the difference between the wedged hepatic venous pressure (WHVP) and the free hepatic venous pressure (FHVP). It represents the pressure difference between the portal vein and IVC.

Interpretation of HVPG values:

HVPG value	Interpretation
1-5 mm Hg	Normal
>5 mm Hg	Portal hypertension
≥10 mm Hg	Clinically significant portal hypertension, varices can develop
≥12 mm Hg	Risk of variceal bleeding and ascites

Solution to Question 4:

Budd-Chiari syndrome is a posthepatic cause of portal hypertension.

Budd Chiari syndrome is a congestive hepatopathy characterized by the obstruction of hepatic venous outflow. The most common cause is due to hepatic vein thrombosis. Obstruction of two or

Sold by @itachibot

more major hepatic veins causes Budd-Chiari syndrome. It is characterized by hepatomegaly, ascites, and abdominal pain.

Solution to Question 5:

Carvedilol is used to reduce the intrahepatic resistance in portal hypertension.

The pharmacologic management of portal hypertension includes:

• Drugs that reduce the portal blood flow

• Non-selective β blockers - Propranolol and nadolol are preferred

• Somatostatin and its analogs - Octreotide, lanreotide, and vapreotide

• Vasopressin and analogs - Terlipressin

• Drugs that decrease the intrahepatic resistance

• Carvedilol - due to its α blockade property

Solution to Question 6:

The serum ascites albumin gradient of ≥1.1g/dL is suggestive of portal hypertension.

SAAG helps distinguish ascites caused by portal hypertension from other causes of ascites.

Solution to Question 7:

The likely diagnosis in the given clinical scenario with serum-ascites albumin gradient (SAAG) >1.1g/dL [calculated as serum albumin level - ascitic fluid albumin level, which in this case is 3-1=2g/dL] and ascitic fluid protein <2.5 g/dL is late Budd Chiari syndrome.

Serum-ascites albumin gradient (SAAG) can be used to differentiate between ascites due to portal hypertension and non-portal hypertensive ascites. It represents the pressure within the hepatic sinusoids and correlates with the hepatic venous pressure gradient.

It is given by the formula: SAAG = Serum albumin level - Ascitic fluid albumin level

Interpretation of SAAG:

• SAAG ≥1.1 g/dL indicates that the ascites is due to increased pressure in the hepatic sinusoids. It reflects the presence of portal hypertension.

• SAAG <1.1 g/dL indicates that the ascites is not related to portal hypertension. It may be due to infection or malignancy.

In cases with high-SAAG (≥1.1), ascitic protein level analysis further helps in identifying the etiology:

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

• ≥2.5 g/dL indicates that the hepatic sinusoids are normal and are allowing the passage of protein into the ascites as in early Budd-Chiari syndrome.

• <2.5 g/dL indicates that the hepatic sinusoids have been damaged and scarred and no longer allow passage of protein as in late Budd-Chiari syndrome

Solution to Question 8:

Refractory ascites is defined as the inability to mobilize ascitic fluid despite:

• Compliance with dietary sodium restriction

• Administration of maximum doses of diuretics -

• 400mg of spironolactone per day and

• 160mg of furosemide per day

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

The diagnosis of spontaneous bacterial peritonitis(SBP) is made when the fluid sample has an absolute neutrophil count >250/µL.

SBP is the spontaneous infection of the ascitic fluid without an intraabdominal source. Infection with E.coli and other gut bacteria is most commonly seen. However, gram-positive bacteria like Streptococcus viridans, Staphylococcus aureus, and Enterococcus sp., can also be found. Abdominal paracentesis is the investigation of choice. It can be managed with antibiotic therapy.

Solution to Question 10:

The given clinical scenario along with ANC ≥ 250 cells/µl in ascitic fluid is suggestive of spontaneous bacterial peritonitis (SBP). Daily antibiotic prophylaxis is recommended in patients who have had an episode of SBP to prevent a recurrence.

SBP is the spontaneous infection of the ascitic fluid without an intraabdominal source. The most common causative organisms are E.coli and other gut bacteria. The diagnosis of SBP is made when the ascitic fluid has an absolute neutrophil count >250/µl.

According to the latest recommendations, weekly antibiotic prophylaxis is no longer recommended. For patients with low protein ascites (<1.5 g/dL [15g/L]) along with either impaired renal function or liver failure, we use prolonged outpatient trimethoprim-sulfamethoxazole (one double-strength tablet daily). Alternatives include ciprofloxacin 500 mg per day or norfloxacin 400 mg per day.

For patients on chronic trimethoprim-sulfamethoxazole therapy, renal function should be monitored by checking serum electrolytes, blood urea nitrogen (BUN) and creatinine every one to six months depending on the stability of patient. If there is worsening renal function,

Sold by @itachibot

trimethoprim-sulfamethoxazole should be discontinued.

Solution to Question 11:

The risk of bleeding in a patient with oesophageal varices cannot be predicted based on the number of varices present.

The factors that are used to predict the risk of variceal bleeding are:

• Severity of cirrhosis (Child's class, MELD score)

• History of a previous variceal bleed

• Location of the varices - varices with a thinner layer of supporting tissue are likely to bleed more

• Size of the varices - larger varices have a higher risk of bleeding

• Endoscopic appearance of varices - Presence of red wale sign, hematocystic spots, diffuse erythema, and cherry red spots are associated with increased risk of bleeding

• Variceal pressure

Solution to Question 12:

Transjugular intrahepatic portosystemic shunt (TIPS) is not recommended in hepatic encephalopathy as it can aggravate its symptoms.

TIPS creates a low-resistance channel between the hepatic vein and an intrahepatic branch of the portal vein. It uses a percutaneous approach under angiographic guidance. This reduces portal pressure.

It is used in the management of:

• Oesophageal varices - in patients who fail medical or endoscopic management and have a poor surgical outcome

• Refractory ascites

• Gastric varices

• Ectopic varices

• Portal hypertensive gastropathy

• Hepatorenal syndrome

• Budd Chiari syndrome

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Solution to Question 13:

The given clinical scenario along with the image showing caput-medusae is suggestive of acute esophageal variceal bleeding due to portal hypertension. Upper GI endoscopy is performed within 12 hours after fluid resuscitation in variceal bleeding.

The management of esophageal variceal bleeding involves:

- Resuscitation with IV fluids, airway protection, and oxygen
- Blood transfusion to maintain Hb between 7-9g/dL
- Antibiotic prophylaxis with ceftriaxone before endoscopy
- An attempt at hemostasis with octreotide or terlipressin or vasopressin
- Upper GI endoscopy within 12 hours after fluid resuscitation to confirm the source of the bleed
- Endoscopic variceal ligation of sclerotherapy
- In case, of failure or rebleeding - TIPS

Nonselective beta-blockers like propranolol and nadolol are used in the prophylaxis of variceal bleeding.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Solution to Question 14:

The given image shows Sengstaken-Blakemore tube. Pressure in the oesophageal balloon should not exceed 45 mm Hg.

It is used for balloon tamponade in variceal hemorrhage when medical therapy fails or is not available. Initially, only the gastric balloon should be inflated, with 200– 250 mL of air, as this will usually control bleeding. The oesophageal balloon is used in continued bleeding, it should be deflated for about 5 minutes every 6 hours to avoid oesophageal mucosal damage.

Endotracheal intubation prior to tube insertion reduces the risk of pulmonary aspiration.



If you purchased this from someone else, they have been scammed.
Sold by @itachibot

Solution to Question 15:

Gross disorientation with an intact response to pain and voice suggest grade 3 hepatic encephalopathy.

Hepatic encephalopathy is the alteration of mental status and cognitive function occurring in liver failure. This occurs due to the failure of removal of gut-derived neurotoxins like ammonia by liver. These neurotoxins enter the brain leading to the development of symptoms.

Solution to Question 16:

The earliest symptom of hepatic encephalopathy is an alteration in sleep pattern.

Other signs and symptoms of hepatic encephalopathy include:

- Asterixis also called the flapping tremor
- Mood changes and inappropriate behavior
- Disorientation and confusion

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 17:

The given clinical scenario is suggestive of hepatic encephalopathy. Antibiotics are unlikely to precipitate hepatic encephalopathy.

The common precipitating factors of hepatic encephalopathy are :

- Gastrointestinal bleeding
- Infections like SBP or UTIs
- Peritoneal tap
- Constipation
- Renal failure
- Hypovolemia
- Hypoxia
- Drugs - sedatives, antidepressants, diuretics
- Hypoglycemia
- Hypokalemia

Solution to Question 18:

Rifaximin can be used in the treatment of hepatic encephalopathy. It lowers blood ammonia levels.

Sold by @itachibot

Treatment of hepatic encephalopathy includes:

• Correction of precipitating factors

• Lactulose - decreased GI transit time due to cathartic actions leading to increased elimination of ammonia

• Rifaximin - lowers the blood ammonia levels

• L-ornithine L-aspartate - stimulate the urea cycle, with resulting loss of ammonia

Solution to Question 19:

The absence of shock is a part of the diagnostic criteria for hepatorenal syndrome.

Hepatorenal syndrome is a progressive renal failure in patients with acute or chronic liver disease.

The diagnostic criteria for hepatorenal syndrome include:

• Cirrhosis with ascites

• Diagnosis of acute kidney injury (AKI)

• Increase in serum creatinine of ≥ 0.3mg/dL in 48 hours, OR

• 50% increase in serum creatinine from baseline within 7 days

• Absence of shock

• No evidence of recent use of nephrotoxic agents

• Absence of intrinsic renal disease

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 20:

Among the given options, platypnea is most likely to be seen in patients with hepatopulmonary syndrome.

Hepatopulmonary syndrome is characterized by a triad of liver disease, hypoxemia, and intrapulmonary vascular dilatations. Most patients have dyspnea on exertion. The classic clinical manifestations are:

• Platypnea - Increase in dyspnea when the patient moves from supine to upright position

• Orthodeoxia - Decrease in arterial oxygen tension when the patient moves from supine to upright position

The hypoxemia is due to a ventilation-perfusion mismatch which causes an increased alveolar-arterial (A-a) gradient. The most sensitive test for diagnosis is contrast bubble echocardiography.

Liver transplantation is the definitive treatment of this condition following which the symptoms subside.

Sold by @itachibot

Solution to Question 21:

The given clinical scenario is suggestive of hepatorenal syndrome. The best treatment modality for it is liver transplantation.

Hepatorenal syndrome is characterised by a rise in:

- AST
- ALT
- Total bilirubin
- Serum creatinine
- Blood urea nitrogen (BUN).

Management of hepatorenal syndrome include:

- Norepinephrine
- Octreotide
- Terlipressin
- Midodrine
- Intravenous albumin
- Liver transplantation- definitive treatment

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

Chronic Hepatitis B is not a MELD exception.

MELD (Model for End-stage Liver Disease) score is used to prioritize patients awaiting liver transplantation. However, there are certain diseases that impair patient survival and are not accounted for in the MELD score. Patients with these diseases are given upgraded MELD points to accurately represent the mortality risk and these conditions are called MELD exceptions.

The MELD exceptions include:-

- Portopulmonary hypertension
- Hepatocellular carcinoma
- Familial amyloid polyneuropathy
- Primary hyperoxaluria (necessitating liver-kidney transplantation)
- Cystic fibrosis with liver disease
- Highly selected cases of hilar cholangiocarcinoma.
- Hepatopulmonary syndrome
- Hepatic artery thrombosis 7–14 days post-liver transplantation.

Sold by @itachibot

Solution to Question 23:

The Child Turcotte Pugh (CTP) score does not include creatinine.

The components of the CTP score are:

• Albumin

• Bilirubin

• International normalized ratio (INR)

• Ascites

• Encephalopathy

The Model for End-stage liver disease (MELD 3.0) is used to prioritize patients awaiting liver transplantation and includes:

• Female sex

• Bilirubin

• Albumin

• Serum sodium (mEq/L)

• INR (International Normalized ratio)

• Creatinine

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 24:

The given patient belongs to Child-Turcotte-Pugh class B.

The score can be calculated as follows:

Serum bilirubin-2.5mg/dl: 2

Serum albumin-3g/dl: 2

Prothrombin time-5 seconds or INR-2: 2

No encephalopathy: 1

Mild ascites: 2

Therefore, the Child-Turcotte-Pugh score is (2+2+2+1+2)= 9 which is included in Class B.

Sold by @itachibot

Peptic Ulcer Disease and Related Disorders

Question 1:

Which of the following is an epithelial mucosal defense mechanism of the gastric mucosa?

a) Layer of mucous

b) Alkaline tide

c) Microvascular bed

d) Restitution

Question 2:

A 50-year-old woman complains about a burning pain in her chest. She reports using NSAIDs for her constant knee pain. This drug impairs her gastric mucosal defense mechanism primarily by:

a) Irritant effect on gastric mucosa

b) Stimulating acid production

c) Inhibiting prostaglandin synthesis

d) Inhibiting nitric oxide synthesis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

All of the following are true about the risk factors of NSAIDs associated peptic ulcer disease except:

a) Smoking increases the risk

b) H pylori infection increases the risk

c) Even low dose of a NSAID can cause ulcers

d) Concomitant use of glucocorticoids decreases the risk

Question 4:

What pattern of gastritis is most common in patients with H.pylori-related duodenal ulcers?

a) Autoimmune gastritis

Sold by @itachibot

b) Non-atrophic pangastritis

c) Granulomatous gastritis

d) Antral predominant gastritis

Question 5:

A patient presents with frequent epigastric pain with burning sensation for the past 2 years. The pain usually occurs 2-3 hours after a meal and is relieved by taking food. He also says that he sometimes wakes up at night because of severe pain. The most probable cause for his symptoms is:

a) Gastric ulcer

b) Duodenal ulcer

c) Esophagitis

d) GERD

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 6:

Which of the following is true about gastric ulcers?

a) Usually occur in 4th decade

b) Rarely malignant

c) Low basal acid production

d) High stimulated acid production

Question 7:

H pylori causes gastric mucosal injury due to the presence of:

a) Cag A

b) VacA

c) LPS

d) Hop proteins

Question 8:

Sold by @itachibot

You are a first-year resident. You are asked to prescribe an H.pylori eradication regimen to a patient with peptic ulcer disease. Which of the following drugs will you not include in it?

a) Bismuth subsalicylate

b) Erythromycin

c) Amoxicillin

d) Metronidazole

Question 9:

In which of the following patients presenting with epigastric pain and discomfort is H.pylori eradication therapy not indicated:

a) 46-year-old man with a family history of gastric cancer in the father

b) 40-year-old woman with rheumatoid arthritis on NSAIDs for 5 years

c) 58-year-old man with 35-pack years history of smoking

d) 36-year-old man with GERD on PPI's for 3 months

Question 10:

Eradication of H.pylori infection in patients with peptic ulcer disease is determined by?

a) Serology

b) Urea breath test

c) Biopsy urease test

d) Culture

Question 11:

H.pylori increases the risk of:

a) GERD

b) Barrett's esophagus

c) Adenocarcinoma of cardia of stomach

d) Gastric MALT lymphoma

Question 12:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

The most common complication in patients with peptic ulcer disease is:

a) Perforation

b) GI bleeding

c) Gastric outlet obstruction

d) Gastrocolic fistula

Question 13:

A 47-year-old woman presented to the ER with an altered sensorium. Her husband reported that she has had abdominal pain, vomiting, and constipation for the past 4 days. She has been self-medicating for gastritis. Her lab investigations are as shown. Which of the following drugs is she most likely on that could have caused her symptoms?

a) Aluminium hydroxide

b) Magnesium hydroxide

c) Calcium carbonate

d) Proton pump inhibitors

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 14:

A patient is admitted to the ICU after he suffered trauma to his head. This patient is prone to develop what type of gastric ulcers?

a) Curling ulcers

b) Cushing ulcers

c) Stress ulcers

d) Refractory ulcers

Question 15:

You would diagnose a patient to have a refractory gastric ulcer if the ulcer fails to heal after:

a) 12 weeks

b) 8 weeks

c) 10 weeks

d) 14 weeks

Sold by @itachibot

Question 16:

A middle-aged woman presents with acidity, abdominal pain, diarrhea, and weight loss for the past few months. Her symptoms are not relieved by taking antacids. An endoscopic evaluation shows ulcers on the second part of the duodenum. Which of the following conditions should be suspected in this patient?

a) Stress ulcer

b) H. pylori infection

c) Malignant duodenal ulcer

d) Zollinger Ellison syndrome

Question 17:

A 35-year-old man presented with recurrent duodenal ulcers with poor response to medical therapy. He had a history of hyperparathyroidism and transsphenoidal prolactinoma resection. Which among the following is the next appropriate step?

a) Urea breath test

b) Secretin test

c) Serum gastrin levels

d) Endoscopic ultrasound

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

Which of the following investigations employs somatostatin analogues to detect gastrinomas?

a) CECT

b) PET/CT with Ga-DOTATATE

c) OctreoScan

d) Endoscopy

Question 19:

Which of the following viruses cause infectious gastritis in immunocompromised patients?

a) Cytomegalovirus

b) Norovirus

Sold by @itachibot

c) Enterovirus

d) Coxsackie virus

Question 20:

An anemic patient on evaluation showed macrocytes with hypersegmented neutrophils on peripheral smear. Serology revealed anti-parietal cell antibodies. Which of the following is most likely to be associated with her condition?

a) Zollinger Ellison syndrome

b) VIPoma

c) Type A gastritis

d) Type B gastritis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

An 84-year-old woman living in an assisted care facility is rushed to the hospital with an upper GI bleed. On arrival, she is febrile and the nurse notes a sudden spurt of blood from her nasogastric tube. The following investigations are done. Which of the following is responsible for this patient's condition?

a) H.pylori

b) E.coli

c) Cytomegalovirus

d) Herpes simplex virus

Answer Key

Question No.	Correct Option
1	d
2	c
3	d
4	d
5	b
6	c

Sold by @itachibot

7	b
8	b
9	d
10	b
11	d
12	b
13	c
14	b
15	a
16	d
17	c
18	c
19	a
20	c
21	b

Sold by ... If you purchased this from ... else, you may have been scam... d.

Detailed Explanations

Solution to Question 1:

Restitution is an epithelial mucosal defense mechanism. It is the process by which healthy surface epithelial cells from the adjoining areas migrate to restore a damaged region of the gastric mucosa.

The gastric mucosa has well-established mucosal defense mechanisms against various assaults like HCl, pepsin, alcohol, etc. These include:

• Pre-epithelial defense: A layer of mucous over the surface epithelium. This is secreted by the surface epithelium and contains HCO_3- which maintains neutral pH at the cell surface

• Epithelial defense: It is constituted by the surface epithelial cells and their intercellular tight junctions that perform the following functions

• Act as a physical barrier

• Restitution is the migration of healthy cells to damaged areas

• Rapid renewal of surface epithelial layer

• Generating factors that prevent cellular damage such as heat shock proteins

• Sub-epithelial defense:

• Rich microvascular bed of the gastric submucosa that provides HCO_3- and washes away the toxic metabolites.

Sold by @itachibot

• Alkaline tide- It is caused by the release of HCO_3^- into the gastric capillaries by the parietal cells. This HCO_3^- helps in the quick neutralization of HCl.

Solution to Question 2:

The NSAIDs impair the mucosal defense mechanism primarily by inhibiting prostaglandin synthesis.

NSAIDs also decrease nitric oxide synthesis and have an irritant effect on the gastric mucosa but inhibition of COX enzymes/ prostaglandin synthesis is the most prominent factor.

Almost all the mucosal defense mechanisms are primarily mediated by prostaglandins synthesized by the gastric mucosal cells. Apart from the prostaglandins, the other factors that play a role in mucosal defense include:

• Nitric oxide (NO)- Causes vasodilation thereby increasing the mucosal blood flow.

• Several growth factors like EGF, TGF, FGF- Play a role in epithelial regeneration and restitution.

Solution to Question 3:

Concomitant use of glucocorticoids with NSAIDs increases the risk of peptic ulcer disease.

Other risks factors for NSAID-induced PUD include:

• H pylori infection

• Serious or multisystem disease

• Alcohol and smoking

• Concomitant use of the following drugs with NSAIDs increases the risk of PUD and its complications:

• Glucocorticoids

• Anticoagulants

• Clopidogrel

• Multiple NSAIDs

PUD may occur even with low doses of NSAIDs. No dose of NSAIDs is considered completely safe.

Solution to Question 4:

Antral predominant gastritis is associated with duodenal ulcers. Antral predominant gastritis is associated with increased acid production by the parietal cells. This subjects the duodenum to increased acid exposure, leading to the formation of duodenal ulcers.

H. pylori infection may result in the following patterns of gastritis:

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Sold by @itachibot

• Non-atrophic pangastritis- mostly asymptomatic and sometimes associated with MALToma.

• Corpus predominant gastritis- affects the corpus and causes atrophy. This leads to the loss of acid-producing parietal cells that are primarily found in the corpus. Hence, it is associated with decreased acid production. This is also a predisposing factor for the development of gastric cancer.

• Antral predominant gastritis- spares the parietal cells. Hence, it is associated with increased acid production.

Option A: Autoimmune gastritis is caused due to autoantibodies against the parietal cells. It is not associated with H. pylori.

Option C: Granulomatous gastritis is characterized by the presence of granulomas. It is associated with Crohn's disease and sarcoidosis.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 5:

Epigastric pain with a burning sensation that usually occurs 2-3 hours after a meal and is relieved by taking food suggestive of duodenal ulcer.

The following patterns of pain are usually seen in peptic ulcer disease:

• Duodenal Ulcer- Pain occurs 1.5 to 3 hours after a meal and is frequently relieved by antacids or food. Pain may awaken the patient from sleep.

• Gastric Ulcer- Pain/discomfort may be precipitated by food. Nausea and weight loss are common.

• Perforated ulcer- The bed of the ulcer tunnels through the entire thickness of the gastric wall and opens into the peritoneum. This presents with sudden onset severe abdominal pain and leads to peritonitis. Radiography shows gas under the diaphragm.

• Penetrating ulcer- It is a type of perforation where the ulcer tunnels into an adjacent organ like the pancreas/liver. In this case, the pain is no longer relieved by food or antacids and might radiate to the back.

Many patients with peptic ulcers, especially elderly patients, may not have these symptoms and may directly present with a complication like bleeding or perforation.

Solution to Question 6:

Gastric ulcers are associated with low basal acid production and low stimulated acid production.

Gastric acid secretion occurs in two patterns:

• Basal acid output- Occurs in a circadian pattern. It is mediated by ACh from the vagus and histamine.

• Stimulated acid production- Occurs in response to food. It is mediated by gastrin in addition to other factors.

Sold by @itachibot

Gastric Ulcer	Duodenal Ulcer
Less common	More common
Occurs later in life, with a pe ak incidence in the 6th decad e	Occurs earlier in life
Basal and stimulated acid pr oduction is low/normal, exce pt in type 3 gastric ulcers	Basal and nocturnal acid pro duction is increased
Can representmalignancyand should be biopsied	Rarely malignant
Most common site:Antrum(di stal to acid-secreting mucosa) on the lesser curvature	Most common site: 1st part o f the duodenum (within 3cm s from the pylorus)
Painafterfood intake	Painbefore food intake, andr eliefontaking food

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

H.pylori produces vacuolating exotoxin (Vac A) which causes gastric mucosal injury. Vac A inhibits CD4+ T-cells and disrupts the function of B cells, macrophages & mast cells.

Cag A positive strains of H pylori cause an increased risk of gastric cancer.

Solution to Question 8:

Erythromycin is not effective in H pylori eradication.

Note:

Vonoprazan, a potassium-competitive acid blocker, has been recently approved by the FDA for the treatment of peptic acid-related disorders.

Solution to Question 9:

The need for H pylori eradication therapy in patients with GERD on long-term PPIs is not well established and is controversial.

H pylori eradication therapy is offered to patients diagnosed with peptic ulcer disease (PUD) as it greatly decreases the recurrence of ulcers. In addition to this, it should also be offered to patients at increased risk of PUD and gastric cancer such as:

• Patients with long-term NSAIDs usage. Eg. Rheumatoid arthritis

• Patients taking gastric acid inhibitors for >1 year

Sold by @itachibot

• Patients with increased exposure to risk factors of gastric cancers. Eg. heavy smokers, exposure to dust, coal

• Patients with previous history/family history of gastric neoplasm

Solution to Question 10:

Eradication of H. pylori infection can be determined by a urea breath test.

A stool antigen test can also be employed.

Solution to Question 11:

H.pylori increases the risk of gastric MALT lymphoma.

H.pylori is protective against the following:

• GERD

• Barrett's esophagus

• Adenocarcinoma of esophagus

• Adenocarcinoma of the cardia of the stomach

H. pylori are inversely related to gastroesophageal reflux disease (GERD) and its sequelae, which include Barrett's esophagus and esophageal adenocarcinoma.

H.pylori strains positive for Cag A may be particularly protective against the development of esophageal adenocarcinoma. Thus, colonization with H. pylori, particularly Cag A strains, may be protective against the more severe forms of GERD.

Eradication of H. pylori is associated with mild worsening of gastroesophageal reflux disease (GERD) in patients with corpus-dominant or pangastritis. It improves symptoms in those with antral-dominant gastritis.

Solution to Question 12:

GI bleeding is the most common complication of peptic ulcer disease.

Other complications:

• Perforation- second most common

• Gastrocolic fistulas

• Gastric outlet obstruction

• Rarely malignancy

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Solution to Question 13:

The given clinical features indicating hypercalcemia and investigations showing renal insufficiency, metabolic alkalosis are suggestive of milk-alkali syndrome. Long-term use of calcium carbonate can lead to this condition.

In susceptible individuals, there is a loss of negative feedback, leading to excess calcium absorption when there is excess intake. It results in the following triad of clinical features:

• Hypercalcemia

• Alkalosis- due to HCO_3 reabsorption caused by PTH suppression, natriuresis, and water depletion

• Renal failure- due to damage to renal tubules

It is reversible once ingestion of antacid is stopped. However, in severe cases, a chronic form called Burnett's syndrome can develop. It is associated with irreversible renal damage.

Solution to Question 14:

The gastric ulcers associated with head trauma are Cushing ulcers. Head trauma leads to vagal stimulation which in turn leads to increased gastric acid production.

Option A: Curling ulcers are seen in severe burns.

Option C: Stress ulcers are seen in conditions like shock, sepsis, etc. They are thought to be caused by local ischemia and splanchnic vasoconstriction.

Option D: Refractory ulcers are those that do not heal even after adequate medical therapy.

Solution to Question 15:

A gastric ulcer is said to be refractory if it fails to heal after 12 weeks.

A refractory duodenal ulcer is that which fails to heal after 8 weeks.

Management of refractory PUD:

• Rule out other diseases or causes

• High dose PPI

• Surgery

Solution to Question 16:

The given scenario of symptoms not improving with antacids, associated diarrhea, and the unusual location of ulcers are highly suggestive of Zollinger Ellison syndrome.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Zollinger Ellison syndrome (ZES) is a peptic ulcer disease caused by unregulated gastrin release from a neuroendocrine tumor/gastrinoma. About 60% of the tumors are malignant, hence, it is important to identify ZES.

The clinical presentation of ZES is very similar to that of peptic ulcer disease. The features which help in suspecting ZES include:

• Unusual locations- the second part of the duodenum and beyond

• Refractory to medical therapy

• Recurrence after acid-reducing surgery

• Presenting with complications like bleeding and perforation

• Not associated with H. pylori or NSAID ingestion

Patients of ZES often have associated diarrhea and involvement of the esophagus (esophagitis, ulceration, or stricture).

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

The given clinical scenario is suggestive of Zollinger Ellison syndrome (ZES). The next appropriate step to evaluate the patients for ZES is by measuring the serum gastrin levels. History of hyperparathyroidism and transsphenoidal prolactinoma resection indicates MEN1 syndrome. ZES is commonly associated with MEN1.

The normal gastrin levels are <150 pg/mL. An increase of >10 times the normal levels is highly suggestive of ZES.

Secretin test is a gastrin provocative test. It is done when the gastrin levels and the acid secretory studies are inconclusive for ZES. It is the most sensitive and specific gastrin provocative test for the diagnosis of gastrinoma.

Endoscopic ultrasound is done to localize the tumor. It is done after the diagnosis of ZES has been established.

Solution to Question 18:

OctreoScan detects gastrinomas using their property of uptake of somatostatin analogues.

OctreoScan detects endocrine tumors that express receptors for somatostatin in conditions such as Zollinger-Ellison syndrome (ZES) and multiple endocrine neoplasia (MEN) syndromes.

Other modalities used for localizing the tumor in ZES include:

• Endoscopic ultrasound (EUS)

• PET/CT with Ga-DOTATATE is the investigation of choice when available.

Note: PET/CT with Ga-DOTATATE is the gold-standard test for localizing tumors in pheochromocytoma.

Sold by @itachibot

Solution to Question 19:

Viruses that cause infectious gastritis in immunocompromised individuals include cytomegalovirus and Herpes simplex virus.

Solution to Question 20:

The given peripheral smear findings and serology are suggestive of pernicious anemia. It is usually seen in type A gastritis.

Type A autoimmune gastritis is caused by circulating antibodies against the parietal cells. The destruction of parietal cells leads to the deficiency of intrinsic factor that is necessary for the absorption of vitamin B12. The resulting B12 deficiency causes pernicious anemia.

Solution to Question 21:

The given clinical scenario along with her endoscopy findings are suggestive of phlegmonous gastritis. It is caused due to E.coli.

Phlegmonous gastritis is a potentially life-threatening disorder characterized by marked and diffuse acute inflammatory infiltrates of the entire gastric wall accompanied by necrosis. It usually affects the immunocompromised and elderly individuals.

Organisms associated with phlegmonous gastritis are:

- E.coli
- Streptococci
- Staphylococci
- Proteus
- Haemophilus species

Other uncommon forms of gastritis are:

- Eosinophilic gastritis: Presents as intestinal obstruction
- Russel's body gastritis: Pseudotumor on endoscopy
- Granulomatous gastritis is seen in Crohn's disease
- Lymphocytic gastritis is seen in celiac disease

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Evaluation of Diarrhea

Question 1:

A patient presents with chronic diarrhea, bloating sensation, and weight loss. History is significant for stools that are bulky, pale, foul-smelling, and difficult to flush. Further testing reveals fecal fat excretion of 8.2g/day. Which of the following would be a differential diagnosis?

a) Short bowel syndrome

b) Primary lactase deficiency

c) Pernicious anemia

d) Wilson's disease

If you purchased this from someone else, you may have been scammed. Sold by @Itachibot

Question 2:

A 22-year-old man presented with diarrhea and intolerance to dairy products. On investigation, he was found to have a lactase deficiency. Which of the following agents is least likely to cause symptoms of lactose intolerance?

a) Condensed Milk

b) Skimmed Milk

c) Yogurt

d) Ice Cream

Question 3:

Which of the following is incorrect about lactose intolerance?

a) It may be seen in celiac disease

b) It can mimic irritable bowel syndrome

c) Delayed gastric emptying decreases the symptoms

d) Rapid small-intestinal transit time decreases the symptoms

Question 4:

Sold by @itachibot

Which of the following is unlikely to be present in a patient with large bowel diarrhea?

a) Steatorrhea

b) Small volume stools

c) Blood and mucus in stools

d) Tenesmus

Question 5:

All are true regarding osmotic diarrhea except :

a) Fasting improves the symptoms

b) Stool osmolar gap >100 mOsm

c) Increased osmolarity of stool is due to unabsorbed dietary nutrients

d) Are associated with enterotoxigenic E. coli

Question 6:

A 32-year-old woman presents with complaints of abdominal pain and recurrent diarrhea for several months. History reveals that she regularly goes on camping trips. Stool osmolality was 200 mOsm/kg H2O and the stool osmolar gap was 20 mOsm/kg H2O. What is the most likely diagnosis?

a) Secretory diarrhea

b) Osmotic diarrhea

c) Factitious diarrhea

d) Infectious diarrhea

Question 7:

All are true regarding bile acid diarrhea except :

a) Seen in limited ileal disease

b) No or mild steatorrhea

c) Normal bile acid pool size

d) Good response to low fat diet

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 8:

Which of the following is the most common viral cause of diarrhea in adults?

a) Enterovirus

b) Rotavirus

c) Norovirus

d) Cytomegalovirus

Question 9:

A father brings his 12-year-old son with complaints of severe abdominal pain and diarrhea. The pain is localized to the right iliac fossa. USG abdomen reveals enlarged mesenteric lymph nodes and a normal appendix. Which of the following is the likely causative organism?

a) Yersinia enterocolitica

b) Campylobacter jejuni

c) Shigella dysenteriae

d) Salmonella enterica

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 10:

Which of the following is not associated with travellers' diarrhoea?

a) Shigella

b) Norovirus

c) Giardia

d) Enterovirus

Question 11:

A child was brought with complaints of watery diarrhea for 48 hours. On examination, tachycardia and decreased skin turgor were present. Stool microscopy revealed the following organisms. Which of the given findings is likely to be seen in this patient?

Sold by @itachibot

a) Hypokalemic acidosis

b) Hypokalemic alkalosis

c) Hyperkalemic acidosis

d) Hyperkalemic alkalosis

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Question 12:

A 19-year-old woman presents to the ER with complaints of acute onset nausea, vomiting, and diarrhea. She gives a history of consumption of street food earlier the same day. Which of the following organisms is likely to be responsible for her condition?

a) Vibrio cholerae

b) Salmonella enterocolitis

c) Enterotoxigenic E. coli

d) Staphylococcus aureus

Answer Key

Question No.	Correct Option
1	a
2	c
3	d

Sold by @itachibot

4	a
5	d
6	c
7	d
8	c
9	a
10	d
11	a
12	d

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

The above scenario is suggestive of steatorrhoea, which is seen in short bowel syndrome.

Steatorrhoea is defined as excess fat in stools (>7g/day) due to poor absorption of lipids. It is characterized by stools are bulky, pale, foul-smelling, and difficult to flush.

Short bowel syndrome (SBS) is a malabsorption disorder caused by a lack of functional small intestine. It can lead to the typical features of malabsorption such as abdominal pain, diarrhoea, steatorrhoea, weight loss and anaemia.

Option B: Primary lactase deficiency leads to defective absorption of lactose. It does not affect lipid absorption.

Option C: Pernicious anemia is due to an absence of gastric parietal cell-intrinsic factor which is necessary for the absorption of vitamin B12.

Option D: Wilson's disease is associated with increased absorption of copper and is not a part of malabsorption syndrome.

Solution to Question 2:

Yogurt is the least likely to cause symptoms of lactose intolerance. It contains lactobacillus that produces lactase and is generally well tolerated.

Lactase is a brush border enzyme that hydrolyzes lactose into glucose and galactose. In lactase deficiency, patients have great variability in clinical symptoms, depending on both the severity of lactase deficiency and the amount of lactose ingested.

The goal of treatment in patients with isolated lactase deficiency is achieving patient comfort through the elimination of dietary lactose.

Foods that are high in lactose include milk (12 g/cup), ice cream (9 g/cup), and cottage cheese (8 g/cup). Aged cheese has lower lactose content. Most people can tolerate up to 12g of lactose a day

Sold by @itachibot

without significant symptoms.

Solution to Question 3:

Rapid small-intestinal transit time increases the symptoms of lactose intolerance.

Lactose intolerance can be due to:

• Primary lactase deficiency- a genetically determined deficiency or absence of lactase enzyme

• Secondary lactase deficiency- due to small intestinal mucosal diseases such as celiac disease

Factors affecting symptoms of lactose intolerance:

• Amount of lactose in the diet.

• Rate of gastric emptying- symptoms are more likely when gastric emptying is rapid (post-gastrectomy and skimmed milk intake).

• Small intestinal transit time- more rapid small intestinal transit time makes symptoms more likely.

• Antibiotic use- results in decreased colonic microflora which in turn decreases the production of easily absorbed short-chain fatty acids from non-absorbed lactose. This can worsen the symptoms.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

Small bowel diarrhea is associated with steatorrhea, not large bowel diarrhea.

Tenesmus refers to the constant urge to defecate with little or no stool passed.

Features	Small bowel	Large bowel
Volume of stools	Large	Small
Frequency of stool	No change	Increased
Blood and mucus in stools	Absent	Present
Tenesmus	Absent	Present
Steatorrhea	Present	Absent
Vomiting	May be present	Infrequent

Solution to Question 5:

Enterotoxigenic E. coli (ETEC) is associated with secretory diarrhea.

Sold by @itachibot

Stool osmolar gap = measured osmolality - calculated osmolality

= (290 to 300) - 2 (stool Na+ + stool K+)

Feature	Osmotic diarrhea	Secretory diarrhea
Definition	Diarrhea due to poorly absorbed osmotically active substances in the colonic lumen.	Diarrhea due to excessive secretion of NaCl, HCO3-, and water into the gut.
Fasting	Improves symptoms	No effect
Stool osmolar gap (mOsm)	>100	<50
Causes	Malabsorption syndrome	Tumors - VIPomaToxins - ETEC, V. cholera

Sold by @itachibot
If you purchased this from someone
you may have been scammed.

Solution to Question 6:

The most likely diagnosis is factitious diarrhea.

Factitious diarrhoea is a form of Münchausen's syndrome. In patients with diarrhea, stool osmolality is always >290-300 mOsm/kg H2O. A stool osmolality of <290 mOsm/kg H2O is suggestive of dilution with urine or water to the sample, by the patient. In this case stool osmolality can be lower than plasma osmolality (<250 mOsm/kg) or very high (>400 mOsm/kg) due to purposeful dilution with hypotonic/concentrated solute by the patient to the stool sample.

The stool osmolar gap is used to differentiate between osmotic and secretory diarrhea. A gap of:

• <50 suggests secretory diarrhoea

• >100 suggests osmotic diarrhoea

Stool osmolar gap = measured osmolality - calculated osmolality

= (290 to 300) - 2 (stool Na+ + stool K+)

Solution to Question 7:

Bile acid diarrhea has no response to a low-fat diet.

Features	Bile acid diarrhea	Fatty acid diarrhea
The extent of ileal disease	Limited	Extensive
Ileal bile acid absorption	Reduced	Reduced
Fecal bile acid loss compensated by hepatic synthesis	Yes	No
Faecal bile acid excretion	Increased	Increased

Sold by @itachibot

Features	Bile acid diarrhea	Fatty acid diarrhea
Steatorrhea	None or mild	>20gm
Response to cholestyramine	Yes	No
Response to a low-fat diet	No	Yes

Solution to Question 8:

The most common viral cause of diarrhoea in adults (and overall) is norovirus (Norwalk virus).

In children, the most common cause is rotavirus.

Solution to Question 9:

Yersinia enterocolitica causes diarrhea with symptoms resembling appendicitis, also called pseudoappendicitis.

The patients present with abdominal pain which can be localized to the right iliac fossa, thus mimicking appendicitis. USG shows a normal appendix and may show enlarged mesenteric lymph nodes.

Appendectomy is not indicated for Yersinia infection pseudoappendicitis.

Causative agent	Special features
Campylobacter jejuni	Triggers Guillain–Barré syndrome,Small bowel lymphoproliferative disorder
Shigella dysenteriae	Severe colitis affecting all 4 layers
Clostridium difficile	Toxic megacolon
Nontyphoid Salmonella	Aortic /Endovascular infection

Solution to Question 10:

Enterovirus is not an etiological agent for travelers' diarrhea.

The various causes of travelers' diarrhea are as follows:

• Bacterial

• Enterotoxigenic E. coli (ETEC)- most common

• Shigella

If you purchased this from someone else, Sold by @itachibot have been scammed.

Sold by @itachibot

• Salmonella

• S. aureus

• B. cereus

• Viral

• Norovirus

• Rotavirus

• Parasite

• Giardia lamblia

• Entamoeba histolytica.

Solution to Question 11:

The image shows the characteristic fish in stream appearance of Vibrio, suggestive of cholera. Diarrhea due to Vibrio cholerae results in hypokalemic acidosis.

Cholera exotoxin activates adenylate cyclase (↑ cAMP) leading to loss of bicarbonate, water, and electrolytes leading to hypokalemic acidosis.

Note: The other cause for hypokalemic acidosis is renal tubular acidosis.

Solution to Question 12:

The above scenario is suggestive of acute onset gastroenteritis. Since the symptoms appeared shortly after ingestion, they are caused due to preformed toxins. From the given options, Staphylococcus aureus is associated with preformed toxins.

Etiology of gastroenteritis are as follows :

1. Preformed toxin:

• Staphylococcus aureus

• Bacillus cereus

• Clostridium perfringens

2. Toxin forming:

• Vibrio cholerae

• Enterotoxigenic E. coli (ETEC)

3. Invasive:

• Shigella

• Salmonella

• Yersinia

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

• Campylobacter

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Irritable Bowel Syndrome

Question 1:

Which of the following is not a diagnostic criterion for irritable bowel syndrome?

a) Abdominal pain related to defecation

b) Persistent diarrhoea after a 48 hour fast

c) Change in frequency of stool

d) Change in form of stool

Question 2:

The most common pattern seen in irritable bowel syndrome patients is _____

a) Diarrhoea only

b) Constipation only

c) Prolonged constipation followed by diarrhoea

d) Constipation alternating with episodes of diarrhoea

Question 3:

A patient presents with a 3-month history of crampy abdominal pain associated with diarrhea. Which of the following features, if present, will support a diagnosis of irritable bowel syndrome?

a) Steatorrhea

b) Nocturnal diarrhoea

c) Persistent diarrhoea after a 48 hour fast

d) Onset of symptoms during a period of stress

Question 4:

Which of the following is not implicated in the pathogenesis of irritable bowel syndrome?

a) Visceral hypersensitivity

b) Activation of mid-cingulate cortex

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Increased serotonin

d) Decreased colonic motor potentials

Question 5:

A 35-year-old housewife has been diagnosed with irritable bowel syndrome - diarrhea predominant. Which neurotransmitter is associated with this condition?

a) Acetylcholine

b) Neuropeptide Y

c) Vasoactive intestinal peptide

d) Serotonin

Question 6:

Which of the following parts of the brain are associated with irritable bowel syndrome?

a) 1 & 2

b) 3 & 4

c) 1 & 4

d) 2 & 3

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following is not a risk factor for post-infectious irritable bowel syndrome?

a) Smoking

b) Depression

c) Male gender

d) Toxicity of infecting bacterial strain

Question 8:

In a suspected case of irritable bowel syndrome, which of the following investigations is not recommended?

a) Complete blood count

Sold by @itachibot

b) Sigmoidoscopy

c) Stool examination for ova and parasites

d) Stool fat test

Question 9:

Which of the following is a marker of inflammatory bowel disease, which helps to rule out irritable bowel syndrome?

a) Faecal fat

b) Faecal lactoferrin

c) Faecal trypsin

d) Faecal chymotrypsin

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following statements is false regarding the FODMAPs diet recommended for a patient with irritable bowel syndrome?

a) It includes fermentable oligosaccharides

b) Diet low in FODMAPs reduces symptoms in IBS patients

c) Sorbitol is an example of FODMAP

d) FODMAPs are highly absorbed by the small intestine

Question 11:

A 35-year-old woman with constipation-predominant irritable bowel syndrome presents with worsening abdominal pain and bloating. Which of the following drugs can be prescribed for her condition?

a) Loperamide

b) Lubiprostone

c) Alosetron

d) Eluxadoline

Question 12:

Sold by @itachibot

You have prescribed paroxetine to a patient with irritable bowel syndrome. This medication is most effective in reducing which of the following symptoms?

a) Diarrhoea

b) Constipation

c) Abdominal pain

d) Bloating

Question 13:

Drug 'X', a 5 HT3 receptor antagonist, is prescribed to treat a patient with diarrhea-predominant irritable bowel syndrome. Identify the drug 'X'.

a) Alosetron

b) Tegaserod

c) Lubiprostone

d) Ondansetron

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

A patient with irritable bowel disease was started on a drug. A few days after administration of the drug, the patient complained of acute GI distress suggestive of ischemic colitis. What was the drug taken by the patient?

a) Lubiprostone

b) Linaclotide

c) Alosetron

d) Tegaserod

Question 15:

A 28-year-old female patient comes to you for advice. She is a known case of irritable bowel syndrome. She asks you when is the best time to take antispasmodics for her postprandial abdominal pain. What would you suggest?

a) 30 minutes before meals

b) 30 minutes after meals

c) Along with meals

Sold by @itachibot

d) When the pain starts

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	d
5	d
6	d
7	c
8	d
9	b
10	d
11	b
12	b
13	a
14	c
15	a

Detailed Explanations

Solution to Question 1:

Persistent diarrhoea after a 48-hour fast is not a diagnostic criteria for IBS.

Solution to Question 2:

The most common pattern in IBS patients is constipation alternating with episodes of diarrhoea.

Based on bowel movement patterns, IBS is classified as:

• IBS-C: Irritable bowel syndrome-constipation predominant

• IBS-D: Irritable bowel syndrome-diarrhoea predominant

• IBS-M: Irritable bowel syndrome-mixed

Sold by @itachibot

The majority of patients change subtypes over the course of illness.

Note: Abdominal pain is a pre-requisite for the diagnosis of IBS. This means that for the diagnosis of IBS, recurrent abdominal pain must be present.

Solution to Question 3:

The onset of symptoms during a period of stress or emotional upset is a feature that supports a diagnosis of irritable bowel syndrome (IBS).

Features in favour of IBS:

• Abdominal pain that improves/worsens with defecation

• Association with stress or emotional symptoms

• Small volume of stools

Features not in favour of IBS:

• Onset of symptoms in old age

• Presence of fever and weight loss

• Persistent diarrhoea after a 48h fast

• Nocturnal diarrhoea

• Steatorrhoea

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 4:

Decreased colonic motor potentials are not implicated in the pathogenesis of irritable bowel syndrome (IBS). On the contrary, increased colonic motor potentials are seen.

The pathogenesis of IBS is poorly understood. However, the following have been implicated in its pathogenesis:

• Increased colonic motor potentials lead to increased high-amplitude propagating contractions (HAPCs) which causes diarrhea.

• Visceral hypersensitivity leads to exaggerated sensory responses to visceral stimulation by food. This explains the postprandial symptoms in IBS.

• CNS dysregulation - dysfunctional activation of the mid-cingulate cortex and prefrontal lobes leads to increased perception of visceral pain.

• Abnormal serotonin pathways - serotonin-containing enterochromaffin cells are increased in IBS patients.

• Altered gut flora

Sold by @itachibot

Solution to Question 5:

Serotonin is associated with irritable bowel syndrome diarrhoea type (IBS-D).

Abnormal serotonin pathways is one of the causes for symptoms in IBS. The serotonin containing enterochromaffin cells are increased in a subset of IBS-D patients. This provides a rationale for the use of serotonin antagonists in the treatment of this condition.

Solution to Question 6:

Mid cingulate cortex and prefrontal lobe are associated with IBS.

Mid cingulate cortex: Modulation of this region is associated with changes in the subjective unpleasantness of pain.

Prefrontal lobe: Dysfunction of this area leads to the increased perception of visceral pain.

Solution to Question 7:

Male gender is not a risk factor for post-infectious IBS.

Post-infectious IBS develops following a gastrointestinal infection with microbes such as Campylobacter, Salmonella and Shigella.

Risk factors for developing post-infectious IBS:

- Young age and female gender
- Prolonged duration of illness
- Toxicity of infecting bacterial strain
- Smoking
- Depression
- Hypochondriasis

Solution to Question 8:

A stool fat test is not recommended in a suspected case of irritable bowel syndrome (IBS).

IBS is a clinical diagnosis that can be made without performing many tests if the patient presents with typical symptoms. However, it is recommended to perform the following tests in most patients to rule out other conditions:

- Complete blood count
- Sigmoidoscopic examination
- Stool examination for ova and parasites

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• 3-week lactose-free diet

In patients >40 years, air-contrast barium enema or colonoscopy should also be performed.

Solution to Question 9:

Fecal lactoferrin is the marker of IBD that is used to rule out irritable bowel syndrome. Fecal lactoferrin is a highly sensitive and specific marker for detecting intestinal inflammation.

Fecal calprotectin is another marker of IBD and it is used to rule out irritable bowel syndrome and bacterial overgrowth. It correlates well with histologic inflammation, predicts relapses, and detects pouchitis.

Option A: Fecal fat is normally digested by pancreatic lipase and absorbed from the jejunum. In disorders of exocrine pancreatic function, fat absorption decreases, resulting in steatorrhea and increased fecal fat.

Options C & D: Fecal trypsin, chymotrypsin are pancreatic enzymes, therefore a decrease in fecal levels below normal suggests a disorder of the exocrine pancreas.

Solution to Question 10:

FODMAPs are poorly absorbed in the small intestine.

FODMAPs stands for Fermentable Oligosaccharides, Disaccharides, Monosaccharides, And Polyols. Foodstuff containing lactose, fructose, and sorbitol are some examples of FODMAPs.

Since FODMAPs are poorly absorbed, they are fermented by gut bacteria in the colon to produce gas and osmotically active carbohydrates. So, a diet low in FODMAPs is recommended to reduce symptoms in IBS patients.

Solution to Question 11:

Lubiprostone is used in the treatment of constipation-predominant IBS.

Lubiprostone is a chloride channel activator that promotes the secretion of water and chloride into the bowel and improves bowel movements.

Loperamide, alosetron, and eluxadoline (opioid mu receptor agonist) are used in the treatment of diarrhea-predominant IBS.

Solution to Question 12:

Paroxetine is most effective in treating constipation. It is a selective serotonin reuptake inhibitor (SSRI) that accelerates oro-caecal transit.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed

Sold by @itachibot

Note: Citalopram is an SSRI that is effective in the management of pain in IBS. It blunts perception of rectal distension and reduces the magnitude of gastrocolic reflex.

Solution to Question 13:

Alosetron is a 5-HT3 receptor antagonist used in the treatment of IBS–D.

Options B and C: Tegaserod and lubiprostone are used in the treatment of constipation-predominant irritable bowel syndrome (IBS–C).

Option D: Ondansetron is used in the treatment of nausea and vomiting.

Solution to Question 14:

Ischaemic colitis is a side-effect of alosetron.

Alosetron is a 5-HT3 receptor antagonist that is used to reduce diarrhoea in irritable bowel syndrome.

Option A: Lubiprostone is associated with nausea and diarrhea.

Option B: Linaclotide is associated with diarrhea.

Option D: Tegaserod is associated with cardiac toxicity.

Solution to Question 15:

The best time to take antispasmodics for postprandial pain is 30 min before meals.

Antispasmodics such as dicyclomine inhibit the gastrocolic reflex and thus reduce pain due to intestinal spasms. It is best taken 30 min before meals so that the effective blood levels are achieved shortly before the anticipated onset of pain.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Inflammatory Bowel Disease - Clinical Features and Diagnosis

Question 1:

Which of the following genes is associated with Crohn's disease?

a) NOD2

b) JAK2

c) BRAF V600E

d) STAT3

Question 2:

Which of the following conditions are not associated with inflammatory bowel disease?

a) Turner's syndrome

b) Wiskott-Aldrich syndrome

c) Hermansky-Pudlak syndrome

d) Klinefelter syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

In patients who have undergone an appendectomy, there is a lower risk for developing which of the following diseases?

a) Ulcerative colitis

b) Crohn's disease

c) Carcinoma colon

d) Diverticulosis

Question 4:

Early-onset inflammatory bowel disease is associated with:

a) JAK2 mutation

Sold by @itachibot

b) FOXP3 mutation

c) IL-10 deficiency

d) NOD 2 mutation

Question 5:

Which of the following is true about immune dysregulation polyendocrinopathy enteropathy X-linked (IPEX) syndrome?

a) B cell defect

b) Due to FOXP3 gene mutation

c) Associated with Crohn's disease-like enteropathy

d) Recurrent staphylococcal skin infections

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which of the following infections should you rule out before diagnosing a patient with IBD on the basis of clinical symptoms and biopsy findings?

a) Mycobacterium Avium

b) Herpes simplex

c) Campylobacter jejuni

d) Cytomegalovirus

Question 7:

Which of the following serological markers is positive in ulcerative colitis?

a) Anti-pancreatic antibody

b) Anti-flagellin antibody

c) ASCA

d) ANCA

Question 8:

Which of the following sites is most commonly involved in Crohn’s disease?

Sold by @itachibot

a) Terminal ileum

b) Rectosigmoid colon

c) Transverse colon

d) Proximal duodenum

Question 9:

A patient presented with acute severe small bowel obstruction to the ER. An emergency small bowel resection was done. The resected specimen is shown below. What could be the likely diagnosis?

a) Ulcerative colitis

b) Crohn's disease

c) Intestinal TB

d) Intestinal lymphoma

Question 10:

A 28-year-old man presented with diarrhea, bloody stools, weight loss, and intermittent fever for the past 2 months. Endoscopy revealed the following. What is the most probable diagnosis?

Sold by @itachibot



a) Crohn's disease

b) Intestinal TB

c) Ulcerative colitis

d) Salmonella infection

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 11:

A 70-year-old chronic smoker is being evaluated for chronic diarrhea, steatorrhea, and weight loss. On examination, pallor was noted. Peripheral smear revealed macrocytes and hypersegmented neutrophils. Stool examination showed elevated calprotectin and a positive guaiac test. What is the most likely diagnosis?

a) Ulcerative colitis

b) Diverticulitis

c) Crohn's disease

d) Irritable bowel syndrome

Question 12:

A 30-year-old woman presented with 7-8 episodes of bloody diarrhea. On examination, she was pale and febrile, with a pulse rate of 104 per minute. As she was not responding to broad-spectrum antibiotics, a colonoscopy was done which revealed ulcerations and pseudopolyps. What is the most probable diagnosis?

a) Moderate Crohn's disease

Sold by @itachibot

b) Severe Crohn's disease

c) Moderate ulcerative colitis

d) Severe ulcerative colitis

Question 13:

All of the following are features of Crohn's disease except:

a) Transmural involvement

b) Lead pipe appearance

c) Rectal sparing

d) Perianal fistula

Question 14:

A young man presents to the OPD with complaints of chronic diarrhea, right lower abdominal pain, and weight loss. On examination, you note a low-grade fever and a palpable mass in the right lower quadrant. Colonoscopy revealed sparing of the rectum. The following endoscopic findings may also be seen in his condition except:

a) Aphthous ulcer

b) Longitudinal ulcers

c) Fine granular mucosa

d) Stricture

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 15:

What is the earliest radiographic feature of Crohn's disease?

a) Ram's horn sign

b) Rose-thorn appearance

c) Aphthous ulcer

d) String sign of Kantor

Question 16:

Sold by @itachibot

Which among the following is the earliest radiologic change seen on barium enema in ulcerative colitis?

a) Collar button ulcer

b) Fine mucosal granularity

c) String sign

d) Ram's horn sign

Question 17:

Which of the following findings is seen in patients with ulcerative colitis?

a) Rose thorn appearance

b) Pseudopolyps

c) Skip lesions

d) Pseudodiverticula

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 18:

A 37-year-old man was referred to medical gastroenterology with complaints of chronic bloody diarrhea. Examination revealed blood on rectal examination. A colonoscopy was done as shown below. What is the likely diagnosis?

a) Ulcerative colitis

b) Crohn's disease

Sold by @itachibot

c) Ishchemic colitis

d) Solitary rectal ulcer syndrome

Question 19:

Which of the following complications is unlikely to occur in a patient with ulcerative colitis?

a) Haemorrhage

b) Toxic megacolon

c) Malignant change

d) Fistula

If you bought this from someone else, it was probably stolen and resold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	c
5	b
6	c
7	d
8	a
9	b
10	a
11	c
12	d
13	b
14	c
15	c
16	b
17	b
18	a
19	d

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The gene associated with Crohn's disease is the NOD2 gene. It is present on chromosome 16.

Nucleotide-binding oligomerization domain 2 (NOD2) acts as a cytoplasmic sensor of microbial products. Defective NOD2 gene results in defective innate immunity. When the innate immunity is unable to clear the bacteria, there is an exaggerated response of the immune system (T-cells) resulting in damage to the host tissue.

Option B: JAK2 mutation is associated with polycythemia vera.

Option C: BRAF V600E mutation is associated with melanoma, papillary thyroid cancer, and serous ovarian cancers.

Option D: STAT3 mutations are associated with the autosomal dominant hyper-IgE syndrome.

Solution to Question 2:

Klinefelter syndrome is not commonly associated with inflammatory bowel disease (IBD).

The genetic disorders associated with IBD are:

- Wiskott-Aldrich syndrome
- Immunodysregulation polyendocrinopathy enteropathy X-linked (IPEX) syndrome
- Turner's syndrome
- Hermansky-Pudlak syndrome
- Glycogen storage disease

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Appendectomy is protective against ulcerative colitis.

Smoking is also seen to have a protective effect on ulcerative colitis but it is a risk factor for developing Crohn's disease.

Solution to Question 4:

Early-onset IBD is due to IL-10 receptor dysfunction and IL-10 deficiency. They are associated with severe refractory IBD in early life.

Solution to Question 5:

Sold by @itachibot

IPEX (immune dysregulation, polyendocrinopathy, enteropathy, and X-linked) syndrome is caused by mutations in the FOXP3 gene, which leads to a deficiency of T-regulatory cells, resulting in immune dysfunction.

IPEX is associated with:

• Ulcerative colitis-like autoimmune enteropathy

• Neonatal type 1 DM

• Thyroiditis

• Dermatitis

Solution to Question 6:

Campylobacter enteritis should be ruled out before diagnosing a patient with IBD.

Campylobacter enteritis can mimic the clinical presentation and biopsy appearance of ulcerative colitis or Crohn's disease (IBD). Hence, it should be ruled out, especially in persons with a history of:

• Foreign travel

• Significant animal contact

• Immunodeficiency

• Exposure incurring a high risk of transmission.

It can also cause a relapse of established ulcerative colitis.

A definitive diagnosis of campylobacter infection can be made by culture or real-time PCR.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 7:

Antineutrophil cytoplasmic antibody (ANCA) positivity is a feature of ulcerative colitis.

ASCA (anti-Saccharomyces cerevisiae antibody), anti-pancreatic antibody, and anti-flagellin antibody positivity are seen in Crohn's disease.

Serological Markers in IBD	
Crohn's disease	Ulcerative colitis
ASCAOMP-CAnti I2Anti-pancreatic antibodyAnti-flagellin antibody	p-ANCAAnti-goblet cell Antibody

Solution to Question 8:

Sold by @itachibot

The most commonly affected site in Crohn's disease is the terminal ileum.

Other commonly affected sites in Crohn's disease include the small intestine and the colon. Although it can affect any part of the GI tract from mouth to anus, the rectum is usually spared.

On the other hand, the rectum is almost always involved in ulcerative colitis.

Solution to Question 9:

The above specimen of the small bowel shows creeping fat which is a feature of Crohn's disease (CD).

Mesenteric fat is normally seen along the longitudinal axis of the intestine. However, in Crohn's disease, excessive fibrofatty proliferation results in mesenteric fat wrapping around the inflamed bowel. This is called creeping fat.

Strictures and fistulae are the commonly associated complications of long-standing Crohn's disease. Strictures can cause bowel obstruction.

Other gross features seen in Crohn's disease are:

• Transmural inflammation- full-thickness inflammation involving all the layers of the bowel

• Aphthous ulcers- small, shallow ulcers surrounded by erythema

• Longitudinal ulcers- linear or serpiginous deep ulcers

• Skip lesions- involvement of discontinuous segments of the bowel with intervening normal mucosa

• Cobblestone appearance- deep, longitudinal ulcerations interlaced with intervening normal mucosa

The image below shows cobblestone appearance in Crohn's disease.

Sold by @itachibot If you purchased this from someone else you may have been scammed

Sold by @itachibot

Solution to Question 10:

The clinical presentation along with the endoscopy image showing a cobblestone appearance are suggestive of Crohn's disease (CD).

Endoscopic findings in Crohn's disease include:

• Aphthous ulcers - small, shallow ulcers surrounded by erythema

• Longitudinal ulcers - linear or serpiginous deep ulcers

• Cobblestone appearance - deep, longitudinal ulcerations interlaced with intervening normal mucosa

• Strictures

• Fistulas

Strictures and fistulae are the commonly associated complications of long-standing Crohn's disease.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 11:

The above scenario is suggestive of Crohn's disease presenting as jejunoileitis. This is evidenced by the presence of features of malabsorption such as steatorrhea, chronic diarrhea, and vitamin B12 deficiency anemia.

The fecal calprotectin and lactoferrin are elevated in inflammatory conditions like inflammatory bowel disease, but not in irritable bowel syndrome.

The stool guaiac test or guaiac fecal occult blood test (gFOBT) reveals the presence of microscopic levels of blood suggesting Crohn's disease. Gross bleeding per rectum is more characteristic of ulcerative colitis.

Additionally, smoking is a risk factor for Crohn's disease. However, it is a protective factor in ulcerative colitis.

Solution to Question 12:

Tachycardia, bloody diarrhea, pseudopolyps, and ulcerations on colonoscopy strongly suggest severe ulcerative colitis.

Given below is the Truelove and Witts severity index for ulcerative colitis.

	Mild	Moderate	Severe
Bowel movements	< 4 per day	4 – 6 per day	> 6 per day
Blood in stool	Small	Moderate	Severe

Sold by @itachibot

	Mild	Moderate	Severe
Fever	None	< 37.5°C	> 37.5°C
Tachycardia	None	< 90 mean pulse	>90 mean pulse
Anemia	Mild	Hb > 75% of normal Hb	Hb < 75% of normal Hb
ESR	< 30mm	< 30mm	> 30mm
Endoscopic appearance	ErythemaDecreased vascular markingsFine granularity	Marked erythemaAbsent vascular markingsCoarse granularityContact bleedingNo ulcerations	Spontaneous bleedingUlcerations

If you @itachibot someone else, ...med.

Solution to Question 13:

Lead pipe appearance is seen in ulcerative colitis (not Crohn's disease). This appearance is due to the loss of haustra in the diseased colonic segment.

The image below shows the lead pipe appearance of the colon on barium enema:



Solution to Question 14:

The above clinical scenario is suggestive of Crohn's disease presenting as ileocolitis. Fine granular mucosa is seen in ulcerative colitis, not Crohn's disease.

Ileocolitis presents as right lower abdominal pain, fever, and diarrhea. Pain is usually colicky and is relieved by defecation. It may mimic acute appendicitis in the early stages.

Endoscopic findings in Crohn's disease include:

• Aphthous ulcers - Small, shallow ulcers surrounded by erythema

Sold by @itachibot

• Longitudinal ulcers - Linear or serpiginous deep ulcers

• Cobblestone appearance - Deep, longitudinal ulcerations interlaced with intervening normal mucosa

• Strictures

• Fistulas.

Solution to Question 15:

Aphthous ulcers seen in Crohn's disease are the earliest radiological signs on small bowel follow-through study.

They are shallow ulcers that are often multiple and separated by normal intervening mucosa. They are also the earliest macroscopic findings of Crohn's disease.

Features on barium small bowel follow-through in Crohn's disease include:

• Mucosal ulcers:

• Aphthous ulcers initially

• Deep ulcers (more than 3 mm depth)

• Longitudinal fissures

• Transverse stripes

• Severe mucosal ulcers lead to cobblestone appearance

• Sinus tracts and fistulae

• Creeping fat: widely separated loops of the bowel due to fibrofatty proliferation

• Thickened folds due to edema

• Pseudo-diverticula formation: due to contraction at the site of ulcer with ballooning of the opposite site

• String sign of Kantor: tubular narrowing due to spasm or stricture depending on the chronicity

• Rose thorn appearance

• Partial obstruction.

CT and MR enterography have a higher sensitivity for the diagnosis of Crohn's disease. They may reveal additional features such as:

• Fat halo sign - submucosal fat deposition

• Comb sign - engorgement of the vasa recta.

In gastric Crohn's disease, conical narrowing of the antrum may be seen. This is called Ram's horn sign.

The image below shows aphthous ulcers (focal contrast with surrounding haloes of mucosal edema)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



The image below shows the string sign of Kantor:



The image below shows rose thorn appearance:

Sold by @itachibot



The image below shows cobblestone appearance:



Solution to Question 16:

The earliest radiological change of ulcerative colitis seen on barium enema is fine mucosal granularity.

Double-contrast barium enema allows for exquisite detail of the colonic mucosa and also allows the bowel proximal to strictures to be assessed. It is however contraindicated if acute severe colitis is present due to the risk of perforation.

Early changes in ulcerative colitis:

• Mucosal granularity

• Fine granular pattern

Sold by @itachibot

• Mucosal stippling

• Collar button ulcers

• Deep collar stud ulcers.

The below contrast radiograph shows granular mucosa typical of ulcerative colitis.



Option A: Collar button ulcers are small deep ulcers in which the button surface corresponds to the eroded ulcer base and the button neck corresponds to the undermined and preserved mucosa. They are also seen in ulcerative colitis.

Option C: String sign represents circumferential inflammation and fibrosis, resulting in long segments of luminal narrowing as shown below. It is more characteristic of Crohn's disease.



Option D: Ram's horn sign is seen in gastric Crohn's disease.

Sold by @itachibot

Solution to Question 17:

Pseudopolyps are seen in ulcerative colitis. They are formed due to alternating inflammation and regeneration of the ulcerated mucosal epithelium in ulcerative colitis.

Other gross findings in ulcerative colitis include:

• Continuous mucosal inflammation of the colon most commonly involving the rectum

• Fine mucosal granularity

• Collar button ulcers

• Pseudopolyps.

Rose thorn appearance, skip lesions, and pseudodiverticula are features of Crohn's disease.

Solution to Question 18:

In the above scenario, colonoscopy shows continuous involvement of erythematous rectosigmoid mucosa and has a fine granular surface resembling sandpaper. This along with the clinical features is suggestive of ulcerative colitis.

The image below shows a gross specimen of the colon with diffuse involvement of mucosa and ulcers (without skip lesions) in ulcerative colitis.

The image below shows a gross specimen of jejunum showing the cobblestone appearance as seen in Crohn's disease.

Sold by @itachibot

Sold by @itachibot

Sold by @itachi... else,
If you purchased this from some...
you may have been scammed.

Solution to Question 19:

Fistula is not a complication of ulcerative colitis. It is seen in Crohn's disease.

Hemorrhage, malignant change, and toxic megacolon are complications of ulcerative colitis.

Note: Stricture formation may be a complication of ulcerative colitis, but it is seen only in 5-10% of patients. They are more commonly seen in Crohn's disease.

Sold by @itachibot

Inflammatory Bowel Disease - Complications and Treatment

Question 1:

A patient with inflammatory bowel disease presents with the following lesion. Which of the following is incorrect regarding this condition?

a) Can occur years before the onset of bowel symptoms

b) Steroids should be avoided

c) Can develop years after colectomy

d) Associated with severe disease

Question 2:

A 37-year-old man presented with hematuria and dysuria for 5 days. He also complains of chronic diarrhea and weight loss for 6 months. His urine microscopy revealed the following stones. What is the most likely disorder associated with this finding?

Sold by @itachibot

a) Crohn's disease

b) Ulcerative colitis

c) Irritable bowel syndrome

d) Lactose intolerance

Sold by @itachi... If you purchased this from some... else, ...may have been scammed.

Question 3:

Perianal examination of a patient revealed the following finding. Which condition is this associated with?

a) Ulcerative colitis

Sold by @itachibot

b) Crohn's disease

c) Ischemic colitis

d) Ileocecal tuberculosis

Question 4:

A 28-year-old man was brought to the casualty with a high fever and abdominal pain for the past 2 days. His history is significant for bloody diarrhea over the past few months. The abdominal radiograph of this patient is given below. Identify the condition.

a) Toxic megacolon

b) Intussusception

c) Sigmoid volvulus

d) Pneumatosis intestinalis

Question 5:

A patient with inflammatory bowel disease presents with ocular pain, photophobia, and blurred vision. Which among the following is the most likely cause?

a) Uveitis

b) Episcleritis

c) Scleritis

d) Conjunctivitis

Sold by @itachibot

Question 6:

A 34-year-old man with Crohn's disease presents with red tender swellings on the lower leg as shown below. What is the likely diagnosis?

a) Pyoderma gangrenosum

b) Erythema nodosum

c) Sweet's syndrome

d) Ecthyma gangrenosum

Question 7:

What are the cardiopulmonary manifestations associated with inflammatory bowel disease?

a) 1,2,5

b) 4,5

c) 2,3

d) 1,3,4

Question 8:

A 36-year-old man presents with worsening fatigue, jaundice, and pruritis for the last 3 months. Past history revealed recurrent episodes of bloody diarrhea and weight loss over the

Sold by @itachibot

past year. Lab investigations show raised ALP, GGT, and bilirubin, and an MRCP is done as shown below. Which of the following is most likely to be associated with his current presentation?



a) Chronic pancreatitis

b) Celiac disease

c) Ulcerative colitis

d) Crohn's disease

Sold by
If you purchased this fro
you may have been scam
else,
d.

Question 9:

After initiating treatment for patients with IBD, in which of the following extraintestinal manifestations would you expect to see an improvement?

a) Arthritis

b) Ankylosing spondylitis

c) Uveitis

d) Pyoderma gangrenosum

Question 10:

A patient is diagnosed with ulcerative colitis on endoscopy. He is hemodynamically stable but currently has 3-4 episodes of blood-tinged loose stools per day. Which of the following drug would you prefer for treating this patient?

Sold by @itachibot

a) 5-aminosalicylic acid

b) Azathioprine

c) Budesonide

d) 6-mercaptopurine

Question 11:

A 36-year-old man presented with complaints of 7-8 episodes of bloody diarrhea per day. He is pale and febrile. Heart rate is 110 bpm and ESR is 50mm. Colonoscopy revealed continuous mucosal involvement with ulcerations and mucosal hemorrhages. Which of the following drugs would you not prefer in the management of this patient?

a) Mesalamine

b) Infliximab

c) IV steroids

d) Cyclosporine

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 12:

Which of the following drugs have no role in the management of patients with ulcerative colitis?

a) Oral salicylates

b) Hydrocortisone

c) Metronidazole

d) Cyclosporine

Question 13:

A patient with Crohn's disease is started on infliximab. Which of the following conditions is the patient is at increased risk of developing?

a) Hodgkin's Lymphoma

b) Non-Hodgkin lymphoma

c) Hairy cell Leukaemia

d) Follicular lymphoma

Sold by @itachibot

Question 14:

Ustekinumab is effective in all of the following conditions except:

a) Rheumatoid arthritis

b) Ulcerative colitis

c) Psoriatic arthirtis

d) Crohn's disease

Question 15:

Which of the following drugs has no role after induction of remission has been achieved in a patient with IBD?

a) 5-ASA agents

b) Glucocorticoids

c) Azathioprine

d) Infliximab

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 16:

A patient with IBD was brought to the ER with acute epigastric pain radiating to the back. His lab investigations revealed elevated serum amylase and lipase. He gives a history of being started on a new drug for maintenance therapy of his IBD 2 weeks back. The causative drug is most likely to be:

a) Methotrexate

b) Tacrolimus

c) 5-ASA

d) Azathioprine

Question 17:

What is nutritional therapy in inflammatory bowel disease used for?

a) Maintaining remission in Crohn's disease

b) Maintaining remission in ulcerative colitis

c) Inducing remission in Crohn's disease

Sold by @itachibot

d) Inducing remission in ulcerative colitis

Question 18:

What is the treatment of choice in a patient with refractory ulcerative colitis?

a) Mucosal proctectomy with ileal pouch–anal anastomosis

b) Proctectomy with ileorectal anastomosis

c) Colectomy with ileostomy

d) Ileorectal anastomosis

@itachibot someone else, med. If yo

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	a
5	a
6	b
7	d
8	c
9	a
10	a
11	a
12	c
13	b
14	a
15	b
16	d
17	c
18	a

Detailed Explanations

Sold by @itachibot

Solution to Question 1:

The given image shows pyoderma gangrenosum. It is very difficult to treat and often requires IV antibiotics and IV glucocorticoids. Other drugs used in its management include dapsone, cyclosporine, and infliximab.

Pyoderma gangrenosum is a dermatological extraintestinal manifestation of inflammatory bowel disease. It begins as a pustule and progresses to an ulcer with necrotic tissue and exudates. It may occur years before the onset of bowel symptoms and is usually associated with severe disease.

Lesions are commonly found on the dorsal surface of the feet and legs but may occur on the arms, chest, and even the face. It responds poorly to colectomy and may even develop years after the surgery.

Other dermatological manifestations associated with IBD include:

- Erythema nodosum
- Sweet syndrome (neutrophilic dermatosis)
- Perianal skin tags
- Psoriasis
- Metastatic Crohn's disease (cutaneous granulomas).

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 2:

The given urinary crystal is the bipyramidal or envelope-shaped crystals of calcium oxalate dihydrate. Calcium oxalate stones are more commonly associated with Crohn's disease.

In Crohn's disease (especially post small bowel resection), fatty acids are not absorbed because of ileal dysfunction. These fatty acids in the gut lumen bind to calcium. Because of this, the dietary oxalates which normally bind to the calcium are now left unbound. This leads to increased absorption of dietary oxalates leading to hyperoxaluria and oxalate stones.

Note: Celiac disease is also associated with calcium oxalate stones.

Solution to Question 3:

The above image shows a large hemorrhoidal perianal skin tag. These are found in patients with Crohn's disease, especially those with colon involvement.

Solution to Question 4:

The clinical scenario and radiograph are suggestive of toxic megacolon. It is seen in severe ulcerative colitis.

Sold by @itachibot

Abdominal radiography in toxic megacolon shows transverse or right colonic dilation >6 cm and loss of haustrations.

Toxic megacolon may present with fever, abdominal pain, and abdominal distension or features of sepsis.

Solution to Question 5:

Uveitis is the most common cause of the given symptoms in a patient with inflammatory bowel disease (IBD).

Uveitis is an ophthalmic manifestation of IBD. It is associated with both ulcerative colitis and Crohn's disease. It can develop during remission and is seen even after bowel resection. Prompt intervention with glucocorticoids is required to prevent scarring and visual impairment.

The other common ophthalmic manifestations in IBD are conjunctivitis and episcleritis. Episcleritis, more common in Crohn's, is a benign disorder presenting with mild ocular burning.

Solution to Question 6:

The given clinical scenario suggests a diagnosis of erythema nodosum seen in inflammatory bowel disease (IBD).

Erythema nodosum is a dermatologic manifestation of IBD. It is seen in both ulcerative colitis and Crohn's disease. It manifests as tender erythematous nodules on the anterior surface of the lower legs, ankles, thighs, and arms.

Option A: Pyoderma gangrenosum is a dermatological manifestation of IBD. It begins as pustular lesions which then ulcerate with necrotic foci and exudation.

Option C: Sweet's syndrome is acute febrile neutrophilic dermatosis, seen in ulcerative colitis.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed!

Sold by @itachibot

Option D: Ecthyma gangrenosum is a skin infection caused by Pseudomonas aeruginosa, often seen in immunocompromised patients. It presents as pustules/vesicles which progress to necrotic ulcers.

Solution to Question 7:

The cardiopulmonary manifestations of inflammatory bowel disease (IBD) are:

- Endocarditis
- Myocarditis
- Pleuropericarditis
- Interstitial lung disease

Sold by @itachibot

Both ulcerative colitis and Crohn's disease have an increased risk of venous and arterial thrombosis.

Solution to Question 8:

The above clinical scenario along with MRCP showing multiple areas of intrahepatic narrowing with alternating areas of normal and dilated segments of bile ducts is suggestive of primary sclerosing cholangitis. PSC is more common in ulcerative colitis than in Crohn's disease.

Additionally, recurrent episodes of bloody diarrhea with weight loss seen in this patient also point toward a possible diagnosis of ulcerative colitis.

Primary sclerosing cholangitis (PSC) is an extraintestinal manifestation of IBD characterized by intrahepatic and extrahepatic duct inflammation/fibrosis. Around 50-75% of patients with primary sclerosing cholangitis (PSC) have inflammatory bowel disease (IBD) whereas, only 5% of IBD cases develop PSC.

Option A: Chronic pancreatitis is associated with a chain of lakes appearance- dilated main duct with clubbing and dilatation of the side branches. Patients can present with steatorrhoea i.e., the passage of offensive-smelling large stools due to fat malabsorption.

The image below shows chain of lakes appearance on MRI:

Option B: Celiac disease may be associated with primary biliary cholangitis (PBC), rather than primary sclerosing cholangitis.

Solution to Question 9:

Arthritis worsens with exacerbation of bowel disease and improves with the treatment of bowel symptoms or surgery.

Sold by @itachibot

Ankylosing spondylitis, uveitis, and pyoderma gangrenosum do not correlate with disease activity in the bowel. They may occur during the remission of disease or even after surgery.

Solution to Question 10:

The above scenario is suggestive of mild ulcerative colitis. The preferred drug for mild ulcerative colitis (UC) is 5-ASA agents (5-aminosalicylic acid).

Given below is the Truelove and Witts severity index for ulcerative colitis.

Note: Mirikizumab (an IL-23 inhibitor) was recently FDA approved for the treatment of moderate to severe ulcerative colitis. It reduces stool frequency, rectal bleeding and bowel urgency.

	Mild	Moderate	Severe
Bowel movements	< 4 per day	4 – 6 per day	> 6 per day
Blood in stool	Small	Moderate	Severe
Fever	None	< 37.5°C	> 37.5°C
Tachycardia	None	< 90 mean pulse	>90 mean pulse
Anemia	Mild	Hb > 75% of normal Hb	Hb < 75% of normal Hb
ESR	< 30mm	< 30mm	> 30mm
Endoscopic appearance	ErythemaDecreased vascular markingsFine granularity	Marked erythemaAbsent vascular markingsCoarse granularityContact bleedingNo ulcerations	Spontaneous bleedingUlcerations

Treatment of Ulcerative Colitis			
	Mild	Moderate	Severe
Induction	5-ASA	5-ASA + Oral steroids	IV steroids + IV anti TNF alpha + IV cyclosporine
Maintainance	5-ASA	5-ASA + Azathioprine/6-MP	Oral cyclosporine or infliximab

Solution to Question 11:

The above scenario is suggestive of severe ulcerative colitis. Mesalamine (5-ASA) is ineffective in the management of severe ulcerative colitis (UC). It is used in the treatment of mild-moderate UC.

Given below is the Truelove and Witts severity index for ulcerative colitis.

Note: Mirikizumab (an IL-23 inhibitor) was recently FDA approved for the treatment of moderate to severe ulcerative colitis. It reduces stool frequency, rectal bleeding and bowel urgency.

Sold by @itachibot

	Mild	Moderate	Severe
Bowel movements	< 4 per day	4 – 6 per day	> 6 per day
Blood in stool	Small	Moderate	Severe
Fever	None	< 37.5°C	> 37.5°C
Tachycardia	None	< 90 mean pulse	>90 mean pulse
Anemia	Mild	Hb > 75% of normal Hb	Hb < 75% of normal Hb
ESR	< 30mm	< 30mm	> 30mm
Endoscopic appearance	ErythemaDecreased vascular markingsFine granularity	Marked erythemaAbsent vascular markingsCoarse granularityContact bleedingNo ulcerations	Spontaneous bleedingUlcerations

Treatment of Ulcerative Colitis			
	Mild	Moderate	Severe
Induction	5-ASA	5-ASA + Oral steroids	IV steroids + IV anti TNF alpha + IV cyclosporine
Maintainance	5-ASA	5-ASA + Azathioprine/6-MP	Oral cyclosporine or infliximab

If you purchased this book you may have been

Solution to Question 12:

Metronidazole has no role in the management of ulcerative colitis. It is an antibiotic that is useful in active inflammatory, fistulous, and perianal Crohn's disease.

Note: Mirikizumab (an IL-23 inhibitor) was recently FDA approved for the treatment of moderate to severe ulcerative colitis. It reduces stool frequency, rectal bleeding and bowel urgency.

Solution to Question 13:

Patients with Crohn's disease on infliximab are at increased risk of non-Hodgkin lymphoma (NHL) and hepatosplenic T-cell lymphoma (HS-TCL).

Infliximab is the first biologic therapy approved for Crohn's disease. It is an IgG1 antibody against TNF-alpha.

Drugs used in treatment of Crohn's disease:

• Oral budesonide (mainly for mild to moderate)

• Oral prednisolone

• Methotrexate

• Anti TNF alpha

Sold by @itachibot

• Azathioprine

• 6-mercaptopurine

• Monoclonal antibodies

• Antibiotics

Infliximab is also used in the treatment of moderate to severely active ulcerative colitis.

Solution to Question 14:

Ustekinumab has no role in rheumatoid arthritis. It is effective against both Crohn's disease and ulcerative colitis.

It can also be used in the management of plaque psoriasis and psoriatic arthritis.

It is an IgG1 monoclonal antibody that blocks the activity of IL-12 and IL-23.

Biological therapy in inflammatory bowel disease:

• Anti-TNF therapies:

• Infliximab

• Adalimumab

• Certolizumab

• Anti-integrins:

• Natalizumab

• Ustekinumab

• Vedolizumab

Note: Mirikizumab (an IL-23 inhibitor) was recently FDA approved for the treatment of moderate to severe ulcerative colitis. It reduces stool frequency, rectal bleeding and bowel urgency.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 15:

Glucocorticoids play no role in maintenance therapy in inflammatory bowel disease.

They are used for induction in Crohn's disease and moderate to severe ulcerative colitis. Once clinical remission has been induced, they should be tapered gradually.

Solution to Question 16:

The above scenario is suggestive of pancreatitis. Azathioprine causes pancreatitis in some patients within the first few weeks of therapy. It is completely reversible on discontinuing the drug.

Sold by @itachibot

Azathioprine and 6-mercaptopurine (purine analogues) also have an increased risk of toxicity as some individuals may lack the enzyme necessary for their metabolism. This may lead to hepatitis, bone marrow suppression, etc. Hence, the patient should be monitored regularly with blood counts and liver function tests.

Solution to Question 17:

Nutritional therapy is used for inducing remission in Crohn's disease.

Dietary antigens may stimulate an immune response. Hence, nutritional therapies like total parenteral nutrition, enteral nutrition (peptide-based preparations), etc. can be used to provide bowel rest in active Crohn's disease. These interventions cannot be used for maintaining remission.

Ulcerative colitis does not respond to dietary intervention.

Drugs used in treatment of Crohn's disease:

- Oral budesonide (mainly for mild to moderate)
- Oral prednisolone
- Methotrexate
- Anti TNF alpha
- Azathioprine
- 6-mercaptopurine
- Monoclonal antibodies
- Antibiotics

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 18:

The treatment of choice in refractory ulcerative colitis (UC) is mucosal proctectomy with ileal pouch–anal anastomosis (IPAA).

As ulcerative colitis is a mucosal disease, the rectal mucosa is dissected and removed down to the dentate line of the anus. The ileum is made into a pouch that serves as neorectum.

Sold by @itachibot

Malabsorption Syndromes

Question 1:

Which of the following is unlikely to be seen in patients with malabsorption syndrome?

a) Anemia

b) Constipation

c) Tetany

d) Abdominal pain

Question 2:

Which of the following tests is useful for the assessment of intestinal mucosal function in a patient presenting with symptoms of malabsorption?

a) D-xylulose test

b) D-xylose test

c) Galactomannan test

d) Schilling test

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

The result of a D-xylose test showed urinary excretion of the monosaccharide at 3.6g after 5 hours. Which of the following can be a differential diagnosis?

a) Irritable bowel syndrome

b) Ulcerative colitis

c) Pancreatic insufficiency

d) Crohn's disease

Question 4:

Patients with which of the following conditions are likely to suffer from malabsorption?

a) Small bowel diverticulum

b) Anterior colon resection

Sold by @itachibot

c) Ulcerative colitis

d) Sigmoid volvulus

Question 5:

What is the most common cause of short bowel syndrome in adults?

a) Crohn's disease

b) Mesenteric vascular occlusion

c) Necrotizing enterocolitis

d) Small intestinal perforation

Question 6:

Patients with which of the following conditions are unlikely to have protein malabsorption?

a) Enterokinase deficiency

b) Hartnup syndrome

c) Cystinuria

d) Homocystinuria

Question 7:

A Schilling test was performed on a patient with vitamin B12 deficiency and the results are given below. What is the inference?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Test	Absorption
58Co-labelled cobalamin	Reduced
With intrinsic factor	Reduced
With pancreatic enzyme	Reduced
After 5 days of antibiotics	Reduced

©MARROW

a) Ileal disease

b) Pernicious anemia

c) Small intestinal bacterial overgrowth

d) Chronic pancreatitis

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Question 8:

A patient presented with chronic diarrhea and steatorrhea. D-xylose test was normal and the Schilling test was abnormal. A duodenal biopsy was normal. What is the most likely diagnosis?

a) Celiac disease

b) Ulcerative colitis

c) Intestinal lymphangiectasia

d) Pancreatic insufficiency

Question 9:

An 11-year-old boy presented with steatorrhea and chronic diarrhea. An evaluation revealed delayed intellectual development and ataxia. Acanthocytes were seen on the peripheral blood smear. Ophthalmological examination showed the following. Which of the following is the most likely diagnosis?

Sold by @itachibot



a) Gaucher's disease

b) Abetalipoproteinemia

c) Whipple's disease

d) Protein losing enteropathy

Sold by @itachi... purchased this from some... else, If yo... ...ay have been scammed.

Question 10:

The findings associated with abetalipoproteinemia are :



a) Figure A,B & C

b) Figure B & C

Sold by @itachibot

c) Figure A & B

d) Figure A & C

Question 11:

All are true about the types of celiac disease except :

a) Latent type is associated with HLA DQ 2,8

b) Atypical type has a negative serology

c) Silent type has a positive serology

d) Classic type presents with diarrhea and steatorrhea

Question 12:

What is the most common extraintestinal manifestation of celiac disease?

a) Dermatitis herpetiformis

b) Infertility

c) Anemia

d) Peripheral neuropathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

A 15-year-old girl presented with fatigue, chronic diarrhoea, weight loss, bone pain, and abdominal distension. Investigations revealed iron deficiency anemia and osteoporosis. Which of the following is the single best test to be done for her evaluation?

a) TSH levels

b) C-peptide levels

c) Urine sugar and ketone

d) IgA tissue transglutaminase antibody

Question 14:

A patient who presented with severe malabsorption is found to be positive for IgA tissue trans-glutaminase antibodies. What will be the next best step?

a) Do an intestinal biopsy

Sold by @itachibot

b) Check for anti-endomysial antibodies

c) Check for anti-gliadin antibodies

d) Start gluten-free diet

Question 15:

A 30-year-old man presented with chronic diarrhea, steatorrhea, and weight loss. IgA tissue trans-glutaminase antibodies were found to be positive. Duodenal biopsy revealed the following. Which of the following complications would you not expect in this patient?



a) Intestinal lymphoma

b) Intestinal ulcers

c) Pancreatic insufficiency

d) Collagenous sprue

Question 16:

A 46-year-old lady with celiac disease is recommended a gluten-free diet. Which among the following can she consume?

a) Wheat

b) Barley

c) Rye

d) Rice

Sold by @itachibot

Question 17:

When will you diagnose a patient with refractory celiac disease?

a) No remission of disease despite gluten free diet

b) Intestinal ulcers on biopsy

c) No response to steroids

d) Development of a subepithelial collagen layer

Question 18:

Match the following disorders with their respective sites of involvement:

a) 1-ii , 2-i , 3-iii

b) 1-iii, 2-ii , 3-i

c) 1-i , 2-iii , 3-ii

d) 1-ii, 2-iii, 3-i

Question 19:

All are true about Whipple's disease except :

a) Caused by gram positive organism Tropheryma whipplei

b) Culture of organism from biopsy specimen helps in early diagnosis

c) Males affected more than females

d) Multiple systems are affected

Question 20:

What is the most common CNS manifestation associated with Whipple's disease?

a) Progressive dementia

b) Supranuclear ophthalmoplegia

c) Opsoclonus myoclonus syndrome

d) Oculomasticatory myorrhythmia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 21:

What is the most common cardiovascular presentation seen in patients with Whipple's disease?

a) Endocarditis

b) Pericarditis

c) Myocarditis

d) Pericardial effusion

Question 22:

A 38-year-old man presented with chronic diarrhea, weight loss, and joint pains. A duodenal biopsy was done. Which of the following drug is used in the maintenance phase of treatment of this condition?



a) Penicillin G

b) Ceftriaxone

c) Meropenem

d) Trimethoprim / Sulfamethoxazole

Question 23:

A 42-year-old man presented with diarrhea, floating stools, and weight loss for 3 months. Celiac serology was negative. Biopsy of the small intestinal mucosa revealed villous atrophy

Sold by @itachibot

and crypt hyperplasia with normal mucosal thickness. What is the most likely diagnosis?

a) Celiac disease

b) Whipple's disease

c) Tropical sprue

d) Radiation enteritis

Question 24:

Which of the following would you prescribe for a patient with tropical sprue?

a) Tertacycline + Vitamin B12

b) Metronidazole + Folic acid

c) Amoxicillin + Vitamin B12

d) Tetracycline + Folic acid

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 25:

Which of the following drug is used for the management of a patient with small intestinal bacterial overgrowth?

a) Ciprofloxacin

b) Azithromycin

c) Rifaximin

d) Streptomycin

Answer Key

Question No.	Correct Option
1	b
2	b
3	d
4	a
5	b
6	d

Sold by @itachibot

7	a
8	d
9	b
10	c
11	b
12	a
13	d
14	a
15	c
16	d
17	a
18	a
19	b
20	a
21	a
22	d
23	c
24	d
25	c

If you purchase... you may have...

Detailed Explanations

Solution to Question 1:

Constipation is not a feature of malabsorption syndrome.

Malabsorption syndrome refers to a spectrum of conditions of various etiologies that arises due to diminished absorption of one or more dietary nutrients.

The general features of malabsorption syndrome are abdominal pain, diarrhea, steatorrhea, and weight loss.

Anemia may be present when there is defective absorption of iron, folate, or vitamin B12. Hypocalcemia and tetany can be seen due to defective absorption of vitamin D.

Solution to Question 2:

D-xylose test is useful for the assessment of intestinal mucosal function.

D-xylose is a monosaccharide that is directly absorbed by the intestinal mucosa without undergoing enzymatic digestion.

Sold by @itachibot

The test is performed by administering 25g of D-xylose orally followed by the collection of urine 5h later. The test is considered abnormal (positive) if urinary D-xylose is <4.5g. A positive D-xylose test indicates abnormal small intestinal mucosa.

Solution to Question 3:

A urinary D-xylose <4.5g after 5 hours of oral D-xylose (abnormal) can be seen in Crohn's disease.

D-xylose is a monosaccharide that is directly absorbed by the small intestinal mucosa without undergoing enzymatic digestion. An abnormal (positive) D-xylose test indicates abnormal small intestinal mucosal absorption.

A positive D-xylose test is seen in:

• Celiac disease

• Whipple's disease

• Tropical sprue

• Short bowel syndrome

• Crohn's disease

The test is of limited clinical value today and mostly has been replaced by small intestinal biopsy.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

A small bowel diverticulum can result in malabsorption.

A small bowel diverticulum is a blind pouch protruding from the small intestine that communicates with the lumen of the gut. It becomes a potential site for bacterial overgrowth which causes malabsorption. The growing gut bacteria depletes vitamin B12 and deconjugates bile acids leading to steatorrhea, diarrhea, and macrocytic anemia.

Anterior colon resection, ulcerative colitis, and sigmoid volvulus affect the large intestine and therefore do not cause malabsorption.

Solution to Question 5:

The most common cause of short bowel syndrome in adults is mesenteric vascular occlusion.

Short bowel syndrome (SBS) is a malabsorption disorder caused by a lack of functional small intestine. In adults, the common causes are :

• Mesenteric vascular disease (e.g: mesenteric vascular occlusion.)

• Primary mucosal and submucosal disease (e.g: Crohn's disease)

• Operations without any preexisting small-intestinal disease (e.g: after trauma)

Sold by @itachibot

In children, necrotizing enterocolitis is the most common cause.

Solution to Question 6:

Homocystinuria is not a protein malabsorption disorder. It is a disorder of methionine metabolism that leads to increased homocysteine levels.

Proteins are broken down by pancreatic enzymes and are absorbed as amino acids and peptides. So, protein malabsorption usually occurs only in pancreatic insufficiency. Additionally, there are three rare genetic disorders that lead to protein malabsorption. They are:

• Enterokinase deficiency- Defective conversion of the proenzyme trypsinogen to trypsin. It is characterised by:

• Diarrhoea

• Growth retardation

• Hypoproteinemia

• Hartnup disease- Defective transport of tryptophan and other neutral amino acids. It is characterised by:

• Pellagra-like rash

• Neuropsychiatric symptoms

• Cystinuria- Defective transport of cystine and other dibasic amino acids. It is characterised by:

• Renal calculi

• Chronic pancreatitis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

The above findings on Schilling test show reduced absorption of vitamin B12 irrespective of all interventions. This is suggestive of an ileal disease.

Solution to Question 8:

The clinical scenario describes a case of malabsorption syndrome (chronic diarrhea and steatorrhea), a normal xylose test and normal duodenal biopsy with abnormal schilling test points towards pancreatic insufficiency.

A normal xylose test and normal duodenal biopsy rules out small intestinal pathologies like celiac disease (option A) and intestinal lymphangiectasia (option C), presence of steatorrhea rules out ulcerative colitis (option B) as it affects the large bowel which is not the main site of absorption of fat.

Sold by @itachibot

Ileal disease is also characterized by normal xylose test, normal duodenal biopsy and steatorrhea and abnormal schilling test but among the given options the most appropriate answer would be pancreatic insufficiency due to chronic pancreatitis.

Pancreatic insufficiency is characterised by the deficiency of exocrine pancreatic enzymes that result in maldigestion and malabsorption.

The features in favor of the diagnosis:

• Steatorrhea - a deficiency of pancreatic lipases can lead to undigested fat in stools.

• D-xylose test - d-xylose is a monosaccharide that is directly absorbed by the intestinal mucosa without undergoing any enzymatic action. Normal d-xylose test denotes intact mucosa.

• Schilling test - it is abnormal as pancreatic enzymes are required for the absorption of vitamin B12.

Diagnosis of various malabsorption syndromes:

Pathologies	D xylose test	Duodenal mucosal biopsy	Schilling test
Celiac disease	Decreased	Abnormal; Short or absent villi; epithelial cell damage; crypts hypertrophy	Normal
Ileal disease	Normal	Normal	Abnormal
Chronic pancreatitis	Normal	Normal	Abnormal; normal after pancreatic enzymes substitution
Intestinal lymphangiectasia	Normal	Abnormal; dilated lymphatics, clubbed villi	Normal
Bacterial overgrowth syndrome	Normal or may be abnormal	Usually normal or can have diffuse non specific damage to villi	Usually abnormal; normal after antibiotics administration

Solution to Question 9:

The image showing retinitis pigmentosa and presence of acanthocytes along with the given clinical features is suggestive of abetalipoproteinemia.

Abetalipoproteinemia is a rare genetic disorder in which there is defective chylomicron formation. Chylomicrons are required for the absorption and transport of fats and fat-soluble vitamins.

It presents in early childhood with steatorrhoea and diarrhea, hypolipidemia, failure to thrive, ataxia, sensory and motor neuropathies.

Solution to Question 10:

Figure A shows vacuoles (unabsorbed fat) in the cytoplasm of the enterocytes. Figure B shows acanthocytes. These findings are associated with abetalipoproteinemia.

Sold by @itachibot

Figure C shows distended macrophages in the lamina propria and periodic acid–Schiff (PAS)-positive granules as seen in Whipple's disease.

Acanthocytes are abnormal red blood cells with spiked cell membranes due to thorn-like projections. Causes of acanthocytes in peripheral smear are:

- Abetalipoproteinemia
- Renal disease
- Liver disease
- Hemolytic anemia





Sold by @itachibot

Sold by @itachi... purchased this from some... else, ...y have been scammed.

Solution to Question 11:

Atypical celiac disease has a positive serology with atypical symptoms.

Serology in celiac disease refers to either anti-tissue transglutaminase antibodies (anti-tTG) or anti-endomysial antibodies.

Note: Tropical sprue and celiac disease have similar clinical, endoscopic and histologic findings. A diagnosis of tropical sprue must be considered with a negative celiac serology.

	Type of celiac disease	Serology	Features
A	Classic	+	Diarrhea, steatorrhea
B	Atypical	+	Iron deficiency anemia, osteopenia, ataxia, peripheral neuropathy, infertility
C	Silent	+	No features
D	Latent	-	HLA DQ 2,8

Solution to Question 12:

The most common extraintestinal manifestation of celiac disease is dermatitis herpetiformis (DH).

DH is characterized by intensely pruritic small papules and vesicles on the extensor surfaces of extremities, buttocks, and neck. It can be considered a celiac disease of the skin.

Other extraintestinal manifestations of celiac disease are:

• Anemia

Sold by @itachibot

• Low bone density

• Ataxia

• Peripheral neuropathy

• Infertility

Celiac disease is often associated with other conditions such as type 1 diabetes mellitus, autoimmune thyroid disease, IgA deficiency, bird fancier's lung, and Down syndrome.

The image below shows dermatitis herpetiformis.

If you purcha... you may have... else,

Solution to Question 13:

The presence of fatigue, chronic diarrhea, weight loss and abdominal distention along with investigations showing features of malabsorption (iron deficiency) lead to the diagnosis of celiac disease or gluten sensitive enteropathy. The single best test for which is checking serum for the presence of IgA tissue transglutaminase (anti-TTG) antibodies (Option D).

Celiac disease is a disorder characterized by an immunological response to gluten causing damage to the proximal intestinal mucosa (villous atrophy) and malabsorption of nutrients.

Symptoms usually begin 6 to 24 months from the introduction of gluten into diet, although most patients go undiagnosed.

The common manifestations are - irritability, abdominal distention, chronic diarrhea, weight loss or muscle wasting and extra-intestinally, the symptoms observed are aphthous ulcers, delayed puberty, iron deficiency anemia and short stature.

Screening for celiac disease is by testing for tissue transglutaminase IgA, endomysial, and deamidated antigliadin antibodies. IgA levels in serum are measured to detect false-negative results because of IgA deficiency.

Small intestinal biopsy shows characteristic villus blunting, crypt hyperplasia, and increased intraepithelial lymphocytes.

Sold by @itachibot

The treatment involves the exclusion of gluten-containing food from diet.

Long term complications observed are osteoporosis, anemia, infertility and malignancy (enteropathy related-T cell lymphoma is the most common association)

The Marsh classification is used to categorize celiac disease into different types and to quantify severity of disease involvement.

Solution to Question 14:

In this patient with positive serology, a biopsy should be done next, to confirm the diagnosis of celiac disease.

Tissue transglutaminase Ab (tTG-IgA) is the best screening test. IgA endomysial antibodies (EMA) are most specific.

But a small-intestinal biopsy is required to establish a diagnosis of celiac disease, before starting treatment.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 15:

The above scenario along with the presence of IgA tissue trans-glutaminase antibodies and biopsy showing villous atrophy, crypt hyperplasia, lamina propria infiltrates indicate a case of celiac disease. Pancreatic insufficiency is not a complication of celiac disease.

The complications of celiac disease are:

• Intestinal lymphomas

• Intestinal ulcers- also known as ulcerative jejunoileitis

• Collagenous sprue- development of a subepithelial collagen layer

Solution to Question 16:

Rice can be consumed on a gluten-free diet.

The primary management of celiac disease is avoiding gluten in the diet.

Diet avoided (gluten)"BROW"	Diet Allowed"So MRP"
Barley	Soya bean
Rye	Maize
Oats	Rice
Wheat	Potato

Sold by @itachibot

Solution to Question 17:

Refractory celiac disease is diagnosed when there is no remission of disease despite a strict gluten-free diet for 6-12 months.

Patients are symptomatic and have persistent small intestinal villous atrophy on biopsy. They respond well to steroids.

Solution to Question 18:

The sites involved in the given malabsorption syndromes are as follows:

Disorder	Site of involvement
Celiac disease	Proximal small intestine
Whipple's disea se	Proximal > Distal small intes tine
Tropical sprue	Whole small intestine (panint estine)

Solution to Question 19:

Tropheryma whipplei is difficult to culture from the biopsy specimen. Hence, it cannot be used for the early diagnosis of Whipple's disease.

Whipple's disease is a chronic multisystem disorder caused by the gram-positive bacteria Tropheryma whipplei. It primarily causes malabsorption along with central nervous system & cardiac problems. It mostly affects men and presents with fever, diarrhea weight loss, and arthralgia.

The diagnosis of Whipple's disease is made through a biopsy which shows the presence of PAS-positive macrophages in the small intestine and other organs with evidence of disease.

Solution to Question 20:

The most common CNS manifestation associated with Whipple's disease is progressive dementia.

Other important CNS manifestations in Whipple's disease are:

• Personality and mood changes

• Progressive supranuclear ophthalmoplegia

• Focal neurologic presentations

Sold by @itachibot

• Oculomasticatory myorrhythmia - an unusual combination of rhythmic eye and jaw movements which is pathognomonic of Whipple's disease.

Whipple's disease associated with CNS symptoms has a poor prognosis.

Note: Opsoclonus myoclonus syndrome is a paraneoplastic syndrome associated with neuroblastoma.

Solution to Question 21:

The most common cardiovascular presentation in Whipple's disease is culture-negative endocarditis.

Other cardiovascular findings include:

• Congestive heart failure

• Pericarditis

• Myocarditis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

The above image shows PAS-positive foamy macrophages present in the lamina propria. This along with the given features suggests Whipple's disease. Cotrimoxazole is used in the maintenance phase of the treatment of Whipple's disease.

Treatment of Whipple's disease:

• Induction phase- ceftriaxone or meropenem or penicillin G + streptomycin - 2 weeks

• Maintenance phase- trimethoprim / sulfamethoxazole (TMP/SMX) for 1 year. It has high CNS penetration so it is used in maintenance to prevent relapse.



Sold by @itachibot

Solution to Question 23:

The history of malabsorptive symptoms, a negative celiac serology, and the biopsy findings are suggestive of tropical sprue.

Improvement in symptoms after folic acid intake is a hallmark of tropical sprue.

Tropical sprue is an acquired malabsorptive condition of probable infectious aetiology. It is seen in people from the tropics such as India, Southeast Asia and South America.

Tropical sprue and celiac disease have similar clinical, endoscopic and histologic findings. However, a positive celiac serology (anti-tissue transglutaminase and anti-endomysial antibodies) can be used to rule out tropical sprue.

Solution to Question 24:

The treatment of tropical sprue is tetracycline + folic acid.

Because of marked folate deficiency, folic acid is most often given together with broad spectrum antibiotics for up to 6 months.

Solution to Question 25:

Rifaximin is the preferred drug for the management of small intestinal bacterial overgrowth (SIBO).

The antibiotics are given for around 3 weeks. The other drugs that can be used are :

- Tetracyclines
- Cephalosporins
- Metronidazole
- Amoxicillin-Clavulanic acid

Rifaximin is the antibiotic used in hepatic encephalopathy to lower blood ammonia levels.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Other Gastrointestinal Conditions

Question 1:

An upper GI endoscopy is performed in a patient with dyspeptic symptoms. It shows the presence of thickened gastric folds. Which of the following differential diagnoses would you not consider in this patient?

a) Zollinger–Ellison syndrome

b) Lymphoma

c) Menetrier's disease

d) Pancreatic rests

Question 2:

Which part of the stomach is usually spared in Menetrier's disease?

a) Body

b) Fundus

c) Antrum

d) Greater curvature

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 46-year-old man presented with a history of diarrhea for 5 months, occasional abdominal pain, and weight loss. Upper GI endoscopy showed the following finding. Which of the following is least likely to occur in this patient?

Sold by @itachibot



a) Hypoproteinemia

b) Hypochlorhydria

c) Edema

d) Hematemesis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 4:

A 6-year-old child with vomiting, abdominal pain, weight loss, and peripheral edema underwent an upper GI biopsy. Histology showed foveolar hyperplasia and glandular atrophy within the gastric mucosa. Which of the following is most commonly associated with this condition?

a) Cytomegalovirus

b) Helicobacter pylori

c) Herpes simplex virus

d) Epstein Barr virus

Question 5:

Overexpression of which of the following causes Menetrier's disease?

a) TNF-α

b) TGF-α

c) TNF-β

Sold by @itachibot

d) TGF-β

Question 6:

A 24-year old woman underwent extensive workup for unexplained weight loss, vomiting, and diarrhea. She was suspected to have Menetrier's disease. Which of the following investigations will confirm the diagnosis?

a) Endoscopic full thickness biopsy

b) CECT abdomen

c) Endoscopic punch biopsy

d) MRI abdomen

Question 7:

A middle-aged man presented with a history of dyspepsia. He has been taking pantoprazole for the last few months, however, his symptoms persisted. Upper GI endoscopy was done, and a full-thickness gastric biopsy was taken. The histopathological appearance is shown below. What is the treatment of choice?



a) Partial gastrectomy

b) Lansoprazole

c) Cetuximab

d) Octreotide

Sold by @itachibot
If you received this from someone else, it has been scammed.

Sold by @itachibot

Question 8:

What is the most common site for a MALToma?

a) Duodenum

b) Stomach

c) Ileum

d) Cecum

Question 9:

Which of the following is false about MALTomas?

a) Associated with autoimmune conditions

b) History of chronic gastritis usually seen

c) Usually diagnosed in elderly patients

d) Arises from T lymphocytes

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 10:

A 62-year-old woman underwent an endoscopic biopsy of a gastric tumor. The biopsy shows dense lymphocytic infiltration with lymphoepithelial lesions in the gastric glands. Which of the following investigations will confirm your diagnosis?

a) Cytogenetic testing

b) Immunophenotyping

c) Electron microscopy

d) Fluorescent microscopy

Question 11:

Immunophenotyping from a gastric tumor in an elderly man showed the following findings. The tumor was found to be limited to the gastric mucosa. What is the treatment of choice?

a) Rituximab and cyclophosphamide

b) Three-dimensional conformal radiotherapy

c) Omeprazole, clarithromycin and amoxicillin

d) Radical gastrectomy

Sold by @itachibot

Question 12:

A 65-year-old smoker presents with postprandial abdominal pain and diarrhea for the past 1 year. On examination, she is malnourished, and a bruit is heard in the epigastrium. What is the most probable diagnosis?

a) Acute mesenteric ischemia

b) Chronic mesenteric ischemia

c) Wilkie's syndrome

d) Nutcracker syndrome

If you bought this from someone else, you've been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	c
3	d
4	a
5	b
6	a
7	c
8	b
9	d
10	b
11	c
12	b

Detailed Explanations

Solution to Question 1:

Pancreatic rests are not a differential diagnosis for hypertrophied gastric folds. These usually present as a single lump in the stomach and are formed by ectopic pancreatic tissue.

Sold by @itachibot

Thickened gastric folds can represent a local malignant process, inflammatory process, or hormonal response. The differential diagnoses of large gastric folds include:

• Menetrier's disease

• Zollinger–Ellison syndrome - hypergastrinemia leads to parietal cell hyperplasia and prominent folds

• Malignancy- lymphoma, infiltrating gastric cancer

• Infections- histoplasmosis, tuberculosis.

Solution to Question 2:

The antrum is usually spared in Menetrier's disease.

Foveolar hyperplasia results in the development of large rugal folds in the body and fundus of the stomach, sparing the antrum. These hypertrophied rugal folds give a cobblestone or cerebriform appearance to the gastric mucosa.

The image given below shows the endoscopic appearance of Menetrier's disease.



Solution to Question 3:

The given case scenario and endoscopy image showing large, rugal gastric folds are suggestive of Menetrier's disease. Among the given options, hematemesis is least likely to occur in this patient.

Large, gastric rugal folds in Menetrier's disease occur due to massive overgrowth of mucous cells (foveolar cells) lining the stomach.

Patients usually present with epigastric pain, nausea, diarrhea, vomiting, anorexia, peripheral edema, and weight loss. Occult gastrointestinal bleeding may occur, but overt bleeding is unusual. If overt bleeding is seen, it is usually due to superficial mucosal erosions.

Sold by @itachibot

They develop protein-losing gastropathy, leading to hypoproteinemia as a result of increased secretion of gastric mucus. There is also a decrease in the number of parietal cells which leads to decreased acid secretion (hypochlorhydria) in the stomach. They are at an increased risk of developing gastric carcinoma.

Solution to Question 4:

The given case scenario is suggestive of Menetrier's disease. In children, it occurs commonly due to the infection by cytomegalovirus.

The histopathological features of Menetrier's disease include:

- Foveolar hyperplasia - hyperplasia of surface and glandular mucosa
- Reduction in oxyntic glands, parietal cells, and chief cells
- Dilated, tortuous gastric glands
- Elongation of pits of gastric glands
- Eosinophilic infiltration in lamina propria

The image given below shows the histopathological appearance of Menetrier's disease.



Solution to Question 5:

Overexpression of TGF-α (transforming growth factor-alpha) causes Menetrier's disease. This overexpression results in overstimulation of the epidermal growth factor receptor (EGFR) pathway and increased proliferation of mucous cells, resulting in mucous cell (foveolar) hyperplasia.

If yo... ...y @itachibot ...from someone else, ...commed.

Sold by @itachibot

Solution to Question 6:

Menetrier's disease is confirmed by endoscopic full-thickness biopsy.

A full-thickness biopsy allows for the visualization of the cells on the surface as well as the pits of the gastric glands. The histopathological specimen shows the following features:

• Foveolar hyperplasia - hyperplasia of surface and glandular mucosa

• Reduction in oxyntic glands, parietal cells, and chief cells

• Dilated, tortuous gastric glands

• Elongation of pits of gastric glands

• Eosinophilic infiltration in lamina propria

A punch biopsy can only sample the surface of the gastric gland and hence is not preferred.

The image given below shows the histopathological appearance of Menetrier's disease.



Solution to Question 7:

The given clinical scenario and the image showing foveolar hyperplasia are suggestive of Menetrier's disease. The treatment of choice is cetuximab.

Increased epidermal growth factor receptor(EGFR) signaling has been implicated in the pathogenesis of Menetrier's disease. Cetuximab is an EGFR inhibitor, and blocks this signaling pathway and helps in the improvement of the symptoms. Therefore, it is the first line of treatment.

Medical therapy with proton pump inhibitors (lansoprazole) and somatostatin analogues (octreotide) may also be tried to manage symptoms. If the patient continues to have protein loss despite medical therapy, then total gastrectomy is indicated.

The image given below shows the histopathological appearance of Menetrier's disease.

Sold by @itachibot



Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Solution to Question 8:

The most common site for mucosa-associated lymphoid tissue (MALT) lymphoma or MALToma is the stomach.

MALToma is a type of extranodal marginal zone lymphoma that most commonly occurs in the stomach and to a lesser extent in the salivary glands, lungs, small intestine, etc. It occurs as a result of the malignant proliferation of B cells in the marginal zone of mucosa-associated lymphoid tissue (MALT).

Solution to Question 9:

MALToma arises from B lymphocytes present in the marginal zone of the lymphoid tissue.

The majority of the cases of MALToma are associated with H. pylori induced chronic gastritis. This infection results in an increased proliferation of lymphocytes in the mucosa of the stomach causing MALToma. It is initially low grade and if the infection is not treated, it later progresses to high grade.

The patients usually present with non-specific symptoms like epigastric pain and discomfort. It is usually diagnosed in patients around 60 years of age. It is also associated with autoimmune diseases like Sjogren's syndrome and Hashimoto's thyroiditis.

Solution to Question 10:

The given clinical scenario and lymphoepithelial lesions seen in the gastric glands is suggestive of MALToma. The diagnosis is confirmed by immunophenotyping.

Sold by @itachibot

Histopathological examination of a MALToma shows the presence of lymphoepithelial lesions. These are caused due to the invasion of gastric glands and crypts by the neoplastic lymphocytes. These lymphocytes are positive for CD19, CD20 and negative for CD5 and CD10. Biopsy or aspiration of the bone marrow is also done to see the bone marrow involvement.

Radiological investigations like CT scans, MRIs, and PET scans are useful in identifying the extent of the tumor and infiltration of the tumor to the adjacent organs.

Note: Cytogenetic testing will be helpful in identifying chromosomal abnormalities.

The image given below shows the histopathological appearance of a MALToma.



The below image shows a sample of a MALToma showing positive immunophenotyping for CD20.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 11:

The given case scenario is suggestive of a low-grade gastric MALToma. The treatment of choice for this condition is H. pylori eradication therapy which can be done with the OCM regimen (omeprazole, clarithromycin and metronidazole).

Immunophenotyping in MALTomas shows positive staining for CD19 and CD20 which is suggestive of a B-cell neoplasm. Negative staining for CD5 and CD10 is seen, suggesting it arises from the marginal zone of lymphoid tissues.

Chronic infection with H. pylori leads to the development of MALTomas. When these lesions are limited to the mucosa, treatment of the infection alone is sufficient to cause regression of the tumour.

In high-grade tumors or the tumors which invade into the adjacent or distant organs, radiotherapy and chemotherapy are required to treat the tumor. Radiotherapy is usually three-dimensional conformal radiotherapy. The chemotherapeutic agents used in the treatment of MALToma are rituximab, cyclophosphamide, fludarabine, doxorubicin, vincristine, and prednisone. Radical gastrectomy has a limited role and is done when the disease is confined to the stomach.

Solution to Question 12:

The given case scenario is suggestive of chronic mesenteric ischemia.

Chronic mesenteric ischemia occurs due to atherosclerosis of the celiac or mesenteric arteries. It is common in older women, and they usually have a history of smoking, coronary artery disease, cerebrovascular disease, or peripheral artery disease.

The important clinical features of chronic mesenteric ischemia are:

Sold by @itachibot

• Postprandial abdominal pain occurs as a result of a mismatch between the increased demand and decreased blood flow to the intestine. This is called intestinal angina

• Weight loss- patients avoid eating due to postprandial pain and this leads to significant weight loss

• Diarrhea- occurs as a result of decreased absorption

• Physical examination may reveal epigastric bruit in almost half of the patients.

The gold standard investigation to diagnose the blockade is mesenteric angiography. Once the disease is confirmed, the patients have to be started with antiplatelets and mesenteric angioplasty and stenting should be performed.

Option A: Acute mesenteric ischemia present with acute abdomen, bloody diarrhea and nausea. Angiography will show occlusion of superior mesenteric artery.

Option C: Wilkie's syndrome occurs due to compression of the third portion of the duodenum by the superior mesenteric artery. It can present similarly, however is associated with vomiting and rapid weight loss. It usually occurs in young, asthenic women.

Option D: Nutcracker syndrome occurs due to compression of the left renal vein, between the aorta and the superior mesenteric artery. It presents with varicoceles and hematuria.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Large and medium vessel vasculitis

Question 1:

Which of the following is an example of large vessel vasculitis?

a) Polyarteritis nodosa

b) Takayasu disease

c) Behcet's disease

d) Cogan's syndrome

Question 2:

Biopsy performed on a patient with suspected vasculitis showed the presence of granulomas. Which of the following is the most likely diagnosis?

a) Microscopic polyangiitis

b) Behcet's disease

c) Polyarteritis nodosa

d) Churg- Strauss syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 42-year-old man presented with complaints of fever, malaise, and weight loss. His lab investigations showed mild anemia, elevated ESR, and serology was positive for cANCA. Which of the following is the most likely diagnosis?

a) Cryoglobulinemic vasculitis

b) Granulomatosis with polyangiitis

c) Microscopic polyangiitis

d) Churg-Strauss syndrome

Question 4:

Which of the following is not true about giant cell arteritis?

a) Seen in young women

Sold by @itachibot

b) May lead to ischemic optic neuropathy

c) Good response to glucocorticoid therapy

d) Anemia is present

Question 5:

A 65-year-old woman came with complaints of headache for the past 1-month. She also reported aching of the jaw during her meals. Over the last week, she has experienced episodes of painless visual loss which lasted for around 10 minutes. Which of the following is the best investigation for the diagnosis of her condition?

a) Temporal artery biopsy

b) Colour doppler of temporal artery

c) CT angiography

d) MR angiography

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 6:

A 50-year-old woman presents to your OPD with complaints of pain and stiffness of the shoulders which is worse in the morning. She works as a bus conductor and tells you that she has difficulty raising her shoulders to ring the bell. On examination, there is diffuse tenderness over the shoulders. Which of the following vasculitis is closely associated with her presentation?

a) Polyarteritis nodosa

b) Granulomatosis with polyangiitis

c) Takayasu arteritis

d) Giant cell arteritis

Question 7:

A 60-year-old woman presents with severe right-sided throbbing headache. She also gives history of pain and stiffness of the shoulders and hips. Her lab values show an ESR of 80 mm/hr. What is the next best step in her management?

a) Start beta-blockers

b) Obtain non contrast head CT

c) Perform temporal artery biopsy

Sold by @itachibot

d) Start steroids

Question 8:

Which is the most common artery involved in Takayasu arteritis?

a) Subclavian artery

b) Common carotid artery

c) Internal carotid artery

d) External carotid artery

Question 9:

A 27-year-old woman presents to the hospital with fever, malaise, and arthralgia for a duration of 1 month. She also complains of pain, numbness, and weakness in her right arm, which has been increasing over the past 1 week. Clinical examination reveals weak radial pulses, and her laboratory investigations show elevated ESR. What is the most likely diagnosis?

a) Coarctation of aorta

b) Giant cell arteritis

c) Takayasu arteritis

d) Syphilitic aortitis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

In a patient with suspected Behcet's disease which of the following tests can be used to confirm the diagnosis?

a) Foshay test

b) Tzanck test

c) Pathergy test

d) Cardiolipin antibody test

Question 11:

Sold by @itachibot

A 35-year-old woman presented with recurrent oral and genital ulcerations. An inflammatory skin reaction was observed after intradermal administration of normal saline. Which of the following statements is incorrect about her condition?

a) Associated with HLA B51

b) Associated with painless scarring oral ulcers

c) Associated with erythema nodosum

d) Associated with peripheral vein thrombosis

Question 12:

Which of the following is a medium vessel vasculitis?

a) Polyarteritis nodosa

b) Behcet's disease

c) Microscopic polyangiitis

d) Cogan's syndrome

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

A 47-year-old man presented to the hospital with the findings shown in the image below. He also complained of pain in the upper and lower limbs. On examination, his BP was 160/100 mmHg and his peripheral pulses were palpable. Serology was negative for ANCA. What is the most probable diagnosis?

Sold by @itachibot

a) Takayasu arteritis

b) Microscopic polyangiitis

c) Cryoglobulinemic vasculitis

d) Polyarteritis nodosa

Question 14:

In a patient diagnosed with polyarteritis nodosa which of the following arteries are characteristically spared?

a) Renal arteries

b) Mesenteric arteries

c) Pulmonary arteries

d) Bronchial arteries

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 15:

Which of the following is true about polyarteritis nodosa?

a) Necrotising inflammation of large vessels

b) Patient has hypogammaglobulinemia

c) 90% is associated with ANCA positivity

d) 30% is associated with Hepatitis B

Question 16:

What is the most common cause of vasculitis in children in India?

a) Henoch-Schonlein purpura

b) Kawasaki disease

c) Juvenile idiopathic arthritis

d) Rheumatic fever

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	b
2	d
3	b
4	a
5	a
6	d
7	d
8	a
9	c
10	c
11	b
12	a
13	d
14	c
15	d
16	b

If you purchased this, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Takayasu disease is an example of large vessel vasculitis.

Solution to Question 2:

Churg-Strauss syndrome is the most likely diagnosis as it is a granulomatous vasculitis.

The vasculitis can be classified as:

Granulomatous vasculitis	Non-granulomatous vasculitis
Giant cell arteritisTakayasu arteritisGranulomatosis with polyangiitisChurg-Strauss syndrome	Polyarteritis nodosaKawasaki diseaseBehcet's diseaseMicroscopic polyangiitisHenoch-Schonlein purpura

Sold by @itachibot

Solution to Question 3:

Granulomatosis with polyangiitis (Wegener's granulomatosis) is the most likely diagnosis as it is associated with cANCA.

Solution to Question 4:

Giant cell arteritis is common in older women (>50 years of age).

Giant cell arteritis is an inflammatory vasculitis of the large vessels, commonly seen in females over 50 years of age. It usually involves the temporal artery.

It is characterized by fever, anemia, and headache. Depending on the artery involved, the patient may have tenderness over the affected artery, scalp pain, tongue and jaw claudication. Untreated patients may develop ischemic optic neuropathy.

Laboratory investigations reveal elevated ESR and normochromic normocytic anemia. It shows good response to glucocorticoid therapy.

Solution to Question 5:

The given clinical presentation of headache, jaw claudication, and painless visual loss is suggestive of giant cell arteritis. The best investigation for the diagnosis of giant cell arteritis is temporal artery biopsy.

Histopathology shows panarteritis with the presence of mononuclear cells and giant cells. The biopsy may not be positive in all patients due to patchy histologic findings. Since involvement of the vessel may be segmental, positive yield is increased by obtaining a biopsy segment of 3–5 cm together with serial sectioning of biopsy specimens.

Note : According to the 21st edition of Harrison's Medicine, as involvement of the vessel may be segmental, getting a biopsy segment of 3–5 cm together with serial sectioning of biopsy specimens would help to increase a positive yield. And Robbin's 10th edition mentions that an adequate biopsy requires at least a 1-cm segment.

Solution to Question 6:

The given clinical presentation of painful, tender, and stiff shoulders with restricted movement points toward a diagnosis of polymyalgia rheumatica, which is closely associated with giant cell arteritis.

Polymyalgia rheumatica is characterized by pain and stiffness (worst in the morning) in the muscles of the neck, shoulder, back, hips, and thigh. On examination, diffuse tenderness is present without localization to any specific structures. Pain restricts the active range of motion but passive range of motion and muscle strength are normal.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

ESR and CRP are typically elevated in polymyalgia rheumatica whereas creatine phosphokinase (CPK) lies within the normal range. It shows a good response to prednisone therapy.

Solution to Question 7:

The given clinical presentation of headache and symptoms suggestive of polymyalgia rheumatica with raised ESR points towards a diagnosis of giant cell arteritis (GCA), for which the next best step is initiation of steroids (prednisone).

Although temporal artery biopsy is the diagnostic investigation for GCA, it should not delay the initiation of steroids as it helps in reducing symptoms and preventing visual loss. The biopsy findings usually remain positive for at least 1 month after initiation of steroids.

Newer drugs used in the treatment of giant cell arteritis are tocilizumab (anti-IL-6 receptor) and abatacept (CTLA4-Ig).

Solution to Question 8:

The most common artery involved in Takayasu arteritis is the subclavian artery.

Patients present with pain, claudication, and weakness of the affected limb. The arterial pulse in the affected limb may be weak or absent.

The order of arteries affected in Takayasu arteritis is subclavian artery > common carotid artery > abdominal aorta > renal artery > aortic arch = vertebral > coeliac axis > iliac > pulmonary > coronary artery.

Solution to Question 9:

The above clinical features with a marked weakening of pulse in the upper extremities and laboratory findings of elevated ESR are suggestive of Takayasu arteritis. It is also known as pulseless disease.

Takayasu arteritis is a systemic disease and may present with generalized and vascular symptoms. It commonly affects adolescents or young females. Generalized symptoms like fever, night sweats, weight loss, malaise, arthralgias, and anorexia may occur long before vascular symptoms.

Vascular symptoms depend on the vessel involved:

• Arch of the aorta and its branches:

• Subclavian artery (most commonly affected) - It leads to weak pulses in the upper extremities. Hence, it is also known as pulseless disease. It may also lead to Raynaud's phenomenon.

• Carotid arteries- It leads to weak pulses, blindness, and neurological deficits.

• Aortic arch- It may lead to congestive heart failure.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• Abdominal aorta, celiac axis, and superior mesenteric artery - leads to abdominal pain, nausea, and vomiting

• Iliac vessels - leads to claudication of the legs

• Renal arteries - leads to systemic hypertension

• Pulmonary artery - leads to pulmonary hypertension

• Coronary arteries (least commonly affected) - lead to myocardial infarction

Investigations:

• Laboratory tests reveal an elevated ESR level, anemia, and elevated immunoglobulin levels.

• Arteriography is done as a confirmatory test.

Treatment includes glucocorticoid therapy, methotrexate, immunotherapy with anti-TNF agents, and surgery.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Pathergy test can be used for confirming Behcet's disease. It is a nonspecific inflammatory skin reaction to scratch or intradermal saline injection and is a specific manifestation.

The diagnostic criteria in Behcet's syndrome are the presence of recurrent oral ulcerations plus two of the following findings:

• Recurrent genital ulcerations

• Ocular lesions

• Skin lesions

• Positive pathergy test

Option A: Foshay test is used in cat-scratch disease.

Option B: Tzanck test is used to diagnose herpes infection.

Option D: The cardiolipin antibody test is used to detect antiphospholipid antibodies.

Solution to Question 11:

Recurrent oral and genital ulcerations with a positive pathergy test are suggestive of Behcet's syndrome. It is associated with oral ulcers that are painful and heal without scarring.

Behcet's syndrome usually affects young individuals and is associated with HLA-B51 and anti-Saccharomyces cerevisiae antibodies.

The clinical features of Behcet's syndrome include:

• Oral ulcers - painful, non-scarring

• Genital ulcers - painful, scarring

• Ocular involvement - scarring and bilateral panuveitis

Sold by @itachibot

• Skin involvement - folliculitis, acne-like rashes, erythema nodosum, and pyoderma gangrenosum

• Arthritis and peripheral vein thrombosis

Solution to Question 12:

Polyarteritis nodosa (PAN) is a medium vessel vasculitis.

Solution to Question 13:

The most probable diagnosis is polyarteritis nodosa.

The image shows the gangrene of the fingers. The presence of myalgia, the involvement of cutaneous vessels (gangrene with palpable peripheral pulses), renal vessels causing hypertension and serology negative for ANCA make polyarteritis nodosa (PAN) the most probable diagnosis.

Polyarteritis nodosa (PAN) is necrotizing arteritis of the medium-sized blood vessels. It can affect the skin, peripheral nerves, renal vessels, mesenteric vessels, heart, and brain. Lungs are usually spared in PAN.

The disease initially starts with non-specific symptoms like fever, malaise, arthralgia, and myalgia. Later, depending on the artery involved, patients may develop the following symptoms:

• Skin findings - ulcers, digital gangrene, and livedo reticularis

• Renal arteries - hypertension and renal failure

• Mesenteric arteries - abdominal pain, nausea, vomiting, bleeding, and a bowel perforation

Other options:

1) Takayasu arteritis- It is a large vessel vasculitis. Also known as pulseless disease as it may be associated with weak upper extremity pulses.

2) Microscopic polyangiitis- It is a small vessel vasculitis. It is positive for p-ANCA. Presentation is similar to granulomatosis with polyangiitis but without upper airway involvement.

3) Cryoglobulinemic vasculitis- It is frequently associated with hepatitis C. Circulating cryoglubulins and rheumatoid factor can be found in blood. Most common clinical manifestations are cutaneous vasculitis, peripheral neuropathy, glomerulonephritis and arthritis.

Solution to Question 14:

Pulmonary arteries are characteristically spared in polyarteritis nodosa (PAN). PAN mainly involves renal and visceral arteries.

Histopathology of the involved arteries usually shows the following findings:

• Transmural necrotizing vasculitis of the arteries and adjacent veins. Venules are spared.

• Renal arteries - arteritis without glomerulonephritis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Sold by @itachibot

• Fibrinoid necrosis

• Absence of granuloma and eosinophilia

Solution to Question 15:

One-third of polyarteritis nodosa (PAN) cases are associated with Hepatitis B.

Other options:

Option A: Polyarteritis nodosa is a systemic vasculitis which is associated with necrotizing inflammation of small and medium-sized muscular arteries.

Option B: Patients show hypergammaglobulinemia.

Option C: PAN is rarely associated with ANCA positivity. There is no serological test for PAN. Most patients exhibit an elevated leukocyte count with a predominance of neutrophils.

PAN is usually seen in young adults, characterized by rapidly accelerating hypertension, abdominal pain, bloody stools, myalgias, and peripheral neuropathy.

Arterial inflammation is segmental, necrotizing, and transmural, leading to aneurysms and thrombosis.

The involved vessels are typically found at arterial bifurcations, spreading to veins but sparing venules. The most commonly affected organs are the kidney, heart, liver, and gastrointestinal tract, in that order. Pulmonary arteries are not involved. Inadequate blood flow leads to ischemia, ulcers, infarcts, atrophy, or hemorrhage.

Diagnosis relies on histopathological findings from organ biopsies or the demonstration of aneurysms in small to medium-sized arteries of the renal, hepatic, and visceral vasculature.

Untreated PAN has an extremely poor prognosis, with a 5-year survival rate of only 10 to 20%. Common causes of death include gastrointestinal complications, such as bowel infarcts and perforation, as well as cardiovascular issues. Timely treatment is vital for improved outcomes.

Combining prednisone and cyclophosphamide has shown significant benefits in survival rates. Glucocorticoids alone may be sufficient for milder cases. In PAN cases associated with hepatitis B or C, antiviral therapy is combined with glucocorticoids and plasma exchange. Managing hypertension effectively can help reduce vascular complications. Despite successful treatment, relapses occur in approximately 10-20% of patients.

A histopathological image of PAN showing necrotizing transmural inflammation is given below:

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Sold by @itachibot



The image below shows aneurysms in renal vasculature:



Solution to Question 16:

The most common cause of vasculitis in children in India is Kawasaki disease.

Kawasaki disease has a higher incidence in children in the Indian subcontinent.

The most common vasculitis worldwide is Henoch-Schonlein purpura, whereas in the Indian subcontinent and other Asian countries the most common vasculitis is Kawasaki disease.

Sold by @itachibot

Small vessel vasculitis

Question 1:

Which of the following findings would you not expect in a patient diagnosed with small-vessel vasculitis?

a) Palpable purpura

b) Hematuria

c) Hemoptysis

d) Visual loss

Question 2:

Which of the following structures is most commonly involved in cutaneous small-vessel vasculitis?

a) Muscular arteries

b) Arterioles

c) Postcapillary venules

d) Capillaries

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A patient presents with cutaneous vasculitis, glomerulonephritis, and peripheral neuropathy. Which investigation should be performed next to help you diagnose the condition?

a) ANCA

b) RA factor

c) HbsAg

d) MIF

Question 4:

Which is the most common form of vasculitis associated with connective tissue disorders?

a) Medium and small vessel vasculitis

Sold by @itachibot

b) Medium vessel vasculitis

c) Small vessel venulitis

d) Small vessel arteritis

Question 5:

A pathology resident is examining biopsy specimens of patients with clinical features of vasculitis. In which of the following conditions would he report granulomatous inflammation of the vessels?

a) 1, 2, and 3

b) 2, 4, and 5

c) 1, 2, and 5

d) 1, 3, 4, and 5

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A 45-year-old woman presents with pain over the forehead aggravated on bending and occasional hemoptysis. Laboratory investigations reveal elevated ESR and red cell casts in the urine. Her serology is positive for cANCA. What is the most likely diagnosis?

a) Goodpasture's syndrome

b) Granulomatosis with polyangiitis

c) Microscopic polyangiitis

d) Henoch-Schonlein pupura

Question 7:

A patient who presented with skin rashes, otitis media and hematuria has been diagnosed with granulomatosis with polyangiitis. Which of the following drugs would you not use in the management of this patient?

a) Rituximab

b) Cyclophosphamide

c) Abatacept

d) Methotrexate

Sold by @itachibot

Question 8:

A middle-aged woman presents with hemoptysis, proteinuria, and the clinical presentation given below. What is the most likely diagnosis?

a) Churg-Strauss syndrome

b) Leprosy

c) Systemic lupus erythematosus

d) Granulomatosis with polyangiitis

If you purchased this ... you may have been sc... else, ...d.

Question 9:

Which of the following is not a feature associated with Wegener's granulomatosis?

a) Pulmonary nodules

b) Venous thrombosis

c) Non-necrotizing vasculitis

d) Glomerulonephritis

Question 10:

A 57-year-old man presented with complaints of fever, weight loss, and hemoptysis. His serology was positive for pANCA and biopsy showed small vessel vasculitis with neutrophil infiltration. Which of the following is a feature associated with his condition?

Sold by @itachibot

a) Non-necrotizing vasculitis

b) Fibrinoid necrosis

c) Intravascular granuloma formation

d) Immune complex mediated vasculitis

Question 11:

Which of the following features is not useful to distinguish between microscopic polyangiitis and granulomatosis with polyangiitis?

a) Upper airway involvement

b) Granulomas

c) Pulmonary nodules

d) Necrotizing vasculitis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

A triad of skin lesions, asymmetric mononeuritis multiplex and eosinophilia is seen in which of the following conditions?

a) Cryoglobulinemic vasculitis

b) Polyarteritis nodosa

c) Churg Strauss syndrome

d) Giant cell arteritis

Question 13:

A 47-year-old housewife presented with recurrent attacks of asthma and pain and numbness along the distribution of the median and ulnar nerve. Her serology was positive for pANCA. Further evaluation is unlikely to reveal which of the following findings?

a) Peripheral blood eosinophilia

b) Tissue eosinophilia

c) Extravascular granuloma

d) Large vessel vasculitis

Question 14:

Sold by @itachibot

A 6-year-old boy presents to the ER with colicky abdominal pain. His mother tells you that he refuses to walk due to knee pain. On examination, palpable purpura is noted. What is the most likely diagnosis?

a) Henoch Schonlein purpura

b) Hemolytic uremic syndrome

c) Juvenile Idiopathic arthritis

d) IgA nephropathy

Question 15:

Cryoglobulinemia is associated with _____.

a) Hepatitis C

b) Ovarian cancer

c) Diabetes

d) Leukemia

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	d
2	c
3	a
4	c
5	c
6	b
7	c
8	d
9	c
10	b
11	d
12	c
13	d
14	a

Sold by @itachibot

15	a

Detailed Explanations

Solution to Question 1:

Ischemic optic neuropathy associated with giant cell arteritis (large vessel vasculitis) can present as visual loss.

Small vessel vasculitis is the inflammation of arterioles, capillaries, or venules. It starts with nonspecific symptoms like fever, fatigue, and arthralgia. Since it mainly affects the small vessels and capillaries present in the lungs, kidney, and skin, the majority of the patients will have:

• Hemoptysis

• Glomerulonephritis (hematuria, proteinuria)

• Palpable purpura

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 2:

The structure most commonly involved in cutaneous small-vessel vasculitis is postcapillary venules.

Cutaneous small-vessel vasculitis is defined as vasculitis which is limited to the small vessels of the skin. There is no systemic vasculitis and glomerulonephritis. It most commonly affects postcapillary venules.

It is characterized by the presence of leukocytoclasis (presence of neutrophil debris in and around the blood vessels).

Solution to Question 3:

The presence of cutaneous vasculitis, glomerulonephritis, and peripheral neuropathy are suggestive of small-vessel vasculitis. Investigating for ANCA (antineutrophil cytoplasmic antibodies) should be the next step to aid in diagnosis.

Small Vessel Vasculitis/ ANCA- associated Vasculitis

Sold by @itachibot

Granulomatosis with Polyangiitis(Wegener's granulomatosis)	Microscopic Polyangiitis	Eosinophilic granulomatosis with Polyangiitis(Churg Strauss syndrome)
Characterized by a triad ofupper and lower respiratory tractinvolvement withglomerulonephritis.	Characterized bynecrotizing vasculitiswithglomerulonephritis(pauci immune, crescentic)	Characterized byasthma,peripheral and tissue eosinophilia,extravascular granulomaformation, andvasculitisof multiple organs.
Nodular cavitary infiltratesin lungs showing necrotizing granulomas, and vasculitis on lung biopsy	Diffuse alveolar hemorrhage, hemoptysis.No URTI, No granulomas, No pulmonary nodules/cavities.	Mononeuritis multiplex- most common and characteristic. Biopsy showing granulomas with eosinophilic infiltrates.
C-ANCA +	P-ANCA +	P-ANCA +

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Solution to Question 4:

The most common form of vasculitis associated with connective tissue disorders is small-vessel venulitis.

Connective tissue disorders that can cause vasculitis are:

- Systemic lupus erythematosus
- Rheumatoid arthritis
- Relapsing polychondritis
- Sjögren's syndrome

Solution to Question 5:

Biopsy findings of giant cell arteritis, Churg-Strauss syndrome, and Takayasu arteritis show granulomatous inflammation of vessels as they are granulomatous vasculitis.

The vasculitis diseases can be classified as:

Granulomatous vasculitis	Non-granulomatous vasculitis
Giant cell arteritisTakayasu arteritisGranulomatosis with polyangiitisChurg-Strauss syndrome	Polyarteritis nodosaKawasaki diseaseBehcet's diseaseMicroscopic polyangiitisHenoch-Schonlein purpura

Solution to Question 6:

Sold by @itachibot

The given clinical scenario with cANCA positivity is suggestive of granulomatosis with polyangiitis (Wegener's granulomatosis). It is characterized by the involvement of the upper respiratory tract (sinusitis), lower respiratory tract (occasional hemoptysis), and kidneys (red cell casts).

It is a granulomatous, necrotizing vasculitis of the small vessels that mainly affects:

• Upper respiratory tract- presenting with sinusitis, otitis media, and nasal septal perforation.

• Lower respiratory tract- presenting with hemoptysis.

• Kidney (glomerulonephritis)- presenting with proteinuria, hematuria, and hypertension.

Apart from these organs, the disease can also affect the eye and skin. Skin lesions present as palpable purpura or subcutaneous nodules. Laboratory investigations show anemia, leucocytosis, elevated ESR, and ANCA positivity.

Options A and C: In Goodpasture's syndrome and microscopic polyangiitis, the symptoms of upper respiratory tract involvement are very rare.

Option D: Henoch-Schonlein purpura is commonly seen in children. It presents with palpable purpura, joint pain, abdominal pain, and glomerulonephritis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

Abatacept is a disease-modifying antirheumatic drug (DMARD) used in the treatment of rheumatoid arthritis.

Important drugs used in the management of granulomatosis with polyangiitis (Wegener's granulomatosis) are:

• Cyclophosphamide

• Rituximab (anti CD20 antibody)

• Glucocorticoids

• Methotrexate

Solution to Question 8:

The given clinical scenario showing saddle nose deformity along with hemoptysis, and proteinuria is suggestive of granulomatosis with polyangiitis (Wegener's granulomatosis).

Involvement of upper respiratory tract in granulomatosis with polyangiitis manifests as:

• Paranasal sinusitis, serous otitis media

• Nasal ulcers, purulent or bloody nasal discharge

• Nasal perforation leading to the destruction of septum and saddle nose deformity

• Subglottic tracheal stenosis

Sold by @itachibot

Solution to Question 9:

Wegener's granulomatosis is associated with necrotizing vasculitis.

Wegener's granulomatosis predominantly affects the small blood vessels of the upper respiratory tract, lungs, and kidneys. Lung involvement will show multiple, bilateral cavitating nodules and pulmonary infiltrates on chest X-ray.

The radiograph below shows cavitating nodules in a patient with granulomatosis with polyangiitis (Wegener's granulomatosis).



Renal involvement initially appears as focal or segmental glomerulonephritis on biopsy. If left untreated, it can lead to rapidly progressive glomerulonephritis (RPGN). Patients with Wegener's granulomatosis have an increased incidence of venous thrombosis, like deep vein thrombosis and pulmonary embolism.

The diagnosis is confirmed by the presence of necrotizing granulomatous vasculitis on biopsy. This finding is commonly seen in the tissues obtained from lungs, skin, upper airways, and rarely kidneys.

Solution to Question 10:

The given clinical presentation along with pANCA positivity and biopsy findings of leukocytoclastic (neutrophil infiltration) vasculitis point towards a diagnosis of microscopic polyangiitis. It is associated with fibrinoid necrosis.

Microscopic polyangiitis, also known as hypersensitivity vasculitis, causes segmental fibrinoid necrosis of small arterioles and venules. It is associated with neutrophil infiltration, and hence also called leukocytoclastic vasculitis.

Histopathology of the affected tissue will show necrotizing vasculitis of arterioles, venules, and capillaries without any granulomas. It is caused by ANCA (antineutrophil cytoplasmic antibodies) and is not mediated by the immune complex. It primarily affects the kidneys and lungs.

Sold by @itachibot

Solution to Question 11:

Both microscopic polyangiitis and granulomatosis with polyangiitis are associated with necrotizing vasculitis. Hence, this feature is not useful to distinguish between them.

Solution to Question 12:

A triad of skin lesions, asymmetric mononeuritis multiplex, and eosinophilia is seen in Churg–Strauss syndrome (eosinophilic granulomatosis with polyangiitis).

It is a necrotizing vasculitis of the small-to-medium-sized vessels and in order to be diagnosed with Churg-Strauss, a patient should have evidence of asthma, peripheral blood and tissue eosinophilia (>1000 cells/µL), extravascular granuloma formation and clinical features consistent with vasculitis.

Common presenting symptoms of this condition are lower respiratory tract symptoms such as severe attacks of asthma, upper respiratory tract symptoms such as allergic rhinitis, sinusitis, and nasal polyps, mono neuritis multiplex (a type of peripheral neuropathy affecting at least two nerves) which presents with a complaint of pain, numbness, or weakness in the affected areas and skin lesions like palpable purpura and subcutaneous nodules.

Renal involvement is less common and less severe than other small vessel vasculitis.

P-ANCA antibodies are present in Churg Strauss syndrome.

• Severe attacks of asthma

• Mononeuritis multiplex: a type of peripheral neuropathy affecting at least two nerves. Patients will complain of pain, numbness, or weakness in the affected areas.

• Skin lesions like palpable purpura and subcutaneous nodules.

Glucocorticoids show effective responses in these patients. Other drugs that have been tried are mepolizumab and rituximab.

For fulminant multisystem disease, daily cyclophosphamide with prednisolone followed by azathioprine or methotrexate is used.

Option A: Cryoglobulinemic vasculitis presents with palpable purpura, arthralgia, neuropathy, and glomerulonephritis. It is most commonly associated with Hepatitis C.

Option B: Polyarteritis nodosa can also present with mononeuritis multiplex and skin lesions, but they do not have eosinophilia, it has predominant neutrophilia.

Option D: Giant cell arteritis manifests with the cardinal symptom of headache localized to the temporal region. Common symptoms include headache, polymyalgia rheumatica , jaw claudication, fever, and decrease in body weight. Laboratory results show increased ESR, CRP levels, normochromic or mildly hypochromic anemia.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Solution to Question 13:

The given clinical scenario of asthma, mononeuritis multiplex, and pANCA positivity is suggestive of Churg-Strauss syndrome (eosinophilic granulomatosis with polyangiitis). It affects small to medium-sized arteries.

Laboratory investigations will show peripheral blood eosinophilia. Some patients may be positive for pANCA (antineutrophil cytoplasmic antibodies). Histopathology will show the presence of necrotizing vasculitis, tissue eosinophilia, and extravascular granulomas.

Histopathology of ANCA associated vasculitides can be remembered as:

• Microscopic polyangiitis: necrotizing vasculitis

• Wegener's granulomatosis: necrotizing vasculitis with granulomas

• Churg Strauss syndrome: necrotizing vasculitis with extravascular granulomas and eosinophilia

Solution to Question 14:

The given clinical presentation of palpable purpura, colicky abdominal pain, and arthritis, point towards a diagnosis of Henoch Schonlein purpura (HSP) also known as IgA vasculitis.

It is a small vessel vasculitis that occurs due to IgA immune complex deposition within the wall of the blood vessels. It is commonly seen in children and is characterized by:

• Palpable purpura without thrombocytopenia and coagulopathy.

• Arthralgia- commonly affecting hip, knee, or ankle joints.

• Abdominal pain

• Glomerulonephritis: proteinuria, microscopic hematuria, RBC casts in urine.

The skin biopsy will show small vessel vasculitis, with IgA and C3 deposition in the affected organs by immunofluorescence.

Option B: Hemolytic uremic syndrome is characterized by a triad of microangiopathic hemolytic anemia, thrombocytopenia, and acute kidney injury.

Option C: Juvenile idiopathic arthritis (JIA) is characterized by fever of at least 2 weeks duration associated with arthritis, evanescent erythematous rash, hepatosplenomegaly, generalized lymphadenopathy, and serositis.

Option D: Patients with IgA nephropathy or Berger's disease present with gross hematuria, often associated with a preceding or ongoing viral infection of the upper respiratory tract (synpharyngitic presentation).

Solution to Question 15:

Cryoglobulinemia is associated with hepatitis C infection.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Following Hepatitis C infection, there is an aberrant immune response leading to an immune complex formation containing hepatitis C antigens, polyclonal IgG specific to Hepatitis C, and monoclonal IgM rheumatoid factor, which triggers an inflammatory cascade results in Cryoglobulinemic vasculitis.

Cryoglobulins are the immunoglobulins that precipitate at temperatures less than 37°C, the presence of these cryoglobulins in serum leads to Cryoglobulinemia, which is characterized by cutaneous vasculitis, palpable purpura, arthritis, peripheral neuropathy, and glomerulonephritis.

Important conditions associated with cryoglobulinemia are:

- Hepatitis C
- Multiple myeloma
- Connective tissue disorders
- Lymphoproliferative disorders

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Crystal Arthropathies

Question 1:

A patient receiving anti-tubercular therapy has been recently diagnosed with gout. Which of the following statements is incorrect about this condition?

a) The crystals are positively birefringent

b) Ethambutol can precipitate gout

c) More common in males

d) Most commonly involves the first MTP joint

Question 2:

An elderly man presented to the OPD with painful swelling of his left big toe. Synovial fluid examination showed negatively birefringent needle-shaped crystals. What are these crystals made of?

a) Calcium apatite

b) Monosodium urate

c) Calcium pyrophosphate

d) Ammonium urate

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following synovial fluid crystals cannot be identified using polarized light microscopy?

a) Monosodium urate

b) Calcium pyrophosphate

c) Calcium apatite

d) Calcium oxalate

Question 4:

Sold by @itachibot

A patient has a history of multiple episodes of painful swelling of ankles and big toes which is relieved with NSAIDs. You would avoid all of the following drugs in her except:

a) Furosemide

b) High dose salicylates

c) Cyclosporine

d) Pyrazinamide

Question 5:

A patient with chronic gout now presented to the emergency room after sustaining trauma to his left foot. Identify the abnormality in his X-ray given below.

a) Podagra

b) Martel's sign

c) Chopart's fracture

d) March fracture

Question 6:

A 52-year-old man presented to the OPD with complaints of severe joint pain in his big toe with history of multiple similar episodes in the past. An ultrasound of the first metatarsophalangeal joint was performed. What is the characteristic finding depicted in the image given below?

Sold by @itachibot



a) Double wall sign

b) Double density sign

c) Double contour sign

d) Double barrel sign

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

A middle-aged man presented to the ER with severe pain and swelling as shown in the image below. He told that he had consumed 2 bottles of beer a few hours before. Which of the following drugs is not used in the management of this patient?

a) Indomethacin

Sold by @itachibot

b) Prednisone

c) Allopurinol

d) Colchicine

Question 8:

Which of the following drugs is the best agent for the long-term treatment of chronic gout in a 50-year-old patient?

a) Probenecid

b) Colchicine

c) Allopurinol

d) NSAIDS

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 58-year-old man with history of gout is now diagnosed with hypertension. Which of the following is the drug of choice for this patient?

a) Hydrochlorothiazide

b) Metolazone

c) Losartan

d) Furosemide

Question 10:

Which joint is most commonly affected in calcium pyrophosphate disease?

a) Wrist

b) Knee

c) Metacarpophalangeal joint

d) Metatarsophalangeal joint

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	a
2	b
3	c
4	b
5	b
6	c
7	c
8	c
9	c
10	b

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Detailed Explanations

Solution to Question 1:

In gout, the crystals are negatively birefringent on polarized light microscopy of the joint aspirate.

Gout is a metabolic disease that is characterized by the deposition of monosodium urate crystals in joints and connective tissue. It is more common in males.

It usually presents as monoarthritis that may be acute and episodic or chronic. Acute gout presents with a joint that is red, hot, swollen, and extremely tender on examination. The metatarsophalangeal joint of the great toe is the most frequently involved joint.

Synovial fluid analysis of joint aspirate is diagnostic and shows the following features:

• Increased leukocyte count with a neutrophilic predominance

• Negatively birefringent crystals on polarized light microscopy of the synovial fluid

Antitubercular drugs like pyrazinamide and ethambutol cause hyperuricemia by inhibiting uric acid excretion in the kidney.

Monosodium urate crystals as seen on polarized light microscopy:

Sold by @itachibot



Solution to Question 2:

The given clinical presentation and polarised microscopy showing negatively birefringent needle-shaped crystals point towards a diagnosis of gout. In gout, the negatively birefringent needle-shaped crystals are due to the deposition of monosodium urate crystals.

Monosodium urate crystals in gout:



Option A: Calcium apatite crystals can accumulate in areas of tissue damage, hypercalcemia, or hyperparathyroid states.

Option C: Calcium pyrophosphate crystals are positively birefringent rhomboid-shaped crystals that are seen in pseudogout.

Calcium pyrophosphate crystals seen in pseudogout:

Sold by @itachibot



Option D: Ammonium urate crystals can cause urinary bladder stones which are radiolucent but may develop a calcific coating making it visible on X-rays.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Calcium apatite crystals cannot be identified using polarized light microscopy.

Polarized light microscopy can identify crystals such as:

- Monosodium urate
- Calcium pyrophosphate
- Calcium oxalate

Solution to Question 4:

The given scenario of multiple painful swelling of ankles and toes that responded to NSAIDs is suggestive of gout. High-dose salicylates (>3g/day) do not precipitate gout.

Although high-dose salicylates inhibit both the secretion and reabsorption of urate, the inhibition of reabsorption is more than the inhibition of secretion, and therefore the net effect is uricosuric.

On the other hand, low-dose salicylates (<2 to 3g/day) are associated with uric acid retention. They inhibit renal tubular secretion of uric acid resulting in hyperuricemia and may precipitate gout.

Aspirin is not used to treat acute attacks of gout due to its uric acid retention property. However, low-dose aspirin used for cardiovascular prophylaxis need not be discontinued during the treatment of a gout flare.

Sold by @itachibot

Solution to Question 5:

The given X-ray of a patient with chronic gout shows Martel's sign.

Martel's sign also known as the G sign, or rat-bite erosion, refers to the punched-out lytic lesion and overhanging margin of the new bone along the edge of erosion seen in chronic tophaceous gout.

Option A: Podagra is the inflammation of the base of the great toe (metatarsophalangeal joint) during flares of gout. It is not a radiological sign.

Option C: Chopart's fracture is a fracture between talonavicular and calcaneocuboid joints.

Option D: March fracture is fracture neck of 2nd or 3rd metatarsal.

X-ray showing Martel's sign:

Solution to Question 6:

The characteristic finding depicted in the given ultrasound image of a patient with chronic gout is the double contour sign.

Double contour sign is characterized by an echogenic line on the outer surface of the joint cartilage, due to the deposition of monosodium urate crystals on the surface of articular cartilage. It is seen in chronic gout.

Option A: Double wall sign also known as Rigler's sign is an abdominal radiograph finding seen in pneumoperitoneum. Here both the inner and outer walls of the bowel are visible.

Sold by @itachibot



Option B: Double density sign on chest X-ray PA view is seen due to left atrial enlargement. Here, the right border of the left atrium becomes laterally displaced and convex producing this appearance.



Option D: Double barrel sign is a USG finding seen in common bile duct (CBD) obstruction causing dilatation of CBD and its lumen size becomes comparable to portal vein giving the appearance of a double-barrel.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

The given scenario is suggestive of an acute attack of gout. Allopurinol is used in the management of chronic gout.

Urate-lowering drugs such as allopurinol, probenecid, and lesinurad are not initiated during acute attacks.

The treatment options for an acute attack of gout include:

• NSAIDs: Indomethacin, naproxen, ibuprofen, and diclofenac are highly effective

• Colchicine: It acts by inhibiting neutrophil migration and reduces inflammation. It is used orally and is most effective if given early in an attack.

• Glucocorticoids: Systemic glucocorticoids may be used in polyarticular involvement. If the disease is limited to one joint, intraarticular triamcinolone may be given.

Solution to Question 8:

Allopurinol is the best agent for the long-term treatment of chronic or recurrent gout.

Hypouricemic drugs that can be used for definitive treatment of chronic gout include:

• Xanthine oxidase inhibitors- allopurinol, febuxostat

• Uricosuric agents- probenecid, lesinurad

• Recombinant uricases- rasburicase and pegloticase

Solution to Question 9:

Sold by @itachibot

Losartan is the drug of choice for the management of hypertension in a patient with gout because of its uricosuric action.

Thiazides (hydrochlorothiazide, metolazone) and loop diuretics (furosemide) cause reduced excretion of uric acid from the renal tubules and result in hyperuricemia. Therefore they are not preferred for management of hypertension in patients with gout.

Solution to Question 10:

The knee joint is most commonly affected in calcium pyrophosphate (CPPD) disease or pseudogout. Other sites include the wrist, shoulder, ankle, elbow, and hands.

Acute CPPD arthropathy is commonly seen in the elderly and may be associated with conditions such as primary hyperparathyroidism, hypothyroidism, and hemochromatosis.

X-ray of the affected joint may reveal chondrocalcinosis, which represents calcium deposition in articular cartilage. Polarised light microscopy of the joint aspirate demonstrates positively birefringent rhomboid crystals, as shown below:



If yo @itachibot someone else,

Sold by @itachibot

Sjogren's Syndrome and Scleroderma

Question 1:

What is the most common cause of secondary Sjogren's syndrome?

a) Systemic lupus erythematosus

b) Rheumatoid arthritis

c) Scleroderma

d) Primary biliary cirrhosis

Question 2:

Which is the most common extraglandular manifestation seen in patients diagnosed with Sjogren's syndrome?

a) Primary biliary cirrhosis

b) Vasculitis

c) Arthritis

d) Renal tubular acidosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A middle-aged woman presented with complaints of persistent dry eyes and dry mouth since past few months. On examination, dry mucous membranes, bilateral parotid region swelling, and conjunctivitis were noted. This condition is associated with which of the following?

a) HLA-DR2

b) HLA-DR4

c) HLA-DR3

d) HLA-B27

Question 4:

A software engineer presented with complaints of dryness and itching of eyes on staring at his laptop screen. On examination, parotid swelling was noted and the test shown below was

Sold by @itachibot

positive. The serology of this patient is likely to be positive for which of the following antibodies?



a) Anti topoisomerase

b) Anti centromere

c) Anti U1-RNP

d) Anti Ro/SS-A

Sold by @ if you purchased this from you may have been scamm... else,

Question 5:

A 55-year-old woman presents with dry mouth, and difficulty swallowing solid food. Physical examination finds enlargement of her parotid glands along with marked dryness of her buccal mucosa. Laboratory examination finds the presence of both SS-A and SS-B antibodies. Which of the following statements is not true about this condition?

a) About 75% of patients have rheumatoid factor

b) Patients are at high risk for development of B-cell lymphomas

c) Glomerular lesions are common

d) Lacrimal gland is a major target for this disease

Question 6:

A 35-year-old woman presented with complaints of thickening of her skin and the finding shown below. On workup, ANA was positive and immunofluorescence showed nucleolar pattern. What is the most common extracutaneous manifestation associated with her

Sold by @itachibot

condition?



a) Pulmonary fibrosis

b) Myositis

c) Scleroderma renal crisis

d) Raynaud's phenomenon

Question 7:

A 14-year-old girl has the pallor of extremities followed by pain and cyanosis on exposure to cold. Which of the following diseases is she likely to develop in the later stages of life?

a) Myasthenia gravis

b) Scleroderma

c) Mikulicz syndrome

d) Dermatomyositis

Question 8:

Which of the following statements is false about Raynaud's phenomenon?

a) Secondary Raynaud's is associated with more severe symptoms

b) Secondary Raynaud's develops predominantly after the age of 30 years

c) Primary Raynaud's is usually ANA negative

d) Primary Raynaud's is associated with digital tissue necrosis and ulcers

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 9:

What is the most common histological pattern of interstitial lung disease seen in patients with systemic sclerosis?

a) Non specific interstitial pneumonia

b) Usual interstitial pneumonia

c) Desquamative interstitial pneumonia

d) Cryptogenic organizing pneumonia

Question 10:

Which of the following drugs are implicated in systemic sclerosis-like illness?

a) 1, 3, and 4

b) 2, 3, and 5

c) 1, 4, and 5

d) 2, 4, and 5

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 11:

A 38-year-old woman presents with ulceration over her fingers. She has history of Raynaud's phenomenon and GERD since 3 years. Which of the following can be used as a screening test for her condition?

a) Antinuclear antibody (ANA)

b) Anti centromere antibody

c) Anti topoisomerase I (Scl-70)

d) Anti-cardiolipin antibody

Question 12:

A 47-year-old woman presents with complaints of swollen fingers, heartburn, and occasional joint pains. She gives a history suggestive of Raynaud's phenomenon. Physical examination reveals taut, thickened skin over her hands and fingers with limited mobility. The presence of which of the following antibodies will confirm her diagnosis?

Sold by @itachibot

a) Antinuclear antibody

b) Anticentromere antibody

c) Anti-topoisomerase I antibody

d) Anti-RNA polymerase III antibody

Question 13:

A 53-year-old woman presented with thickening of skin over the trunk and proximal parts of the limbs. On workup, her serology was found to be positive for anti-topoisomerase I. Which of the following features is not likely to be associated with her condition?

a) Severe arthralgia

b) Critical limb ischemia

c) Interstitial lung disease

d) Carpal tunnel syndrome

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 14:

Which among the following is more common in limited cutaneous systemic sclerosis when compared to diffuse cutaneous systemic sclerosis?

a) Esophageal dysmotility

b) Myopathy

c) Interstitial lung disease

d) Scleroderma renal crisis

Question 15:

Which of the following is not a component of CREST syndrome?

a) Calcinosis cutis

b) Renal failure

c) Esophageal dysmotility

d) Sclerodactyly

Question 16:

Sold by @itachibot

Which of the following antibodies is likely to be positive in patients who develop scleroderma renal crisis?

a) Anti centromere

b) Anti Ku

c) Anti U1-RNP

d) Anti RNA polymerase III

Question 17:

An elderly man presents to the ER with severe headache, blurring of vision, and oliguria. Examination reveals BP of 200/100mmHg and papilledema. Skin thickening is noted over his hands, with contractures in the wrist and fingers. Investigations reveal serum creatinine of 3mg/dL and peripheral smear shows schistocytes. What is the drug of choice?

a) Direct renin inhibitors

b) ACE inhibitors

c) Steroids

d) Calcium channel blockers

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	b
2	c
3	c
4	d
5	c
6	d
7	b
8	d
9	a
10	d
11	a
12	b
13	b

Sold by @itachibot

14	a
15	b
16	d
17	b

Detailed Explanations

Solution to Question 1:

The most common cause of secondary Sjögren's syndrome is rheumatoid arthritis.

Sjögren's syndrome is a chronic autoimmune disorder that affects the lacrimal and salivary glands. It also affects other organ systems, such as lungs, kidneys, and nervous system. It is seen in association with:

• Rheumatoid arthritis

• Systemic lupus erythematosus

• Scleroderma

• Mixed connective tissue disease

• Primary biliary cirrhosis

• Vasculitis

• Chronic active hepatitis C

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 2:

The most common extraglandular manifestation seen in patients with Sjögren's syndrome is arthritis or arthralgia.

The extraglandular manifestations in Sjögren's syndrome are:

• Arthralgia or arthritis - most common manifestation. It is usually non-erosive and leads to Jaccoud's arthropathy. Jaccoud's arthropathy is a deforming non-erosive arthropathy characterized by the ulnar deviation of the second to 5th fingers with metacarpophalangeal joint subluxation.

• Raynaud's phenomenon

• Interstitial lung disease - most common type is non-specific interstitial pneumonia (NSIP)

• Renal tubular acidosis - Type 1 (distal) RTA

• Primary biliary cirrhosis

• Lymphoma - MALToma

The give X-ray shows Jaccoud's arthropathy:

Sold by @itachibot

The given image shows a patient with Jaccoud's arthropathy of the hands:

Solution to Question 3:

Dry eyes and dry mouth along with parotid swelling and conjunctivitis are suggestive of Sjögren's syndrome. It is associated with HLA-DR3 allele.

HLA allele	Disease
HLA DR2, HL A-DR3	SLE
HLA-DR3	Sjogren's syndrome
HLA-DR4	Rheumatoid arthritis

Sold by @itachibot

HLA allele	Disease
HLA-B27	Seronegative arthritis

Solution to Question 4:

The given scenario of dry eyes, parotid swelling, and a positive Schirmer test is suggestive of Sjögren's syndrome. It is associated with anti-Ro/SS-A and anti-La/SS-B autoantibodies.

The presence of anti-Ro/SS-A and anti-La/SS-B indicate extra glandular involvement and poor prognosis.

Note: Anti-Ro and anti-La in SLE indicate a good prognosis, due to decreased risk of nephritis and vasculitis.

Solution to Question 5:

The given clinical scenario is suggestive of Sjogren's syndrome. In this condition, glomerular lesions are extremely rare.

Defects of tubular function, including renal tubular acidosis, uricosuria, and phosphaturia, are often seen and are associated histologically with tubulointerstitial nephritis.

Lacrimal and salivary glands are the major targets of this disease, while the respiratory and gastrointestinal tracts and the vagina may also be involved. The characteristic decrease in tears and saliva (sicca syndrome) is the result of lymphocytic infiltration and fibrosis of the lacrimal and salivary glands. The infiltrate contains predominantly activated CD4+ helper T cells and some B cells, including plasma cells.

Antibodies detected are:

• Rheumatoid factor - seen in 75% of patients, whether or not coexisting rheumatoid arthritis is present

• Anti-nuclear antibodies (ANAs)

• SS-A (Ro) - associated with early disease onset, longer disease duration, and extra-glandular manifestations, such as cutaneous vasculitis and nephritis

• SS-B (La)

These patients are at high risk for the development of B-cell lymphomas. About 60% of patients have another accompanying autoimmune disorder, such as rheumatoid arthritis, and these patients also have the symptoms and signs of that disorder.

Biopsy of the lip (to examine minor salivary glands) is essential for the diagnosis of Sjögren's syndrome.

The image shows an enlargement of the salivary gland with intense lymphocytic and plasma cell infiltration with ductal epithelial hyperplasia in the salivary gland.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot



Note: Mikulicz syndrome is characterized by lacrimal and salivary gland enlargement from any cause, including sarcoidosis, lymphomas, and other tumors.

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 6:

The given image is suggestive of microstomia. The findings of skin thickening, microstomia, positive ANA, and immunofluorescence showing nucleolar pattern point towards a diagnosis of systemic sclerosis. The most common extracutaneous manifestation of systemic sclerosis (scleroderma) is Raynaud's phenomenon.

Raynaud's phenomenon is characterized by episodes of reversible vasoconstriction in the fingers and toes on exposure to cold. Rarely, it affects the lips, tip of the nose, and earlobes. These episodes classically occur in the following sequence:

- Affected part turning pale due to vasoconstriction.
- Blue due to ischemia, often associated with numbness or pain.
- As blood flow returns, the area turns red with a burning sensation.

Sold by @itachibot

Sold by @itachi... else,
...purchased this from some...
...may have been scammed.

Solution to Question 7:

The above clinical scenario suggests that the patient has Raynaud phenomenon, which is associated with the risk of developing scleroderma in later stages of life.

Raynaud phenomenon is manifested as episodic vasoconstriction of the arteries and arterioles of the extremities, causing pallor of extremities followed by pain and cyanosis. It is seen in virtually all patients with systemic sclerosis or scleroderma and precedes other symptoms in 70% of cases.

The image shows features of Raynaud's phenomenon.

Systemic sclerosis is characterized by progressive fibrosis involving the skin, gastrointestinal tract, and other tissues. Fibrosis may be the result of the activation of fibroblasts by cytokines produced by T cells.

Sold by @itachibot

Focal, and sometimes diffuse, subcutaneous calcifications may develop, especially in patients with CREST syndrome. The most ominous manifestation is malignant hypertension.

Solution to Question 8:

Primary Raynaud's phenomenon is not associated with tissue necrosis and ulcers.

Features	Primary Raynaud's phenomenon	Secondary Raynaud's phenomenon
Age & gender	Young women (<30y)	Older women (>30y)
Associations	No underlying cause	Scleroderma, Sjögren's syndrome and SLE
Family history	Yes	No
ANA test	Negative	Positive
Severity - duration, frequency & pain	Less severe	More severe
Digital tissue necrosis, ulceration & gangrene	No	Yes
Nailfold capillaroscopy(Image below)	Normal capillaries in evenly spaced parallel loops	Distorted capillaries with widened & irregular loops



Solution to Question 9:

The most common histological pattern of interstitial lung disease in systemic sclerosis (scleroderma) is non-specific interstitial pneumonia (NSIP). It commonly affects the lower lobes and is characterised by ground-glass opacities.

Sold by @itachibot

Note: Usual interstitial pneumonia (UIP) is the most common type of ILD in rheumatoid arthritis.

Solution to Question 10:

Drugs implicated in systemic sclerosis-like illness are:

• Bleomycin

• Pentazocine

• Cocaine

• L-tryptophan-containing dietary supplements

Chemicals implicated in systemic sclerosis-like illness are:

• Heavy metals

• Polyvinyl chloride (PVC)

• Toluene

• Trichloroethylene

Cyclophosphamide and bosentan are used in the treatment of systemic sclerosis (SSc). Cyclophosphamide reduces the progression of ILD associated with SSc. Bosentan reduces the risk of new ulcers in Raynaud's phenomenon associated with SSc.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

The presence of digital ulcers, Raynaud's phenomenon, and GERD are suggestive of systemic sclerosis (scleroderma). Antinuclear antibody (ANA) is the preferred screening test as it is the most sensitive test for both limited and diffuse variants of systemic sclerosis.

Option B: Anti centromere antibody is the most specific test for CREST syndrome.

Option C: Anti topoisomerase I (Scl-70) is the most specific test for diffuse cutaneous systemic sclerosis.

Option D: Anti-cardiolipin antibody is seen in SLE and APLA syndrome.

Solution to Question 12:

The given scenario of skin thickening in the hands and fingers (sclerodactyly), Raynaud's phenomenon, heartburn (GERD), and occasional arthralgia is suggestive of limited cutaneous systemic sclerosis (lcSSc). The presence of anticentromere antibody (ACA) will confirm the diagnosis of limited cutaneous systemic sclerosis.

Antinuclear antibody (ANA) is positive in most cases of systemic sclerosis. The absence of this antibody suggests considering an alternate diagnosis.

Sold by @itachibot

Solution to Question 13:

The presence of skin thickening over the trunk and proximal parts of the limbs along with positive anti-topoisomerase I antibodies is suggestive of diffuse cutaneous systemic sclerosis (dcSSc). Critical limb ischemia is commonly associated with limited cutaneous systemic sclerosis (lcSSc).

Solution to Question 14:

Esophageal dysmotility is more commonly seen in limited cutaneous systemic sclerosis (lcSSc) compared to diffuse cutaneous systemic sclerosis.

Systemic sclerosis is divided into two based on the patterns of involvement - diffuse and limited.

Limited cutaneous systemic sclerosis is characteristically associated with CREST syndrome, which includes calcinosis cutis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyly and telangiectasia.

Skin involvement, interstitial lung disease (Option C), myopathy (Option B), and scleroderma renal crisis (Option D) are more common in diffuse cutaneous systemic sclerosis (dcSSc) than limited cutaneous systemic sclerosis.

The incidence of ischemic digital ulcers, CREST features, and primary biliary cirrhosis is more in limited cutaneous systemic sclerosis. The frequency of pulmonary arterial hypertension (PAH) is equal in both types.

Solution to Question 15:

Renal failure is not a component of CREST syndrome.

CREST syndrome is seen in limited cutaneous systemic sclerosis (lcSSc) and consists of the following:

- Calcinosis cutis
- Raynaud's phenomenon
- Esophageal dysmotility
- Sclerodactyly
- Telangiectasia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



Solution to Question 16:

Anti-RNA polymerase III antibody is associated with scleroderma renal crisis.

Scleroderma renal crisis is scleroderma with accelerated hypertension, acute kidney injury and renal failure. The antibodies associated with it are:

- Anti-RNA polymerase III
- Topoisomerase I
- U3-RNP

Target Antigen	Characteristic Clinical Association
Topoisomerase I	Early ILD, cardiac involvement, scleroderma renal crisis
RNA Polymerase III	Scleroderma renal crisis, increased cancer risk, GAVE (gastric antral vascular ectasia)
U3-RNP	PAH, ILD, scleroderma renal crisis, and myositis
Th/To	Interstitial lung disease (ILD) and Pulmonary artery hypertension (PAH)
Ku	Overlap syndrome

Solution to Question 17:

If you purchased this from someone else, it may have been scammed. Sold by @itachibot

The given clinical scenario is suggestive of scleroderma renal crisis. The drug of choice is ACE inhibitors.

Scleroderma renal crisis is a medical emergency characterized by acute onset malignant hypertension and kidney failure, usually in the absence of previous evidence of significant kidney disease. The use of steroids is associated with an increased risk of scleroderma renal crisis.

Clinical features of scleroderma renal crisis are:

• Accelerated hypertension - headache, vomiting, blurring of vision, hypertensive retinopathy, and hypertensive encephalopathy.

• Acute kidney injury - oliguria, mild proteinuria, microscopic hematuria, renal failure.

• Pulmonary edema, congestive heart failure.

• Microangiopathic hemolytic anemia (schistocytes), thrombocytopenia.

In patients with persistent hypertension, the addition of angiotensin II receptor blockers, calcium channel blockers, and direct renin inhibitors should be considered.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Sarcoidosis

Question 1:

In patients with sarcoidosis, which of the following is the most commonly affected organ?

a) Lungs

b) Eyes

c) Liver

d) Skin

If you bought this from someone else, it was scammed. Sold by @itachibot

Question 2:

A 35-year-old woman with history of sarcoidosis presents with skin lesions shown in the image below. Identify the condition.

a) Acne rosacea

b) Lupus pernio

c) Heliotrope rash

d) Malar rash

Question 3:

Sold by @itachibot

A 24-year-old man presented with fever, productive cough, and weight loss. On examination, the following skin lesions were seen. His chest X-ray showed hilar lymphadenopathy. Which is the next best investigation for the management of this patient?

a) Sputum AFB smear

b) Serum ACE levels

c) CD4/CD8 ratio in BAL

d) Lymph node biopsy

Sold by
if you purchased this fro
you may have been scam
else,
d.

Question 4:

A man came to the clinic with fatigue, cough, and breathing difficulty. His chest X-ray showed bilateral hilar lymphadenopathy and lung biopsy revealed non-caseating granuloma. What is the most common ocular manifestation associated with this condition?

a) Anterior uveitis

b) Scleritis

c) Episcleritis

d) Conjunctivitis

Question 5:

Which is the most common cranial nerve to be involved in sarcoidosis?

a) Optic nerve

Sold by @itachibot

b) Facial nerve

c) Oculomotor nerve

d) Trigeminal nerve

Question 6:

Which of the following features would you not expect in a patient diagnosed with Lofgren's syndrome?

a) Arthritis

b) Parotid enlargement

c) Bilateral hilar adenopathy

d) Erythema nodosum

Question 7:

A 44-year-old construction worker, who is a known case of sarcoidosis, has now presented with polyuria, constipation, and myalgia for the past 5 days. An ECG was taken which showed a short QT interval. Which of the following is responsible for this condition?

a) Increased bone resorption

b) Decreased renal excretion of calcium

c) Increased production of vitamin D

d) Increased production of PTH related peptide

Question 8:

A patient was evaluated for unexplained cough and shortness of breath. Investigations revealed elevated serum calcium and a chest X-ray showed hilar lymphadenopathy. A lymph node biopsy was taken to confirm the diagnosis. Which of the following is least likely to be seen in the histopathological examination?

a) Non-caseating granuloma

b) Asteroid bodies

c) Schaumann bodies

d) Masson bodies

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 9:

What is the most common abnormality of liver function in sarcoidosis?

a) Elevation of transaminases

b) Elevation of alkaline phosphatase

c) Coagulopathy

d) Elevation of bilirubin

Question 10:

A middle-aged woman presented to you with shortness of breath and dry cough for the past 3 months. On examination, the following skin lesions were seen. You will order all of the following investigations except:



a) Serum angiotensin converting enzyme levels

b) CD4/CD8 counts in the blood

c) Biopsy of the lymph nodes

d) Gallium scan

Question 11:

Sold by @itachibot

The chest radiograph of a young man with sarcoidosis shows pulmonary infiltrates in the upper lobes of his lungs without hilar lymphadenopathy. What is the grade of lung involvement according to Scadding scoring system?

a) Grade I

b) Grade II

c) Grade III

d) Grade IV

Question 12:

Which of the following lymph nodes is not involved in the Garland triad?

a) Right paratracheal nodes

b) Right hilar nodes

c) Left hilar nodes

d) Left paratracheal nodes

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

Which of the following is the first-line drug for a patient diagnosed with sarcoidosis with multiple organ involvement?

a) Prednisolone

b) Hydroxychloroquine

c) Methotrexate

d) Infliximab

Question 14:

Which of the following conditions is not associated with a significant elevation of angiotensin-converting enzyme levels?

a) Sarcoidosis

b) Hyperthyroidism

c) Diabetes

d) Miliary TB

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	b
3	a
4	a
5	b
6	b
7	c
8	d
9	b
10	b
11	c
12	d
13	a
14	c

if you purcha... you may have ... else, ...d.

Detailed Explanations

Solution to Question 1:

Sarcoidosis most commonly affects the lungs.

Sarcoidosis is a multisystem inflammatory disease characterized by non-caseating granulomas. It commonly involves the lungs, skin, liver, and eyes.

Solution to Question 2:

The given condition is lupus pernio which is a cutaneous manifestation of sarcoidosis.

Lupus pernio is a chronic inflammatory skin lesion, which is diagnostic for a chronic form of sarcoidosis. It involves the bridge of the nose, the area beneath the eyes, and the cheeks and often is purple in color.

Other options:

Sold by @itachibot

Option A: Acne rosacea is a skin disorder with onset in middle■aged adults (30–50 years old) and involves the convexities of the face.

Option C: Heliotrope rash presents as violaceous lesions around eyelids. It is seen in dermatomyositis.

Option D: Malar rash or butterfly rash is classically seen in SLE.

Solution to Question 3:

The given clinical features along with examination findings of lupus vulgaris is suggestive of tuberculosis. The next best investigation is AFB smear microscopy of the sputum sample.

Lupus vulgaris is a manifestation of cutaneous tuberculosis. It has a characteristic slow enlarging plaque with an elevated border and central atrophy.

Solution to Question 4:

The given clinical features along with bilateral hilar lymphadenopathy and non-caseating granuloma are suggestive of sarcoidosis. The most common ocular manifestation in sarcoidosis is anterior uveitis.

Anterior uveitis presents with photophobia, blurry vision, and lacrimation. Other ocular manifestations include retinitis, pars planitis, and dry eyes.

Solution to Question 5:

The most common cranial nerve to be involved in sarcoidosis is the facial nerve leading to facial palsy.

Neurosarcoidosis frequently manifests as follows:

• Cranial nerve involvement- optic neuritis (CN II), extraocular muscles involvement (CN III, IV, VI) and facial palsy (CN VII)

• Space occupying lesions

• Diabetes insipidus - due to anterior hypothalamus involvement

• Basilar meningitis

Solution to Question 6:

Parotid enlargement is not a feature of Lofgren's syndrome. Lofgren's syndrome is an acute form of sarcoidosis that has a good prognosis, with most of the patients experiencing disease resolution within 2 years.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

It is characterized by:

• Arthritis

• Erythema nodosum

• Bilateral hilar lymphadenopathy

Heerfordt–Waldenström syndrome also known as uveoparotid fever is a variant of sarcoidosis characterized by:

• Fever

• Parotid enlargement

• Facial palsy

• Anterior uveitis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

The given clinical features along with a short QT interval points towards hypercalcemia. Hypercalcemia in a patient with sarcoidosis is due to the increased production of vitamin D by the non-caseating granulomas.

Hypercalcemia and hypercalciuria are seen in some patients with sarcoidosis. The increased production of vitamin D (1,25-dihydoxyvitamin D) results in increased intestinal absorption of calcium, leading to hypercalcemia.

Vitamin D supplementation and increased sun exposure in patients with sarcoidosis can precipitate hypercalcemia.

Solution to Question 8:

The given clinical scenario is suggestive of sarcoidosis, in which the histopathological examination of lesions does not show Masson bodies. Masson bodies are polypoid plugs of loose organizing connective tissue seen in cryptogenic organizing pneumonia.

The cardinal histopathologic feature of sarcoidosis, irrespective of the organ involved, is the non-caseating granuloma. Two other microscopic features are sometimes seen in the granulomas:

• Schaumann bodies- inclusion bodies with concentric rings of calcification

• Asteroid bodies- stellate inclusions in the giant cells

Solution to Question 9:

The most common abnormality of liver function in sarcoidosis is the elevation of alkaline phosphatase (ALP) levels.

Sold by @itachibot

Sarcoidosis of the liver presents as a granulomatous disease that causes intrahepatic cholestasis. As the disease advances it leads to elevated liver enzymes followed by elevated bilirubin levels.

Solution to Question 10:

The given clinical scenario and examination findings of erythema nodosum are suggestive of sarcoidosis. CD4/CD8 ratio in bronchoalveolar lavage (BAL) fluid, not blood, is used for diagnosis of sarcoidosis.

Erythema nodosum is inflammation of the subcutaneous fat (panniculitis) seen in sarcoidosis.

The investigations used in the diagnosis of sarcoidosis include:

• Serum ACE (angiotensin-converting enzyme) levels - elevated

• BAL- lymphocytosis and increased CD4/CD8 ratio. A ratio of >3.5 in BAL is strongly supportive of sarcoidosis.

• PET-CT scan - to identify areas of granulomatous disease

• Gallium scan shows the following signs:

• Panda sign- due to bilateral involvement of parotid and lacrimal glands in sarcoidosis

• Lambda sign- due to bilateral hilar and right paratracheal lymph nodes which resembles lambda (λ)

• Biopsy showing non-caseating granulomas in a patient with features of sarcoidosis confirms the diagnosis.

The image given below shows the Panda sign:



If yo... purchased this from someone else, Sold by @itachibot ...have been scammed.

Solution to Question 11:

Sold by @itachibot

According to the Scadding scoring system, the presence of pulmonary infiltrates alone represents grade III.

The Scadding scoring system is used to grade lung involvement in sarcoidosis. It is based on the chest radiograph findings:

• Grade I: Hilar lymphadenopathy, often with right paratracheal involvement. No pulmonary infiltrates.

• Grade II: Hilar lymphadenopathy + pulmonary infiltrates

• Grade III: Pulmonary infiltrates only

• Grade IV: Fibrosis

Sarcoidosis predominantly involves the upper lobes of the lungs.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 12:

Garland triad does not involve left paratracheal lymph nodes.

Garland triad also known as the 1-2-3 sign or Pawnbrokers sign is a lymph node enlargement pattern in sarcoidosis that involves:

• Right paratracheal nodes (1)

• Right hilar nodes (2)

• Left hilar nodes (3)



Solution to Question 13:

The first-line drug for sarcoidosis with multiple organ involvement is prednisolone.

Management of sarcoidosis:

Sold by @itachibot

• Asymptomatic patients- monitored for disease progression

• Localized/ limited cutaneous and ocular lesions- treated using topical steroids

• Multisystem involvement:

• Systemic steroids

• Other drugs like hydroxychloroquine, methotrexate, and infliximab are considered when the steroids are ineffective or not tolerated

Solution to Question 14:

Diabetes is associated with a mild elevation of angiotensin-converting enzyme (ACE) levels.

Significant elevation of ACE levels is seen in:

• Sarcoidosis

• Silicosis

• Leprosy

• Gaucher's disease

• Hyperthyroidism

• Disseminated granulomatous infections such as miliary tuberculosis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Dermatomyositis and Related Disorders

Question 1:

What is the most common inflammatory myopathy seen in the elderly?

a) Polymyositis

b) Dermatomyositis

c) Inclusion body myositis

d) Myositis secondary to overlap syndrome

Question 2:

An IT professional complained of weakness of fingers while using the keyboard. It has been progressing over the past 2 months. It is more severe on his right hand than his left hand. Which of the following is the most likely diagnosis?

a) Polymyositis

b) Dermatomyositis

c) Inclusion body myositis

d) Necrotizing myopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Patients with which of the following myopathies are generally resistant to immunosuppressive therapies?

a) Dermatomyositis

b) Polymyositis

c) Inclusion body myositis

d) Necrotizing myopathy

Question 4:

Which of the following group of muscles are unlikely to be involved in a patient diagnosed with an inflammatory myopathy?

Sold by @itachibot

a) Ocular muscles

b) Pharyngeal muscles

c) Neck flexors

d) Quadriceps muscles

Question 5:

Which inflammatory myopathy is most frequently associated with other autoimmune diseases?

a) Inclusion body myositis

b) Polymyositis

c) Dermatomyositis

d) Necrotizing autoimmune myopathy

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A physician is suspecting dermatomyositis in a 27-year-old woman with complaints of proximal myalgia and dysphagia. The presence of which of the following signs would be pathognomonic of this condition?

a) Gottron's papules

b) V sign

c) Gottron's sign

d) Shawl sign

Question 7:

A 50-year-old woman presented with difficulty in activities like climbing stairs, getting up from a chair, and combing hair. She is a known case of ovarian cancer. The following finding was noted on examination. What is the most probable diagnosis?

Sold by @itachibot

a) Polymyositis

b) Dermatomyositis

c) Inclusion body myositis

d) Discoid lupus erythematosus

Question 8:

Which of the following would help you to differentiate between dermatomyositis and polymyositis in a patient presenting with muscle weakness?

a) Elevated creatine kinase

b) Interstitial lung disease

c) Erythematous rash

d) Involvement of proximal muscles

Question 9:

A 50-year-old woman was recently diagnosed with dermatomyositis. She should be screened for which of the following conditions?

a) Diabetes mellitus

b) Malignancy

c) HIV

d) Hypertension

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 10:

A 30-year-old woman complained of difficulty reaching high cabinets and carrying grocery bags. It has been progressing over the past 3 months. Which is the initial investigation of choice in this patient?

a) ESR

b) Creatine kinase

c) Electromyogram

d) hs-CRP

Question 11:

A middle-aged woman presented with muscle weakness and pain over her shoulders. On examination, the following rash and mechanic's hands were noted. Positive titers of which of the following antibodies are associated with her condition?

a) Anti-Mi-1

b) Anti-Jo1

c) Anti-PM/Scl

d) Anti-SRP

Question 12:

Sold by @itachibot

A patient with dermatomyositis presented with cough, hemoptysis, and weight loss. After a detailed evaluation, she was diagnosed with lung cancer. Which of the following autoantibodies is strongly associated with this condition?

a) Anti-Mi-2

b) Anti- Jo1

c) Anti PM/SCL

d) Anti 155/140

Question 13:

A bodybuilder complains that he is facing difficulty in doing squats and is too weak to climb stairs. He denies any headaches or changes in vision. A quadriceps muscle biopsy reveals perimysial inflammation and perifascicular atrophy. Which of the following is the most likely diagnosis?

a) Inclusion body myositis

b) Dermatomyositis

c) Polymyositis

d) Necrotizing autoimmune myopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 14:

A housemaid presents to you with generalized skin thickening and numbness of fingers on exposure to cold. She also complains of muscle pain, and difficulty while walking and climbing stairs. Investigations reveal the following. The presence of which of the following antibodies is diagnostic of this condition?

a) Anti – Jo 1

b) Anti PM/Scl

c) Anti centromere

d) U3- RNP

Question 15:

A 35-year-old woman presents with severe generalized body ache, fatigue, sleep disturbances, and low mood for the past 4 months. On examination, increased pain sensitivity and tender bony prominences are noted. Laboratory and radiological investigations are normal. What is

Sold by @itachibot

the most probable diagnosis?

a) Polymyositis

b) Dermatomyositis

c) Depression

d) Fibromyalgia

Question 16:

Which of the following statements is false about fibromyalgia?

a) More common in females

b) Aerobic exercise improves symptoms

c) SPECT studies show hyperperfusion of the thalamus

d) Opioid analgesics are ineffective in pain management

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	c
3	c
4	a
5	b
6	a
7	b
8	c
9	b
10	b
11	b
12	d
13	b
14	b
15	d
16	c

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The most common inflammatory myopathy seen in elderly patients (>50 years) is inclusion body myositis.

Inclusion body myositis is a type of inflammatory myopathy. It is characterized by progressive, asymmetric muscle weakness most commonly involving the wrist, finger flexors, and quadriceps. Muscles of the face and neck may also be involved, presenting as dysphagia.

Inclusion body myositis is named so because of the presence of inclusion bodies containing amyloid-like protein seen on microscopy.

Solution to Question 2:

The given history of progressive and predominantly asymmetric muscle weakness is most likely suggestive of inclusion body myositis.

Inclusion body myositis is a type of inflammatory myopathy. It is seen in people older than 50 years of age. It is characterized by progressive, asymmetric muscle weakness most commonly involving the wrist, finger flexors, and quadriceps. Muscles of the face and neck might also be involved, leading to dysphagia.

Polymyositis, dermatomyositis, and necrotizing myopathy have predominantly symmetric muscle involvement.

If you purchased this from someone else
Sold by @itachibot
you may have been scammed.

Solution to Question 3:

Patients with inclusion body myositis are generally resistant to immunosuppressive therapies.

Solution to Question 4:

Ocular muscles are unlikely to be involved as they are spared in inflammatory myopathy. They are not involved even in advanced, untreated cases.

The pattern of muscle involvement in various inflammatory myopathies is:

• Proximal muscles are most commonly involved in dermatomyositis and polymyositis, causing difficulty in getting up from a chair, brushing hair, etc.

• Proximal and distal muscles are involved in inclusion body myositis, such as wrist, finger flexors, and quadriceps.

Sold by @itachibot

• Pharyngeal and neck-flexor muscles are often involved in all forms of inflammatory myopathy, causing dysphagia or difficulty in holding up the head. Respiratory muscles may also be affected in advanced cases.

Solution to Question 5:

Inflammatory myopathy most frequently associated with other autoimmune diseases is polymyositis.

Solution to Question 6:

The presence of Gottron's papules is pathognomonic of dermatomyositis. They are erythematous to violaceous papules that occur symmetrically over the extensor aspects of metacarpophalangeal and interphalangeal joints.

Gottron's sign, on the other hand, is erythematous rash over the extensor surfaces of joints such as the knuckles, elbows, knees, and ankles.

V sign is confluent macular erythema over the lower anterior neck and upper anterior chest.

Shawl sign is the confluent macular erythema over the upper back and lateral aspect of arms.

All these signs are present in dermatomyositis, but Gottron's papules and heliotrope rash are pathognomonic.

Solution to Question 7:

The given image shows upper eyelid lilac erythema or heliotrope rash. The given clinical presentation of proximal myopathy with the characteristic heliotrope rash and history of an internal malignancy point towards a diagnosis of dermatomyositis.

Classic cutaneous signs in dermatomyositis:

• Heliotrope rash/ eyelid erythema- violaceous erythema of the upper eyelids. Pathognomonic of dermatomyositis.

• V sign- confluent macular erythema over the lower anterior neck and upper anterior chest

• Shawl sign- confluent macular erythema over the upper back and lateral aspect of arms

• Gottron’s papules- erythematous papules and plaques over the extensor surface of the metacarpophalangeal and interphalangeal joints. Pathognomonic of dermatomyositis.

• Gottron's sign- erythematous rash over the extensor surfaces of joints such as the knuckles, elbows, knees, and ankles

• Mechanic’s hands- rough, hyperkeratotic lines on the lateral and palmer areas of the fingers

• Scalp erythema and hair loss

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

• Facial rash- confluent erythema in the malar distribution involving the cheeks, nasal bridge, and nasolabial folds

• Nail fold changes

• Flagellate erythema- linear erythema on the upper back with pruritus

• Poikiloderma- mottled pattern of hyperpigmented and hypopigmented macules with telangiectasias

• Cutaneous calcinosis- accumulation of calcium deposits in the skin

Facial rash in dermatomyositis-

Eyelid erythema/ heliotrope rash in dermatomyositis-

V- sign in dermatomyositis-

Sold by @itachibot

Shawl sign in dermatomyositis-

Flagellate erythema in dermatomyositis-

Sold by @itachibot

Gottron papules in dermatomyositis-

Poikilodermatous changes in dermatomyositis-

If yo ltachi some else, ed.

Sold by @itachibot

(a)

(b)

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 8:

An erythematous rash helps in differentiating between the two conditions as it is seen only in dermatomyositis.

Option A: Elevated levels of creatine kinase are seen in all the inflammatory myopathies.

Options B and D: The involvement of proximal muscles more than the distal muscles and interstitial lung disease is seen in both polymyositis and dermatomyositis.

Solution to Question 9:

This patient should be screened for malignancy as dermatomyositis is often associated with underlying malignancies.

The most common cancer associated with dermatomyositis is ovarian cancer. Other malignancies that are commonly associated with dermatomyositis are:

- Breast cancer
- Melanoma
- Colon cancer
- Non-Hodgkin's lymphoma

Solution to Question 10:

The given scenario is suggestive of an inflammatory myositis for which the initial investigation of choice is creatine kinase (CK).

Sold by @itachibot

Creatine kinase is a marker of muscle breakdown. It is increased in dermatomyositis and polymyositis. However, it may be normal in inclusion body myositis.

The most specific investigation for inflammatory myositis is a muscle biopsy.

Solution to Question 11:

The given clinical scenario and findings of Gottron papules (image) are suggestive of dermatomyositis. Positive titers of Anti-Jo-1 (antisynthetase) antibodies are associated with interstitial lung disease, arthritis, and mechanic's hands in patients with dermatomyositis.

Other autoantibodies seen in dermatomyositis are:

• Anti-Mi-2 - highly specific to dermatomyositis

• Anti-SRP (anti-signal recognition protein) - associated with severe myositis

• Anti–PM-Scl - associated with overlapping myositis and scleroderma

• Anti-p155/140 or anti-transcription intermediary factor (TIF)-1γ - seen in cancer-associated myositis

• Anti-MDA5 (anti-CADM 140) - associated with rapidly progressive lung disease and cutaneous ulcer

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

Anti-155/140 antibody is strongly associated with dermatomyositis with malignancy.

Adenocarcinomas of the cervix, lung, ovaries, and pancreas account for the majority of cancers associated with inflammatory myopathies.

Antibody	Clinical association
Anti-Jo-1 and other antisynthetase antibodies	PolymyositisDermatomyositis with interstitial lung disease, arthritis, and mechanic's hands
Anti-Mi-2	Dermatomyositis with rash more than myositis
Anti-MDA5(anti-CADM 140)	Dermatomyositis with rapidly progressive lung disease and cutaneous ulcers
Anti- 155/140 orAnti-transcription intermediary factor (TIF)-1γ	Cancer-associated myositis
Anti-SAE	Cancer-associated dermatomyositis, dermatomyositis with rapidly progressive lung disease

Sold by @itachibot

Solution to Question 13:

The clinical presentation and muscle biopsy finding of perimysial inflammation and perifascicular atrophy makes dermatomyositis the most likely diagnosis.

Muscle biopsy findings in various inflammatory myositis are as follows:

Dermatomyositis	Polymyositis	Inclusion body myositis
Perimysialinfiltrates	Endomysialinfiltrates	Endomysialinfiltrates
Perivascular infiltrates andpe rifascicularatrophy	Perivascular infiltrates	Perivascular infiltrates with a myloid inclusion bodies and r immed vacuoles
CD4+cellular infiltrates	CD8+cellular infiltrates	CD8+cellular infiltrates

If you purchased this from so
you may have been scammed.
Sold by @ita

Solution to Question 14:

The given clinical scenario having features of both systemic sclerosis and dermatomyositis points towards a diagnosis of overlap syndrome. The presence of anti-PM/Scl antibodies is diagnostic of overlap syndrome.

Anti-PM/Scl is a specific antinuclear antibody directed against a nucleolar-protein complex.

Overlap syndrome also known as mixed connective tissue disease usually involves a combination of three connective tissue diseases:

• Systemic lupus erythematosus (SLE)

• Progressive systemic sclerosis (PSS)

• Polymyositis (PM) or dermatomyositis (DM)

Solution to Question 15:

The most probable diagnosis for the given clinical scenario is fibromyalgia.

Fibromyalgia is characterized by chronic widespread musculoskeletal pain and tenderness accompanied by other non-specific somatic symptoms like fatigue and sleep disturbance, for >three months.

It is more common in females. On examination, they have tenderness in multiple soft tissue anatomical locations. Laboratory findings are normal.

Patients with fibromyalgia may have an increased prevalence of other syndromes associated with pain and fatigue, like chronic headaches, chronic fatigue syndrome, irritable bowel syndrome, interstitial cystitis/ painful bladder syndrome, and pelvic pain syndromes.

Sold by @itachibot

Options A and B: Polymyositis and dermatomyositis have positive laboratory findings such as specific antibodies and radiological findings of multiorgan involvement (ex: interstitial lung disease in polymyositis).

Option C: Although the given clinical scenario closely resembles clinical depression, the examination findings showing tenderness and pain sensitivity favor the diagnosis of fibromyalgia.

Solution to Question 16:

Single-photon emission computed tomography (SPECT) studies done on fibromyalgia patients show hypoperfusion of the thalamus.

Fibromyalgia is characterized by chronic widespread musculoskeletal pain and tenderness. It is more common in females.

Patients with fibromyalgia have an increased prevalence of other syndromes associated with pain and fatigue, including chronic fatigue syndrome, chronic headaches, irritable bowel syndrome, interstitial cystitis/ painful bladder syndrome, and pelvic pain syndromes.

Non-pharmacological treatment strategies for fibromyalgia include physical conditioning and aerobic exercise. Opioid analgesics should be avoided in fibromyalgia as they are not effective and are associated with opioid-induced hyperalgesia that can worsen both the symptoms and function.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Seronegative Spondyloarthritides

Question 1:

Seronegative spondyloarthritides are associated with which of the following?

a) HLA-B27

b) HLA-B16

c) HLA-B47

d) HLA-DR4

Question 2:

A 26-year-old man presented with pain and stiffness in his lower back. His X-ray spine is given below. Which of the following is true about his condition?

a) Steroids are the first-line therapy

b) Symptoms improve with activity

c) It is associated with HLA B47

d) Associated with low serum IgA levels

Question 3:

If yo... bought by @itachibot ...s from someone else, ...scammed.

Sold by @itachibot

What are the key cytokines involved in the pathogenesis of ankylosing spondylitis?

a) Interferon γ & IL-23

b) Interferon γ & IL-17

c) TNF-α & IL-13

d) TNF-α & IL-17

Question 4:

A patient with Crohn's disease presents with back pain and stiffness. The physician suspects ankylosing spondylitis and orders an MRI for evaluation. Which of the following joints will be the earliest to be involved?

a) Costovertebral joint

b) Lumbosacral joint

c) Sacroiliac joint

d) Sacrococcygeal joint

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

A 28-year-old male patient with low backache and early morning stiffness for the past 1 year is diagnosed with ankylosing spondylitis. Which of the following is not associated with the haplotype related to this condition?

a) Behcet's disease

b) Reiter's syndrome

c) Reactive arthritis

d) Psoriatic spondylitis

Question 6:

The following test is performed in a patient to evaluate back pain associated with stiffness. Identify the test.

Sold by @itachibot



©MARROW

a) Gaenslen test

b) Thomas test

c) Adam's test

d) Schober's test

Question 7:

Which is the most common extra-articular manifestation seen in patients diagnosed with ankylosing spondylitis?

a) Anterior uveitis

b) Cauda equina syndrome

c) Inflammatory bowel disease

d) Acne fulminans

Question 8:

Which of the following conditions predominantly involve the upper lobe of the lung?

a) Rheumatoid arthritis

b) Asbestosis

c) Scleroderma

d) Ankylosing spondylitis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 9:

A patient who presented with chronic lower back pain has been diagnosed with ankylosing spondylitis. What is the first line of pharmacological therapy?

a) TNF α inhibitors

b) Opioids

c) NSAIDS

d) Steroids

Question 10:

A 27-year-old man with ankylosing spondylitis is not responding to standard therapy. His physician decides to prescribe him an anti-IL-17 monoclonal antibody. Which of the following drugs would he prescribe?

a) Ustekinumab

b) Secukinumab

c) Inflximab

d) Adalimumab

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Question 11:

A 15-year-old boy presents with generalized malaise and painful swelling of his right knee joint since 10 days. He also complains of pain and discomfort during urination. He was treated for diarrhea last month. What is the most likely diagnosis?

a) Reactive arthritis

b) Enteropathic arthritis

c) Hemolytic uremic syndrome

d) Pyogenic arthritis

Question 12:

A patient presents with painful knee swelling, dysuria, and conjunctivitis. Which of the following infectious agents is least likely to cause this?

a) Shigella dysenteriae

b) Neisseria gonorrhoea

Sold by @itachibot

c) Campylobacter jejuni

d) Chlamydia trachomatis

Question 13:

A middle-aged man presented with pain and swelling of his right knee for one week. It started progressing to the ankle joints and feet for the past 2 days. He gives a history of diarrhea and burning micturition 4 weeks ago. On examination, his soles showed the following lesions. What is the most likely diagnosis?

a) Syphilitic arthritis

b) Enteropathic arthritis

c) Psoriatic arthritis

d) Reactive arthritis

Question 14:

An elderly man presents with pain and deformity of his toes. The clinical findings are shown in the image below. What is the most likely diagnosis?

Sold by @itachibot

a) Rheumatoid arthritis

b) Osteo arthritis

c) Psoriatic arthritis

d) Septic arthritis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 15:

A 16-year-old girl is being evaluated for arthritis and uveitis. Which of the following would you not consider as a differential diagnosis?

a) Psoriatic arthritis

b) Enteropathic arthritis

c) Juvenile onset spondyloarthritis

d) Reactive Arthritis

Question 16:

Which of the following is not a part of the SAPHO syndrome?

a) Synovitis

b) Acne

c) Psoriasis

d) Hyperostosis

Sold by @itachibot

Question 17:

Uveitis is an uncommon manifestation in patients with which of the following conditions?

a) Ankylosing spondylitis

b) Psoriatic arthritis

c) Reiter's syndrome

d) Rheumatoid arthritis

If you bought this from someone else, it was stolen.

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	c
5	a
6	d
7	a
8	d
9	c
10	b
11	a
12	b
13	d
14	c
15	c
16	c
17	d

Detailed Explanations

Solution to Question 1:

Seronegative spondyloarthritides are associated with HLA-B27.

Sold by @itachibot

The spondyloarthritides are a group of overlapping disorders that share certain clinical features and genetic associations (HLA-B27). These disorders include:

- Psoriatic arthritis
- Ankylosing spondylitis
- Reactive arthritis
- Enteropathic arthritis
- Juvenile onset spondyloarthritis
- Undifferentiated spondyloarthritis

Usually, rheumatoid factor (RF) assay is negative in spondyloarthritides. Hence, they are also referred to as seronegative arthritis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

The given clinical features along with the classical bamboo-spine appearance on X-ray point towards a diagnosis of ankylosing spondylitis (AS). In ankylosing spondylitis, the symptoms improve with activity.

Ankylosing spondylitis (AS) is an inflammatory disorder that affects the axial skeleton primarily, and may also involve peripheral joints. It is strongly associated with HLA-B27. It usually presents in the mid-twenties.

It presents with lower back pain that improves with activity and worsens with rest or inactivity. Early morning stiffness is a characteristic feature. It is associated with elevated serum IgA levels. NSAIDs are the first line of pharmacological therapy.

It is also associated with extra-articular manifestations like uveitis, interstitial lung disease, aortic regurgitation, etc.

NSAIDs are the first-line treatment for ankylosing spondylitis.

Solution to Question 3:

The key cytokines involved in the pathogenesis of ankylosing spondylitis are TNF-α and IL-17. IL-23 also plays a role in pathogenesis.

Solution to Question 4:

The earliest joint to be involved in ankylosing spondylitis is the sacroiliac joint.

Ankylosing spondylitis (AS) is an inflammatory disorder that primarily affects the axial skeleton. It is more common in males, usually presenting between 20-30 years of age.

Sold by @itachibot

Ankylosing spondylitis (AS) and peripheral arthritis can occur as extra-intestinal manifestations of inflammatory bowel disease (IBD). They are more common in Crohn's disease than in ulcerative colitis.

Solution to Question 5:

Ankylosing spondylitis is related to the HLA-B27 haplotype. This haplotype is not associated with Behcet's disease. It is associated with HLA B51.

Ankylosing spondylitis is a chronic inflammatory disease that mainly affects the spine and sacroiliac joint. Individuals who inherit this class I HLA allele (HLA-B27) have a 100–200 fold greater chance (odds ratio or relative risk) of developing the disease compared with those who do not carry HLA-B27. Many autoimmune diseases are associated with different class II HLA alleles.

Solution to Question 6:

The test performed is modified Schober's test. This test is done in patients with suspected ankylosing spondylitis.

It is used to detect reduced flexion of the spine. The test involves marking two points- at the lumbosacral junction and 10 cm above it. The patient is then asked to bend forward with knees fully extended and the distance between the two points is measured.

This distance normally increases by ≥2 cm. An increase of <2 cm indicates reduced flexion.

In the image given below the distance increases by 5cm which is normal.



Option A: Gaenslen's test is used to detect primary chronic inflammation of the sacroiliac joint.

Option B: Thomas test is used to detect fixed flexion deformity of the hip.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Sold by @itachibot

Option C: Adam's forward bending test is performed on a patient to differentiate structural from non-structural scoliosis.

Solution to Question 7:

The most common extraarticular manifestation of ankylosing spondylitis (AS) is anterior uveitis.

It usually presents with unilateral pain, photophobia and increased lacrimation. It is also associated with other spondyloarthritides like reactive arthritis, psoriatic arthritis and enteropathic arthritis.

Other extra-articular manifestations of AS include:

• GIT: ileitis, colitis, inflammatory bowel disease

• Cardiovascular: increased risk of IHD, aortic insufficiency, heart block

• Respiratory: upper lobe fibrosis

• Neurological: cauda equina syndrome

• Skin: acne fulminans, hidradenitis suppurativa

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

Ankylosing spondylitis predominantly involves the upper lobe of the lung leading to pulmonary fibrosis.

Rheumatoid arthritis, scleroderma, and asbestosis predominantly involve the lower lobes of the lungs.

Solution to Question 9:

Nonsteroidal anti-inflammatory drugs (NSAIDs) are the first line of pharmacological therapy for ankylosing spondylitis (AS).

Management of ankylosing spondylitis includes:

• First-line management:

• Physiotherapy to maintain posture and range of motion.

• NSAIDs- reduce pain and tenderness, increase mobility, and slow down the disease progression.

• TNF α inhibitors- in patients who do not respond to NSAID therapy. It is not used in all patients because of side effects like reactivation of TB, hypersensitivity reactions, pancytopenia, drug-induced psoriasis, etc.

• Anti-IL-17 antibodies- in patients with failed anti-TNF therapy.

• Surgery- improves mobility in patients with severe arthritis and joint stiffness.

Sold by @itachibot

Solution to Question 10:

Secukinumab is an anti-IL-17 monoclonal antibody used in patients with ankylosing spondylitis who are not responding to NSAIDs and anti-TNF therapy.

Management of ankylosing spondylitis includes:

• First-line management:

• Physiotherapy to maintain posture and range of motion.

• NSAIDs- reduce pain and tenderness, increase mobility, and slow down the disease progression.

• TNF α inhibitors- in patients who do not respond to NSAID therapy. It is not used in all patients because of side effects like reactivation of TB, hypersensitivity reactions, pancytopenia, drug-induced psoriasis, etc.

• Anti-IL-17 antibodies- in patients with failed anti-TNF therapy.

• Surgery- improves mobility in patients with severe arthritis and joint stiffness.

Option A: Ustekinumab is an anti-IL-12/23 monoclonal antibody used in the treatment of psoriasis.

Options C and D: Infliximab and adalimumab are anti-TNF-α monoclonal antibodies used in the treatment of ankylosing spondylitis.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 11:

The given scenario describes an acute arthritis following an infection which is also associated with urethritis and generalized malaise. These features are suggestive of reactive arthritis.

Reactive arthritis also known as Reiter's syndrome is a seronegative spondyloarthritis that occurs as a consequence of a recent enteric or urogenital infection. It is associated with HLA B27. It is also associated with extra-articular manifestations such as:

• Urogenital- urethritis, prostatitis, cervicitis

• Ocular- conjuctivitis, anterior uveitis

• Mucocutaneous- oral ulcers, keratoderma blennorrhagica, circinate balanitis

In some patients it presents as a triad of:

• Arthritis

• Urethritis

• Conjunctivitis

Solution to Question 12:

Sold by @itachibot

The given scenario of arthritis, urethritis, and conjunctivitis is suggestive of reactive arthritis. It is not caused by Neisseria gonorrhea.

Disseminated gonococcal infection causes gonococcal arthritis. It comprises localized septic arthritis and arthritis-dermatitis syndrome.

Reactive arthritis is usually caused due to LPS (lipopolysaccharide) component of the gram-negative organisms or cross-reactivity with synovial antigens. The common causative organisms include:

• Chlamydia trachomatis (most common)

• Salmonella

• Shigella

• Yersinia

• Campylobacter

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 13:

The given scenario of acute arthritis following a recent infection and additive pattern (involvement of new joints within few days) and examination finding of keratoderma blennorrhagica (image) are suggestive of reactive arthritis.

Keratoderma blennorrhagica is a mucocutaneous manifestation of reactive arthritis. It presents as vesicles and pustules that lead to hyperkeratotic lesions or crusting on the palms and soles.

Other mucocutaneous manifestations of reactive arthritis include:

• Oral ulcers - superficial, transient, and often asymptomatic

• Circinate balanitis - consists of vesicles that rupture to form painless superficial erosions on the glans penis

Solution to Question 14:

The given clinical image showing dactylitis, nail changes and telescoping of toes is suggestive of psoriatic arthritis.

Clinical features of psoriatic arthritis include:

• Psoriatic nail changes- yellowing and hypertrophy

• Dactylitis- inflammation of the toes

• Telescoping (shortening) of toes due to osteolysis

• Severe deformity- suggestive of arthritis mutilans

• Enthesitis- heel pain at the sites of ligament or tendon attachment

Psoriatic arthritis of the hands commonly involves proximal and distal interphalangeal joints (PIP, DIP) and metacarpophalangeal joints (MCP). The image below shows psoriatic arthritis of

Sold by @itachibot

the hands:

Sold by @ita... If you purchased this from so... else, you may have been scammed.

Solution to Question 15:

Juvenile onset spondyloarthritis (JSpA) would not be considered as a differential diagnosis as it has no associated extra-articular manifestations like uveitis, lung involvement, etc.

It is a seronegative spondyloarthritis with onset between 7-16 years of age. It usually presents as asymmetric arthritis predominantly involving the lower extremities.

Solution to Question 16:

Psoriasis is not a part of SAPHO syndrome.

The SAPHO syndrome is characterized by:

- Synovitis- arthritis
- Acne- acne conglobata
- Pustulosis- palmoplantar pustulosis, hidradenitis suppurativa
- Hyperostosis- sternoclavicular and spinal hyperostosis
- Osteitis- sterile osteomyelitis

Solution to Question 17:

Uveitis is a rare manifestation of rheumatoid arthritis (RA). Ocular manifestation in RA includes scleritis, episcleritis, and keratoconjunctivitis.

Sold by @itachibot

Uveitis is a common ocular manifestation of seronegative arthritis e.g. Reiter's syndrome, ankylosing spondylitis, psoriatic arthritis, and juvenile rheumatoid arthritis (JRA).

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

SLE and APS

Question 1:

What type of hypersensitivity reaction is seen in systemic lupus erythematosus?

a) Type I

b) Type II

c) Type III

d) Type IV

Question 2:

Patients with which of the following complement deficiencies are at an increased risk of developing systemic lupus erythematosus?

a) C1q

b) C5

c) C3

d) C1 inhibitor

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Anti-nuclear antibodies are required for the diagnosis of which of the following?

a) Scleroderma

b) Systemic lupus erythematosus

c) Rheumatoid arthritis

d) Sjogren's syndrome

Question 4:

A 25-year-old marathon runner presents with complaints of joint pain and rashes over the face. She says the rashes appear on exposure to the sun. On physical examination, a butterfly rash is seen over her face sparing the nasolabial folds. Which of the following antibodies will you evaluate to screen this patient?

Sold by @itachibot

a) Anti-dsDNA antibodies

b) Anti-Sm antibodies

c) Antinuclear antibodies

d) Anti-histone antibodies

Question 5:

A school teacher presented with painful oral ulcers, photosensitive rashes, and arthralgia. Presence of which of the following antibodies would be diagnostic of her condition?

a) Anti Ro/SS-A antibodies

b) Anti-Sm antibodies

c) Antinuclear antibodies

d) Anti-dsDNA antibodies

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 6:

An SLE patient is found to have a butterfly rash. Which of the following is an incorrect statement regarding this finding?

a) Most common acute rash in SLE

b) Spares the nasolabial folds

c) Increased risk of squamous cell carcinoma

d) Associated with scaling

Question 7:

What is the most common pulmonary manifestation of SLE?

a) Shrinking lung syndrome

b) Pleuritis

c) Intra alveolar hemorrhage

d) Interstitial inflammation

Question 8:

Sold by @itachibot

A 40-year-old man with history of SLE presented to your OPD with shortness of breath. It is associated with chest pain that worsens on inspiration. On auscultation, you notice a pleural friction rub. Investigations reveal the following. What is the most likely diagnosis?

a) Lupus pneumonitis

b) Interstitial fibrosis

c) Shrinking lung syndrome

d) Intra-alveolar hemorrhage

Question 9:

What is the most common neurological manifestation seen in patients with systemic lupus erythematosus?

a) Seizures

b) Depression

c) Cognitive impairment

d) Polyneuropathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A young girl is being evaluated for low mood, anhedonia, weight loss, and insomnia since past 1 month. On examination, multiple hesitation marks were seen on her left forearm. She was diagnosed with systemic lupus erythematosus (SLE) one year back. Which of the following autoantibodies is associated with this presentation?

a) Anti-ribosomal P antibodies

b) Anti-Ro/SSA antibodies

c) Anti-neuronal antibodies

d) Anti-histone antibodies

Question 11:

What is the most common cardiac manifestation seen in patients with systemic lupus erythematosus?

a) Pericarditis

b) Myocarditis

Sold by @itachibot

c) Libman Sack endocarditis

d) Myocardial infarction

Question 12:

Which of the following is a feature of arthritis in SLE?

a) Non-deforming arthritis

b) Predilection for large joints

c) Asymmetric involvement

d) Non-erosive arthritis

Question 13:

A young woman who is a known case of SLE develops proteinuria. The doctor suspects lupus nephritis and performs a renal biopsy. It shows the presence of wire loop appearance without glomerular proliferation. In which class of lupus nephritis will you categorize this patient's biopsy findings?

a) Mesangial proliferative lupus nephritis

b) Focal lupus nephritis

c) Diffuse lupus nephritis

d) Membranous lupus nephritis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

A 40 year old female presented to the OPD for her annual routine check-up. she is a patient of SLE diagnosed a decade ago, among the given options what is the most common cause of death in such patients?

a) Cardiovascular disease

b) Renal failure

c) Infections

d) Disseminated intravascular coagulation

Question 15:

Sold by @itachibot

Which of the following drugs is recommended for all patients with SLE?

a) Hydroxychloroquine

b) NSAIDs

c) Methotrexate

d) Systemic steroids

Question 16:

A patient with a history of SLE presented with hematuria and proteinuria. Investigations revealed elevated creatinine and low C3, C4 levels. He was unresponsive to steroid therapy and his physician prescribed him belimumab for further management. What is the mechanism of action of this drug?

a) TNF inhibitor

b) IL-1 antagonist

c) Anti CD-20 antibody

d) BAFF inhibitor

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

A 60-year-old man presented with fever, malaise, skin rashes, and arthralgia. On further evaluation, his serology was positive for ANA and anti-histone antibodies. Use of which of the following drugs could have led to this condition?

a) Rifampin

b) Bupivacaine

c) Warfarin

d) Hydralazine

Question 18:

What is the ratio of occurrence of drug-induced lupus erythematosus in females and males?

a) 9:1

b) 3:1

c) 2:1

d) 1:1

Sold by @itachibot

Question 19:

Which of the following statements is true regarding drug-induced lupus erythematosus?

a) Renal involvement is rare

b) Commonly associated with anti ds-DNA

c) Common in females

d) Reduced C3 and C4 levels

Question 20:

Antiphospholipid syndrome is associated with all of the following except:

a) Recurrent fetal loss

b) Thrombocytosis

c) Arterial thrombosis

d) Libman-Sacks endocarditis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

A 34-year-old woman comes for evaluation of recurrent fetal losses. She has had 4 spontaneous abortions, all of them at <10 weeks of gestational age, and has a history of deep vein thrombosis. Which of the following antibodies is not useful in making a diagnosis in this condition?

a) Lupus anticoagulant

b) Anti-cardiolipin

c) Anti-beta-2 glycoprotein

d) Anti-beta-2 microglobulin

Question 22:

A 40-year-old man presented with painful swelling of his right calf muscle. The clinical test shown below was positive. Investigations revealed elevated D-dimer, high titres of lupus anticoagulant and anti-cardiolipin antibodies. You would use all of the following drugs for his management except:

Sold by @itachibot



a) Warfarin

b) Glucocorticoids

c) Aspirin

d) Hydroxychloroquine

Sold by @itachi... purchased this from some... else, If you may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	a
3	b
4	c
5	b
6	c
7	b
8	c
9	c
10	a
11	a
12	d
13	c
14	a

Sold by @itachibot

15	a
16	d
17	d
18	d
19	a
20	b
21	d
22	b

Detailed Explanations

Solution to Question 1:

Type III hypersensitivity reaction (immune complex deposition) is seen in systemic lupus erythematosus.

Systemic lupus erythematosus (SLE) is an autoimmune condition in which autoantibodies are directed against a part of the nucleus such as double-stranded DNA, histone, ribonuclear proteins, etc. The presence of these autoantibodies leads to the deposition of immune complexes and activation of the complement system.

Solution to Question 2:

C1q complement deficiency is a risk factor for the development of systemic lupus erythematosus (SLE).

Risk factors for SLE are:

- Genetic
- Genetic defects resulting in C1q, C2, C4 complement deficiencies
- Presence of HLA DR3
- Mutation in the TREX1 gene
- Environmental
- Epstein-Barr virus (EBV) infection
- Oral contraceptive pills/ hormone replacement therapy
- Smoking
- Prolonged exposure to silica

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Solution to Question 3:

Antinuclear antibodies (ANAs) are required for the diagnosis of systemic lupus erythematosus.

ANAs are autoantibodies that bind to the contents of the cell nucleus. They are present in > 95% of patients with SLE. A positive antinuclear antibody (ANA) test with a titer of at least 1:80 is the obligatory entry criterion along with the additive criteria in 7 clinical and 3 immunologic domains. If a total of >10 points achieved, it is indicative of SLE according to 2019 EULAR/ACR Classification Criteria for Systemic Lupus Erythematosus (SLE)

Most sensitive antibody for SLE: ANA

Most specific antibody for SLE: Anti smith > Anti dsDNA

ANA may also be positive in the other options but it is not a criterion for diagnosis of these conditions.

The following antibodies are specific for their respective autoimmune conditions.

• Scleroderma (option A): Anti-topoisomerase I(Scl-70) and anti-centromere antibodies highly specific for Systemic sclerosis.

• Rheumatoid arthritis (option C): RF and anti-CCP antibodies

• Sjogren syndrome (option D): Anti-Ro, Anti-La

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

The given clinical scenario is suggestive of systemic lupus erythematosus (SLE). The best screening test in this condition is to evaluate for the presence of antinuclear antibodies (ANAs).

Solution to Question 5:

The given clinical scenario of oral ulcers, photosensitive rash, and arthralgia is suggestive of systemic lupus erythematosus (SLE). Anti-Sm antibodies are most specific for SLE and hence their presence is diagnostic of this condition.

Solution to Question 6:

Butterfly rash or malar rash in a patient with SLE is not associated with an increased risk of squamous cell carcinoma.

It is the most common type of acute skin rash seen in SLE. It is a photosensitive, erythematous, and scaly rash that is prominent over the cheeks and the nose. It spares the nasolabial folds.

Common causes of malar rash are as follows:

• Erysipelas

• Cellulitis

Sold by @itachibot

- Systemic lupus erythematosus
- Rosacea
- Pellagra
- Dermatomyositis

Solution to Question 7:

The most common pulmonary manifestation of SLE is pleuritis, with or without pleural effusion.

The most common manifestations of different systems involved in SLE are:

- Musculoskeletal- arthritis and arthralgia
- Skin- photosensitivity
- Kidneys- proteinuria
- Neurologic- cognitive impairment
- Cardiac- pericarditis
- Pulmonary- pleuritis
- Hematologic- anemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

The given clinical scenario along with elevated hemidiaphragm and no evidence of ILD on chest CT is suggestive of shrinking lung syndrome (SLS), which is a rare complication of systemic lupus erythematosus (SLE).

It is characterized by dyspnea and episodes of pleuritic chest pain. These symptoms occur in the absence of pleural or parenchymal involvement.

PFT shows decreased lung volume with a restrictive pattern, and DLCO is often reduced. Radiological investigations show an elevated hemidiaphragm with no evidence of ILD.

The following X-ray shows shrinking lung syndrome.

Sold by @itachibot

Pulmonary manifestations seen in SLE are:

• Pleuritis with or without pleural effusion- most common pulmonary manifestation.

• Lupus pneumonitis - It is the inflammation of the lung tissue (pneumonitis) due to SLE (lupus).

• Interstitial lung fibrosis

• Shrinking lung syndrome

• Intra-alveolar hemorrhage

If you purchased this from someone else, you may have been scammed. Sold by @itachi

Solution to Question 9:

The most common neurological manifestation seen in patients with systemic lupus erythematosus (SLE) is cognitive impairment.

It can manifest as impaired memory and reasoning. Other common neurological manifestations in SLE are psychosis, depression, and seizures.

The most common manifestations of different systems involved in SLE are:

• Musculoskeletal- arthritis and arthralgia

• Skin- photosensitivity

• Kidneys- proteinuria

• Neurologic- cognitive impairment

• Cardiac- pericarditis

• Pulmonary- pleuritis

• Hematologic- anemia

Sold by @itachibot

Solution to Question 10:

The given clinical scenario is suggestive of depression in a patient with SLE. It is associated with anti-ribosomal P antibodies.

Antiribosomal P antibody is associated with psychosis or depression in SLE.

Solution to Question 11:

Pericarditis is the most common cardiac manifestation seen in patients with systemic lupus erythematosus.

Other cardiac manifestations include:

• Myocarditis

• Libman-Sacks endocarditis.

Patients with SLE are at an increased risk for myocardial infarction, usually due to accelerated atherosclerosis.

The most common manifestations of different systems involved in SLE are:

• Musculoskeletal- arthritis and arthralgia

• Skin- photosensitivity

• Kidneys- proteinuria

• Neurologic- cognitive impairment

• Cardiac- pericarditis

• Pulmonary- pleuritis

• Hematologic- anemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

Non-erosive arthritis is a common feature of arthritis in SLE.

Features of arthritis in SLE are:

• Symmetrical and polyarticular arthritis

• A predilection for the small joints of the hand (the proximal interphalangeal and carpal joints)

• Non-erosive and deforming arthritis (reducible)

In contrast to rheumatoid arthritis which is characterized by erosive, deforming joint involvement, SLE has non-erosive and deforming arthritis (reduciable).

Solution to Question 13:

Sold by @itachibot

Wire loop appearance without glomerular proliferation is seen in diffuse lupus nephritis (Class IV).

The capillaries are thickened due to the subendothelial deposits and appear as wire loops. It is the most common and most severe form of lupus nephritis.

Renal involvement is common in patients with SLE. The majority of the patients will present with

- Proteinuria, nephrotic syndrome
- Microscopic hematuria, hypertension
- Urinary casts
- Abnormal renal function tests- elevated serum creatinine
- Low C3 and C4 levels

Sold by @itachibot
If you purchased this from else, you may have been scammed.

Solution to Question 14:

The most common cause of death in patients with SLE after 10 years of disease is accelerated atherosclerosis.

The leading causes of death in the first 10 years of the disease are infections and renal failure; subsequently, Cardiovascular events become increasingly frequent causes of mortality.

Solution to Question 15:

Hydroxychloroquine is recommended for all patients with SLE. It decreases the symptoms, reduces flares and organ damage.

Important drugs used in the management of SLE are:

Sold by @itachibot

• NSAIDs- used for a shorter period to reduce pain and fever. They are not used in all patients because of the increased risk of atherosclerosis and renal injury.

• Hydroxychloroquine or chloroquine- should be used in all patients with SLE

• Glucocorticoids

• topical glucocorticoids- in mild skin lesions

• systemic steroids- in severe skin lesions and organ involvements.

• Immunosuppressive agents- used in combination with glucocorticoids in severe skin lesions and organ involvement. They help to reduce the dose and toxicity of glucocorticoids. Important agents are:

• Methotrexate

• Azathioprine

• Mycophenolate mofetil and cyclophosphamide- immunosuppressive agents of choice in renal involvement.

• Biologic drugs- like belimumab and rituximab are used when there is no response to conventional drugs.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

The given clinical scenario is suggestive of lupus nephritis. Belimumab is a BAFF inhibitor used in patients with lupus nephritis who do not show response to initial therapy with steroids.

Belimumab is a human monoclonal antibody that inhibits the soluble form of a B cell survival factor (known as BLyS or BAFF) which promotes the formation of memory B cells and autoantibodies. Belimumab inhibits this factor and therefore is used in the management of SLE.

It is used in combination with either cyclophosphamide or MMF in patients with lupus nephritis who do not show response to initial therapy with steroids.

CD20 is an antigen present on the B cells. Rituximab is an anti-CD20 monoclonal antibody that is used in patients who are resistant to the standard therapies.

Solution to Question 17:

The given clinical scenario with positive ANA and anti-histone antibodies point towards a diagnosis of drug-induced lupus erythematosus. Hydralazine which is a vasodilator is associated with the development of drug-induced lupus erythematosus.

Drug-induced lupus erythematosus is an autoimmune condition caused due to exposure to certain medications. It completely resolves after the causative drug is discontinued. Important drugs associated with this condition are:

• Sulfonamides

• Hydralazine

Sold by @itachibot

- Isoniazid
- Infliximab
- Procainamide
- Propylthiouracil
- ACE inhibitors
- Thiazides
- TNF-alpha antagonists

Solution to Question 18:

The female to male ratio in drug-induced lupus erythematosus (DILE) is 1:1.

The female to male ratio in systemic lupus erythematosus (SLE) is 9:1.

	Systemic lupus erythematosus (SLE)	Drug-induced lupus erythematosus (DILE)
Age	Common in young individuals	Common in elderly (50-70 years)
Female: male ratio	9:1	1:1
Organs involved	Can affect many organs	Skinis mainly involved.Renal and brain involvement arerare
Complement levels	Reduced C3, C4 levels	Normal C3, C4 levels
Antibodies		
-Antinuclear	Common	Common
-Antihistone	Rare	Common
-Anti ds-DNA	Common	Rare

Solution to Question 19:

Renal involvement is rare in drug-induced lupus erythematosus.

	Systemic lupus erythematosus (SLE)	Drug-induced lupus erythematosus (DILE)
Age	Common in young individuals	Common in elderly (50-70 years)

Sold by @itachibot

	Systemic lupus erythematosus (SLE)	Drug-induced lupus erythematosus (DILE)
Female: male ratio	9:1	1:1
Organs involved	Can affect many organs	Skinis mainly involved.Renal and brain involvement arerare
Complement levels	Reduced C3, C4 levels	Normal C3, C4 levels
Antibodies		
-Antinuclear	Common	Common
-Antihistone	Rare	Common
-Anti ds-DNA	Common	Rare

Sold by @itachibot
If you purchased this from someone
you may have been scammed.

Solution to Question 20:

Thrombocytosis is not associated with antiphospholipid syndrome (APS). On the contrary, thrombocytopenia is seen.

APS is an autoantibody-mediated condition that is characterized by arterial thrombosis, venous thrombosis, and recurrent fetal loss. These autoantibodies bind to the phospholipids present on the endothelial cells and activate coagulation and thrombus formation.

Important manifestations of venous thrombosis are:

• Deep vein thrombosis of legs

• Pulmonary embolism

• Livedo reticularis

• Cerebral venous thrombosis

Important manifestations of arterial thrombosis are:

• Stroke, transient ischemic attacks

• Retinal artery thrombosis

• Myocardial ischemia

• Libman-Sacks endocarditis- due to the vegetations formed by platelet fibrin microthrombi

• Leg ulcers, gangrene of the digits

Important obstetric manifestations are:

• Preeclampsia

• Recurrent fetal loss

• Premature birth

Important hematological abnormalities are:

Sold by @itachibot

• Thrombocytopenia- occurs due to increased platelet aggregation and thrombus formation

• Coombs positive hemolytic anemia- seen when autoantibodies attack red blood cells

• Thrombotic microangiopathic syndromes- thrombotic thrombocytopenic purpura (TTP) and hemolytic uremic syndrome (HUS)

Solution to Question 21:

The given clinical history of >3 spontaneous abortions at <10 weeks of gestational age along with a history of deep vein thrombosis suggests a diagnosis of antiphospholipid syndrome (APS). Anti beta-2 microglobulin is not useful in diagnosing this condition.

For the diagnosis of antiphospholipid syndrome, high titers of one or more of the following three antibodies should be demonstrated along with at least one clinical symptom of thrombosis or pregnancy complications:

• Lupus anticoagulant (LA) - most specific

• Anticardiolipin (aCL) antibodies - most sensitive

• Anti-beta-2 glycoprotein antibodies

Solution to Question 22:

The given findings of positive Moses' sign with elevated D-dimer, lupus anticoagulant, and anti-cardiolipin antibodies are suggestive of antiphospholipid syndrome (APS). Glucocorticoids are not useful in the general management of APS, they are used only in the treatment of catastrophic APS.

Squeezing of the relaxed calf muscles from side to side is painful as the thrombosed deep veins in the calf are always tender. This test is known as Moses' sign.

The clinical manifestations in APS occur due to the activation of coagulation and thrombus formation. Therefore, anticoagulants and antiplatelet agents are the mainstays in the treatment. Important drugs used are:

• Warfarin- is an anticoagulant that inhibits vitamin K dependant epoxidase.

• Low molecular weight heparins (LMWH)- warfarin cannot be used in pregnancy due to its teratogenicity. In these situations, LMWH is used for anticoagulation.

• Aspirin- is an antiplatelet agent.

• Hydroxychloroquine- is an immunomodulatory agent that can be used to prevent recurrent thrombosis.

• Statins

Note: The only condition where corticosteroids are beneficial is catastrophic APS. It is a life-threatening rapidly progressive thromboembolic disease involving simultaneously three or more organs. It is seen in very few patients.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Rheumatoid Arthritis

Question 1:

Which of the following is associated with rheumatoid arthritis?

a) HLA B27

b) HLA DR4

c) HLA DR3

d) HLA DR5

Question 2:

Which part of the spine is most commonly affected in patients with rheumatoid arthritis?

a) C1 - C2

b) C6 - C7

c) L4 - L5

d) T1 - T2

Question 3:

Identify the following deformity seen in a patient complaining of pain and swelling of her hands.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Swan neck deformity

b) Boutonnière deformity

c) Bouchard's nodes

d) Hitchhiker's deformity

Question 4:

A patient presents to your clinic with the following deformity. In which of the following conditions is this deformity seen?

a) Osteoarthritis

b) Gout

c) Rheumatoid arthritis

d) Psoriatic arthritis

Question 5:

An elderly woman presents with a chronic history of pain in the small joints of hands with stiffness of joints in the early hours of the day. The image of the patient's hands is given below. What is the most likely diagnosis?

Sold by @itachibot

a) Rheumatoid Arthritis

b) Osteoarthritis

c) Complex Regional Pain Syndrome

d) Villonodular synovitis

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 6:

What is the most common extra-articular manifestation of rheumatoid arthritis?

a) Subcutaneous nodule

b) Sjogren's syndrome

c) Felty's syndrome

d) Vasculitis

Question 7:

Which is the most common cardiac valvular lesion seen in patients with rheumatoid arthritis?

a) Mitral regurgitation

b) Aortic stenosis

c) Aortic regurgitation

d) Mitral stenosis

Sold by @itachibot

Question 8:

Which of the following would you not expect to see in a patient diagnosed with Felty syndrome?

a) Splenomegaly

b) Neutropenia

c) Hepatomegaly

d) Rheumatoid arthritis

Question 9:

Which is the most common type of lymphoma seen in patients with rheumatoid arthritis?

a) Mantle cell lymphoma

b) Hodgkin's lymphoma

c) Diffuse large B cell lymphoma

d) Marginal zone lymphoma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A patient complains about difficulty doing chores due to morning stiffness of the joints. Investigations show elevated ESR, CRP, and positive Rheumatoid factor. Which of the following is not likely to be seen in this condition?

a) Hyperandrogenism

b) Hip fractures

c) Keratoconjunctivitis sicca

d) Congestive heart failure

Question 11:

All of the following characteristics are found in the pleural effusion fluid of a rheumatoid arthritis patient except:

a) RA factor

b) High glucose

c) Cholesterol crystals

Sold by @itachibot

d) High LDH

Question 12:

Which of the following radiological features would help to differentiate between rheumatoid arthritis and SLE?

a) Juxta-articular osteoporosis

b) Ulnar deviation

c) Articular erosions

d) Swelling of PIP joint

Question 13:

What is the most common cause of death in patients with rheumatoid arthritis?

a) Interstitial lung disease

b) Coronary artery disease

c) Atlantoaxial subluxation

d) Gastrointestinal bleeding

Question 14:

In a suspected case of rheumatoid arthritis, which of the following markers should be used to diagnose this condition?

a) Anti-nuclear antibodies

b) Rheumatoid factor

c) Anti- CCP antibody

d) Anti dsDNA antibody

Question 15:

Which of the following is the drug of choice for the treatment of a patient with rheumatoid arthritis?

a) Hydroxychloroquine

b) Sulfasalazine

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Methotrexate

d) Infliximab

Question 16:

A patient with rheumatoid arthritis is being treated with methotrexate and low-dose corticosteroids for the past 4 months. However, the disease is still progressing. What would be your recommendation for the further management of this patient?

a) Stop oral methotrexate and start parenteral methotrexate

b) Add sulfasalazine and hydroxychloroquine

c) Continue corticosteroids and methotrexate

d) Start only anti-TNF alpha agents

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

Which of the following is a Janus kinase inhibitor used in patients with rheumatoid arthritis?

a) Anakinra

b) Tocilizumab

c) Abatacept

d) Tofacitinib

Question 18:

A 28-year-old pregnant lady visits your OPD with complaints of painful swelling of her hands since past 2 months. Investigations reveal positive rheumatoid factor and anti-CCP antibodies. Which of the following DMARDs should not be prescribed to her?

a) Rituximab

b) Leflunomide

c) Hydroxychloroquine

d) Sulfasalazine

Question 19:

What is the isotype of the rheumatoid factor most commonly measured in the laboratory?

Sold by @itachibot

a) Ig M

b) Ig G

c) Ig A

d) Ig D

Question 20:

Iritis is seen in all except:

a) SLE

b) Behcet's disease

c) Ulcerative colitis

d) Rheumatoid arthritis

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	b
2	a
3	b
4	c
5	a
6	a
7	a
8	c
9	c
10	a
11	b
12	c
13	b
14	c
15	c
16	b
17	d
18	b

Sold by @itachibot

19	a
20	d

Detailed Explanations

Solution to Question 1:

Rheumatoid arthritis is associated with HLA DR4.

Solution to Question 2:

The atlantoaxial joint (C1 & C2) is the most commonly affected part of the spine in patients with rheumatoid arthritis.

The joints in rheumatoid arthritis are bilaterally and symmetrically involved. The commonly affected joints are:

• Metacarpophalangeal (MCP) joints

• Wrists

• Proximal interphalangeal (PIP) joints

• Metatarsophalangeal joints (MTP) joints

• Atlantoaxial joint

There is relative sparing of the DIP joint in rheumatoid arthritis. The involvement of DIP joints should direct suspicion towards osteoarthritis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

The image is showing flexion of the PIP joint with hyperextension of the DIP joint is suggestive of Boutonnière deformity seen in patients with rheumatoid arthritis.

Heberden's and Bouchard's nodes are nodules in osteoarthritis that are palpable in the distal interphalangeal joint and proximal interphalangeal joints respectively.

Solution to Question 4:

The given image shows a swan-neck deformity seen in a patient with established rheumatoid arthritis(RA). Swan-neck deformity is hyperextension at the proximal interphalangeal (PIP) joint and flexion at the distal interphalangeal (DIP) joint.

Some of the other small joint deformities of the hand in a patient with RA include:

Sold by @itachibot

Boutonnie■re deformity with flexion at the PIP joint and hyperextension at the DIP joint.

Z-line deformity with hyperextension at the DIP joint and subluxation of the 1st metacarpophalangeal (MCP) joint.

Solution to Question 5:

The above image shows flexion at the PIP joint and extension in the DIP joints which is known as boutonniere deformity which is seen in rheumatoid arthritis(RA).

If the patient gives a history of pain/stiffness in the joints in the early hours of the day which usually decreases as the day progresses suggesting inflammatory arthritis and if it presents with predominant involvement of smaller joints, it is most likely a case of RA(option A).

It occurs due to the accumulation of inflammatory mediators which on increased joint mobility are flushed out due to increased blood supply.

Pain and stiffness in osteoarthritis (Option B) usually increase as the day progresses due to an increase in the activity of the joint. They typically present with nodules around the distal interphalangeal joint (Heberden nodes) and proximal interphalangeal joint (Bouchard nodes).

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 6:

The most common extra-articular manifestation of rheumatoid arthritis(RA) is subcutaneous nodules.

The nodules are generally benign, non-tender, and firm, and occur in areas subjected to repeated trauma such as the forearm, sacral prominences, and Achilles tendon. They can also occur in the lungs, pleura, pericardium, and peritoneum.

Patients who have a history of cigarette smoking had early onset arthritic manifestations and tested positive for RA factor or ACPA are more likely to develop extra-articular manifestations. Other common extra-articular manifestations include secondary Sjogren's syndrome, interstitial lung disease, pulmonary nodules, and anemia. FELTY'S syndrome is a triad of neutropenia, splenomegaly, and nodular RA.

Solution to Question 7:

The most common cardiac valvular lesion seen in patients with rheumatoid arthritis is mitral regurgitation.

Solution to Question 8:

Hepatomegaly is not seen in Felty syndrome.

Sold by @itachibot

Felty syndrome is defined by the clinical triad of:

• Neutropenia

• Splenomegaly

• Rheumatoid arthritis

It occurs in the late stages of rheumatoid arthritis (RA).

Other syndromes associated with rheumatoid arthritis are:

• Caplan's syndrome- characterized by both rheumatoid arthritis and pneumoconiosis. It is associated with development of multiple peripheral basilar nodules in the lungs.

• Secondary Sjogren's syndrome- development of keratoconjunctivitis sicca or xerostomia in association with connective tissue disorder.

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Solution to Question 9:

The most common histopathologic type of lymphoma seen in patients with rheumatoid arthritis is diffuse large B cell lymphoma (DLBCL).

The risk of developing lymphoma increases if the patient has Felty's syndrome or high levels of disease activity.

Solution to Question 10:

The given clinical scenario is suggestive of rheumatoid arthritis, which is associated with hypoandrogenism.

It is characterized by low levels of serum testosterone, LH, and DHEA.

The conditions associated with rheumatoid arthritis are:

• Cardiovascular disease, congestive heart failure

• Osteoporosis - hip fractures are more likely to occur

• Ocular - keratoconjunctivitis sicca, episcleritis, scleritis, keratitis

• Secondary Sjogren's syndrome

• Hypoandrogenism in men and postmenopausal women

Solution to Question 11:

Low glucose (<60 mg/dL) is seen in the pleural effusion fluid of a patient with rheumatoid arthritis (RA).

RA causes exudative pleural effusion and fulfills the Light's criteria. High LDH and RA factor are found in the pleural fluid of RA patients. Less commonly, patients with RA may have a cholesterol

Sold by @itachibot

effusion. It is sterile and has a milky or opaque appearance. The milky appearance is due to the presence of cholesterol crystals and an elevated cholesterol level.

Light's criteria is used to distinguish between exudative and transudative pleural effusions. An exudative pleural effusion satisfies at least one of the following criteria:

• Pleural fluid protein/serum protein > 0.5

• Pleural fluid LDH / serum LDH > 0.6

• Pleural fluid LDH more than 2/3rds upper limit of normal for serum



Solution to Question 12:

Articular erosions on X-ray is seen in rheumatoid arthritis (RA), but not in SLE.

Juxta-articular osteoporosis, swelling of PIP joint, and ulnar deviation can be seen in both SLE and RA.

Jaccoud arthropathy is a deforming non-erosive arthropathy characterized by the ulnar deviation of the second to 5th fingers with metacarpophalangeal joint subluxation seen in SLE.

The give X-ray shows Jaccoud's arthropathy:

If you purchas... you may have... else,

Sold by @itachibot

The given image shows a patient with Jaccoud's arthropathy of the hands:

Solution to Question 13:

The most common cause of death in patients with rheumatoid arthritis is cardiovascular disease.

Solution to Question 14:

The most appropriate marker to diagnose rheumatoid arthritis (RA) is the anti-CCP (cyclic citrullinated peptide) antibody as it is the most specific marker for RA.

Anti-CCP, ESR, and CRP values correlate with the disease activity.

Sold by @itachibot

Rheumatoid factor can occur in other autoimmune diseases as well as in chronic infections. Therefore it is not considered specific.

Solution to Question 15:

Methotrexate which is a DMARD is the drug of choice for the treatment of rheumatoid arthritis (RA).

A combination regimen is used when there is a failure to achieve adequate improvement with methotrexate only.

Methotrexate in low doses causes adenosine release and suppresses joint inflammation. However, in high doses, it causes inhibition of dihydrofolate reductase (DHFR).

Solution to Question 16:

For further management of a patient with rheumatoid arthritis (RA) with worsening disease activity, the preferred therapy is a step-up to oral triple therapy with methotrexate, sulfasalazine, and hydroxychloroquine. The addition of leflunomide or a biological (anti - TNF) to methotrexate is another alternative.

Ideally, the approach to a patient with early RA is aggressive. Methotrexate is always the first DMARD that the patient is started on, and if an adequate response is not observed within 3-6 months, a step up in therapy is made by adding on drugs.

For assessment of progression, composite indices like Disease Activity Score - 28 (DAS-28) can used.

Other options:

Option A: Changing the route of administration of methotrexate from oral to parenteral is not a part of step up

Option C: Priority is given to minimising the duration the patient is on steroids.

Option D: Anti -TNF agents (adalimumab, etanercept) are not recommended as a monotherapy, and are reserved for treatment in combination with DMARDs. This is due to the fact that the efficacy of monotherapy is lesser than that of DMARDs, and the adverse effects include increased risk of serious infections, re-activation of latent TB and opportunistic fungal infections. They are reserved for cases where there is a lack of response with the first line agents.

Solution to Question 17:

The Janus kinase inhibitor used in the treatment of rheumatoid arthritis is tofacitinib.

Tofacitinib, baricitinib, and upadacitinib are Janus kinase inhibitors used as monotherapy or combination therapy in rheumatoid arthritis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option A: Anakinra is an IL-1 antagonist.

Option B: Tocilizumab is an IL-6 antagonist.

Option C: Abatacept acts by inhibiting co-stimulation of T cells.

Solution to Question 18:

The given clinical scenario is suggestive of rheumatoid arthritis (RA). Leflunomide should not be prescribed in pregnancy as it is a category X drug.

DMARDs classified as Category X in pregnancy (contraindicated in pregnancy) are:

- Methotrexate
- Leflunomide

Solution to Question 19:

IgM isotype is the rheumatoid factor (RF) most commonly measured in laboratories.

IgM RF has been found in 75–80% of patients with rheumatoid arthritis. IgG and IgA isotypes of RF have also been identified.

However, RF is not specific for rheumatoid arthritis and can also be found in the following conditions:

- Primary Sjögren's syndrome
- Systemic lupus erythematosus
- Subacute bacterial endocarditis
- Hepatitis B and C

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 20:

Iritis is seen in all the above conditions except rheumatoid arthritis.

Iritis or iridocyclitis is also known as anterior uveitis.

Anterior uveitis is seen in:

- Ankylosing spondylitis
- Juvenile rheumatoid arthritis
- Psoriasis
- Reactive arthritis
- Systemic lupus erythematosus
- Leprosy

Sold by @itachibot

Posterior uveitis is seen in:

• Vogt-Koyanagi Harada syndrome

• Diseases like toxoplasmosis, cysticercosis, coccidioidomycosis, toxocariasis, histoplasmosis

• Infections- Candida, Pneumocystis carinii, Cryptococcus, Aspergillus, and Cytomegalovirus

• Cat-scratch disease, Whipple's disease, and brucellosis

Both anterior and posterior uveitis is seen in:

• Sarcoidosis

• Behcet's disease

• Inflammatory bowel disease

• Herpes infections, Syphilis, Lyme disease, TB, onchocerciasis

Intermediate uveitis/pars planitis is seen in multiple sclerosis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Asthma & COPD

Question 1:

Which among the following about idiosyncratic asthma is false?

a) Presents in childhood

b) Normal IgE levels

c) Commonly have concomitant nasal polyps

d) Negative skin test for common inhalant allergens

Question 2:

A 36-year-old woman presented with wheezing after taking aspirin for a headache. Which of the following is true regarding this condition?

a) Raised serum IgE levels

b) Positive skin test for inhaled allergens

c) Associated with nasal polyps

d) A type of extrinsic asthma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

What is the best way to diagnose patients with suspected bronchial asthma?

a) Chest X-ray

b) Skin test for inhaled allergens

c) FEV1 values

d) Demonstration of reversible obstruction

Question 4:

A young woman is rushed to ED due to wheezing and gasping for breath. She is unable to speak in full sentences. Her husband reports that she has had a cough with fever for 3 days. On examination, PR- 121/min, RR- 32/min, and has cyanosis. Her ABG analysis and chest X-ray are given below. What is the likely diagnosis?

Sold by @itachibot



a) Acute respiratory distress syndrome

b) Asthma with impending failure

c) Acute pneumothorax

d) Acute severe asthma

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 5:

Which of the following drugs is not recommended in the treatment of patients with bronchial asthma?

a) Sodium cromoglycate

b) Zafirlukast

c) Roflumilast

d) Dupilumab

Question 6:

Which of the following is false regarding leukotriene modifiers in the management of bronchial asthma?

a) Ketotifen belongs to this group

b) Associated with Churg- Strauss syndrome

c) Protect against exercise-induced asthma

d) Diminishes nocturnal symptoms of asthma

Sold by @itachibot

Question 7:

Treatment of choice for patients with type 2 brittle asthma is:

a) Inhaled Salbutamol

b) Subcutaneous salbutamol

c) Subcutaneous terbutaline

d) Subcutaneous adrenaline

Question 8:

Which of the following class of drugs cannot be used in the treatment of patients with bronchial asthma?

a) IL-4 receptor antagonists

b) Anti–IL-5 antibodies

c) Anti IgE antibody

d) IL- 1 antagonist

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 6-year-old asthmatic boy is rushed to the casualty. He is not responding to the commands and is having fast shallow breathing. On auscultation, the chest is silent and paradoxical breathing is present. ABG analysis shows respiratory acidosis. What is the best step in management for him?

a) Hyperbaric oxygen therapy

b) MDI with salbutamol and corticosteroids

c) Immediate endotracheal intubation

d) Intravenous magnesium sulfate

Question 10:

Which of the following statements is false regarding COPD?

a) Increased resistance is seen in airways ≤2 mm diameter

b) Changes in large airways cause cough and sputum

Sold by @itachibot

c) Ventilation-perfusion mismatching is seen

d) Rise in post bronchodilator FEV1>12%

Question 11:

What happens to Reid index in patients with chronic bronchitis?

a) Increases

b) Decreases

c) Remains same

d) Increases initially and then decreases

Question 12:

Which of the following findings is least likely to be seen in patients with COPD?

a) Barrel chest

b) Hoover's sign

c) Cyanosis

d) Clubbing

Question 13:

A 28-year-old coal miner presented with breathing difficulties. He reported that his symptoms get aggravated at work. His chest X-ray showed features of emphysema. On evaluation, he was found to have an α1-antitrypsin deficiency. Which pathological type of emphysema is most likely present in this patient?

a) Centri-acinar

b) Pan-acinar

c) Irregular

d) Para-septal

Question 14:

An elderly man is being evaluated for exertional dyspnea, cough, and weight loss. His chest CT is given below. What is it suggestive of?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) Emphysematous lung

b) Atypical pneumonia

c) Calcified granulomas

d) Interstitial fibrosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 15:

A 55-years old chronic smoker presented with chronic cough with expectoration. He has been having similar episodes for the past 2 years. He also reported that he had to rest every few minutes while walking due to breathlessness. Pulmonary function tests showed FEV1/FVC of 0.5 and FEV_1 of 50%. Which of the following is not a feature of the given condition?

a) Irreversible airflow limitation

b) Affects small airway disease

c) Decrease in residual volume

d) Associated with ineffective repair of elastin

Question 16:

Chest X-ray of an emphysematous patient is as shown. What is the best investigation to measure his lung volume?

Sold by @itachibot



a) Body plethysmography

b) Gas dilution

c) Fractional exhaled NO

d) DLCO

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

A 66-year-old chronic smoker presents with progressive shortness of breath and chronic cough. On examination, he has a barrel chest and has pursed-lip breathing. Hyperresonance and inspiratory wheezes are heard. Pulmonary functions are likely to reveal all of the following, except:

a) Decreased vital capacity

b) Increased diffusion capacity for carbon monoxide

c) Increased total lung capacity

d) Decreased FEV1/FVC ratio

Question 18:

The Gold's criteria for severe COPD are:

a) FEV1/FVC <0.7 and FEV1 ≥ 30%

b) FEV1/FVC <0.7 and FEV1 < 30%

c) FEV1/FVC<0.7 and FEV1< 20%

Sold by @itachibot

d) FEV1/FVC<0.7 and FEV1 ≥ 50%

Question 19:

A patient presents to the OPD with shortness of breath. He reports walking slower than his peers due to breathlessness. He has had two hospitalizations for similar complaints this year. His post-bronchodilator PFTs are FEV1/FVC < 0.55 and FEV1 55%. Under what category would he be classified based on GOLD criteria?

a) Moderate COPD, Group C

b) Moderate COPD, Group E

c) Severe COPD, Group A

d) Severe COPD, Group D

Question 20:

Which of the following is false regarding acute exacerbations in COPD patients?

a) May not be accompanied with fever

b) Viral infection is a common precipitating factor

c) Glucocorticoids are contraindicated

d) May present with congestive heart failure

Question 21:

A 60-year-old male patient who is a known case of COPD presented with acute exacerbation and is admitted to the ICU. Which of the following statements is correct with regards to the initial management of this patient?

a) Non-Invasive PPV should be given

b) Invasive PPV should be given

c) IV corticosteroids should be administered

d) Permissive hypercapnia is allowed

Question 22:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

A 45-year-old obese chronic smoker is rushed to the ER with difficulty in breathing and a productive cough. He has a past history of two similar episodes this year. On examination, RR- 34/min, with use of accessory muscles and central cyanosis is present. ABG is as shown. Chest X-ray shows increased vascular markings. What is the best initial step of management?

a) Low flow humidified oxygen

b) High flow humidified oxygen

c) Salbutamol and budesonide nebulisation

d) Slow infusion of theophylline

Question 23:

In patients with which of the following conditions is NIPPV contraindicated?

a) COPD

b) Central hypoventilation syndrome

c) Cardiovascular instability

d) Chest wall disorders

Question 24:

Which of the following is not used in the long-term treatment of COPD patients?

a) Theophylline

b) Azithromycin

c) Oral glucocorticoids

d) N Acetylcysteine

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	c
3	d
4	d

Sold by @itachibot

5	c
6	a
7	d
8	d
9	c
10	d
11	a
12	d
13	b
14	a
15	c
16	a
17	b
18	a
19	b
20	c
21	a
22	a
23	c
24	c

If you purch... else, you may hav... d.

Detailed Explanations

Solution to Question 1:

Idiosyncratic or intrinsic asthma does not present in childhood. It typically shows a later onset of disease (adult-onset asthma).

Solution to Question 2:

The given clinical scenario is suggestive of aspirin-sensitivity asthma. It is closely associated with nasal polyps.

Aspirin-exacerbated respiratory disease (AERD), also known as Samter's triad or aspirin-sensitive asthma. It is a chronic medical condition that consists of:

• Intrinsic asthma

• Recurrent sinus disease with nasal polyps

• Sensitivity to aspirin and other NSAIDs

Sold by @itachibot

It has normal serum IgE levels and a negative skin test for inhaled allergens.

Solution to Question 3:

The best way to diagnose bronchial asthma is by the demonstration of reversible obstruction of the airways. Following the administration of an inhaled beta 2 agonist, a 12% and 200 ml increase in the FEV1 demonstrates the reversibility of airway obstruction.

Options A and B: Chest X-ray and skin test for inhaled allergens are not diagnostic of bronchial asthma.

Option C: FEV1 values are measured to calculate FEV1/FVC ratio to diagnose any obstructive pathology and not specifically for asthma alone.

Solution to Question 4:

The given clinical scenario of dyspnea, tachycardia along with the ABG analysis suggestive of respiratory alkalosis and chest-Xray showing hyperinflated lungs is pointing to a diagnosis of acute severe asthma. A history of cough with fever is indicating a probable respiratory infection to have triggered the asthma exacerbation.

Features of acute severe asthma vs asthma with impending respiratory failure:

Note: The X-ray is not showing the features of ARDS and pneumothorax, hence they are ruled out as the right options.

Acute severe asthma	Asthma with impending respiratory failure
Dyspneic, speaks in words	Mute patient
Agitated patient	Silent / confused patient
Heart rate > 120 /min, tachycardia	Relative bradycardia
Pulsus paradoxus present	Pulsus paradoxus absent (due to respiratory muscle fatigue)
Accessory muscles of respiration used	Paradoxical breathing is seen
Loud wheeze throughout the respiration	Silent chest
Peak expiratory flow rate < 40 ml/min	Peak expiratory flow rate < 25 ml/min
Respiratory alkalosis	Respiratory acidosis
Hyperventilation leading to low or normal $PaCO_2$	Normal or rising $PaCO_2$

If you purchased this from someone else, Sold by @itachibot may have been scammed.

Sold by @itachibot

Solution to Question 5:

Roflumilast is a phosphodiesterase-4 inhibitor used in the treatment of COPD, not asthma.

Drugs used in the treatment of bronchial asthma

Bronchodilators

• Beta-2 sympathomimetics: Salmeterol, formoterol

• Methylxanthines: Theophylline, aminophylline

• Anticholinergics: Ipratropium bromide, tiotropium bromide

Anti-inflammatory

• Corticosteroids (inhalational): Beclomethasone, budesonide, fluticasone

• Corticosteroids (systemic): Prednisolone, hydrocortisone

• Mast cell stabilizers: Sodium cromoglycate, ketotifen

• Leukotriene antagonists: Montelukast, zafirlukast

• Anti-IgE: Omalizumab

• Anti-IL5: Mepolizumab, reslizumab

• Anti-IL4 receptor: Dupilumab

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

Ketotifen is an example of a mast cell stabilizer, not leukotriene modifiers.

Leukotriene modifiers are a group of agents used for mild persistent asthma. They block LT1-receptors thus causing reduced eosinophilic inflammation and causing bronchodilator. They include montelukast, zafirlukast, and zileuton.

Leukotriene modifiers are less effective than inhaled corticosteroids (ICS) in controlling asthma. They are used as add-on therapy in some patients whose asthma is not controlled with low doses of ICS.

Churg–Strauss syndrome (CSS) is a rare granulomatous small vessel vasculitis, that affects the heart, kidneys, peripheral nerves, etc. It occurs against a background of long-standing asthma and marked peripheral eosinophilia. Several cases of CSS are associated with the use of montelukast and zafirlukast.

Solution to Question 7:

The treatment of choice for type II brittle asthma is subcutaneous adrenaline.

Sold by @itachibot

Brittle asthma is a type of severe asthma where patients have very serious and often, life-threatening attacks despite appropriate therapy. It is of two types:

Type 1	Type 2
A persistent pattern of variability in PEFR (peak expiratory flow rate) readings	Generally normal or near-normal PEFR readings but few unpredictable exacerbations occur without an obvious trigger
Oral steroids, infusion of bronchodilators, and dietary exclusion of allergens may be helpful	Respond only to subcutaneous epinephrine

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 8:

The treatment options for asthma do not include IL-1 antagonists (anakinra). Anakinra is used in the treatment of rheumatoid arthritis.

Cytokines mediating airway inflammation in asthma: IL-4, IL-5, IL-13, TGF-β, and chemokines.

Omalizumab is a monoclonal antibody against IgE. It is available for the treatment of children ≥6 years of age with documented allergen-induced asthma that is inadequately controlled by inhaled corticosteroids.

The new treatment options for asthma under clinical trial are:

- Mepolizumab: anti–IL-5 antibody
- Benralizumab: anti–IL-5 receptor antibody
- Dupilumab: anti–IL-4 receptor antibody

Solution to Question 9:

The given clinical scenario is suggestive of severe asthma with impending respiratory failure. The best step in the management is immediate endotracheal intubation.

Features of asthma with impending respiratory failure are:

- Mute or confused patient
- Paradoxical breathing
- Silent chest on auscultation
- Relative bradycardia
- Peak expiratory flow rate (PEFR) <25ml/min
- Arterial blood gas analysis shows respiratory acidosis.

Sold by @itachibot

Option A: Hyperbaric oxygen is given in CO poisoning and high altitude pulmonary edema.

Option B: MDI with salbutamol and corticosteroids is only useful for acute severe asthma.

Option D: Intravenous magnesium sulfate is not routinely used.

Solution to Question 10:

The rise in post-bronchodilator FEV1>12% is a feature of asthma, not COPD. It suggests reversible airflow obstruction.

Chronic obstructive pulmonary disease (COPD) is a chronic respiratory disease characterized by the presence of airflow obstruction that is partially reversible.

Cigarette smoking is the most common cause of COPD. Cigarette smoke exposure affects large airways, small airways, and alveoli:

• Changes in large airways cause cough and sputum

• Changes in small airways (≤2mm) and alveoli are responsible for physiologic alterations

The major site of increased resistance in most individuals with COPD is in airways ≤2 mm diameter.

Characteristic cellular changes include goblet cell metaplasia. Goblet cells are mucus-secreting cells that replace surfactant-secreting Club cells.

Nonuniform ventilation and ventilation-perfusion mismatching are characteristic of COPD.

Solution to Question 11:

Reid index is increased in chronic bronchitis.

Reid index is the ratio of the thickness of the mucous gland layer to the thickness of the bronchial wall.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

Clubbing is not a sign of COPD. Its presence should trigger the investigation of the causes of clubbing, particularly lung cancer.

Signs of COPD:

• Signs of respiratory distress

• Tripod posture - facilitates the use of accessory muscles of respiration

• Pursed lip breathing - seen in emphysema

• Central cyanosis

• Signs of cachexia

• Severe weight loss

• Bitemporal wasting

• Signs of hypercapnia (type 2 respiratory failure) - more common in chronic bronchitis

• Barrel-shaped chest - increase in the anteroposterior diameter

• Enlarged lung volumes

• Positive Hoover's sign - paradoxical inward movement of the chest during inspiration

• Early inspiratory coarse crackles

• Normal vesicular breath sounds of reduced-intensity with bilaterally prolonged expiration

• Blue bloaters - cyanotic (blue) and edematous (bloaters), chronic bronchitis predominant features

• Pink puffers - thin, non-cyanotic (pink) at rest, and severely dyspneic (puffers), emphysema predominant features

Sold by @itachibot

Causes of clubbing:

• Pulmonary
• Bronchogenic carcinoma
• Lung abscess, empyema, bronchiectasis
• Chronic interstitial pneumonitis
• Cystic fibrosis
• Sarcoidosis
• Gastrointestinal
• Inflammatory bowel disease
• Sprue
• Neoplasms - esophagus, liver, bowel
• Cardiovascular
• Cyanotic congenital heart disease
• Subacute bacterial endocarditis
• Aortic aneurysm
• Arteriovenous fistula
• Endocrine
• Hyperthyroidism (Grave's disease)

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

The image below shows the characteristic tripod posture in a patient with COPD.

Solution to Question 13:

Sold by @itachibot

The pathological type most likely to be present in this patient is pan-acinar emphysema. This type is commonly seen in patients with α1-antitrypsin deficiency.

Solution to Question 14:

The CT scan shows emphysematous lungs, with reduced parenchymal markings and focal areas of low attenuation. This represents the emphysematous destruction of the lung, indicative of hyperinflation.

Radiographic studies in COPD show:

• Emphysematous bullae

• A paucity of parenchymal markings

• Hyperlucency suggests the presence of emphysema

• Increased lung volumes and flattening of the diaphragm suggest hyperinflation

Option B: Atypical pneumonia shows patchy bilateral ground-glass opacities.

Option C: Calcified granulomas are seen as dense, bright nodules.

Option D: Interstitial fibrosis shows a honeycomb pattern of the lung parenchyma as shown below.



If yo... sold by @itachibot ...this from someone else, ...n scammed!

Solution to Question 15:

The clinical picture is highly suggestive of chronic obstructive pulmonary disease(COPD). An increase in the residual volume is a feature of COPD.

COPD is defined as a chronic respiratory disease characterized by the presence of airflow obstruction that is partially reversible.

Sold by @itachibot

COPD is the clinical expression of abnormalities of airways and/or alveoli and includes:

• Emphysema is defined pathologically as the abnormal and permanent dilation of airways distal to the terminal bronchiole, along with the destruction of their walls.

• Chronic bronchitis is defined clinically as the presence of persistent cough with sputum production on most of the days for ≥3 months in 2 consecutive years.

• Small airways diseases are conditions affecting airways of ≤2 mm, resulting in inflammation and narrowing of the same.

Solution to Question 16:

Body plethysmography is the best investigation to measure lung volume in an emphysematous patient with bullous lesions as seen on the X-ray.

To measure absolute lung volumes, two techniques are used:

• Body plethysmography: It is used to measure the volume of any gas containing space in the thorax including bullae. It is accurate in both obstructive and restrictive diseases.

• Inert gas dilution technique: It is used to measure the volume of regions that are well ventilated which excludes trapped gas or bullae. It is accurate in restrictive diseases only.

The amount of gas trapped in bullae can be measured by subtracting lung volumes measured by inert gas dilution technique from lung volume measured by body plethysmography.

Fractional exhaled nitric oxide (FENO) is a non-invasive test used in the diagnosis and monitoring of bronchial asthma. NO levels are increased in asthmatics and related to the eosinophilic inflammation in the airways.

Solution to Question 17:

The given clinical diagnosis is suggestive of COPD and the underlying pathological change is likely to be emphysema. It is associated with a decreased diffusion capacity for carbon monoxide (DLCO).

Emphysema is defined pathologically as the abnormal and permanent dilation of airways distal to the terminal bronchiole, along with the destruction of their walls. This leads to hyperinflation due to air trapping and impaired diffusion of gases.

Pulmonary function tests in COPD:

• TLC: Normal or increased

• FEV1: Decreased (<80% predicted)

• FVC: Decreased

• FEV1/FVC: Decreased

• DLCO: Decreased

Diseases associated with lowered DLCO:

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

• Emphysema

• Interstitial Lung disease

• Primary pulmonary hypertension

Note - Bronchial asthma is associated with an increased DLCO.

Solution to Question 18:

Gold's criteria for severe COPD (stage III) are FEV1/FVC <0.7 and FEV1 ≥ 30% but <50%

The severity of COPD is assessed by Global Initiative for Chronic Obstructive Lung Disease (GOLD) classification:

Note: FEV1/FVC and FEV1 predicted are measured after the administration of a bronchodilator.

GOLD Stage	Severity	Spirometry
I	Mild	FEV1/FVC <0.7 and FEV1 ≥80% predicted
II	Moderate	FEV1/FVC <0.7 and FEV1 ≥50% but <80% predicted
III	Severe	FEV1/FVC <0.7 and FEV1 ≥30% but <50% predicted
IV	Very severe	FEV1/FVC <0.7 and FEV1 <30% predicted

Recent update: PRISm refers to a group of COPD patients with apparent normal spirometry, but with emphysema (as seen on radiographic evaluation) and a wide range of lung function impairment. These patients have a higher mortality risk as compared to other COPD patients. The possible risk factors are exposure to smoking, high BMI, and reduced total lung capacity. Spirometry findings are low to normal FEV1, and normal FEV1/FVC ratio.

Solution to Question 19:

This patient belongs to moderate COPD, group E category.

The severity of COPD assessed by the Global Initiative for Lung Disease (GOLD) is classified as:

According to GOLD 2023 guidelines, COPD patients are classified into Group A, B, or E based on the severity of dyspnoea, frequency of exacerbations and hospitilization. Dyspnoea may be assessed using the modified MRC Dyspnoea scale or COPD Assessment Test (CAT score)

* single inhaler therapy maybe more convenient and effective than multiple inhalers

ICS - inhaled corticosteroids

LAMA - Long-acting Muscarinic Antagonists

Sold by @itachibot

LABA - Long-acting Beta Agonists

Modified MRC (mMRC) dyspnoea scale:

• 0- Breathless only with strenuous exercise

• 1- Breathless when hurrying on a level or walking up a hill

• 2- Walks slower than peers or has to stop for breath when walking on level

• 3- Stops for breath after walking about 100m or for a few minutes

• 4- Breathless at rest

GOLD Stage	Severity	Spirometry
I	Mild	FEV1/FVC <0.7 and FEV1 ≥80% predicted
II	Moderate	FEV1/FVC <0.7 and FEV1 ≥50% but <80% predicted
III	Severe	FEV1/FVC <0.7 and FEV1 ≥30% but <50% predicted
IV	Very severe	FEV1/FVC <0.7 and FEV1 <30% predicted

	>=2 moderate exacerbations or >=1 leading to hospitalisation	Group E:LABA+LAMAconsider LABA+LAMA+ICS* if blood eosinophil count >=300
0 or 1 moderate exacerbation (not leading to hospital admissions)	Group A: mMRC 0-1, CAT<10Bronchodilator	Group B: mMRC >=2, CAT >=10LABA + LAMA*

Solution to Question 20:

Glucocorticoids are used during an exacerbation of COPD. They hasten recovery and reduce the chances of future exacerbations.

Exacerbations are episodes of increased dyspnoea and cough, with a change in the amount and character of sputum. They are often triggered by an infection with one-third of cases having a viral etiology. They may or may not have other signs of illness such as fever, myalgias, and sore throat. In severe cases, they may present with pneumonia and congestive heart failure.

Note: While oral glucocorticoids are contraindicated in long-term treatment due to side effects, they are used for acute exacerbation of COPD.

Sold by @itachibot

Solution to Question 21:

Non-invasive positive pressure ventilation (NIPPV) should be the first mode of ventilation used in COPD patients with acute respiratory failure provided there are no absolute contraindications. This approach improves gas exchange, reduces work of breathing and the need for intubation, decreases hospitalization duration and enhances overall survival outcomes.

Invasive positive pressure ventilation is only considered after failure of NIPPV, resulting in severe respiratory distress, life threatening hypoxemia, severe hypercarbia and/or acidosis, markedly impaired mental status, respiratory arrest or hemodynamic instability (Option B).

Apart from NIPPV, short-acting inhaled beta 2 agonists (albuterol, salbutamol) with or without short-acting anticholinergics (ipratropium) and Long-acting anticholinergics (tiotropium) are recommended as the initial bronchodilators. Maintenance therapy with long-acting beta 2 agonists (formoterol) must be started as soon as possible.

Systemic corticosteroids are useful in the treatment of acute exacerbations of COPD as they reduce airway inflammation and improve lung function over time. However, the effects are not immediate. NIPPV on the other hand provides immediate relief to the patient and reduces the work of breathing. Therefore, it would not be the best choice in this case. (Option C)

Antibiotics can also be considered, as they reduce duration of hospitalisation. Treatment is given for 5 days.

Permissive hypercapnia refers to a ventilation strategy where tidal volume is reduced in order to avoid hyperinflation-associated lung trauma. It is not preferred at the initial presentation. It is only permitted once invasive PPV is started following NIPPV failure (Option D)

An exacerbation of COPD is defined as an event characterized by dyspnea and/or cough and sputum that worsen over < 14 days. Exacerbations of COPD are often associated with increased local and systemic inflammation caused by airway infection, pollution, or other insults to the lungs.

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Solution to Question 22:

The given clinical scenario along with the ABG and X-ray findings is suggestive of acute exacerbation of COPD with respiratory failure. The best initial step of management is low-flow humidified oxygen.

Oxygen supplementation is a crucial component of acute therapy. Administering O2 should target a SpO2- 88-92% or PaO2- 60-70 mmHg. This helps to minimize the risk of worsening hypercapnia with excess supplemental oxygen.

The given ABG shows respiratory acidosis as PaCO2 >45 mmHg, and pH<7.35. In respiratory acidosis, a trial of non-invasive ventilation is given. Hence, low-flow humidified oxygen is the first step in management.

Option B: High flow oxygen causes dryness, crusting epistaxis, and damages the lung hence not used.

Option C: Salbutamol and budesonide nebulization are not effective as initial management in exacerbation of COPD with respiratory failure.

Sold by @itachibot

Option D: Methylxanthines are not recommended in the management of acute exacerbations due to their side effect profiles.

Solution to Question 23:

Noninvasive positive-pressure ventilation (NIPPV) is contraindicated in cardiovascular instability. NIPPV can impede venous filling of the heart, decrease cardiac output, and worsen/precipitate hypotension. Moreover, in unstable states with impending cardiovascular collapse, invasive ventilation is preferred.

Indications of NIPPV:

• Apnea - central and obstructive

• Central hypoventilation syndrome

• Neuromuscular and chest wall disorders

• Respiratory failure - PaCO2 >45 mmHg

Contraindications of NIPPV:

• Cardiovascular instability

• Impaired mental status or inability to cooperate

• Copious secretions or the inability to clear secretions

• Craniofacial abnormalities

• Trauma precluding effective fitting of the mask

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 24:

Oral glucocorticoids are not used in the long-term treatment of COPD because of an unfavorable benefit/risk ratio.

Chronic use of glucocorticoids is associated with significant side effects, including osteoporosis, weight gain, cataracts, glucose intolerance, and increased risk of infection.

Treatment of COPD:

• Smoking cessation

• Anticholinergic agents- Ipratropium bromide

• Beta Agonists- Long-acting inhaled β agonists, such as salmeterol or formoterol are commonly used.

• Methylxanthines- theophylline produces improvements in expiratory flow rates and vital capacity.

• Phosphodiesterase inhibitors - roflumilast is used when FEV1<50%.

• Antibiotics- macrolides like azithromycin reduce exacerbations when used over one year

Sold by @itachibot

• N-acetylcysteine has been used in patients with COPD for both its mucolytic and antioxidant properties

• Oxygen therapy

Non-pharmacologic therapies:

• Lung volume reduction surgery

• Lung transplantation

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Pneumonia

Question 1:

Which of the following organisms can cause both typical and atypical pneumonia?

a) Legionella

b) Pneumococcus

c) Mycoplasma

d) Staphylococcus aureus

Question 2:

There is an outbreak of legionella pneumonia in the hospital. Which of the following is least likely to have played a role in the transmission of the organism?

a) Home split air conditioners

b) Cooling towers

c) Evaporative condensers

d) Humidifiers

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 55-year-old smoker presents with high-grade fever, shortness of breath, diarrhea, and confusion. Gram-staining of the sputum revealed neutrophilia with no organisms. Chest x-ray showed bilateral multilobar infiltrates. Lab values are given below. Which pathogenic agent is suspected?

a) Legionella

b) Pneumocystis jirovecii

c) Klebsiella pneumoniae

d) Streptococcus pneumoniae

Question 4:

Urinary antigen detection test is recommended for diagnosing patients with?

Sold by @itachibot

a) Mycoplasma pneumonia

b) Chlamydia pneumonia

c) Streptococcus pneumonia

d) Legionella pneumonia

Question 5:

A middle-aged man presents with fever, cough with scanty sputum, and fatigue. He had a throat infection a week ago. His blood investigations and chest X-ray are given below. Which is the most likely causative agent for his condition?



a) Pneumococcus

b) Mycoplasma

c) Legionella

d) Chlamydia

Question 6:

Which of the following is not true for patients with Mycoplasma pneumonia?

a) Causative organism cannot be cultured routinely

b) Diffuse infiltrates in the lung are seen

c) Cefotaxime is the drug of choice

d) Serology is useful in diagnosis

Sold by @itachibot

Question 7:

A chronic alcoholic presents to the OPD with fever, shortness of breath, chest pain, and productive cough. On examination, there is bronchial breathing on the right side and a dull note on percussion. A chest x-ray is done. What is the most likely causative agent?

a) Streptococcus pneumoniae

b) Klebsiella pneumoniae

c) Pseudomonas aeruginosa

d) Pneumocystis jirovecii

Question 8:

A 3-month-old male child was brought with a history of fever and respiratory distress. Suspecting pneumonia, a chest X-ray was ordered. Which of the following is the most likely causative organism?

Sold by @itachibot

a) Klebsiella pneumoniae

b) Streptococcus pneumoniae

c) Mycoplasma pneumoniae

d) Staphylococcus aureus

Question 9:

Which of the following is the best biomarker for lobar pneumonia?

a) C-reactive protein

b) Procalcitonin

c) Fibrinogen

d) Serum amyloid A

Question 10:

All are components of CURB65 except

a) Confused state

b) Respiratory rate of > 30 /min

c) Blood urea nitrogen value greater than 7 mmol/L

d) Systolic blood pressure less than 100 mm Hg

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 11:

A 71-year-old man presents to the OPD with high-grade fever, productive cough. On examination, the respiratory rate was found to be 32/ min, heart rate of 102/ min, and blood pressure of 90/60 mmHg. How will you manage this patient?

a) Admit to ICU and treat with azithromycin and ceftriaxone

b) Admit to the ward and treat with levofloxacin and doxycycline

c) Treat him on OP basis with azithromycin and ceftriaxone

d) Admit to the ward and treat with azithromycin and levofloxacin

Question 12:

A patient in the ICU is diagnosed with community-acquired-MRSA. What is the best line of management?

a) Ceftriaxone + Azithromycin + Ampicillin

b) Ceftriaxone + Moxifloxacin + Linezolid + Clindamycin

c) Ceftriaxone + Carbenepem + Moxifloxacin

d) Ceftriaxone + Linezolid

Question 13:

An epileptic patient presented with fever, breathlessness, and productive cough for 2 days. The last episode of seizures occurred a week ago. On examination, PR-102/min, RR-30/min, and oral cavity revealed a gingival infection. Bronchial breath sounds were heard on auscultation on the right side. What is the drug of choice for his condition?

a) Clindamycin

b) Azithromycin

c) Moxifloxacin

d) Ceftriaxone

Question 14:

A 30-year-old HIV-positive man presents with fever for 3 weeks, dry cough, and significant weight loss. He complains of retrosternal chest pain that is worse on inspiration. His chest X-ray is given below. What is the most likely diagnosis?

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot



a) Staphylococcal pneumonia

b) Pneumococcal pneumonia

c) Tuberculosis

d) Pneumocystis jirovecii pneumonia

Question 15:

In an immunocompromised patient, which of the following is the drug of choice for prophylaxis of pneumocystis jirovecii?

a) Clotrimazole

b) Sulfadiazine-Pyrimethamine

c) Trimethoprim-Sulfamethoxazole

d) Sulfadoxine-Pyrimethamine

Question 16:

Which of the following is false regarding patients with Pneumocystis jirovecii pneumonia?

a) Glucocorticoids are indicated if Pao2 <70 mmHg

b) Recent oropharyngeal candidiasis is an indication for prophylaxis

c) Elevation of LDH is a common finding

d) Aerosolized pentamidine is used in the treatment

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 17:

A 36-year-old male HIV patient presents to the ED with a cough with bloody sputum. He also has night sweats, chills, and difficulty in breathing. A chest X-ray and examination of the sputum show the following findings. Which of the following is the drug of choice for his condition?





a) Amikacin

b) Cotrimoxazole

c) Minocycline

d) Linezolid

Question 18:

Which of the following drugs is commonly used for treating community-acquired pneumonia in OPD?

a) Vancomycin

b) Ceftriaxone

c) Azithromycin

d) Streptomycin

Question 19:

Sold by @itachibot

What is the most likely causative organism of ventilator-associated pneumonia among the following?

a) Clostridium

b) Klebsiella

c) Acinetobacter

d) Hemophilus

Question 20:

Which of the following statements is false regarding patients with parapneumonic effusion?

a) Diabetes is a risk factor

b) pH <7.2 may be a finding in pleural fluid analysis

c) Pleural fluid protein >4 gm/dL is an indication of tube thoracostomy

d) USG is helpful in characterizing the effusion

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	a
4	d
5	b
6	c
7	b
8	d
9	b
10	d
11	a
12	b
13	a
14	d
15	c

Sold by @itachibot

16	d
17	b
18	c
19	c
20	c

Detailed Explanations

Solution to Question 1:

Legionella pneumonia presents as both typical and atypical pneumonia.

Pneumonia is an infection of the lung parenchyma. Pneumonia that develops outside the hospital setting is commonly referred to as community-acquired pneumonia (CAP). There are two types of community-acquired pneumonia.

• Typical Pneumonia

• Atypical Pneumonia

Solution to Question 2:

Home split air-conditioning units do not use water to cool the air, so they are not a risk factor for legionella transmission.

The following are the risk factors for transmission of legionella:

• Contamination of water supplies in large facilities such as hospitals, hotels, apartment buildings

• Contaminated water sources such as hot tubs, birthing pools, fountains, whirlpools

• Water using machinery- cooling towers, evaporative condensers, humidifiers

Solution to Question 3:

The given clinical scenario is highly suggestive of atypical pneumonia caused by legionella.

Clinical features of Legionnaire's disease:

• High-grade-fever(>104 F)

• Cough with scanty sputum

• Extrapulmonary manifestations involve

• gastrointestinal- diarrhea, abdominal pain, nausea, vomiting

• neurological- most common-confusion, headache, encephalopathy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Relative bradycardia

• Occurrence of illness in an environment in which water supply is at risk of contamination

• Failure to respond to beta-lactams and aminoglycoside antibiotics

Investigations:

• On gram-staining of the sputum, numerous neutrophils with no organisms

• Hyponatremia

• Elevated liver enzymes

Solution to Question 4:

A urinary antigen detection test is recommended for the diagnosis of Legionella pneumonia. The sensitivity (70%) and specificity (99%) of the test are high.

The test can detect antigen even after the initiation of appropriate antibiotic therapy. Urinary antigen detection test can also be used for Streptococcus pneumonia but is not used due to the availability of better alternatives.

Other investigations for Legionella:

• DFA (direct fluorescent antibody)- is highly specific but not sensitive

• Culture- Gold standard, BCYE media is used

• DNA detection with PCR

Treatment: Drugs of choice are azithromycin and respiratory fluoroquinolones (levofloxacin, moxifloxacin).

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

The given clinical scenario along with chest X-ray and lab values is suggestive of atypical pneumonia caused by Mycoplasma pneumoniae. The blood investigations show decreased hemoglobin, elevated LDH, and elevated unconjugated bilirubin suggestive of hemolytic anemia.

Mycoplasma is the most common bacteria causing atypical pneumonia and the second most common cause of pneumonia.

Clinical Presentation:

• Pulmonary manifestations:

• URTI- tracheobronchitis/ pharyngitis

• Pneumonia(seen in only 10% patients)

• Extrapulmonary manifestations:

• Cold agglutinin disease(IgM mediated autoimmune hemolytic anemia)

• Erythema multiforme or Steven Johnson syndrome

Sold by @itachibot

• Arthralgia with adenopathy

• Guillain Barre Syndrome

• Myocarditis

• Pericarditis

• Encephalitis

• Bullous myringitis

Chest X-ray shows patchy consolidation at the hilum and fanning to the periphery.

Azithromycin is the drug of choice. Doxycycline is the second-best drug for the management of atypical pneumonia.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

Cefotaxime is not the drug of choice for the treatment of Mycoplasma pneumonia. Cefotaxime is a cell-wall synthesis inhibitor, hence it is not effective on M.pnemoniae as they lack a cell wall.

Investigations:

• PCR of respiratory secretions

• Chest X-ray shows-

• Patchy consolidation at the hilum and fanning to the periphery

• Diffuse peribronchial and perivascular interstitial infiltrates

• Airspace consolidation

• Reticulonodular opacification

• HRCT shows- areas of ground-glass attenuation

• On PPLO agar (Dienes method) shows- fried-egg appearance.

The diagnosis is made by serology/PCR. Azithromycin is the drug of choice for management. Doxycycline is the second-best drug for the management of atypical pneumonia.

Solution to Question 7:

The given clinical scenario and chest X-ray are suggestive of lobar pneumonia caused by Klebsiella pneumoniae.

Klebsiella pneumonia is associated mostly with alcoholics and shows upper lobe involvement. The patients present with red-currant jelly sputum.

Chest X-ray shows:

• Cavitation

• Pneumatocele (also seen in S. aureus)

Sold by @itachibot

• Bulging fissure sign: Classically seen in Klebsiella pneumonia. It represents expansive lobar consolidation causing fissural bulging; or displacement of copious amounts of inflammatory exudates within the affected parenchyma. It is seen in any space-occupying process in the lung such as pulmonary hemorrhage, lung abscess, and lung tumors.



Sold by
purchased this fr
you may have been scan
else,

Solution to Question 8:

The most likely causative agent of pneumonia in an infant with X-ray showing pneumatoceles (air-filled cysts within lung parenchyma) is Staphylococcus aureus.

In newborns and infants, S. aureus is a significant cause of severe respiratory tract infections, presenting with dyspnea, fever, and respiratory failure. Community-acquired infections are often preceded by viral infections, most commonly influenza. Among adults, nosocomial S. aureus pulmonary infections are frequently observed in intubated patients within intensive care units.

S. aureus often causes necrotizing pneumonitis that may be associated with complications such as pneumothorax and empyema.

Pneumatocoele can also be caused by Streptococcus pneumoniae, Haemophilus influenzae, E. coli, group A streptococci, Klebsiella pneumoniae, etc. However, Staphylococcus aureus is most commonly implicated.

Other options:

Option A: Klebsiella pneumoniae causes lobar pneumonia in alcoholics and diabetics. Patients present with fever, dyspnea, and a productive cough with red currant jelly sputum. Classic chest X-ray findings include upper lobe consolidation, classically on the right side, cavitation, and bulging fissure sign.

The following image shows bulging fissure sign:

Sold by @itachibot

Option B: Streptococcus pneumoniae causes lobar pneumonia. Patients with pneumococcal pneumonia present with fever, dyspnea, and productive cough. On examination, the involved lobes are dull to percussion, bronchial breath sounds, and crackles are heard on auscultation.

The following image shows right lower lobe consolidation:

Option C: Mycoplasma pneumoniae causes atypical pneumonia. Patients present with dry cough, fever, chills, malaise, and upper respiratory tract symptoms. On auscultation, wheezes or rales are heard. Chest X-ray shows peribronchial pneumonia with prominent bronchial markings, interstitial infiltration, and atelectasis.

The below chest X-ray shows patchy consolidation at the hilum and fanning to the periphery:

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

The best biomarker for lobar pneumonia is procalcitonin.

The most commonly used biomarkers of severe inflammation are C-reactive protein (CRP) and procalcitonin (PCT). Levels of these acute phase reactants increase in the presence of bacterial pathogens. PCT testing distinguishes between bacterial (mostly lobar pneumonia) and viral infection to decide the need for antibacterial therapy or discontinuation of treatment.

CRP, fibrinogen, and serum amyloid A are other acute phase reactants.

Solution to Question 10:

Among the options, systolic blood pressure less than 100 mm Hg (Option D) is not a component of CURB-65. CURB65 includes blood pressure, systolic ≤90 mmHg or diastolic ≤60 mmHg.

CURB-65 criteria is used to assess the severity of illness in a case community-acquired pneumonia. It includes confusion (C), urea >7 mmol/L (U), respiratory rate ≥30/min (R), blood pressure systolic ≤90 mmHg or diastolic ≤60 mmHg (B) and an age of ≥65 years.

Pneumonia severity index (PSI) is another index to assess the severity of illness and the risk of mortality in the case of pneumonia. It includes 20 variables and depending on the score, patients are assigned into one of the 5 classes.

Class, I, II, III - at low risk of death and can be treated on an outpatient basis

Class IV & V require hospitalization

Solution to Question 11:

Sold by @itachibot

The given clinical scenario is highly suggestive of lobar pneumonia and the patient belongs to a CURB-65 score of 3. The patient can be best managed by a combination of azithromycin and ceftriaxone in an ICU setting. After culture sensitivity, specific management is done.

Solution to Question 12:

The Empiric treatment of choice for ICU in patients with suspected CA-MRSA is:

• β-lactam (e.g., ceftriaxone) + Fluoroquinolone/Azithromycin

PLUS

• Linezolid / Vancomycin with or without Clindamycin to cover CA-MRSA

Solution to Question 13:

The given clinical scenario is highly suggestive of lobar pneumonia due to aspiration. The drug of choice is clindamycin as anaerobes are implicated.

Anaerobes play a significant role only when an episode of aspiration has occurred days to weeks before presentation for pneumonia. The combination of an unprotected airway (e.g., in patients with alcohol or drug overdose or a seizure disorder) and significant gingivitis constitutes the major risk factor.

Anaerobic pneumonia is often complicated by abscess formation and significant empyemas or parapneumonic effusions.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 14:

The most likely diagnosis in a case of pneumonia in a HIV positive patient, along with the presence of fever, and non-productive cough with radiographic findings of diffuse interstitial infiltrates and symmetric perihilar involvement is Pneumocystis jirovecii pneumonia (PCP). A characteristic feature of PCP is retrosternal chest pain, which worsens on inspiration and is described as sharp or burning.

It is important to note that even though the incidence of PCP is the highest at CD4+ counts of <200/µL (80%), it can be seen in patients with higher counts.

Other options:

Option D: Tuberculosis (TB) should be considered as a possibility in cases of pneumonia in people living with HIV. However, the absence of hilar lymph node involvement, non-productive cough, and no extrapulmonary (pleural, pericardial, lymphatic) features are not consistent with tuberculosis.

TB can occur at any stage of HIV infection. In the early stages, pulmonary TB appears typically with upper-lobe infiltrates and cavitation. In advanced HIV infection, with CD4+ T-cell count below 200/µL, TB presents with diffuse interstitial infiltrates, minimal cavitation, pleural effusion,

Sold by @itachibot

and lymphadenopathy. Extrapulmonary TB is fairly common in HIV, with it being documented in 40-60% of cases.



Option A: Staphylococcal pneumonia most commonly presents with a productive cough, and radiologically presents with pneumatoceles (cavitations)

Option B: Patients with pneumococcal pneumonia also present with fever, dyspnea, and productive cough. Imaging usually shows lobar involvement.

Sold by @itachibot



Solution to Question 15:

Trimethoprim-Sulfamethoxazole combination (commonly known as cotrimoxazole) is the first-line drug for both prophylaxis and treatment of Pneumocystis jirovecii infections in immunocompromised patients.

Cotrimoxazole acts by causing a sequential block of folate metabolism. CD4+ T cell counts below 200 cells/µL are an indication to start prophylaxis.

Solution to Question 16:

Aerosolised pentamidine is not used in the treatment of PCP. It is used in PCP prophylaxis.

Diagnosis of PCP:

• Histopathological staining of induced sputum examination

• Bronchoscopy with BAL (most sensitive)

• Elevated LDH

Treatment and Prophylaxis:

• Drug of choice- trimethoprim/sulfamethoxazole (TMP-SMX)

• IV pentamidine- treatment of choice for severe disease in the patient unable to tolerate TMP-SMX

• Glucocorticoids- indicated if PaO2 <70 mmHg or with an A–a gradient >35 mmHg

• Other drugs used- dapsone/pyrimethamine + leucovorin, clindamycin/primaquine, and atovaquone

• Aerosolised pentamidine is an alternative used in PCP prophylaxis.

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Sold by @itachibot

Prophylaxis is indicated when CD4+ T cell count <200/uL (CD4 cell %<15), recent oropharyngeal candidiasis, and prior PCP infection.

Solution to Question 17:

The given clinical scenario along with the chest X-ray and gram-staining is suggestive of Nocardiosis. The drug of choice is cotrimoxazole.

Nocardia is a gram-positive and acid-fast bacteria. Nocardia pneumonia has a sub-acute presentation with mostly cough with thick sputum and fever. Generally, hemoptysis is uncommon. Chest radiography shows nodular infiltrations involving bilateral lung fields.

Extrapulmonary manifestations are common, the brain is the most commonly involved organ. It is associated with dense bilateral infiltrates with cavitating nodules and subacute brain abscesses.

The drug of choice is cotrimoxazole. Severe disease is treated with a combination of cotrimoxazole, amikacin, and ceftriaxone, or imipenem.

Note: Mnemonic SNAP:

Sulfamethoxazole-trimethoprim (Cotrimoxazole) is the drug of choice for Nocardia.

For Actinomyces, Penicillins are the drugs of choice.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 18:

Azithromycin is the drug of choice for treatment of community-acquired pneumonia (CAP) in an OPD setup.

Outpatient management of CAP:

1. For previously healthy patients who have not taken antibiotics within the past 3 months:

• A macrolide – Azithromycin, 500 mg orally as a first dose and then 250 mg orally daily for 4 days, or 500 mg orally daily for 3 days, or

• Clarithromycin, 500 mg orally twice a day; or

• Doxycycline 100 mg orally twice a day

2. For patients with such comorbid medical conditions as chronic heart, lung, liver, or kidney disease, diabetes mellitus, alcoholism, malignancy, asplenia, immunosuppressant conditions or use of immunosuppressive drugs or use of antibiotics within the previous 3 months:

• A respiratory fluoroquinolone (moxifloxacin, 400 mg orally daily; gemifloxacin, 320 mg orally daily; levofloxacin, 750 mg orally daily) or

• A macrolide (as above) plus a beta-lactam (amoxicillin, 1 g orally three times a day; amoxicillin-clavulanate, 2 g orally twice a day; cefuroxime, 500 mg orally twice a day).

3. In regions with a high rate (> 25%) of infection with high level (MIC ≥ 16 mcg/mL) macrolide-resistant Streptococcus pneumoniae, consider the use of alternative agents listed in (2) for patients with comorbidities.

Sold by @itachibot

Other options: Vancomycin, Ceftriaxone, and Streptomycin are administered parenterally, and hence, not appropriate for outpatient management.

Solution to Question 19:

Among the given options, Acinetobacter species is the most commonly isolated organism in cases of ventilator-associated pneumonia (VAP). VAP develops in a mechanically ventilated patient >48 hours after endotracheal intubation.

The organisms causing ventilator-associated pneumonia (VAP) significantly vary from one hospital to another. However, the organisms that frequently cause VAP in hospital settings are Staphylococcus aureus, Pseudomonas aeruginosa, Acinetobacter sp., Stenotrophomonas maltophilia, and gram-negative rods (Enterobacter species, K. pneumoniae, and Escherichia coli).

Three factors distinguish nosocomial pneumonia from community-acquired pneumonia:

• Different infectious agent

• Different antibiotic susceptibility patterns, specifically, a higher incidence of drug resistance

• Poorer underlying health status of patients putting them at risk for more severe infections

Within 48 hours of admission, 75% of seriously ill hospitalized patients have their upper airway colonized with organisms from the hospital environment. Impaired cellular and mechanical defense mechanisms in the lungs of hospitalized patients raise the risk of infection after aspiration has occurred.

VAP patients may be infected with Acinetobacter species and Stenotrophomonas maltophilia. Anaerobic organisms (Bacteroides, anaerobic streptococci, Fusobacterium) may also cause pneumonia in the hospitalized patient.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 20:

Pleural Fluid Protein >4 gm/dL is not an indication of tube thoracostomy.

Parapneumonic effusion is pleural effusion associated with bacterial pneumonia, lung abscess, or bronchiectasis. When it worsens, it can result in the collection of frank pus in the pleural space. This is known as empyema.

Risk factors include diabetes, immunosuppression, alcoholism, cancer, and poor dental hygiene.

Investigations:

• Chest X-ray:

• Loculations or encystments (complicated parapneumonic effusion)

• Intrapleural air-fluid level (bronchopulmonary fistula)

• Parenchymal infiltrations (underlying pneumonia)

• Ultrasound: Shows echogenic pleural fluid with septations and helps in characterizing the effusion

Sold by @itachibot

• Pleural fluid analysis:

• Exudative neutrophilic effusion

• Low pH <7.2, low glucose <60 mg/dL, high LDH >1000 IU/L in complicated parapneumonic effusion

• CECT: Split pleura sign seen in empyema

Treatment:

• Antibiotics: Tailored according to the microorganism

• Therapeutic thoracocentesis

• Tube thoracostomy is indicated if:

• Pleural fluid pH <7.2

• Frank pus

• Microorganism isolated in Gram stain or culture of pleural fluid

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Interstitial Lung Diseases and pneumoconiosis

Question 1:

Which of the following is the most common interstitial lung disease?

a) Non-specific interstitial pneumonia

b) Desquamative interstitial pneumonia

c) Idiopathic pulmonary fibrosis

d) Respiratory bronchiolitis associated lung disease

Question 2:

An 80-year-old woman was found to have a restrictive disease with reduced lung volumes on a pulmonary function test. A high-resolution CT of the thorax was done and is shown below. Which of the following is not a feature of this condition?



a) Clubbing of the digits

b) Inspiratory coarse crackles

c) Breathlessness on exertion

d) Non-productive cough

Question 3:

Sold by @itachibot

Sold by @itachibot

A 70-year-old woman presents with breathlessness on exertion and dry cough for 4 weeks. She has swollen joints and early morning stiffness for which she is on treatment for a few years. On auscultation, inspiratory fine crackles were heard. An HRCT was ordered and it showed bilateral, subpleural and basilar honeycombing pattern. PFT of the patient is likely to show which of the following patterns?

a) FEV1/FVC ratio <0.7 and DLCO increases

b) FEV1/FVC ratio <0.7 and DLCO decreases

c) FEV1/FVC ratio >0.7 and DLCO decreases

d) FEV1/FVC ratio >0.7 and DLCO increases

Question 4:

Which of the following is not a characteristic HRCT finding of usual interstitial pneumonia (UIP)?

a) Honeycombing and loss of lung architecture

b) Ground-glass opacities

c) Traction bronchiectasis

d) Bibasilar subpleural reticular opacities

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 5:

Which of the following conditions is also known as Hamman-Rich syndrome?

a) Usual interstitial pneumonia

b) Non-specific interstitial pneumonia

c) Acute interstitial pneumonia

d) Cryptogenic organising pneumonia

Question 6:

A 52-year-old woman presented with dyspnoea. General physical examination revealed deformity of the hand as shown below. There is restricted mobility in the small joints and loss of wrinkle lines on the facial skin. An HRCT of the thorax showed diffuse subpleural, symmetric, ground-glass opacities of the lungs. Which of the following histopathological finding is likely to be seen in the lung biopsy of this patient?

Sold by @itachibot

a) Homogenous interstitial involvement

b) Temporal heterogeneity of lesions

c) Spatial heterogeneity of lesions

d) Honeycombing pattern

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 7:

Which of the following is not a smoking-related interstitial lung disease (ILD)?

a) Desquamative interstitial pneumonia

b) Respiratory bronchiolitis associated ILD

c) Pulmonary langerhans cell histiocytosis

d) Pulmonary lymphangioleiomyomatosis

Question 8:

A man presented with progressively worsening cough with sputum, breathlessness, easy fatiguability, and loss of appetite for 6 months. PFT was suggestive of a restrictive disorder. Trans-bronchial biopsy showed cells positive for CD1a and langerin (CD207). Which of the following is not a feature of this condition?

a) They may present with spontaneous pneumothorax

b) Cysts and nodules are seen on HRCT

c) It is a smoking related interstitial lung disease

Sold by @itachibot

d) It typically affects elderly smokers

Question 9:

Which of the following statements is false regarding pulmonary alveolar proteinosis?

a) Acquired due to antibodies formed against GM-CSF

b) Copious, watery PAS-negative sputum

c) X-ray shows bilateral diffuse opacities

d) Treatment is whole lung lavage

Question 10:

Farmer's lung is associated with which of the following organisms?

a) Histoplasmosis

b) Coccidiomycosis

c) Thermophilic actinomycetes

d) Cryptococcosis

Question 11:

A daily-wage worker from the cement industry got a chest X-ray as a part of his routine health check-up. What could be the cause of the findings shown below?



If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

a) Berylliosis

b) Anthracosis

c) Asbestosis

d) Silicosis

Question 12:

A woman has been working for a company that manufactures roofing sheets for 12 years, but the company does not provide her with personal protective equipment. Which of the following cancers is commonly seen in such individuals?

a) Pleural mesothelioma

b) Peritoneal mesothelioma

c) Adenocarcinoma of lung

d) Laryngeal carcinoma

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 13:

Which among the following is a common concern for a patient presenting with chronic dry cough, working in the granite mining industry?

a) Asbestosis

b) Silicosis

c) Farmer's lung

d) Anthracosis

Question 14:

A 45-year-old man comes to the emergency room with an acute onset of dyspnoea. He tells you that he has been working in the sandblasting industry for about a year and the company does not provide personal protective equipment. He also gives a history of on and off dry cough. CT scan shows the following. What is the most likely diagnosis?

Sold by @itachibot



a) Complicated silicosis

b) Acute silicosis

c) Simple silicosis

d) Pulmonary Alveolar Proteinosis

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 15:

A 30-year-old man has recently started work in the stone cutting industry. The company provides him with a respirator mask but they do not use dust control measures. If he works in similar conditions for 20 more years, which of the following will you most likely see in his HRCT scan image?

a) A typical crazy pavement pattern

b) Small round opacities in upper lobes

c) Thin-walled cysts surrounded by normal lung parenchyma

d) Thick calcified pleural plaques

Question 16:

Which of the following pneumoconiosis is associated with an increased incidence of tuberculosis?

a) Asbestosis

b) Silicosis

c) Coal worker's pneumoconiosis

Sold by @itachibot

d) Farmer's lung

Question 17:

Choose the correct option with respect to Caplan's syndrome.

a) Rheumatoid arthritis + pneumoconiosis

b) Rheumatoid arthritis + anterior uveitis

c) Hilar adenopathy + polyarthritis + erythema nodosum

d) Arthritis + urethritis + conjunctivitis

Question 18:

A factory worker presents to the clinic with fever, tiredness, and a chronic dry cough. He says his symptoms are worst on the first day of the week, improve by the end of the week, and are almost non-existent over the weekends. What could be the most likely diagnosis?

a) Bagassosis

b) Byssinosis

c) Berylliosis

d) Bornholm's disease

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

Which interstitial lung disease is likely to be seen in a patient suffering from systemic sclerosis?

a) Respiratory bronchiolitis associated ILD

b) Non-specific interstitial pneumonia

c) Acute interstitial pneumonia

d) Bronchiolitis obliterans organising pneumonia

Question 20:

A 45-year-old man who has worked as a miner for the past 25 years, comes to the OPD with complaints of dyspnoea and cough. Pulmonary function tests and a lung biopsy are done. The histopathology of the sample is shown below. Which type of emphysema can this condition

Sold by @itachibot

give rise to?



a) Panacinar emphysema

b) Centrilobular emphysema

c) Paraseptal emphysema

d) Irregular emphysema

Sold by @itachibot If you purchased this from somewhere else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	b
3	c
4	b
5	c
6	a
7	d
8	d
9	b
10	c
11	c
12	c

Sold by @itachibot

13	b
14	b
15	b
16	b
17	a
18	b
19	b
20	b

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Detailed Explanations

Solution to Question 1:

The most common interstitial lung disease (ILD) is idiopathic pulmonary fibrosis (IPF).

ILD refers to the remodeling of interstitial space, the region between the epithelium and endothelium. The main pathology is fibrosis, mediated by TGF-ß and platelet-derived growth factor.

Classification of Interstitial Lung Disease (ILD)
ILDs of known etiologyOccupational: Coal worker's pneumoconiosis, silicosisTreatment-related: Radiation, drug-inducedConnective tissue disease-related: Rheumatoid arthritis, Sjogren's syndrome, scleroderma
ILDs of unknown etiologyIdiopathic pulmonary fibrosis (IPF):Most commontypeNon-specific interstitial pneumonia (NSIP)Cryptogenic organizing pneumonia (COP)Acute interstitial pneumonia (AIP)/Hamman-Rich syndromeRespiratory bronchiolitis associated ILD (RBILD)Desquamative interstitial pneumonia (DIP)Lymphocytic interstitial pneumonia (LIP)Pleuroparenchymal fibroelastosis
GranulomatousSarcoidosisHypersensitivity pneumonitis

Sold by @itachibot

Classification of Interstitial Lung Disease (ILD)
Other formsLymphangioleiomyomatosis (LAM)Pulmonary Langerhans' cell histiocytosis (PLCH)Pulmonary alveolar proteinosis (PAP)

Solution to Question 2:

The above HRCT shows a honeycomb appearance of lung parenchyma due to interstitial fibrosis in a restrictive lung disorder that is suggestive of interstitial lung disease (ILD). Inspiratory coarse crackles are not a feature of interstitial lung disease.

Auscultating a patient with ILD will reveal bilateral basal end-inspiratory fine crackles also called velcro crackles or rales.

Patients with interstitial lung disease usually present with a non-productive cough and breathlessness on exertion. Clubbing may be seen on examination, especially in advanced cases.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

The given clinical scenario along with the HRCT findings are suggestive of interstitial lung disease (ILD). FEV1/FVC ratio >0.7 and decreased DLCO are likely to be seen in the PFT. A history of swollen joints and early morning stiffness hints at rheumatoid arthritis. It is associated with the usual interstitial pneumonia (UIP) pattern of ILD.

In ILD, restrictive pattern is seen on pulmonary function test, which is characterized by:

- Decreased total lung capacity (TLC)
- Decreased forced vital capacity (FVC)
- Decreased residual volumes (RV)
- Decreased DLCO (diffusing capacity of the lungs for carbon monoxide)
- FEV1/FVC ratio - normal or increased, >0.7

Solution to Question 4:

Ground-glass opacities on HRCT are not a characteristic finding in UIP.

Honeycombing, loss of lung architecture, traction bronchiectasis, and basilar subpleural reticular opacities are seen in UIP pattern of ILD on HRCT.

Sold by @itachibot

Solution to Question 5:

Acute interstitial pneumonia (AIP) is also known as Hamman–Rich syndrome.

AIP is a rare, idiopathic, fulminant interstitial lung disease (ILD) that is similar in presentation to ARDS. It is often preceded by a viral upper respiratory tract infection.

Histologically, it shows diffuse alveolar damage. HRCT shows bilateral patchy, ground-glass opacities. Lung biopsy shows diffuse alveolar damage with hyaline formation.

Solution to Question 6:

The given clinical scenario and the image showing sclerodactyly suggestive of scleroderma along with the HRCT findings point to a diagnosis of non-specific interstitial pneumonia (NSIP). It is characterized by homogeneous interstitial involvement.

NSIP is often associated with connective tissue disease but can also be drug-induced. The common drugs implicated in NSIP are:

- Bleomycin
- Busulfan
- Methotrexate
- Amiodarone

It usually affects middle-aged woman who have never smoked. There is uniformity of interstitial involvement across the biopsy section, and this may be predominantly cellular or fibrosing. HRCT shows homogenous involvement with symmetrical ground-glass opacities.

Non-specific interstitial pneumonia (NSIP) responds to steroids. Other treatment modalities include cytotoxic drugs like azathioprine or cyclophosphamide and also biologics like rituximab.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

Pulmonary lymphangioleiomyomatosis (LAM) is not related to smoking.

It is usually seen in young females with tuberous sclerosis. LAM can lead to pneumothorax or chylous effusions.

Pulmonary function testing usually reveals an obstructive or mixed obstructive-restrictive pattern. HRCT shows thin-walled cysts surrounded by normal lung parenchyma.

The smoking-related ILDs are:

- DIP - Desquamating interstitial pneumonia.
- RB-ILD - Respiratory bronchiolitis associated ILD.
- LCH - Langerhans cell histiocytosis.

Sold by @itachibot

Solution to Question 8:

The given scenario is suggestive of pulmonary Langerhans cell histiocytosis (LCH). It is a smoking-related ILD that typically affects young adults. Smoking cessation leads to improvement in the condition.

The specific immunohistochemical markers for LCH are CD1a and langerin (CD207). On electron microscopy, the presence of Birbeck granules in the cytoplasm is characteristic (as shown below).



HRCT shows bilateral irregularly shaped cysts with nodules, predominantly in the upper lobe. There is a risk of spontaneous pneumothorax in this condition due to cyst rupture.

Solution to Question 9:

Chunky, gelatinous, PAS-positive sputum is a feature of pulmonary alveolar proteinosis (PAP).

PAP is a condition characterized by the accumulation of surfactants in alveoli and terminal airways. It can be due to impaired clearance or abnormal production of surfactants.

PAP can be of three forms:

• Primary (most common): There is impaired GM-CSF-dependent surfactant clearance by the alveolar macrophages. It can be due to:

• Autoantibodies against GM-CSF (acquired/idiopathic/autoimmune PAP)

• Mutation in GM-CSF receptors (hereditary PAP)

• Secondary: There is a reduction in the number or ability of alveolar macrophages that causes impaired surfactant clearance. It can be due to:

• Silicosis

• Immunodeficiency disorders

• Hematopoietic disorders and malignancies

If you purchased this book from anywhere else, you may have been sold.

Sold by @itachibot

• Congenital: It affects neonates and children. There is abnormal surfactant production due to mutation in the genes coding for:

• Surfactant protein B

• Surfactant protein C

• ATP binding cassette protein member 3

About 30% of patients are asymptomatic. Symptomatic patients can have persistent dry cough, progressive dyspnea, and pleuritic chest pain

X-ray shows bilateral, diffuse opacities and HRCT shows a crazy-paving pattern (as shown below). The treatment is whole lung lavage.



Solution to Question 10:

Farmer's lung is a hypersensitivity pneumonitis caused due to exposure to mouldy hay, which contains thermophilic actinomycetes, principally Thermopolyspora polyspora.

Solution to Question 11:

The given X-ray shows irregular opacities, which appear to be pleural thickenings i.e., pleural plaques. The sign along with a history of working in the cement industry suggests a diagnosis of asbestosis.

Workers of the cement, shipping, and textile industries are at a higher risk of developing asbestosis. Asbestos tends to involve the pleura first.

The earliest manifestation is benign asbestosis-related pleural effusion. With time, the plaques calcify resulting in the formation of thick pleural plaques. The pleural thickenings can be seen on the X-ray. This is the "holly leaf sign".

Sold by @itachibot

Lung parenchymal involvement occurs after years of exposure (at least 10 years). They present with cough and exertional dyspnoea. It predominantly affects the lower lobes of the lung, in contrast to other forms of pneumoconiosis, which affect the upper lobes.

Pulmonary function testing in asbestosis typically shows a restrictive pattern with a decrease in both lung volumes and diffusing capacity.

The image below shows thick plueral plaques.



If you purchased this from elsewhere, you may have been

Solution to Question 12:

Workers in the roofing industry are exposed to asbestos. Lung cancer (of all histologic types) is the most common cancer associated with asbestos exposure.

Smoking along with asbestos exposure multiplies the risk to a great extent. Cough and hemoptysis are common presentations.

Pleural mesothelioma is the most specific cancer associated with asbestos exposure, but not the most common. Smoking does not seem to be associated with mesothelioma. Pleuritic chest pain or a mass on the radiographic image are common presentations.

Solution to Question 13:

Silicosis is a common concern for people working in granite mining industries.

Silica is found in rock, granite, slate, and sandstone. People working in mining, stone cutting, sandblasting, glass and cement manufacturing, and quarrying are exposed to silica. The active molecule that causes silicosis is free silica or crystalline silica (quartz).

Silicosis can present as:

• Acute silicosis - due to heavy exposure in closed spaces and they manifest within 2 years.

Sold by @itachibot

• Chronic silicosis (simple silicosis) - due to long-term (10 to 30 years), less intense exposure.

• Complicated silicosis- simple silicosis may progress to complicated silicosis that presents with chronic cough and progressive dyspnoea on exertion.

Note: Silicosis is the most common pnemoconiosis in the world.

Solution to Question 14:

A history of working in the sandblasting industry for about a year without PPE suggests a heavy exposure to silica. This along with the CT finding of bilateral nodules, interstitial fibrosis, and crazy pavement pattern suggests a diagnosis of acute silicosis.

Acute silicosis is seen rarely. It manifests within 2 years of heavy exposure. Common presentations are dyspnoea and hypoxemia.

The X-ray may show diffuse miliary infiltrates or consolidation. HRCT may also show bilateral centrilobular nodules and ground-glass opacification. Bronchoalveolar lavage yields a PAS-positive milky white fluid.

Solution to Question 15:

In this scenario, the man works in the stone cutting industry, indicating exposure to silica. He is expected to have less intense, long term exposure owing to a few but substandard measures to control exposure to silica. Such individuals are more likely to develop simple silicosis. These patients are asymptomatic, and HRCT shows small round opacities in the upper lobes.

This can progress to complicated silicosis. Here, the nodules coalesce and form masses leading to impairment of lung function. They present with chronic cough and progressive dyspnoea on exertion. This may progress further to cause progressive massive fibrosis.

PFT shows a mixed obstructive and restrictive pattern. Calcification of hilar lymph nodes is seen in a few cases, called eggshell calcification. This is depicted in the image below.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option A: A crazy pavement pattern is seen with acute silicosis.

Option C: Thin-walled cysts surrounded by normal lung parenchyma are seen in pulmonary lymphangiomyomatosis (LAM).

Option D: Pleural plaques are seen in patients suffering from asbestosis.

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

Silicosis is associated with an increased incidence of tuberculosis.

Silica causes pulmonary alveolar macrophage dysfunction and hence, patients with silicosis have a higher risk of developing lung infections from organisms like M. Tuberculosis, atypical mycobacteria and fungi.

Radiographic imaging of silicotuberculosis shows cavitation.

Solution to Question 17:

Caplan's syndrome also called rheumatoid pneumoconiosis, is a combination of rheumatoid arthritis and pneumoconiosis.

It manifests as pneumoconiotic nodules along with features of seropositive rheumatoid arthritis. Caplan syndrome was first described in coal miners and later in patients with silicosis too.

Option C: Hilar adenopathy, polyarthritis, and erythema nodosum are characteristic of Lofgren syndrome.

Option D: Arthritis, urethritis, and conjunctivitis are characteristic of Reiter syndrome.

Sold by @itachibot

Solution to Question 18:

The above characteristic pattern of symptoms is called Monday fever or Monday chest tightness and is due to byssinosis.

Byssinosis is seen commonly in cotton factory workers due to cotton dust exposure. It is also caused due to flax, hemp, and jute dust.

It is due to occupational hypersensitivity reaction with symptoms occurring (or recurring) on the first day of the workweek and diminishing by the weekend due to development of tolerance.

Solution to Question 19:

Non-specific interstitial pneumonia (NSIP) pattern of ILD is the most common pulmonary manifestation in systemic sclerosis.

The CT scan of NSIP shows ground-glass opacities. It responds to steroids given along with azathioprine or cyclophosphamide.

Other connective tissue diseases associated with ILD:

- Rheumatoid arthritis
- Sjögren syndrome
- Dermatopolymyositis
- Systemic lupus erythematosus (SLE).

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 20:

The given clinical scenario and the microscopic image showing abundant black-colored pigment within the macrophages of lung parenchyma is characteristic of pulmonary anthracosis/coal worker's pneumoconiosis. It may give rise to centrilobular emphysema.

The inhaled carbon particles are engulfed by the lung macrophages. They accumulate in the connective tissue along lymphatics or in lymphoid tissue along the bronchi or the hilum, which causes the blackening of the tissues. Aggregates of carbon dust are present as coal macules or nodules which consist of carbon-laden macrophages.

It is commonly seen in people who work in environments where there is high air pollution with carbon particles, such as coal mines. It is usually benign and less than 10% of the cases develop progressive massive fibrosis which may lead to pulmonary dysfunction, pulmonary hypertension, and cor pulmonale. Coal workers may also develop emphysema or chronic bronchitis, independent of smoking.

Unlike asbestosis, this condition does not predispose to cancer.

Sold by @itachibot

Bronchiectasis and Lung Abscess

Question 1:

An old woman presents with a persistent productive cough and clubbing. On auscultation, wheeze and crackles are present. There is no history of fever. Which of the following is the investigation of choice to confirm the diagnosis?

a) HRCT

b) CECT

c) Helical CT

d) Spiral CT

If you purchased this from someone else, you have been scammed. Sold by @itachibot

Question 2:

A 30-year-old woman is admitted to the hospital with a fever and cough. She has a history of repeated chest infections for several years and coughs up yellow-green purulent sputum with some streaky hemoptysis. What could be the underlying condition based on the CT given below?



a) Cystic fibrosis

b) Tuberculosis

c) Amyloidosis

d) Lung abscess

Sold by @itachibot

Question 3:

Which of the following is an unlikely complication in patients with bronchiectasis?

a) Cor pulmonale

b) Amyloidosis

c) Lung cancer

d) Empyema

Question 4:

A 58-year-old white man presents with a 6-month history of productive cough, weight loss, fever, and night sweats. HRCT shows a cluster of grapes appearance. A biopsy shows features of a non-tuberculous mycobacterial infection. Cystic fibrosis is ruled out. Which of the following drugs is not to be used in the treatment regimen?

a) Macrolide

b) Ethambutol

c) Bronchodilators

d) Mucolytic dornase

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 5:

A college student presents with complaints of fever, chronic productive cough, steatorrhea. There is a past history of recurrent lung infections and multiple hospitalizations with similar complaints. Which of the following is false about this condition?

a) An autosomal-recessive disorder

b) Abnormality in CFTR gene leads to defective calcium transport

c) Most common mutation is F508 deletion

d) Males have infertility in adult life

Question 6:

A 17-year-old boy presents to the clinic with an acute respiratory tract infection. He has a history of asthma associated with recurrent respiratory tract infections and sinusitis. Genetic testing in childhood had revealed CFTR mutations. Which of the following is likely to be

Sold by @itachibot

decreased in this patient?

a) Sweat chloride

b) Serum bicarbonate level

c) Sperm count

d) Immunoreactive trypsinogen

Question 7:

A young boy is being evaluated for cystic fibrosis. Sweat chloride shows values of 38 and 44 mEq/L (normal<40). What is the next best test to confirm the diagnosis of cystic fibrosis?

a) 72-hour fecal fat

b) Immunoreactive trypsinogen levels

c) DNA analysis

d) Trans epithelial nasal potential difference

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

A 15-year-old boy has been having recurrent hospitalizations for infections, since 2 years of age. After a detailed evaluation, a diagnosis of Kartagener's syndrome was made. Which of the following features is not a component of the triad seen in this condition?

a) Situs inversus totalis

b) Chronic sinusitis

c) Male infertility

d) Bronchiectasis

Question 9:

Infertility in Kartagener’s syndrome is due to which of the following?

a) Oligospermia

b) Asthenospermia

c) Undescended testes

d) Epididymis obstruction

Sold by @itachibot

Question 10:

Which is the most common cause of lung abscess?

a) Tuberculosis

b) Pneumonia

c) Aspirated oropharyngeal secretion

d) Sepsis

Question 11:

A bed-ridden patient with a Ryles tube tried to pull out the tube. This led to aspiration following which he developed fever with chills and cough. X-ray findings were suggestive of lung abscess. Which of the following pathogens is unlikely to have caused it?

a) Bacteroides

b) Streptococcus milleri

c) Microaerophilic streptococci

d) Staphylococcus aureus

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Question 12:

A 64-year-old man was admitted to the emergency department after being found unconscious due to alcohol intoxication. On physical examination, his dental hygiene was poor. Chest auscultation revealed fixed amphoric breath sounds over the right hemithorax. Which of the following lung lobes is most likely to be affected?

a) Posterior upper lobe

b) Anterior upper lobe

c) Apical upper lobe

d) Posterior lower lobe

Question 13:

A middle-aged man presents with persistent fever and productive cough for 6 weeks with night sweats and anemia. The following is his CXR. What is the likely diagnosis?

Sold by @itachibot



a) Carcinoma Lung

b) Tuberculosis

c) Emphysematous bullae

d) Lung abscess

Sold by @itachi. If you purchased this from someone else, you may have been scammed.

Question 14:

A 60-year-old male patient presented with 2 weeks of productive cough, fever, and shortness of breath. A chest radiogram was done as shown below. Which of the following is the most appropriate therapy for him?



a) Cephalexin

Sold by @itachibot

b) Ciprofloxacin

c) Clindamycin

d) Vancomycin

Question 15:

A patient with fever, productive cough, night sweats, and clubbing is diagnosed with lung abscess. Which of the following is an unlikely complication of this condition?

a) Bronchiectasis

b) Brain abscess

c) Lung carcinoma

d) Hemoptysis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

What is the most common presentation of inhalational anthrax?

a) Atypical pneumonia

b) Hemorrhagic mediastinitis

c) Lung abscess

d) Broncho pulmonary pneumonia

Question 17:

Which of the following is not a common feature of allergic bronchopulmonary aspergillosis?

a) Distal bronchiectasis

b) Cough

c) Wheezing

d) Raised serum IgE levels

Question 18:

Which of the following tests is not used to diagnose a patient with suspected allergic bronchopulmonary aspergillosis?

Sold by @itachibot

a) Peripheral eosinophilia

b) Serum IgE > 1000 IU/ml

c) Positive skin prick test

d) Culture of Aspergillus fumigatus from sputum

Question 19:

A patient with acute myeloid leukemia underwent induction therapy. Within the next few days, he developed a cough and fever. On evaluation, HRCT showed a halo sign. Eventually, the patient deteriorated and died due to respiratory compromise. On autopsy, lung histopathology showed infarction and invasion of blood vessels by Aspergillus hyphae. Which of the following is not true about this condition?

a) Fungal hyphae are broad and aseptate

b) Yeast forms are not seen in infected tissue

c) Most of the patients are asymptomatic

d) Exposure to hay barns is a risk factor.

If you purchased this from someone else, Sold by @itachibot
they have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	d
5	b
6	c
7	c
8	c
9	b
10	c
11	d
12	a
13	d
14	c

Sold by @itachibot

15	c
16	b
17	a
18	d
19	a

Detailed Explanations

Solution to Question 1:

The clinical scenario is suggestive of bronchiectasis. HRCT is the investigation of choice - most specific.

Bronchiectasis is a suppurative lung disease with abnormal and irreversible dilatation of the bronchi. It is an end-stage condition of various diseases (infections, congenital or autoimmune disorders) that result in the destruction of the bronchial wall and its surrounding supporting tissues.

Ried's classification divides bronchiectasis into 3 types based on the phenotype of the affected bronchi:

• Cylindrical or tubular (most common)

• Saccular or varicose

• Cystic

Traction bronchiectasis is a type of bronchiectasis seen in conditions causing extensive pulmonary fibrosis (tuberculosis, idiopathic pulmonary fibrosis). The mechanical traction due to fibrosis pulls apart the adjacent bronchi, making them distorted and abnormally dilated. There is no destruction of the bronchial wall.

Clinically, they have a chronic productive cough, clubbing, crackles, and rhonchi.

Solution to Question 2:

The given clinical scenario and the CT showing signet ring sign is suggestive of bronchiectasis which can be due to cystic fibrosis.

A history of repeated chest infections is an indication of cystic fibrosis and a major risk factor for the development of bronchiectasis.

Solution to Question 3:

Lung cancer is not a known complication of bronchiectasis.

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

Complications of bronchiectasis include:

• Massive hemoptysis

• Empyema and lung abscess

• Respiratory failure

• Pulmonary hypertension and cor pulmonale

• Secondary amyloidosis (AA type)

Chronic and recurrent infections lead to secondary amyloidosis (AA type) and involve the kidneys.

Solution to Question 4:

The clinical scenario is suggestive of bronchiectasis secondary to a non-tuberculous mycobacterial (NTM) infection. Mucolytic dornase is recommended routinely in cystic fibrosis (CF)-related bronchiectasis but not in non-CF bronchiectasis.

Macrolides (azithromycin or clarithromycin) combined with rifampicin and ethambutol are used in the treatment of non-CF bronchiectasis.

Treatment of bronchiectasis:

• Antibiotic therapy: This is used to treat exacerbations and is given for 10-14 days. Commonly administered antibiotics include:

• Amoxicillin or clarithromycin: First-line drugs

• Ciprofloxacin: For Pseudomonas aeruginosa

• Airway clearance:

• Postural drainage: This is done by gravity-assisted positioning of the patient and chest physiotherapy.

• Mucolytic treatment: only in cystic fibrosis patients

• Normal or hypertonic saline (≥3%) nebulization

• Humidified air inhalation

• Drugs like N-acetylcysteine or carbocysteine

• Others:

• Inhaled bronchodilators like long-acting β2 agonists and anticholinergics

• Long-term oxygen therapy: In patients with chronic respiratory failure

• Pulmonary rehabilitation

• Adequate hydration

• Resection of a focal area of suppuration, lung transplantation: In refractory cases

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 5:

The given clinical scenario is suggestive of cystic fibrosis. Abnormalities in cystic fibrosis transmembrane conductance regulator protein (CFTR) gene lead to defective chloride transport and not calcium transport.

Cystic fibrosis is an autosomal recessive disorder affecting the epithelial cells of various exocrine glands of the body. It is caused by mutations in the gene coding for CFTR protein. The most common CFTR mutation is F508 deletion.

The most prominent clinical features of cystic fibrosis are chronic respiratory diseases (fever, productive cough, recurrent lung infections) and pancreatic insufficiency (steatorrhea).

Male patients with cystic fibrosis usually have infertility due to congenital absence of bilateral vas deferens and defective sperm transport, resulting in azoospermia.

Solution to Question 6:

The given clinical scenario is suggestive of cystic fibrosis. In these patients, sperm count is most likely to be decreased.

Male patients with cystic fibrosis usually have infertility due to congenital absence of bilateral vas deferens and defective sperm transport, resulting in obstructive azoospermia.

Option A: Sweat chloride levels are elevated (≥60 mEq/L) due to abnormalities in the CFTR protein leading to defective chloride transport.

Option B: Serum bicarbonate levels are increased due to excessive loss of salt in the sweat resulting in hypochloremic alkalosis.

Option D: Immunoreactive trypsinogen levels are increased due to exocrine pancreatic insufficiency. Measurement of these levels is a diagnostic screening test in newborns.

Solution to Question 7:

DNA analysis is used to confirm the diagnosis of cystic fibrosis in patients with equivocal (one normal and one abnormal value) or frankly normal sweat chloride values.

The diagnosis is confirmed if both of the following criteria are met:

• Presence of

• Clinical symptoms affecting at least one organ system (respiratory, gastrointestinal, or genitourinary) or

• Positive newborn screening or

• A sibling with cystic fibrosis

• Evidence of dysfunction of the CFTR gene by

• Two positive sweat chloride test results obtained on separate days or

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Sold by @itachibot

• Identification of two CFTR mutations by genetic testing or

• An abnormal measurement of nasal potential difference

Option A: 72-hour fecal fat is not used in the diagnosis of CF.

Option B: Measuring the levels of immunoreactive trypsinogen is a screening test in newborns.

Option D: Trans epithelial nasal potential difference measurement can be used as an adjunct to sweat test, but it has limited availability and is predominantly used as a research tool.

Solution to Question 8:

Male infertility is typical in Kartagener syndrome, but not a component of the triad.

Kartagener's syndrome is primary ciliary dyskinesia characterized by the classical triad of:

• Situs inversus

• Chronic sinusitis

• Bronchiectasis

Primary ciliary dyskinesia (also called immotile cilia syndrome) is an autosomal recessive disorder affecting ciliary motility. It occurs due to ultrastructural defects in the dynein arm of cilia. This causes the retention of secretions due to a failed mucociliary clearance mechanism. Clinically, this manifests as chronic sinusitis and recurrent infections that ultimately lead to bronchiectasis.

As ciliary function is necessary to ensure proper rotation of the developing organs in the chest and abdomen, its absence, will lead to situs inversus or a partial lateralizing abnormality.

Male patients with this condition tend to be infertile, due to asthenospermia (immotile spermatozoa due to impaired sperm flagella function).

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 9:

Kartagener's syndrome is a type of primary ciliary dyskinesia characterized by the classical triad of

• Situs inversus

• Chronic sinusitis

• Bronchiectasis

Infertility in Kartagener's syndrome is due to asthenospermia (immotile spermatozoa due to impaired sperm flagella function), not azoospermia (complete absence of spermatozoa in ejaculate).

Primary ciliary dyskinesia (also called immotile cilia syndrome) is an autosomal recessive disorder affecting ciliary motility. It occurs due to ultrastructural defects in the dynein arm of cilia. This causes the retention of secretions due to a failed mucociliary clearance mechanism. Clinically, this manifests as chronic sinusitis and recurrent infections that ultimately lead to bronchiectasis. As ciliary function is necessary to ensure proper rotation of the developing organs in the chest and abdomen, its absence will lead to situs inversus or a partial lateralizing abnormality.

Sold by @itachibot

Approximately half of the patients with primary ciliary dyskinesia have Kartagener syndrome.

Solution to Question 10:

The most common cause of lung abscess is aspirated oropharyngeal secretion.

Lung abscess represents necrosis and cavitation of the lung following microbial infection. They are broadly divided into two categories:

• Primary lung abscesses: usually arise from aspiration, often caused principally by anaerobic bacteria (most common)

• Secondary lung abscesses: arise in the setting of an underlying condition, such as a post-obstructive process (e.g., a bronchial foreign body or tumor) or a systemic process (e.g., HIV infection or another immunocompromising condition)

If you purchased this from someone else, it has been scammed. Sold by @itachibot

Solution to Question 11:

The given clinical scenario is suggestive of a primary lung abscess. Staphylococcus is involved in the formation of secondary lung abscess, not primary.

Microbial pathogens causing lung abscess:

Clinical conditions	Pathogens
Primary lung abscess (due to aspiration)	Anaerobes (e.g.,Peptostreptococcusspp.,Prevotellaspp., Bacteroides spp.,Streptococcus milleri), microaerophilic streptococci
Secondary lung abscess (due to underlying conditions)	Staphylococcus aureus, gram-negative rods (e.g.,Pseudomonas aeruginosa,Enterobacteriaceae),Nocardiaspp.,Aspergillusspp.,Legionellaspp.,Pneumocystis jirovecii
Embolic lesions	Staphylococcus aureus(often from endocarditis),Fusobacterium necrophorum(Lemierre's disease)

Solution to Question 12:

The given clinical scenario is suggestive of a primary lung abscess due to aspiration. Most common lobes involved in primary lung abscess are posterior upper lobes and superior lower lobes, given the predisposition of aspirated materials to be deposited in these areas.

Sold by @itachibot

The right lung is affected more commonly than the left because the right mainstem bronchus is less angulated.

In secondary abscesses, the location of the abscess may vary with the underlying cause.

Solution to Question 13:

The clinical scenario described along with the chest X-ray image showing air-fluid level is suggestive of a lung abscess.

The patients usually present with complaints of fever, cough with sputum production, and chest pain. Patients with putrid lung abscesses may report foul-smelling sputum. Patients with chronic lung abscesses usually have a history of night sweats, fatigue, and anemia.

On physical examination:

• Poor dentition/gingival disease

• Amphoric and/or cavernous breath sounds on lung auscultation

• Digital clubbing

• The absence of a gag reflex

CXR features of lung abscess:

• A lung cavity with an air-fluid level

• The cavity wall is thick and irregular, and a surrounding pulmonary-infiltrate is often present

• Computed tomography (CT) is more sensitive than chest radiography and is used to detect small cavities

The two x-ray images given below show thick-walled cavities with air-fluid levels on lateral and PA views respectively.



If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot



Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

In the above clinical scenario, the patient most likely has a lung abscess as depicted by the radiograph. Clindamycin is the drug of choice in the treatment of lung abscesses.

When there are contraindications to clindamycin, a combination of penicillin and metronidazole is likely to be as effective as clindamycin.

Anaerobes of oral origin are the most common causative organism and require broad-spectrum antibiotic coverage. Vancomycin, ciprofloxacin, and cephalexin have no significant activity against anaerobes.

Below is a chest X-ray showing a lung abscess (yellow dotted circle) with an air-fluid level (arrow).

Sold by @itachibot

Solution to Question 15:

Lung carcinoma is unlikely in patients with lung abscesses. Lung carcinoma may cause lung abscess (in 10-15% of cases) but lung abscess is not a premalignant condition.

Complications of lung abscess:

- Empyema
- Pneumatocele
- Bronchopleural fistula
- Bronchiectasis
- Metastatic infections like brain abscess
- Life-threatening hemoptysis
- Massive aspiration of lung abscess contents

Solution to Question 16:

The most common presentation of inhalation anthrax is hemorrhagic mediastinitis, which is seen in 38% of the patients.

Inhalational anthrax also known as wool sorter's disease is caused by Bacillus anthracis. Infection usually occurs among workers handling contaminated animal wool, hair, and skin in textile and tanning industries.

Patients present with malaise, fever, cough, shortness of breath & headache. B. anthracis spores are phagocytosed by alveolar macrophages and transported to mediastinal lymph nodes. Hemorrhagic necrosis of the thoracic lymph nodes draining the lung results in hemorrhagic mediastinitis. Pulmonary infiltrates, mediastinal widening, and pleural effusion are other findings.

Treatment of inhalation anthrax includes a combination of intravenous ciprofloxacin and clindamycin for at least 14 days, antitoxins (Raxibacumab, Obiltoxaximab), anthrax immunoglobulins and steroids (if associated meningitis is present).

Prophylaxis with ciprofloxacin (500mg twice daily for 2 months) is recommended for anyone at high risk of inhalational exposure to anthrax spores.

Solution to Question 17:

Central bronchiectasis is a typical feature of allergic bronchopulmonary aspergillosis (ABPA), not distal bronchiectasis.

ABPA is an eosinophilic pulmonary disorder caused by a hypersensitivity reaction to the colonization of airways by the fungus Aspergillus fumigatus. It occurs almost exclusively in patients with asthma (~2.5% of patients) or cystic fibrosis (up to 15%). In chronic cases, repeated episodes of bronchial obstruction, inflammation, and mucoid impaction can lead to

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

bronchiectasis, fibrosis, and respiratory compromise. While central bronchiectasis is characteristic, some patients can develop chronic cavitary lesions.

With the development of ABPA, asthma or cystic fibrosis typically worsens and may manifest with a new or worsening cough or an increase in sputum production or wheezing. Many patients report coughing up thick sputum casts. The presence of hyperattenuating mucus in the airways is highly specific.

ABPA is diagnosed serologically, based on the presence of:

Serum IgE > 1000 IU/ml

Positive skin prick test or IgE levels against A. fumigatus extract

Detection of Aspergillus-specific antibodies

Eosinophilia >500 cells/µL

Treatment - The use of systemic glucocorticoids is indicated for ABPA, but the course should be tapered by 3-6 months. Antifungal agents like itraconazole, voriconazole, and newer azoles can be used for over 4 months to control disease activity. Monoclonal antibodies against IgE, IL-4, and IL-5 receptors can be used for severe refractory cases.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

The culture of Aspergillus fumigatus from sputum is not a diagnostic criterion for allergic bronchopulmonary aspergillosis (ABPA). The sensitivity of sputum culture in ABPA is low.

ABPA is diagnosed serologically, based on the presence of:

- Serum IgE > 1000 IU/ml.
- Positive skin prick test or IgE levels against A. fumigatus extract.
- Detection of Aspergillus-specific antibodies.
- Eosinophilia >500 cells/µL.

Solution to Question 19:

The given clinical scenario is suggestive of aspergillosis. Fungal hyphae of Aspergillus species are narrow, septate, and have branching at 45o.

Yeast forms of Aspergillus are not seen in infected tissue.

Exposure to the high number of conidia that are found in construction sites, hay barns, moldy barks, and composts is a risk factor.

Clinical features:

- Most of the patients are asymptomatic
- Fever
- Cough

Sold by @itachibot

• Chest discomfort

• Trivial hemoptysis

• Shortness of breath

Halo sign-on HRCT refers to a localized ground glass appearance representing an area of hemorrhagic infarction surrounding a nodule. It is seen in invasive aspergillosis, a distinct entity where there is a direct invasion of hyphae into the lung tissue.

The air crescent sign describes the crescent of air that can be seen in invasive aspergillosis. It usually heralds recovery and is the result of increased granulocyte activity.

Monod sign refers to the gas that surrounds a mycetoma (most commonly an aspergilloma) in a pre-existing pulmonary cavity.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Pulmonary Function Tests

Question 1:

On performing a pulmonary function test, reduction in FEV1/FVC is characteristic of:

a) Restrictive disease

b) Obstructive disease

c) Normal lung function

d) Interstitial lung disease

Question 2:

A 66-year-old smoker presented with chronic productive cough and shortness of breath. Physical examination revealed a barrel chest and an expiratory wheeze on auscultation. All of the following findings would be seen in his pulmonary function tests except?

a) Normal residual volume

b) Decreased FVC

c) Normal vital capacity

d) Decreased FEV1/FVC

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

An 8-year-old boy presents with intermittent wheeze and cough. His mother reports that these episodes are precipitated by physical exercise. There is a history of atopic dermatitis in his mother. Pulmonary function tests are performed. Which of the following findings are least likely to be seen in him?

a) Decreased forced expiratory volume in one second

b) Increased residual volume

c) Decreased mean expiratory flow rate

d) Decreased diffusion capacity

Question 4:

Sold by @itachibot

Decreased maximum mid-expiratory flow rate is seen in obstruction of:

a) Medium airways

b) Large airways

c) Small airways

d) Trachea

Question 5:

Patients with which of the following conditions is likely to have pulmonary function test findings as given below:

a) Bronchiectasis

b) Bronchiolitis

c) Pleural effusion

d) COPD

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 6:

A 76-year-old retired worker from a granite quarry presents with progressive dyspnea and a chronic cough. He has never smoked cigarettes. His chest X-ray is shown below. Which of the following patterns would you see on his pulmonary function tests?



a) FEV1/FVC decreases, compliance decreases

b) FEV1/FVC increases and compliance increases

Sold by @itachibot

c) FEV1/FVC decreases and compliance increases

d) FEV1/FVC increases and compliance decreases

Question 7:

An elderly woman presents with breathlessness on exertion. On auscultation, bilateral basal rales are heard. A chest CT is done and is shown below. Which of the following is likely to be increased in her?

a) Total vital capacity

b) Forced vital capacity

c) Lung compliance

d) A-a gradient

Question 8:

The diffusing capacity of the lungs for carbon monoxide is decreased in patients with:

a) Alveolar haemorrhage

b) Congestive heart failure

c) Asthma

d) Emphysema

Question 9:

Sold by @itachibot

A 30-year-old woman presents with shortness of breath and fatigue. She has a medical history of limited cutaneous systemic sclerosis. Her pulmonary function test results are given below. What is the most probable diagnosis?

PFT	Observed	Predicted
FVC	2.63	2.82
FEV1%	88%	80%
DLCO	5.26	16.3

©MARROW

a) Pulmonary artery hypertension

b) Interstitial lung disease

c) Congestive heart failure

d) Bronchiectasis

Question 10:

A flow-volume loop of a patient is given below. What is the likely diagnosis?



©MARROW

Sold by @itachibot

a) Pulmonary edema

b) Interstitial lung disease

c) Tracheal stenosis

d) Emphysema

Question 11:

The flow-volume curve after spirometry in patient 1 is given below. The most likely diagnosis in this patient is:



©MARROW

a) Endobronchial neoplasm

b) Bronchial Asthma

c) Myasthenia gravis

d) Emphysema

Answer Key

Question No.	Correct Option
1	b
2	a
3	d

Sold by @itachibot

4	c
5	c
6	d
7	d
8	d
9	a
10	c
11	c

Detailed Explanations

Solution to Question 1:

Reduction in FEV1/FVC is characteristic of obstructive lung disease.

Forced vital capacity (FVC) is the volume of expired air recorded when a person inhales to the total lung capacity and exhales as hard and as completely as possible. The volume of air expired in the first second is called the forced expiratory volume 1 (FEV1).

Given below is a graphical representation of FEV and FEV1:



FVC is reduced in both obstructive and restrictive lung diseases. In obstructive diseases, the slope of FVC is steady, therefore the FEV1 value is low and the FEV1/FVC ratio decreases. In restrictive diseases, the airflow is fast in the beginning leading to a higher value of FEV1 and FEV1/FVC ratio increases.

Solution to Question 2:

Sold by @itachibot
If you got this from someone else, then it is stolen and tampered.

Sold by @itachibot

The given clinical scenario is suggestive of chronic obstructive pulmonary disease (COPD). It is characterized by an increased residual volume.

According to Uptodate, decreased vital capacity and increased total lung capacity is indicative of hyperinflation seen in COPD. FVC can be used a proxy for VC in normal individuals but in COPD FVC is less than VC due to air trapping, airflow limitation, and small airway collapse.

Solution to Question 3:

The given clinical scenario is suggestive of asthma. The diffusing capacity of the lungs for carbon monoxide (DLCO) tends to be normal or high in these patients.

Solution to Question 4:

Decreased maximum mid-expiratory flow rate (MMFR) is seen in obstruction of small airways.

MMFR is the average expiratory flow rate during the middle 50% of the vital capacity i.e., forced expiratory flow rate between 25 and 75% of the vital capacity.

This is the most sensitive measurement of early airflow obstruction, particularly in small airways.

Solution to Question 5:

Low FEV1 and FVC with a high FEV1/FVC ratio are characteristics of restrictive lung disease. Among the options, pleural effusion is a restrictive lung disease.

- Obstructive lung diseases
- Asthma
- COPD
- Bronchiectasis
- Bronchiolitis
- Restrictive lung diseases
- Intrinsic
- Interstitial lung disease
- Extrinsic
- Myasthenia gravis
- Kyphoscoliosis
- Pleural effusion

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 6:

The chest X-ray shows calcification of the hilar lymph nodes called eggshell calcification. The clinical scenario along with the chest X-ray is suggestive of silicosis. It is a restrictive lung disease characterized by increased FEV1/FVC and decreased compliance.

It is an occupational interstitial lung disease.

Solution to Question 7:

The given clinical scenario along with the honeycomb pattern of lung parenchyma due to interstitial fibrosis is suggestive of interstitial lung disease (ILD). The alveolar-arterial gradient (PAO2-PaO2) is increased in ILD. It is a restrictive lung disease.

In ILD, the alveolar-capillary bed is damaged and thus there is a decrease in diffusion capacity. Therefore, while the PAO2 (Alveolar O2) remains normal, the PaO2 (arterial) decreases, and the alveolar-arterial difference increases.



Solution to Question 8:

Diffusing capacity of the lungs for carbon monoxide (DLCO) is decreased in patients with emphysema.

DLCO measures the ability of the lungs to transfer gas from inhaled air in the alveoli to the red blood cells in pulmonary capillaries.

In general, the DLCO is used to identify the cause of dyspnea or hypoxemia. It is also used to monitor disease progression in interstitial lung disease, and identify pulmonary hypertension in patients at risk, such as those with systemic sclerosis.

DLCO is decreased in patients with:

Sold by @itachibot

• Interstitial lung disease

• Emphysema (COPD usually have normal DLCO)

• Bronchiolitis obliterans

• Pulmonary hypertension

• Thromboembolic disease

• Chronic congestive Heart Failure

DLCO is increased in patients with:

• Alveolar hemorrhage

• Acute Congestive heart failure

• Asthma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

The above PFT pattern shows near-normal FVC, increased FEV1%, and markedly decreased DLCO. This along with the given history is suggestive of pulmonary artery hypertension(PAH). It is more commonly associated with limited cutaneous systemic sclerosis.

Option B: Interstitial lung disease is a restrictive lung disease. In ILD, the FVC is usually,<80% of the predicted value. It is more commonly associated with diffuse cutaneous systemic sclerosis.

Options C and D: Congestive heart failure and bronchiectasis have obstructive pathology. They would have decreased FEV1% and normal DLCO.

Solution to Question 10:

The likely diagnosis is tracheal stenosis. The flow-volume loop shows flattened inspiratory and expiratory curves, suggestive of fixed upper-airway obstruction.

Sold by @itachibot



A normal flow-volume loop begins on the X-axis (Volume axis); at the start of the test, both flow and volume are equal to zero. After the starting point, the curve rapidly mounts to a peak: Peak expiratory flow (PEF).

Then, the PEF the curve descends (=the flow decreases) as more air is expired. A normal, non-pathological F/V loop will descend in a straight or a convex line from the top (PEF) to the bottom (FVC).

The forced inspiration that follows the forced expiration has roughly the same morphology, but the PIF (Peak Inspiratory Flow) is not as distinct as PEF.

Solution to Question 11:

The given flow-volume curve shows decreased lung volumes (total lung capacity, vital capacity, and residual volume) with normal shape, suggestive of a restrictive pattern. The curve is a miniature version of the normal curve with shift towards the right. It is seen in myasthenia gravis.

Myasthenia gravis presents with ventilatory restriction due to reduced respiratory muscle strength.

Asthma, emphysema, and endobronchial neoplasms present with obstructive flow patterns.

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Sold by @itachibot



Sold by @itachi... else,
If you purchased this from some...
you may have been scammed.

Sold by @itachibot

Respiratory Failure and ARDS

Question 1:

Which of the following is seen in patients with brain stem injury?

a) Low pO_2 and high pCO_2

b) Normal pO_2 and high pCO_2

c) Low pO_2 and low pCO_2

d) Low pO_2 and normal pCO_2

Question 2:

A 45-year-old man underwent emergency surgery following a road traffic accident. The surgery was performed under general anesthesia. Two days later, he developed features of respiratory failure. Which of the following types is likely to be present in him?

a) Type 1

b) Type 2

c) Type 3

d) Type 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A patient is admitted for myocardial infarction complicated by severe mitral regurgitation. He suddenly develops shortness of breath, decreased urine output, and drowsiness. Blood pressure is 86/40 mmHg, pulse 120 bpm, and respiratory rate 24 per minute. Which of the following statements is false?

a) Up to 40% of cardiac output may be distributed to the respiratory muscles

b) Patient will develop type 2 respiratory failure

c) Intubation and mechanical ventilation are treatment of choice

d) Batwing appearance can be expected on an x-ray

Question 4:

Sold by @itachibot

A 47-year-old construction site worker is admitted for fever, nausea, vomiting, and pain abdomen radiating to the back. The next day he develops sudden onset breathlessness and hypoxia. He has been drinking daily for the past 10 years. He had a myocardial infarction 6 months back. His chest X-ray is given below. What is the most probable diagnosis?

a) Interstitial lung disease

b) ARDS

c) Hospital acquired pneumonia

d) Left ventricular failure

Sold
If you purchased this
you may have been sc
else,
d.

Question 5:

According to the berlin definition, moderate ARDS is characterized by all, except

a) PaO2/FiO2 200 - 300mm/Hg

b) Bilateral interstitial infiltrates

c) Symptom onset within a week

d) No cardiac failure on echocardiography

Question 6:

Which of the following is least likely to be present in patients with ARDS?

a) Decreased pulmonary compliance

b) Increased pulmonary artery pressure

Sold by @itachibot

c) Increased left atrial pressure

d) PCWP <18 mm Hg

Question 7:

A fisherman was rushed to the ED following a near-drowning incident. On arrival, he was unconscious and was not breathing. He was resuscitated and stabilized. A few hours later, he developed sudden onset breathlessness and low saturation. Chest X-ray showed diffuse alveolar opacities. He was started on mechanical ventilation. What will be the recommended ventilator setting for him?

a) Low tidal volume and high PEEP

b) Low tidal volume and low PEEP

c) High tidal volume and high PEEP

d) High tidal volume and low PEEP

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	b
5	a
6	c
7	a

Detailed Explanations

Solution to Question 1:

Patients with brain stem injury develop type 2 respiratory failure. It is characterized by low pO2 and high pCO2.

Acute respiratory failure refers to acute impairment in gas exchange between the lungs and the blood causing hypoxia with or without hypercapnia.

Sold by @itachibot

Solution to Question 2:

The features of respiratory failure in the perioperative period of this patient is most likely due to lung atelectasis. Type 3 respiratory failure is most likely to be present in him.

Type 3 respiratory failure is also known as perioperative respiratory failure.

After general anesthesia, there is a decrease in functional residual capacity due to a reduction in muscular tone. This leads to airway closure and the collapse of the dependent lung units.

Atelectasis can be treated by frequent changes in position, chest physiotherapy, upright positioning, and control of incisional and abdominal pain. Non-invasive positive-pressure ventilation may also be used to reverse regional atelectasis.

Solution to Question 3:

The given clinical scenario is suggestive of cardiogenic shock. He is likely to develop type 4 respiratory failure, not type 2.

Patients in cardiogenic shock often have respiratory distress secondary to acute pulmonary edema (batwing appearance on x-ray). This increases the requirement of oxygen delivery to the respiratory muscles from <5% to up to 40% of cardiac output.

Hypoperfusion and increased oxygen requirement lead to inadequate perfusion of respiratory muscles which causes type 4 respiratory failure.

Intubation and mechanical ventilation are the treatment of choice.

Solution to Question 4:

The given clinical scenario along with the chest X-ray finding is suggestive of ARDS (acute respiratory distress syndrome) secondary to acute pancreatitis. Acute pancreatitis gives rise to inflammatory processes and endothelial damage that results in ARDS.

CXR features in ARDS:

• Early findings - normal or diffuse alveolar opacities (consolidation), bilateral and which obscure the pulmonary vascular markings

• Late findings - these opacities progress to more extensive consolidation that is diffuse

Clinical disorders commonly associated with ARDS:

• Direct lung injury:

• Pneumonia

• Aspiration of gastric contents

• Pulmonary contusion

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Sold by @itachibot

• Near-drowning

• Toxic inhalation injury

• Indirect lung injury:

• Sepsis

• Severe trauma- multiple fractures, flail chest, head trauma, burns

• Multiple transfusions

• Drug overdose

• Pancreatitis

• Post-cardiopulmonary bypass

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

Moderate ARDS is characterized by PaO2/FiO2: 101 - 200 mmHg. PaO2/FiO2: 200 - 300 mm/Hg is classified as mild ARDS.

Diagnostic criteria in ARDS—Berlin definition:(requires each of the following criteria to be present to diagnose ARDS)

• Onset - Acute: Respiratory symptoms must have begun < 1 week of a known clinical insult

• Bilateral alveolar or interstitial infiltrates: on a chest X-ray image or CT scan

• Respiratory failure must not be fully explained by cardiac failure or fluid overload. Objective assessment e.g Echo should be done in cases of no known cause.

• Impairment of oxygenation: The severity of hypoxemia defines the severity of ARDS.

Note: American-European Consensus Conference (AECC) classification of ARDS is no longer used.

Image: Alveolar/interstitial infiltrates in ARDS patient

Severity	PaO2/FiO2(On ventilator settings that include PEEP ≥5 cm H2O)
Mild	> 200 mmHgbut≤ 300 mmHg
Moderate	> 100 mmHgbut≤ 200 mmHg
Severe	≤ 100 mmHg

Sold by @itachibot



Sold by @itachi... else, If you purchased this from some... may have been scammed.

Solution to Question 6:

Increased left atrial pressure is not likely to be present in ARDS.

ARDS is characterized by

- PCWP of ≤18 mmHg with normal LA pressure
- Decreased pulmonary compliance
- Increased pulmonary artery pressure
- Severe hypoxemia



Sold by @itachibot

Solution to Question 7:

The given clinical scenario is suggestive of ARDS secondary to drowning. The ventilator setting recommended is low tidal volume and high PEEP.

Drowning causes direct lung injury. The aspiration of water into the lungs damages the surfactant. This disrupts the alveolar-capillary membrane and leads to alveolar edema, resulting in ARDS.

The ventilator settings recommended in ARDS are:

• Low tidal volume: To avoid barotrauma due to overdistention of the alveoli.

• High PEEP (positive end-expiratory pressure): This prevents the collapse of alveoli at the end of the expiration and improves oxygenation.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Neoplasms of the Lung

Question 1:

Which of the following is false regarding lung cancer?

a) It is the most common cancer in Indian men

b) Tobacco smoking is the most common cause

c) Adenocarcinoma is the most common histologic type

d) It is the most common cause of cancer death

Question 2:

What is the most common type of lung cancer in light smokers?

a) Squamous cell carcinoma

b) Adenocarcinoma

c) Small cell carcinoma

d) Large cell carcinoma

Question 3:

A 60-year-old man with a 35 pack-year history of smoking presented for a routine check-up. His brother was diagnosed with lung cancer at a similar age. Which of the following would be preferred to screen him for lung cancer?

a) Chest X-Ray

b) Low dose CT

c) High resolution CT

d) CECT

Question 4:

Which of the following is false about hypertrophic osteoarthropathy?

a) Platelets have a role in the development of HOA

b) Primary HOA disappears in adulthood

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) HOA is commonly associated with lung malignancies

d) Skin changes are more common in secondary HOA

Question 5:

A middle-aged non-smoker presented with a dry persistent cough and hemoptysis. A CT scan showed a mass in the periphery of his right lung. A biopsy was taken, cells showed an acinar pattern with mucin formation. The cells were positive for thyroid transcription factor 1 and napsin A. What other feature is most likely to be seen in this patient?

a) Hypertrophic osteoarthropathy

b) Lambert-Eaton myasthenic syndrome

c) Hypercalcemia

d) SIADH

If you purchased this from someone else, you have been scammed. Sold by @Itachibot

Question 6:

A female patient comes with complaints of unintentional weight loss and right shoulder pain for the past 6 months. She reports that the pain radiates to her arm and is not relieved by NSAIDs. Chest X-ray is given below. Which nerve roots are likely to be compressed?

a) C5, C6

b) C8, T1, T2

c) C6. C7

d) C3, C4, C5

Sold by @itachibot

Question 7:

A COPD patient presented to the emergency with an acute exacerbation. On evaluation, a chest X-ray revealed a solitary pulmonary nodule. Which of the following patterns of calcification would suggest a likely malignancy in this patient?

a) A dense central nidus

b) Popcorn calcification

c) Multiple punctate foci

d) Asymmetric calcification

Question 8:

What is the treatment of choice in patients with adenocarcinoma of the lung with EGFR mutation?

a) Bevacizumab

b) Paclitaxel

c) Imatinib

d) Gefitinib

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 55-year-old man presents with persistent cough and hemoptysis. The intern notices that he has moon-like facies and purple striae on his abdomen. On evaluation, a chest X-ray revealed a mass in his right lung. Which of the following type of lung carcinoma could be the cause of this patient's features?

a) Adenocarcinoma

b) Bronchoalveolar carcinoma

c) Small cell carcinoma

d) Squamous cell carcinoma

Question 10:

An elderly man was rushed to the ED with confusion and multiple episodes of vomiting. His wife reported that he had increased frequency of urination and constipation for the past 3

Sold by @itachibot

days. He is a known case of lung cancer. ECG showed a shortened QT interval. What could be the primary cause of his symptoms?

a) Ectopic secretion of ACTH

b) Syndrome of inappropriate antidiuretic hormone secretion

c) Ectopic secretion of PTH related peptide

d) Tumor lysis syndrome

Question 11:

A 62-year-old woman presents with vomiting, diarrhea, and decreased urine output since morning. On the previous day, she received chemotherapy for her stage 4 lung cancer. On examination, she is drowsy, HR-110/min and RR-30/min. All of the following biochemical findings can be seen in this patient except:

a) Hyperuricemia

b) Hypercalcemia

c) Hyperkalemia

d) Hyperphosphatemia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Which of the following is not used in the management of patients with tumor lysis syndrome?

a) Sodium bicarbonate

b) Rasburicase

c) Aggressive hydration

d) Febuxostat

Question 13:

Which of the following patients with non-small lung carcinoma is a suitable candidate to undergo a potentially curative surgical resection:

a) 60-year-old smoker with a tumor within 2cm of carina

b) 45-year-old woman with malignant pleural effusion

c) 55-year-old non-smoker with tumor involving the main pulmonary artery

d) 57-year old light smoker with ipsilateral mediastinal node metastases

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	b
3	b
4	d
5	a
6	b
7	d
8	d
9	c
10	c
11	b
12	a
13	d

If you purchased this, you may have received it for free.

Detailed Explanations

Solution to Question 1:

The most common cancer seen in Indian men is lip and oral cavity cancer, not lung cancer.

Most common cancer:

• In men globally- Lung cancer

• In Indian men- Lip and oral cavity cancer

• In women globally- Breast cancer

• In Indian women- Breast cancer

Tobacco smoking is a primary cause of the development of lung cancer. It increases 10 fold greater risk of developing lung cancer compared to those who have never smoked.

It is the most common cause of cancer death.

Solution to Question 2:

Sold by @itachibot

The most common lung cancer in light smokers (<10 pack-year history) is adenocarcinoma.

WHO classification of lung carcinomas:

The WHO classification system divides epithelial lung cancers into four major cell types:

• Small-cell lung cancer (SCLC)

• Non-small-cell lung carcinomas (NSCLCs)

• Adenocarcinoma

• Squamous cell carcinoma

• Large-cell carcinoma

Adenocarcinoma is most commonly seen in:

• Lifetime non-smokers

• Light smokers (<10 pack-years history)

• Women

• Younger adults (<60 years)

In smokers, the most common types are small-cell carcinoma and squamous cell carcinomas.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Low-dose CT/ thin-slice spiral chest CT would be preferred to screen him for lung cancer. He belongs to the high-risk population to develop lung cancer.

The high-risk population is individuals whose age is ≥60 years and who have a smoking history of ≥10 pack-years.

Low dose CT scans are shown to detect more lung nodules and cancers than the standard CXR in selected high-risk populations. Also, up to 85% of cancers thus detected are classified as stage I. Early staged cancers are considered potentially curable with surgical resection.

Here, the aim is to reduce medical radiation exposure as much as possible, low dose CT is effective in screening for lung cancer, and is thus preferred.

Solution to Question 4:

Skin changes are more common in primary HOA, not secondary HOA.

Hypertrophic osteoarthropathy is a condition characterized by the clubbing of the digits. In advanced stages, it is characterized by the periosteal new bone formation and synovial effusion. The exact mechanism of its development is unclear. However, platelets are said to have a role in the development of HOA via platelet-derived growth factor.

There are two types of HOA:

Sold by @itachibot

Primary HOA	Secondary HOA
Inherited autosomal dominant condition	Acquired condition, secondary toLung neoplasmsMesotheliomaSuppurative lung infectionsCongenital heart diseaseThyroid acropachyInflammatory bowel disease
Skin changeslike thickening, seborrhea, excessive sweating are commonly seen	Skin changes are unusual
The symptomsdisappearinadulthood	The symptoms persist till the underlying cause is treated

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

The given clinical scenario along with the biopsy findings is suggestive of lung adenocarcinoma. It is most commonly associated with hypertrophic osteoarthropathy.

Adenocarcinomas are mainly seen in the periphery of the lung. They grow in various patterns like acinar, lepidic, papillary, micropapillary, and solid with mucin formation. Cells are positive for markers like thyroid transcription factor 1 and napsin A.

Option C: Hypercalcemia due to ectopic production of PTH-related peptide is seen in squamous cell carcinoma.

Options B and D: However, SIADH, Cushing's syndrome, Lambert-Eaton myasthenic syndrome are associated with small-cell lung carcinoma.

Solution to Question 6:

The given clinical scenario along with the X-ray findings is suggestive of Pancoast syndrome / superior sulcus tumor. This affects the C8, T1, and T2 nerve roots.

It presents with pain that radiates to the ulnar distribution of the arm, often with the destruction of the 1st and 2nd ribs. Patients with the central or endobronchial growth of the primary tumor may present with a cough, hemoptysis, wheeze, stridor, dyspnea, or post-obstructive pneumonitis.

Regional spread can cause

• Tracheal obstruction

• Esophageal compression with dysphagia

• Recurrent laryngeal paralysis with hoarseness

• Phrenic nerve palsy with the elevation of the hemidiaphragm and dyspnea

• Sympathetic nerve paralysis with Horner's syndrome (enophthalmos, ptosis, miosis, and anhydrosis)

Sold by @itachibot

Solution to Question 7:

Asymmetric calcification is suggestive of a malignant lesion.

Benign lesion	Malignant lesion
Lack of growth over a period of >2 years	The growth of the lesion is ev ident in comparison with an old X-ray
Patterns of calcification:Dens e central nidusMultiple punct ate fociBull's eye granulomaP opcorn ball hamartomas	Patterns of calcification:Asy mmetricLack of calcification
Associated chest symptoms a re uncommon	chest symptoms like cough, h emoptysis are common
Complications are uncommo n	Complications like atelectasis, pneumonitis, and effusion ar e common

Sold by
If you purchased this fr
you may have been scan
else,
d.

Solution to Question 8:

Adenocarcinomas with EGFR mutation are exquisitely sensitive to tyrosine kinase inhibitors of the EGFR signal pathway such as gefitinib and erlotinib.

Option A: Bevacizumab is a monoclonal antibody against VEGF.

Option B: Paclitaxel- a taxane, is a plant alkaloid and chemotherapeutic agent.

Option C: Imatinib is a tyrosine kinase inhibitor effective in the treatment of CML and GIST.

Solution to Question 9:

In the given clinical scenario the patient has features of Cushing's syndrome, secondary to small-cell lung cancer. This occurs due to ectopic ACTH production. It can also seen in pulmonary carcinoids.

Most effective management is treating underlying SCLC or carcinoid.

Solution to Question 10:

The primary cause of his symptoms is ectopic secretion of PTH related peptide. Lung cancers are associated with several paraneoplastic syndromes. Squamous cell lung cancer is associated with hypercalcemia due to ectopic PTH related peptide. Hypercalcemia may present as vomiting,

Sold by @itachibot

abdominal pain, constipation, increased frequency of urination, kidney stones, bone pain and shortened QT interval.

Solution to Question 11:

The given clinical scenario is suggestive of tumor lysis syndrome (TLS). It is associated with hypocalcemia, not hypercalcemia.

TLS is caused by the destruction of a large number of rapidly proliferating neoplastic cells. It is usually seen within 1-5 days of receiving chemotherapy. It has also been observed with the administration of glucocorticoids, letrozole, tamoxifen, rituximab and gemtuzumab. Rarely, spontaneous necrosis of malignancies causes TLS.

Clinical features:

• Nausea, vomiting, diarrhea, lethargy

• Pruritus, renal colicky pain, oliguria

• Muscle cramps, carpopedal spasm, tetany

Biochemical findings:

• Hyperuricemia

• Hyperkalemia

• Hypocalcemia

• Hyperphosphatemia

Due to these biochemical changes, there is an increased risk of arrhythmias, acidosis, and acute kidney injury.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

Sodium bicarbonate is not recommended in the treatment of tumor lysis syndrome (TLS).

Urinary alkalinization by sodium bicarbonate decreases the solubility of xanthine, hypoxanthine, and calcium phosphate. This potentially increases the risk of intratubular crystallization.

Management of TLS includes:

• Aggressive hydration

• Allopurinol

• Rasburicase - recombinant urate oxidase

• Febuxostat - non-purine selective xanthine oxidase inhibitor

Solution to Question 13:

Sold by @itachibot

Patients with ipsilateral mediastinal node metastases are suitable candidates for curative resection.

In patients with NSCLC, the following are contraindications to potentially curative resection:

- Extra-thoracic metastases
- Vocal cord involvement
- Superior vena cava syndrome
- Involvement of the main pulmonary artery
- Cardiac tamponade
- Phrenic nerve paralysis
- Malignant pleural effusion
- Tumor within 2 cm of the carina
- Metastasis to the contralateral lung
- Metastases to supraclavicular lymph nodes
- Contralateral mediastinal node metastases

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Sleep apnea

Question 1:

Sleep apnea is defined as a temporary pause in breathing during sleep lasting at least :

a) 30 seconds

b) 20 seconds

c) 15 seconds

d) 10 seconds

Question 2:

A patient is referred to the sleep clinic for polysomnography. Which of the following is not a component of it?

a) EEG

b) Skeletal muscle activation

c) Peripheral pulse oximetry

d) Arterial blood gases

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A middle-aged obese patient presented with complaints of daytime tiredness and decreased exercise tolerance. Echocardiology revealed reduced ejection fraction. The sleep study showed periods of apnea without respiratory efforts. Which of the following is not true regarding his condition?

a) Transient cessation of neural drive to all respiratory muscles

b) Associated with Cheyne Stokes breathing

c) Is due to decreased sensitivity to pCO_2

d) Can be induced by CPAP at high pressure

Question 4:

Which of the following drugs is not a risk factor for the development of central sleep apnea?

Sold by @itachibot

a) Opioids

b) Strychnine

c) Barbiturates

d) Gelsemium

Question 5:

A patient comes to the sleep clinic and the following pattern is seen on polysomnography. What type of breathing pattern is it?



a) Hyperventilation respiration

b) Cheyne-Stokes respiration

c) Kussmaul's respiration

d) Biot's respiration

Question 6:

A 40-year-old man presents with daytime sleepiness and impaired concentration and memory. On examination, his BMI is 41 kg/m2, blood pressure is 160/100 mm of Hg. His awake arterial blood gas analysis is given below. What is the most likely diagnosis?

a) Obstructive sleep apnea

b) Narcolepsy

c) Obesity hypoventilation syndrome

Sold by @itachibot

d) Central sleep apnea

Question 7:

A 55-year-old man is brought to the OPD by his wife for gasping episodes during sleep and excessive daytime sleepiness. She reported that he even dozed off while driving yesterday and nearly caused an accident. He is being treated for hypertension, gastro-oesophageal reflux, and type 2 diabetes. What else would be the ideal therapy for his symptoms?

a) Supplemental oxygenation therapy

b) Uvulopalatopharyngoplasty

c) Continuous positive airway pressure therapy

d) Implantable neuro-stimulation of the hypoglossal nerve

Question 8:

Which of the following is not a component of Syndrome Z?

a) Obstructive sleep apnea

b) Blood pressure more than 130/85 mmHg

c) Fasting triglyceride more than150mg/dl

d) LDL more than 100mg/dl

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	d
3	c
4	b
5	b
6	c
7	c
8	d

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Sleep apnea is defined as a temporary pause in breathing during sleep lasting at least 10 seconds.

Sleep apneas can be further divided into:

• Obstructive apneas - Sleep apnea with persistent respiratory effort.

• Central apneas - Sleep apnea with an absence of respiratory effort.

The gold standard for the diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS) is an overnight polysomnogram (PSG).

Solution to Question 2:

Arterial blood gases are not part of polysomnography. However, they are usually measured to rule out alveolar hypoventilation secondary to lung disease or chest wall abnormalities.

An overnight polysomnogram is a multi-parametric test and is the gold standard for the diagnosis of obstructive sleep apnea. It includes:

• EEG

• Eye movements (EOG)

• Muscle activity or skeletal muscle activation (EMG)

• ECG

• Respiratory airflow and respiratory effort indicators

• Peripheral pulse oximetry

• Sleep latency

• Sleep efficiency

• Arousal index

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



REOG, LEOG	-	Electro-oculography
C3A2, C4A1, O1A2, O2A1	-	Electroencephalography
Chin	-	Chin electromyography
ECG1	-	Electrocardiography
SpO2	-	Pulse oximetry
HR	-	Heart rate
MSnore	-	Oronasal airflow
THO and ABD	-	Thoracic and Abdominal movements

©MARROW

Other sleep disorders diagnosed by polysomnography:

- Narcolepsy
- Idiopathic hypersomnia
- Periodic limb movement disorder
- REM behavior disorder
- Parasomnias

If you purchased this from some else, you may have been scammed. Sold by @itachi

Solution to Question 3:

The given clinical scenario is suggestive of central sleep apnea (CSA), and it is caused by increased sensitivity to pCO_2. Thus changes in pCO_2 levels during sleep lead to an unstable breathing pattern that manifests as hyperventilation alternating with apnea. Crescendo-decrescendo breathing pattern known as Cheyne-Stokes respiration is seen.

Risk factors for CSA:

- Heart failure
- Opioid medications
- Hypoxia (e.g., breathing at high altitude)
- Continuous positive airway pressure (CPAP) at high pressures seems to induce central apnea; this condition is referred to as complex sleep apnea.

Solution to Question 4:

Strychnine is not a risk factor as it does not cause respiratory depression. It is a CNS stimulant. Only drugs that cause respiratory centre depression are risk factors for the development of central

Sold by @itachibot

sleep apnea.

Option A: Opioids depress the respiratory centre in a dose-dependent manner.

Option C: Barbiturates cause respiratory depression in high doses.

Option D: Gelsemium is an ergot-type alkaloid and is a CNS depressant. Often used in homoeopathy, in large doses, paralysis and death from respiratory failure can occur.



The pink shaded area shows an apneic period without any respiratory efforts.

Solution to Question 5:

The breathing pattern shown in the image is suggestive of Cheyne-Stokes respiration. It is characterized by progressively deeper and sometimes faster breathing, followed by a gradual decrease that results in a temporary stop in breathing called apnea.

It is caused by an increased sensitivity of the respiratory center to arterial PCO_2. There is an apneic phase, during which arterial PO_2 falls and arterial PCO_2 rises. These changes in the arterial blood gas content stimulate the respiratory center, resulting in hyperventilation and hypocapnia, followed by recurrence of apnea. It is present in 40% of patients with advanced HF and usually is associated with low cardiac output.

Sold by @itachibot



Solution to Question 6:

The most probable diagnosis of this patient is obesity hypoventilation syndrome.

Obesity hypoventilation syndrome (OHS) or Pickwickian syndrome is defined as the presence of awake alveolar hypoventilation in an obese individual which cannot be attributed to other conditions associated with alveolar hypoventilation.

Patients with OHS are obese and give a history of daytime hypersomnolence, loud snoring, choking during sleep, fatigue, impaired concentration, and memory. On examination usually, there is a small oropharynx and a thick neck.

OHS may be associated with the following findings:

• Elevated serum bicarbonate (>27 mEq/L)

• Hypercapnia (PaCO2 >45 mmHg)

• Hypoxemia (PaO2 <70 mmHg)

• Polycythemia

Options A and D: In obstructive sleep apnea and central sleep apnea, the awake ABG values are usually normal.

Option B: There are no ABG changes in narcolepsy.

Solution to Question 7:

The given clinical scenario suggests a diagnosis of obstructive sleep apnea/hypopnea syndrome (OSAHS). The ideal therapy is continuous positive airway pressure therapy (CPAP).

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option A: Supplemental oxygenation can improve oxygen saturation but there is little evidence that it improves OSAHS symptoms.

Option B: Uvulopalatopharyngoplasty is the most common surgery done in OSAHS but it is reserved for patients who cannot tolerate CPAP.

Option D: Unilateral neuro-stimulation of the hypoglossal nerve through a surgically implanted device is a recently tested therapy for OSAHS. It can be used as an alternative in selected patients who cannot tolerate CPAP.

Solution to Question 8:

Syndrome Z is defined as a combination of obstructive sleep apnea syndrome with metabolic syndrome. LDL more than 100 mg/dL is not a criterion for the diagnosis of metabolic syndrome.

Metabolic syndrome is diagnosed based on The NCEP: ATP III Criteria. Three or more of the following must be present :

• Central obesity: waist circumference >102 cm (M), >88 cm (F)

• Hypertriglyceridemia: triglyceride level ≥150 mg/dL or specific medication

• Low HDL cholesterol: <40 mg/dL and <50 mg/dL for men and women, respectively, or specific medication

• Hypertension: blood pressure ≥130 mmHg systolic or ≥85 mmHg diastolic or specific medication

• Fasting plasma glucose level ≥100 mg/dL or specific medication or previously diagnosed type 2 diabetes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Urine Analysis

Question 1:

Urine sample of a patient has been collected for analysis. How soon should it be analyzed?

a) Within 6-8 hours

b) Within 2-4 hours

c) Within 4-6 hours

d) Within 1-2 hours

Question 2:

All of the following determine the color of urine except:

a) Chemical content

b) Concentration

c) pH

d) Solubility of solutes

Question 3:

Which of the following drugs can cause green-colored urine in a patient?

a) Triamterene

b) Rifampicin

c) Desferoxamine

d) Imipenem

Question 4:

What value of urine specific gravity is called isosthenuria?

a) 1.020

b) 1.010

c) 1.015

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) 1.025

Question 5:

A patient is being evaluated for the following appearance of his urine. A urine dipstick test is performed and the result is negative. What is the likely diagnosis?

a) Myoglobinuria

b) Haemoglobinuria

c) Porphyria

d) Urolithiasis

If you purchase
you may have
else,

Question 6:

Which of the following patients are considered to have significant hematuria based on their urine analysis findings?

a) 1,2,3,4 and 5

b) 2,4 and 5

c) 1 and 4

d) 2, 3 and 5

Question 7:

Sold by @itachibot

A patient with hematuria, generalized edema, and abdominal distension is suspected to have a glomerular pathology. Which of the following is likely to be seen in the urine?

a) Bright red cells

b) Acanthocytes

c) Beta-2 microglobulin

d) Waxy casts

Question 8:

An 18-year-old boy presents to the clinic with a history of malaise. He was diagnosed with mild sensorineural hearing loss nearly 10 years ago. He also has a history of recurrent hematuria. What is the likely diagnosis?

a) Fanconi's anemia

b) Alport syndrome

c) Autosomal dominant polycystic kidney disease

d) IgA nephropathy

Question 9:

Which of the following is true about proteinuria?

a) Proteinuria is always due to glomerular pathology

b) Albuminuria upto 150 mg/day is considered normal

c) Nephrotic range proteinuria is > 3.5 g/day

d) Proteins are reabsorbed in the DCT

Question 10:

All of the following findings in a patient are suggestive of non-glomerular proteinuria except:

a) Proteinuria < 1 g/day

b) Type of protein excreted is β2-microglobulin

c) Albumin : β2-microglobulin ratio is reduced

d) Absence of N-acetyl glucosaminidase

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Question 11:

A patient presents with recurrent kidney stones. The microscopic examination of urine is shown below. What is the type of stone seen in this patient?

a) Struvite

b) Oxalate

c) Uric acid

d) Cystine

Question 12:

Which of the following is incorrect about casts in urine?

a) Width of cast depends on the size of tubule in which it is formed

b) Casts in urine are specific to kidneys

c) Site of formation is proximal convoluted tubule

d) Casts are made up of uromodulin

Question 13:

A patient with cryptococcal meningitis was started on amphotericin B. Two days later, he is found to have decreased urine output. Urinalysis reveals granular casts. What could be the possible diagnosis?

Sold by @itachibot

a) Mycotoxic nephropathy

b) Acute tubular necrosis

c) Acute interstitial nephritis

d) Papillary necrosis

Question 14:

Which of the following casts are associated with a poor prognosis?

a) Epithelial casts

b) WBC casts

c) Hyaline casts

d) Broad casts

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	b
2	d
3	a
4	b
5	c
6	b
7	b
8	b
9	c
10	d
11	d
12	c
13	b
14	d

Detailed Explanations

Sold by @itachibot

Solution to Question 1:

Urine samples should be analyzed within 2-4 hours after collection to avoid cell lysis and precipitation of solutes. If urine analysis is not possible in the given time, then the sample can be stored at 4-8 oC for a maximum of 12 hours.

Urine samples can be collected using the following techniques:

• Midstream sample during spontaneous voiding

• Fresh specimen from an indwelling catheter

• Suprapubic needle aspiration of the bladder, especially in infants

Solution to Question 2:

The solubility of solutes is not a determining factor of the color of urine. Determinants of the color of urine are:

• Chemical content

• Concentration

• pH (Normal pH is 4.5-8)

The normal color of urine is pale straw due to the presence of urochrome.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 3:

Green urine can be caused by triamterene, a potassium-sparing diuretic.

Solution to Question 4:

Isosthenuria is defined as a urine specific gravity of 1.010, which is similar to that of plasma. The normal specific gravity of urine is around 1.020 to 1.030.

Specific gravity and osmolality are measures of the concentration or relative density of urine.

• Specific gravity: The weight of a solution relative to that of equal amounts of water. It is measured by refractometry or using reagent strips.

• Osmolality: The gold standard for relative density. It is defined as the number of osmoles of solute per kilogram of solvent. It is measured directly with an osmometer.

Isosthenuria means that the kidney is neither concentrating nor diluting the urine. This is seen in chronic renal failure (CRF) and acute tubular necrosis (ATN).

Abnormality	Causes
Isosthenuria	CRF, ATN

Sold by @itachibot

Abnormality	Causes
Hyposthenuria (low specific gravity)	Diabetes insipidus
Hypersthenuria (high specific gravity)	SIADH, Severe dehydration

Solution to Question 5:

The given image shows reddish discoloration of the urine sample. A negative urine dipstick indicates absence of RBCs and is suggestive of porphyria.

A urine dipstick test is the first step to be done in a patient with high coloured urine to help identify true hematuria. A positive dipstick test identifies heme that is present in red blood cells, haemoglobin or myoglobin. A negative dipstick test implies that the reddish discolouration is due to pigments from certain food items (beetroot), drugs or porphyrias.



Solution to Question 6:

Hematuria is defined as the presence of 2-5 RBCs / high-powered field (HPF). It is considered significant or persistent in the presence of one of the following:

• RBCs > 3 / HPF on 3 or more occasions

• RBCs > 100 / HPF on 1 occasion

• Gross hematuria

Sold by @itachibot

Solution to Question 7:

Acanthocytes are a subtype of dysmorphic red blood cells with protruding blebs that are seen in glomerular disease.

The 40-5-1 rule can be used to identify glomerular diseases based on urinalysis:

• Dysmorphic RBCs - 40% (can range from 10-80%)

• Acanthocytes - 5% (can range from 2-5%)

• Even a single RBC cast

Hematuria can be either due to systemic or renal causes. Renal causes can be further subdivided into glomerular and non-glomerular causes as given below:

Glomerular	Non-glomerular
Dysmorphic RBCs seen (abnormal shape and size)	Isomorphic RBCs seen (normal shape and size)
RBC casts may be present	RBC casts always absent
MCV is reduced (<72 fl)	MCV is normal
Protein excretion > 1g /day	Protein excretion < 1g /day
The cause is usually medical	The cause is usually surgical

If you purchased this, you may have been scammed.

Solution to Question 8:

This clinical scenario is suggestive of Alport syndrome. The clinical features include:

• Recurrent hematuria and nephritic syndrome

• Sensorineural hearing loss

• Anterior lenticonus

Other causes of recurrent hematuria include:

• Alport syndrome

• IgA nephropathy

• Autosomal dominant polycystic kidney disease (ADPKD)

• Papillary necrosis

• Renal cell carcinoma

Solution to Question 9:

Nephrotic-range or heavy proteinuria refers to excretion of > 3.5 g/day or > 1000 mg/m2/day in children.

Sold by @itachibot

Normally, the following mechanisms prevent proteinuria:

• The glomerular basement membrane acts as a barrier to prevent the loss of negatively charged and large proteins.

• Reabsorption of small proteins in the proximal convoluted tubule (PCT).

The various types of proteinuria are:

• Glomerular - increased filtration of proteins due to disruption of basement membrane

• Tubular - damage to tubules prevents protein reabsorption in the PCT

• Overflow - excess production of low molecular weight proteins such as light chains

• Postrenal - infections or stones can lead to secretion of IgA and IgG by the urinary tract

Type of Proteinuria	Total proteinuria (mg/day)	Albuminuria(mg/day)
Normal-range	<150MainlyTamm-Horsfallprotein (70-75 mg/day)	< 30
Moderately increased	150-500	30-300
Severely increased	> 500	> 300
Orthostatic proteinuria	150-500	—

Solu... purchased this... you may have been...

Solution to Question 10:

Non-glomerular proteinuria is characterized by the presence of N-acetyl glucosaminidase (NAG).

This form of proteinuria occurs as a result of tubular injury. N-acetyl glucosaminidase (NAG) is a lysosomal enzyme present in the proximal tubular cells and is released into the urine following any injury to the tubules.

Low-molecular-weight proteins such as β2-microglobulin are normally reabsorbed in the PCT. When there is tubular injury, the proportion of these proteins in urine increases. This lowers the albumin: β2-microglobulin ratio in urine.

In glomerular proteinuria, there is increased filtration of macromolecules like albumin across the glomerular capillary wall. NAG is absent as there is no tubular injury.

	Glomerular proteinuria	Non-glomerular proteinuria
Type of protein excreted	High molecular weight (Albumin)	Low molecular weight (β2-microglobulin)
24 hr protein excretion	> 1-2 g/day	< 1 g/day
RBC casts	Present	Absent
Albumin: β2microglobulin ratio (Normal 100:1)	1000:1	10:1
N-acetyl glucosaminidase (NAG)	Absent	Present

Sold by @itachibot

Solution to Question 11:

The image shows a hexagonal crystal, suggesting cystine stones. These stones are a recurrent feature in patients with cystinuria.

Crystals in urine are usually associated with kidney stones. Crystals form in the urine depends upon a variety of factors including:

• The degree of concentration of constituent molecules

• The urine pH

• The presence of inhibitors of crystallization

Calcium oxalate and uric acid crystals can sometimes be found in normal urine that has been left to stand, due to crystal formation ex vivo.

Solution to Question 12:

The sites of formation of casts are the collecting ducts and DCTs (distal convoluted tubules).

Casts are cylindrical structures formed by the aggregation of Tamm-Horsfall mucoprotein or uromodulin. This protein is secreted in the thick ascending loop of Henle. They assume the shape and size of the lumen in which they are formed. Several factors favour cast formation, such as urine stasis, low pH and high concentration. Casts are the only elements in urine sediments that are specifically of renal origin.

Solution to Question 13:

Muddy-brown granular casts are seen in acute tubular necrosis, a type of intrinsic renal failure. ATN can occur secondary to nephrotoxicity or hypovolemia.

Solution to Question 14:

Broad casts are associated with a poor prognosis.

They refer to wider waxy casts that are seen in advanced chronic kidney disease. It implies dilation of the tubules.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Chronic Kidney Disease

Question 1:

Which of the following is not a risk factor for chronic kidney disease?

a) Childhood obesity

b) Diabetes mellitus

c) Family history of kidney disease

d) Large for gestational age at birth

Question 2:

What is the most common cause of chronic kidney disease?

a) Diabetic nephropathy

b) Glomerulonephritis

c) Hypertension

d) Adult polycystic kidney disease

Question 3:

Patients with which of the following conditions are unlikely to develop chronic renal failure?

a) Post-streptococcal glomerulonephritis

b) Membranous glomerulonephritis

c) Membranoproliferative glomerulonephritis

d) Focal segmental glomerulosclerosis

Question 4:

Which of the following parameters are used in the staging of chronic kidney disease?

a) Serum albumin and creatinine

b) Serum urea and creatinine

c) Urine albumin and eGFR

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) Serum creatinine and eGFR

Question 5:

In patients with chronic kidney disease, at which stage do the complications of kidney failure start appearing?

a) Stage 1

b) Stage 2

c) Stage 3-4

d) Stage 4-5

Question 6:

A 66-year-old woman is being evaluated for new-onset uremic encephalopathy. What is her GFR likely to be?

a) < 50 mL/min/1.73 m2

b) < 15 mL/min/1.73 m2

c) < 40 mL/min/1.73 m2

d) < 30 mL/min/1.73 m2

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following is incorrect about uremic syndrome?

a) It occurs due to accumulation of toxins

b) There is loss of kidney function and hormone regulation

c) Systemic inflammation is seen

d) Can be eliminated by removing urea through intensive dialysis

Question 8:

A 60-year-old man with CKD has been on dialysis for 3 months. He now presents with intense body pains and muscle pain. He suffered from 2 fractures over the last 2 months. On evaluation, he was found to have decreased PTH levels and normal calcium levels. He has been regularly taking oral vitamin D and calcium tablets. What is the likely diagnosis?

Sold by @itachibot

a) Osteomalacia

b) Adynamic bone disease

c) Osteitis fibrosa cystica

d) Tumoral calcinosis

Question 9:

A middle-aged woman with chronic kidney disease presents to the OPD with exercise intolerance, fatigue, and shortness of breath. On examination, she is found to have severe pallor. Which of the following is least likely to be a cause for these findings?

a) Deficiency of erythropoietin

b) Iron deficiency

c) Hyperparathyroidism

d) Sickle cell anemia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following is not seen in patients with chronic renal failure?

a) Peripheral neuropathy

b) Hyperthyroidism

c) Decreased vitamin D levels

d) Osteomalacia

Question 11:

What is the most common neurological disorder seen in patients with CKD?

a) Dementia

b) Restless leg syndrome

c) Peripheral neuropathy

d) Encephalopathy

Question 12:

Sold by @itachibot

A 27-year-old woman goes to her family doctor as she is unable to fall asleep at night. She complains of tingling and crawling sensations over her legs at night. She also feels a strong urge to move her legs, which relieves the symptoms for a while. Which of the following is incorrect about this condition?

a) Autosomal dominant inheritance

b) Associated with low serum ferritin

c) Most common cause is chronic kidney disease

d) Most common movement disorder in pregnancy

Question 13:

What is the most likely cause of convulsions in a patient with CKD?

a) Hypokalemia

b) Hyperkalemia

c) Hypercalcemia

d) Water intoxication

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

What is most common cause of mortality in patients with chronic kidney disease?

a) Cardiovascular disease

b) Sepsis

c) Uremia

d) Dialysis-related complications

Question 15:

Which of the following would be elevated in patients with chronic kidney disease?

a) Fibrinogen

b) Fetuin

c) Albumin

d) Globulin

Sold by @itachibot

Question 16:

What is the predominant electrolyte abnormality seen in patients with advanced CKD?

a) Hyponatremia

b) Hypokalemia

c) Non-anion gap metabolic acidosis

d) High anion gap metabolic acidosis

Question 17:

Which among the following is the most life-threatening manifestation of CKD?

a) Decreased GFR

b) Increased creatinine

c) Hyperkalemia

d) Proteinuria

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

Patients with CKD have increased levels of all of the following except:

a) Insulin

b) PTH

c) Estrogen

d) FGF-23

Question 19:

A patient presents to the emergency with chest pain that is radiating to his shoulder, shortness of breath, and diaphoresis. He is a known case of chronic kidney disease. Which of the following investigations is best for diagnosis?

a) Serial troponin levels over 6 hours

b) Troponin levels on admission and 1 hour later

c) Troponin and CK-MB levels at admission

d) Serial CK-MB levels over 6 hours

Sold by @itachibot

Question 20:

Which of the following is the most sensitive test to detect early diabetic nephropathy in a patient?

a) Urine albumin

b) Renal ultrasound

c) Creatinine clearance test

d) Serum BUN level

Question 21:

A patient with chronic kidney disease is found to have normal-sized kidneys on ultrasonography. Which of the following diagnoses is least likely?

a) Diabetic nephropathy

b) HIV nephropathy

c) Amyloidosis

d) Hypertensive nephropathy

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 22:

A 65-year-old hypertensive patient has recently been diagnosed with stage 3 chronic kidney disease. According to the ESC 2018 guidelines, what target blood pressure would you choose for this patient?

a) 140/90 mm Hg

b) 140/80 mm Hg

c) 130/90mmHg

d) 120/80 mmHg

Question 23:

A 50-year-old man who has been on dialysis for 2 years presents with bone pain and bowing of legs. Further evaluation reveals hyperparathyroidism and hyperphosphatemia. What is the initial treatment of choice in this patient?

a) Sevelamer

b) Cinacalcet

Sold by @itachibot

c) Pamidronate

d) Calcitriol

Question 24:

Which of the following is the most appropriate definitive treatment for anemia in patients with chronic kidney disease?

a) IV iron sucrose

b) Darbepoetin

c) Blood transfusion

d) Oral iron supplements

Question 25:

Which of the following should not be used in a patient with CKD?

a) Ramipril

b) Diclofenac

c) Loop diuretics

d) Insulin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 26:

A 50-year-old man with type 2 diabetes has developed stage 4 CKD secondary to diabetic nephropathy. He currently takes insulin for his diabetes. What changes need to be made to his regimen?

a) No adjustment required

b) Decreased insulin dose

c) Increased insulin dose

d) Add DPP 4 inhibitor

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	d
2	a
3	a
4	c
5	c
6	b
7	d
8	b
9	d
10	b
11	c
12	c
13	d
14	a
15	a
16	d
17	c
18	c
19	a
20	a
21	d
22	b
23	a
24	b
25	b
26	b

Detailed Explanations

Solution to Question 1:

Small for gestational age (SGA) at birth is a risk factor for chronic kidney disease (CKD), not LGA.

Important risk factors for CKD or chronic renal failure are:

- SGA
- Childhood obesity

Sold by @itachibot

- Hypertension
- Diabetes mellitus
- Autoimmune disease
- Family history of kidney disease
- Previous episode of acute kidney injury
- Structural abnormalities of the urinary tract
- Autosomal dominant polycystic kidney disease

Solution to Question 2:

Diabetic nephropathy is the most common cause of chronic kidney disease (CKD).

The leading causes of CKD in decreasing order of frequency are as follows:

- Diabetic nephropathy
- Glomerulonephritis
- Hypertension-associated CKD
- ADPKD (Adult or autosomal dominant polycystic kidney disease)

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 3:

Post-streptococcal glomerulonephritis (PSGN) is least likely to cause chronic renal failure. It is a rapidly progressive condition and a cause of acute renal failure.

Solution to Question 4:

Stages of chronic kidney disease (CKD) are stratified by estimated GFR (eGFR) and degree of albuminuria in order to predict the risk of progression of CKD. Serum creatinine is not directly used in the staging of chronic kidney disease.

Stages of Chronic Kidney Disease based on GFR		
Stage	GFR(ml/min/1.73sq. mt)	Description
1	≥90	Normal or increased GFR with other evidence of Kidney Damage

Sold by @itachibot

Stages of Chronic Kidney Dis ease based on GFR		
2	60-89	Slight decrease in GFR, with other evidence of kidney dam age
3A	45-59	Moderate decrease in GFR wi th or without other evidence of Kidney Damage
3B	30-44	
4	15-29	Severe decrease in GFR, with or without other evidence of kidney damage
5	<15	Established renal failure, also known asEnd-Stage Renal D isease



Solution to Question 5:

The features of kidney failure become prominent as the GFR progressively declines to stages 3 and 4. The following complications occur as a result of loss of kidney function:

- Volume overload
- Hyperkalemia
- Metabolic acidosis
- Hyperphosphatemia
- Mineral and bone disorders
- Hypertension

Sold by @itachibot

• Anemia

• Dyslipidemia

Solution to Question 6:

Clinical features of uremia start to appear in patients with GFR < 15 mL/min/1.73 m2 or end-stage renal disease (ESRD). By this stage, renal clearance is severely impaired and the accumulation of uremic toxins leads to several complications including:

• Uremic encephalopathy

• Pruritus

• Malnutrition

• Anorexia, nausea, fatigue

• Platelet dysfunction

• Sexual dysfunction

• Pericarditis, pleuritis

• Peripheral neuropathy

Sold by @itachibot
this from someone else,
scammed.

Stages of Chronic Kidney Disease

Stage	GFR(ml/ min/ 1.73 m2)	Description
1	≥90	Normal or increased GFR with other evidence of Kidney Damage
2	60-89	A slight decrease in GFR, with other evidence of kidney damage
3A	45-59	Moderate decrease in GFR with or without other evidence of Kidney Damage
3B	30-44	
4	15-29	Severe decrease in GFR, with or without other evidence of kidney damage
5	<15	End-stage renal disease

Sold by @itachibot



Sold by @itachi... If you purchased this from some... you may have been scammed. ...else,

Solution to Question 7:

Uremic syndrome cannot be eliminated by removing urea alone. It is a complex condition occurring due to the accumulation of multiple toxins.

The pathophysiology of uremic syndrome involves the following:

• Accumulation of toxins (uremia) - urea, creatinine, other products of protein metabolism like urates, hippurates, polyamines, indoles.

• Loss of kidney function leading to fluid and electrolyte imbalance and hormone dysregulation.

• Progressive systemic inflammation with vascular and nutritional impairment.

Clinical features secondary to loss of kidney function tend to develop in parallel with declining GFR. However the features of uremia and, hence, uremic syndrome, usually occur only in stage 5 or ESRD.

Solution to Question 8:

This clinical scenario in a CKD patient is suggestive of adynamic bone disease.

In CKD, metabolic bone disorders (renal osteodystrophy) can be of three types:

Osteitis fibrosa cystica	Osteomalacia	Adynamic bone disease
High bone turnover	Low bone turnover	Low bone turnover
Renal calcium loss	Decreased calcitriol formatio n leads to low calcium	It is aniatrogenicconditionInc reased intake ofvitamin Dorc alcium-containingphosphate binders or dialysis.

Sold by @itachibot

Osteitis fibrosa cystica	Osteomalacia	Adynamic bone disease
High PTH occurs in response to low calcium (secondary hyperparathyroidism)	Low vitamin D leads todefective mineralizationDecreased calcium leads to a mild increase in PTH, which stimulates osteoblasts. Thus thebone volumeismaintained.	High vitamin D decreases PTH levels, which leads todecreased bone volume.Mineralizationisdefectivebecause of aluminium, which comes from dialysis.
Treatment:Vitamin D + calcium	Treatment:Vitamin D + calcium	Treatment:Withdrawal of offending agent

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

Sickle cell anemia is not a feature of chronic kidney disease (CKD).

Common causes of anemia in CKD include:

- Erythropoietin deficiency
- Diminished red blood cell survival
- Bleeding diathesis
- Iron deficiency
- Vitamin B12 and folate deficiency
- Hyperparathyroidism (due to bone marrow fibrosis)
- Hemoglobinopathy

Solution to Question 10:

Hyperthyroidism is not a feature of chronic renal failure.

Option A: Peripheral neuropathy can occur secondary to uremia.

Options C and D: In chronic renal failure, calcitriol (vitamin D3) production is decreased, leading to metabolic bone disease, which in turn leads to low serum calcium levels, secondary hyperparathyroidism, and osteomalacia.

Solution to Question 11:

The most common neurological disorder seen in patients with CKD is peripheral neuropathy.

The various neuromuscular manifestations of CKD include:

- Memory and concentration impairment
- Dialysis dementia - due to aluminum intoxication

Sold by @itachibot

- Restless leg syndrome
- Peripheral neuropathy
- Hiccups
- Cramps
- Uremic encephalopathy
- Confusion and lethargy
- Tremors
- Myoclonus
- Asterixis
- Seizures
- Coma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

This clinical scenario is suggestive of restless legs syndrome (RLS) or Willis-Ekbom disease. Chronic kidney disease (CKD) is not the most common cause of this condition.

RLS is a neurological sleep disorder that can be primary or secondary. The primary form is more common. It shows autosomal dominant inheritance in most cases, but can be idiopathic. Secondary RLS is associated with the following:

- Iron deficiency
- Pregnancy - most common movement disorder
- Peripheral neuropathy
- Diabetes mellitus
- CKD
- Parkinson's disease
- Multiple sclerosis

It is characterized by an uncontrollable, restless urge to move the legs. It tends to worsen during periods of inactivity and at evening or night. It is accompanied by crawling, tingling or burning sensations. These symptoms show partial or complete improvement with movement. This cycle may continue for a while and result in insomnia.

Primary RLS is managed by improving sleep hygiene and dopamine agonists (pramipexole or ropinirole) if symptoms persist. Iron supplements can also be given. In secondary RLS, the underlying cause must be treated.

Solution to Question 13:

Sold by @itachibot

The most likely cause of convulsions in a patient with CKD is water intoxication. Volume overload in these patients as well as damage to the blood-brain barrier by uremic toxins causes cerebral edema and seizures.

Solution to Question 14:

Cardiovascular disease is the leading cause of morbidity and mortality in patients at every stage of CKD.

Solution to Question 15:

Chronic kidney disease (CKD) is associated with elevated levels of fibrinogen.

CKD is a pro-inflammatory state characterized by altered levels of acute-phase reactants (APR)

- Elevated levels of positive APR like CRP and fibrinogen (associated with high ESR).
- Reduced levels of negative APR like albumin and fetuin.

Solution to Question 16:

High anion gap metabolic acidosis is the predominant electrolyte abnormality in a patient with advanced CKD. In advanced CKD, impaired renal functioning causes decreased urinary excretion of acids. These retained organic acids result in a high anion gap.

In early CKD, potassium retention is associated with bicarbonate loss. This gives rise to hyperchloremic metabolic acidosis with a normal anion gap in early CKD.

The following electrolyte abnormalities are seen in CKD:

- Hyperkalemia
- Hypocalcemia
- Hyperphosphatemia
- Hyperuricemia
- Hypermagnesemia

Hypokalemia is uncommon in CKD and usually reflects markedly reduced dietary potassium intake, especially in association with excessive diuretic therapy or concurrent GI losses.

Solution to Question 17:

Of the given options, hyperkalemia is the most life-threatening complication of chronic kidney disease (CKD) as it can result in cardiac arrhythmias and sudden death.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

CKD is staged based on declining glomerular filtration rate (GFR) and albuminuria. The various clinical manifestations occur due to loss of kidney function and accumulation of various uremic toxins (urea, creatinine, and other products of protein metabolism).

Complications of declining kidney function include:

- Volume overload
- Hyperkalemia
- Metabolic acidosis
- Hyperphosphatemia
- Mineral and bone disorders
- Hypertension
- Anemia
- Dyslipidemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

Serum estrogen levels are low in CKD patients. This leads to menstrual abnormalities and infertility.

Option A: Plasma levels of insulin are elevated in CKD patients due to impaired clearance by the affected kidneys. Hence, insulin dose must be reduced in these patients.

Option B: Elevated parathyroid hormone levels in CKD are secondary to derangements in the homeostasis of calcium, phosphate, and vitamin D.

Option D: FGF-23 (fibroblast growth factor) is part of a family of phosphatonins that promotes renal phosphate excretion. The levels of this hormone increase early in the course of CKD.

Solution to Question 19:

This clinical scenario is suggestive of acute myocardial infarction (MI) in a patient with chronic kidney disease (CKD). A serial rise in cardiac troponin levels over 3-6 hours following presentation is needed to make a diagnosis. An interval of 1-hour is not considered sufficient.

Minor chronic elevations in serum troponins are commonly observed in patients with CKD even in the absence of cardiovascular disease. Hence, a single elevated troponin value in such patients cannot be used to diagnose MI.

CK-MB is far less sensitive and specific than troponins. Hence, it is not preferred.

Solution to Question 20:

Sold by @itachibot

Microalbuminuria is the most sensitive and earliest finding of diabetic nephropathy. It refers to urine albumin of 30–300 mg/24-hour urine. It is seen in the stage of incipient diabetic kidney disease.

However, microalbuminuria is a very variable finding and is not required to diagnose nephropathy. Non-albuminuric diabetic nephropathy may occur in some patients and regression of albuminuria may be seen in type 1 diabetics.

Diabetic nephropathy can be diagnosed if the following are satisfied:

• Reduced GFR (< 60 mL/min/1.73 m2) with or without persistent albuminuria (> 30 mg/day or urine PCR > 30 mg/g)

• Long duration of diabetes or established diabetic retinopathy

• Other etiologies are ruled out



Solution to Question 21:

Hypertensive nephropathy is unlikely in this patient as it usually presents with shrunken kidneys.

Longstanding CKD due to most causes results in bilaterally small kidneys. Causes of CKD with normal size kidneys are:

• Diabetic nephropathy (kidney size may be increased at the onset)

• Amyloidosis

• HIV nephropathy

Solution to Question 22:

Sold by @itachibot

According to the ESC 2018 guidelines, the target blood pressure in hypertensive patients with CKD is < 140/ 80 mm Hg regardless of age.

Solution to Question 23:

This clinical scenario is suggestive of secondary hyperparathyroidism (HPT) and hyperphosphatemia in a patient with CKD. Sevelamer is the best initial treatment for this patient. It is a phosphate binder that reduces the absorption of phosphate from the gastrointestinal tract.

Other phosphate binders that can be used include:

• Calcium containing

• Calcium acetate

• Calcium carbonate

• Non-calcium containing

• Sevelamer

• Lanthanum

Calcitriol and cinacalcet are used to decrease PTH levels in secondary hyperparathyroidism. However, they also cause increased serum phosphate levels. Hence, they should not be given as initial treatment in patients with hyperphosphatemia.

Bisphosphonates like pamidronate are used in the treatment of primary hyperparathyroidism.

Solution to Question 24:

Anemia in patients with chronic kidney disease (CKD) is primarily due to decreased erythropoietin production. Hence, human recombinant erythropoietin (EPO) such as darbepoetin is the most appropriate treatment and is considered the mainstay.

However, an optimal response to treatment with EPO occurs only when iron stores are adequate. Hence, iron supplements (oral or IV) must be given first to ensure the effectiveness of EPO therapy.

Other treatment options for anemia in CKD are:

• Vitamin B12 and folate supplementation

• Blood transfusion - in severe anemia and failure to EPO therapy

Solution to Question 25:

Diclofenac should not be used in a patient with CKD as it is nephrotoxic.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Renal blood flow is regulated by prostaglandin E2. Aspirin, diclofenac and other NSAIDs inhibit cyclooxygenase. This decreases prostaglandin E2 concentration, thereby causing a reduction in blood flow and GFR.

Nephrotoxic drugs include the following:

- NSAIDs
- Antifungals (amphotericin B)
- Antibiotics
- Aminoglycosides
- Vancomycin
- Cephalosporins
- Sulfonamides
- Streptomycin
- Antivirals
- Acyclovir
- Foscarnet
- Chemotherapy drugs
- Methotrexate
- Platins
- Immunosuppressants (cyclosporine)

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 26:

In CKD patients with diabetes, the dose of insulin must be reduced.

In CKD patients, the plasma levels of insulin are elevated due to diminished renal degradation of insulin by the impaired kidney. This impaired renal clearance of insulin can lead to hypoglycemia. Hence, the dose of insulin must be reduced to prevent this.

Sold by @itachibot

Acute Kidney Injury

Question 1:

Which of the following criteria is diagnostic of acute kidney injury in a patient?

a) Rise in creatinine 1.5 times the baseline value within 7 days

b) Rise in creatinine ≥ 0.3mg/dl from baseline within 7 days

c) Rise in creatinine ≥ 0.5 mg/dl from baseline within 48 hours

d) Urine volume <0.5 mL/kg/h for 24 hours

Question 2:

AKIN and RIFLE criteria are used to classify

a) Acute kidney injury

b) Chronic renal failure

c) Acute glomerulonephritis

d) Nephrotic syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Patients with urine output below what level are said to have anuria?

a) 50 mL/day

b) 70 mL/day

c) 150 ml/day

d) 100 mL/day

Question 4:

Which of the following causes prerenal AKI?

a) 1, 3, and 4

b) 1, 2 and 4

c) 2, 3 and 5

Sold by @itachibot

d) 3, 4 and 5

Question 5:

A 48-year-old woman presented with severe abdominal pain, vomiting, and dark-coloured urine for the past 3 days. She has been on a combination painkiller for rheumatoid arthritis for 10 years. On examination, severe pallor was noted and BP was 180/120 mmHg. CT abdomen is shown below. What is the diagnosis?



a) Septic acute tubular necrosis

b) Chronic kidney disease

c) NSAID-induced nephropathy

d) Acute interstitial nephritis

Question 6:

A 38-year-old woman presents with nausea, vomiting, fatigue, and oliguria for the past three days. On further investigation, urine osmolality was 800 mOsm/kg, urinary sodium 10 meq/L, and BUN: creatinine ratio was 20:1. What is the most likely diagnosis?

a) Pre-renal AKI

b) Renal AKI

c) Post renal AKI

d) Acute cortical necrosis

Sold by @itachibot

Question 7:

A 35-year-old man was admitted following a crush injury to his left leg. Two days later, his urine/plasma creatinine levels are < 20. What is the likely cause?

a) Rhabdomyolysis

b) Congestive heart failure

c) Hepatorenal syndrome

d) Hypovolemia

Question 8:

A 38-year old woman presents with severe muscle pain and high-coloured urine for the past 3 days. She joined a gym 2 weeks ago and has been strenuously working out. She was diagnosed with hypothyroidism 8 years back. Work-up reveals high serum creatine kinase, uric acid, and creatinine. Which of the following is likely to be present in this patient?

a) Hypophosphatemia

b) Hypomagnesemia

c) Hypocalcemia

d) Hypokalemia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 54-year-old man presented with reduced urine output, swelling of his feet, and eyelid puffiness for 1 week. He also complains of back pain for the past 3 months, unresponsive to painkillers. Investigations reveal the following. What is the possible cause of renal failure in this patient?

a) Hypokalemia

b) Hyperkalemia

c) Hypercalcemia

d) Hypocalcemia

Question 10:

Non-oliguric renal failure may be seen in patients with which of the following conditions?

a) 2, 3, and 5

Sold by @itachibot

b) 2 and 3

c) 1, 3, and 5

d) 2, 4, and 5

Question 11:

A 47-year-old electrician was rushed to the hospital following an electrical shock. Examination reveals electrical burns. During the course of treatment, he developed an arrhythmia and was managed appropriately. What arterial blood gas finding would you expect in this patient?

a) Hyperkalemic alkalosis

b) Hypokalemic alkalosis

c) Hyperkalemic acidosis

d) Hypokalemic acidosis

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 12:

A patient with acute renal failure was found to have metabolic acidosis on evaluation. What is the likely cause?

a) Decreased excretion of ammonium

b) Loss of HCO_3

c) Increased aldosterone levels

d) Increased production of H^+

Question 13:

A 75-year-old obese woman, who is a known diabetic presented with chest pain. Coronary angiography was done. Her S.creatinine levels before and after the procedure are given below. Which of the following is not a risk factor for the complication she developed?

a) Age

b) Diabetic nephropathy

c) Recent myocardial infarction

d) Obesity

Sold by @itachibot

Question 14:

Which of the following is false regarding contrast-induced nephropathy?

a) It resolves within a week

b) N-acetyl cysteine can be used

c) Dialysis-requiring AKI is very common

d) Hypoxia due to nephrotoxic agents is the cause

Question 15:

Which of the following is not a marker of kidney injury?

a) Osteopontin

b) Cystatin C

c) Alanine aminopeptidase

d) Acid phosphatase

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

Which intravenous fluid is preferred for the hydration of patients with pre-renal acute kidney injury?

a) Hydroxyethyl starch

b) Dextran

c) 0.9% NaCl

d) Ringer's lactate

Question 17:

Continuous renal replacement therapy is preferred in patients with which of the following features?

a) Severe hypotension

b) Hyperkalemia

c) Metabolic acidosis

d) Uremic symptoms

Sold by @itachibot

Question 18:

Which of the following is not seen in tumor lysis syndrome?

a) Hyperphosphatemia

b) Hyperuricemia

c) Hypercalcemia

d) Hyperkalemia

If you bought this from anyone else, it is stolen.

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	d
5	b
6	a
7	a
8	c
9	c
10	a
11	c
12	a
13	d
14	c
15	d
16	c
17	a
18	c

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

A rise in creatinine 1.5 times the baseline value within 7 days is diagnostic of acute kidney injury (AKI).

AKI is an abrupt decrease in kidney function within hours to days. According to the KDIGO definition, it is defined as any of the following:

• Increase in serum creatinine by ≥0.3 mg/dL within 48 hours OR

• Increase in serum creatinine to ≥1.5 times the baseline within 7 days OR

• Urine volume <0.5 mL/kg/h for 6 hours.

Solution to Question 2:

AKIN and RIFLE criteria are used to classify acute kidney injury (AKI).

The various classifications used in AKI are:

• RIFLE criteria (Risk, Injury, Failure, Loss, End-stage)

• AKIN (AKI Network) Classification

• KDIGO (Kidney Disease Improving Global Outcomes) Classification

RIFLE, AKIN, KDIGO definitions of AKI:

RIFLE, AKIN, KDIGO staging of AKI:

Sold by @itachibot
Stolen this from someone else,
it has been scammed.

	RIFLE	AKIN	KDIGO
S. creatinine	>50% increase that develops in <7 days	>0.3 mg/dl increase or >50% increase that develops in <48 hr	>0.3 mg/dl increase or >50% increase that develops in <7 days
Urine output	<0.5ml/kg/hr for >6 hr	<0.5ml/kg/hr for >6 hr	<0.5ml/kg/hr for >6 hr

RIFLE	Increase in S.creatinine	AKIN	Increase in S.creatinine	KDIGO	Increase in S.creatinine	Urine output
Risk	≥50%	Stage 1	≥0.3mg/dl or ≥50%	Stage 1	≥0.3mg/dl or ≥50%	<0.5mL/kg/hr for >6h
Injury	≥100%	Stage 2	≥100%	Stage 2	≥100%	<0.5mL/kg/hr for >12 hr
Failure	≥200%	Stage 3	≥200%	Stage 3	≥200%	<0.3mL/kg/hr for >24hr or anuria >12hr
Loss	Need for renal replacement therapy (RRT)>4 weeks					
End-stage	Need RRT for >3 months					

Sold by @itachibot

Solution to Question 3:

Anuria is defined as urine output less than 100 mL/day. Oliguria is defined as urine output less than 400 mL/day.

Acute kidney injury (AKI) can have varied clinical features depending on the etiology. The history and physical examination in AKI aims mainly at:

• Finding the etiology

• Identifying the occurrence of any complications

The following clinical features may be seen:

• Oliguria or anuria

• Gross hematuria

• Renal colic pain (in renal stones)

• Loss of skin turgor, dry mouth, hypotension, tachycardia (in hypovolemia)

• Prostatomegaly and/or a pelvic mass (in post-renal obstruction)

• Dysuria, fever with chills and rigors, increased frequency of micturition, renal angle tenderness (in pyelonephritis)

• Edema, hypertension (volume overload)

• Encephalopathy, seizures, platelet dysfunction (uremia)

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 4:

NSAIDs, Rhabdomyolysis and ACE inhibitors cause prerenal acute kidney injury (AKI).

The use of NSAIDs causes reduced production of vasodilatory prostaglandins. This leads to decreased renal perfusion and glomerular capillary pressure. NSAIDs can also lead to acute interstitial nephritis, a cause of renal AKI. However, this is relatively uncommon.

ACE inhibitors cause inhibition of efferent arteriole vasoconstriction, leading to reduced renal perfusion.

Rhabdomyolysis causes AKI by both prerenal mechanisms (due to volume sequestration) and ATN (due to precipitation of myoglobin in the renal tubules).

Nephrotoxins like myeloma light chains, snake venom, myoglobin (due to rhabdomyolysis) cause acute tubular necrosis (ATN). This leads to renal or intrinsic AKI.

Sold by @itachibot



Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

This clinical scenario and CT scan showing papillary calcifications are highly suggestive of chronic kidney disease caused by analgesic nephropathy.

This condition is more common in females secondary to excessive prolonged use of a combination of analgesics. The mechanism of injury is medullary hypoxia leading to papillary necrosis and finally causes chronic tubulointerstitial nephritis.

Patients can remain asymptomatic for a long time. Sloughing or obstruction of papillae can cause abdominal pain and hematuria. It increases the risk of urothelial malignancies. The following CT findings can be seen:

• Papillary calcifications

• Small kidneys

• Irregular contour

Note: NSAID-induced nephropathy causes acute interstitial nephritis. It is a type IV hypersensitivity reaction. The features include non-oliguric AKI, fractional excretion of sodium (FeNa) < 1, and hyperkalemia. It is seen within a few days of use and is reversible.

The image below shows papillary calcifications:

Sold by @itachibot



The given image shows papillary calcification on ultrasonography:



Solution to Question 6:

The given clinical scenario is highly suggestive of prerenal AKI based on the following:

• Urine osmolality >500 mOsm/kg

• Urinary sodium < 20 meq/l

• BUN: Creatinine ratio of 20:1

Differences between prerenal, and renal AKI:

Sold by @itachibot

Indices	Prerenal	Renal
Spot urine sodium (mEq/L)	<20	>40
Fractional excretion of sodium (FENa)	<1%	>2%
Fractional excretion of urea (FEUrea)	<35%	>35%
Urine osmolality (mOsm/kg)	>500	<350
BUN to serum creatinine ratio	>20:1	<10:1
Urine creatinine/plasma creatinine	>40:1	<20:1
Urinary sediment	Nil or hyaline casts only	Renal tubular epithelial casts, muddy brown granular casts

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

This clinical scenario and urine/ plasma creatinine ratio of <20 is suggestive of renal AKI due to rhabdomyolysis-induced acute tubular necrosis.

Rhabdomyolysis is a serious complication of crush injuries. It leads to muscle breakdown, releasing a high amount of myoglobin in the blood. Myoglobinuria can lead to acute kidney injury by the following mechanisms:

- Direct toxic effect of myoglobin and heme on the PCT
- Tubular obstruction by precipitation of pigmented casts
- Renal vasoconstriction can cause FeNa < 1 (unlike other renal causes of AKI)

Indices	Prerenal	Renal
Spot urine sodium (mEq/L)	<20	>40
Fractional excretion of sodium (FENa)	<1%	>2%
Fractional excretion of urea (FEUrea)	<35%	>35%
Urine osmolality (mOsm/kg)	>500	<350
BUN to serum creatinine ratio	>20:1	<10:1
Urine creatinine/plasma creatinine	>40:1	<20:1
Urinary sediment	Nil or hyaline casts only	Renal tubular epithelial casts, muddy brown granular casts

Sold by @itachibot



Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 8:

The given clinical scenario is suggestive of rhabdomyolysis. Hypocalcemia is seen in this condition.

The causes of rhabdomyolysis include:

• Hypothyroidism (precipitated by exertion, alcohol, infection)

• Crush injury

• Electric shock

• Prolonged seizures

• Malignant hyperthermia

• Exertion

• Drugs like fibrates, statins

• Polymyositis / Dermatomyositis

• McArdle's disease (GSD-5) causes recurrent rhabdomyolysis.

Patients often present with muscle pain and high-colored urine. The following features are seen:

• High serum CK (> 5 times normal)

• High serum myoglobin

• High serum LDH

• Hyperkalemia

• Hypermagnesemia

• Hyperphosphatemia

• Hypocalcemia

Sold by @itachibot

• Hyperuricemia

• Once AKI develops, high serum creatinine and urea

The treatment includes administration of large amounts of IV fluids (5-6 liters), alkalization of urine, and hemodialysis.

Solution to Question 9:

Bence-Jones proteins on urine analysis are suggestive of multiple myeloma. It causes acute kidney injury (AKI) by the following mechanisms:

• Hypercalcemia - leads to intense renal vasoconstriction and reduced perfusion

• Myeloma light chains - cause direct tubular toxicity.

Solution to Question 10:

Non-oliguric renal failure may be seen in the following conditions:

• Acute interstitial nephritis

• Acute tubular necrosis (aminoglycoside toxicity, contrast-induced nephropathy)

All the causes of acute tubular necrosis (ATN) cause non-oliguric AKI except acyclovir (oliguric AKI).

The difference in urine output between oliguric and non-oliguric AKI may be due to variations in GFR or in the rate of tubular reabsorption. Non-oliguric forms may be associated with less severe injury.

Other options: Prerenal causes such as congestive heart failure cause oliguric AKI. In post-streptococcal glomerulonephritis, oliguria occurs as a result of decreased GFR.

Solution to Question 11:

Electrical burns result in rhabdomyolysis. It presents with hyperkalemic metabolic acidosis.

Muscle breakdown in rhabdomyolysis leads to hyperkalemia which causes arrhythmias. It also causes high anion gap metabolic acidosis due to the increase in H+ ions and liberation of organic anions.

Solution to Question 12:

In acute renal failure, metabolic acidosis occurs due to decreased excretion of ammonium. This impairs the excretion of H+ ions as NH_4 ions.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Normally, H+ are excreted via the kidney in combination with ammonia which acts as a urinary buffer.

In AKI, there is a decreased capacity of tubules to secrete ammonia. Due to reduced amounts of ammonia, the production of NH4 is impaired. Hence, H+ ions cannot be excreted, which leads to metabolic acidosis.

Solution to Question 13:

The given clinical scenario shows >0.5 mg/dL increase in S.creatinine levels, which is indicative of contrast-induced nephropathy for which obesity is not a risk factor.

Contrast-induced nephropathy is defined as a 25% or 0.5 mg/dL increase from the baseline value of serum creatinine within 48-72 hours of intravenous contrast administration. It occurs due to tubular damage and as a result, the patient develops non-oliguric transient nephropathy.

Risk factors for contrast medium-induced nephropathy:

Patient-related:

• eGFR < 60 mL/min/1.73 m2 before intra-arterial administration

• eGFR < 45 mL/min/1.73 m2 before intravenous administration

• Patient conditions that increase the risk:

• Diabetic nephropathy

• Dehydration

• Congestive heart failure (NYHA grade 3–4) and low LVEF

• Recent myocardial infarction (<24 h)

• Intra-aortic balloon pump

• Peri-procedural hypotension

• Low hematocrit level

• Age over 70

• Concurrent administration of nephrotoxic drugs

• Known or suspected acute renal failure

Procedure-related:

• Intra-arterial administration of contrast medium

• High-osmolality agents

• Large doses of contrast medium

• Multiple contrast medium administrations within a few days

Side effects of iodinated contrast material include:

• Idiosyncratic anaphylactoid reaction - patient develops bronchospasm and hypotension

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• Dose-dependent side effect – contrast-induced nephropathy seen especially with the use of high osmolar agents

Prevention of contrast-induced nephropathy:

• IV fluids (isotonic saline or sodium bicarbonate) started 1 hour before and continued for at least 3-6 hours after the procedure

• Low volume low osmolar contrast should be used

• N-acetyl cysteine may be given prior to contrast exposure

Solution to Question 14:

Dialysis-requiring AKI is uncommon in contrast-induced nephropathy.

Contrast-induced AKI is generally associated with the use of high osmolar contrasts such as:

• Iodinated contrast agents

• High-dose gadolinium

Mechanism of contrast-induced injury includes:

• Medullary ischemia and hypoxia: By inducing intense vasoconstriction

• Toxic tubular damage: By direct injury

It usually develops within 24-48 hrs. Serum creatinine peaks by 3-5 days and normalizes by 7-10 days. It manifests mainly as a non-oliguric renal failure and has an excellent prognosis.

Contrast-induced AKI can be prevented by:

• Use of iso-osmolar or non-ionic contrast agents (least toxic)

• Administering minimum required quantity of contrast

• Pre- and post-hydration (IV normal saline 1 mL/kg/h x 12 h prior and post-exposure)

• N-acetyl cysteine may be given prior to contrast exposure.

Solution to Question 15:

Acid phosphatase is not a marker of kidney injury.

The various markers of kidney injury are as follows:

• IL-18

• Cystatin C

• NGAL (neutrophil gelatinase-associated lipocalin)

• KIM-1 (kidney injury molecule-1)

• LFABP (liver fatty acid-binding protein)

•

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

• NHE3 (sodium hydrogen exchanger 3)

• NAG (N-acetyl glucosaminidase)

• Alanine aminopeptidase

• Osteopontin

• ALP

Solution to Question 16:

Isotonic crystalloids like 0.9% NaCl are preferred for the correction of hypovolemia in pre-renal AKI.

Management of AKI:

1. AKI care bundle:

• Nephrotoxic drugs should be avoided and the dosage of other drugs must be adjusted according to kidney function

• Creatinine levels and urine output must be closely monitored

2. Hemodynamic management:

• If hypovolemic - fluid replenishment using isotonic crystalloids (normal saline)

• If euvolemic - fluid intake = urine output + additional ~500 mL

• If hypervolemic - diuretics are administered. If the response is unsatisfactory, dialysis is required.

3. Management of complications:

• Pulmonary edema - Treated with diuretics, non-invasive ventilation, and dialysis if refractory

• Metabolic acidosis - Treated when pH <7.2 with oral or IV sodium bicarbonate

• Hyperkalemia - Treated by restriction of dietary potassium, administration of sodium polystyrene sulfonate, insulin, glucose, inhaled β-agonist, and calcium gluconate

Options A and B: Hydroxyethyl starch solutions and dextrans increase the risk of severe AKI and are contraindicated.

Option D: Ringer's lactate is preferred only when a large volume of fluids is required.

Solution to Question 17:

Continuous renal replacement therapy (CRRT) is a modality of dialysis that involves hemodialysis and hemofiltration. It is the preferred option in AKI patients with severe hypotension.

It is preferred in the following cases:

• Acute renal failure complicated by hemodynamic instability or heart failure

• Severe volume overload unresponsive to diuretic therapy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Brain edema

• Hypercatabolism state

• Acute or chronic liver failure as a renal support measure

It is an extracorporeal blood purification therapy intended to substitute for impaired renal function over an extended period of time, aimed at being applied for 24 hours a day. There is no rapid shift of volume and osmolarity with this process as in intermittent hemodialysis.

The other types of dialysis are:

• Intermittent hemodialysis - may be done in hemodynamically stable patients with AKI

• Peritoneal dialysis - used when hemodialysis is not available.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

Hypocalcemia, not hypercalcemia, is seen in tumor lysis syndrome (TLS).

TLS is a potentially life-threatening condition characterized by the rapid destruction of a large number of tumor cells. The release of calcium phosphate from lysed cells contributes to hyperphosphatemia and subsequent reciprocal depression in serum calcium levels. The consumption of ionized calcium during the formation of calcium phosphate crystals leads to hypocalcemia.

TLS is commonly observed after initiation of chemotherapy in hematologic malignancies characterized by high cell turnover rates, such as Burkitt's lymphoma and acute lymphoblastic leukemia.

The destruction of neoplastic cells results in the release of nucleic acids, which are broken down into uric acid, leading to hyperuricemia. The acidic environment in the renal tubules can cause uric acid to precipitate, leading to acute renal failure. Potassium is released in large quantities during tumor cell lysis leading to hyperkalemia.

TLS is diagnosed using the Cairo-Bishop criteria which includes both laboratory and clinical criteria.

Prevention is the mainstay of the management of TLS, which involves includes allopurinol, which inhibits the production of uric acid, and aggressive hydration to promote urine flow. Urinary alkalization is not recommended. Febuxostat, a selective xanthine oxidase inhibitor, can be used as an alternative.

In severe cases or when renal failure is present, rasburicase, a recombinant urate oxidase, is used for treatment. Rasburicase rapidly lowers uric acid levels by converting it into a highly soluble compound called allantoin. Renal replacement therapy is often required and should be considered in case of acute renal failure.

Sold by @itachibot

Renal Replacement Therapy

Question 1:

Dialysis is not an absolute indication for patients who present with which of the following?

a) Asterixis

b) Pericarditis

c) Uremic encephalopathy

d) Hyperkalemia

Question 2:

Which of the following is the most commonly used therapeutic modality for patients with end-stage renal disease?

a) Hemodialysis

b) Renal transplant

c) Continuous ambulatory peritoneal dialysis

d) Continuous cycling peritoneal dialysis

Question 3:

Which of the following is incorrect about hemodialysis?■

a) Dialyzers are made up of cellulose membranes

b) Bicarbonate is commonly used as a buffer

c) Glutaraldehyde is used as a reprocessing agent

d) Urea is cleared more efficiently than creatinine

Question 4:

A 50-year-old man with stage 5 chronic kidney disease is planned for hemodialysis. Which of the following is incorrect regarding the vascular access for this procedure?

a) Tunneled catheters are associated with lower infection rates

b) Access of choice is radiocephalic arteriovenous fistula

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

c) Most common complication of arteriovenous grafts is infection

d) Fistulas have higher long term patency rates

Question 5:

What is the rate of blood flow in the hemodialysis machine?

a) 150-250 mL/min

b) 200-300 mL/min

c) 250-450 mL/min

d) 500-800 mL/min

Question 6:

In a patient undergoing hemodialysis, which of the following can be used to monitor the adequacy of dialysis?

a) Urea reduction ratio

b) Blood urea / creatinine ratio

c) Urinary protein / creatinine ratio

d) Creatinine reduction ratio

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following has been most commonly implicated as a cause of dialysis-induced hypotension in patients?

a) Bicarbonate

b) Aluminium

c) Acetate

d) Phosphate

Question 8:

What is the most common acute complication in patients who undergo hemodialysis?

a) Anaphylaxis

Sold by @itachibot

b) Dialysis disquillibrium syndrome

c) Hypotension

d) Muscle cramps

Question 9:

A 61-year-old man is brought to the casualty following a seizure episode. His wife says that he has been very drowsy and has not passed urine for the past 2 days. Investigations reveal serum urea of 71 mg/dl and creatinine of 5.6 mg/dl. He is referred for urgent hemodialysis. You would take all of the following precautions except:

a) Warming the dialysate before dialysis

b) Removing more fluid during the beginning than the end of the procedure

c) Maintaining fluid removal rate below 13 mL/kg/hour

d) Careful evaluation of dry weight

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A 67-year-old woman with CKD underwent her first session of hemodialysis. Towards the end of the session, she became drowsy and had a sudden seizure episode. What is the likely cause for this presentation?

a) Hypotension

b) Aluminium toxicity

c) Anaphylaxis

d) Reverse urea effect

Question 11:

Which of the following is true about anaphylaxis associated with hemodialysis?

a) Occur due to lack of cellulose in the membranes

b) Ethylene oxide is the cause of Type B reaction

c) Dialysis is discontinued in Type A reaction

d) Type A reaction associated with complement activation

Question 12:

Sold by @itachibot

A patient on hemodialysis may develop all of the following chronic complications except:

a) Amyloidosis

b) Dementia

c) Bone disease

d) Hypovolemia

Question 13:

A physician is evaluating a patient with carpal tunnel syndrome. He has been undergoing dialysis for the past 7 years. Deposition of which of the following is responsible for this condition?

a) AA

b) Amyloid beta 1

c) Amyloid beta 2

d) Amyloid transthyretin

Question 14:

A 57-year-old man is referred urgently for management of pulmonary edema. He was diagnosed with stage-5 CKD 3 weeks ago but was reluctant to start dialysis. On examination, he is confused and has a flapping tremor. Investigations are as follows. Which of his clinical features will not improve with dialysis?

a) Pulmonary edema

b) Metabolic acidosis

c) Uremic encephalopathy

d) Renal osteodystrophy

Question 15:

What is the most common cause of death in patients undergoing chronic dialysis?

a) Cardiovascular causes

b) Electrolyte abnormality

c) Metabolic acidosis

d) Encephalopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 16:

You are educating a patient about performing peritoneal dialysis at home. During this process, how much fluid is infused into the peritoneum?

a) 1.5-3 L

b) 2-4 L

c) 3-5 L

d) 0.5-1.5 L

Question 17:

A 56-year-old man presents with fever and severe abdominal pain for 2 days. He has been on peritoneal dialysis for the past 1 year. On examination, there is abdominal distension, guarding, and rebound tenderness. Which of the following is the most common cause of this condition?

a) Gram-positive cocci

b) Gram-positive bacilli

c) Gram-negative cocci

d) Gram-negative bacilli

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A woman is started on continuous ambulatory peritoneal dialysis. Which of the following complications is she unlikely to develop?

a) Residual uremia

b) Metabolic disturbances

c) Insulin resistance

d) Weight loss

Question 19:

Which of the following is incorrect about renal transplants?

a) ABO and HLA mismatch are relative contraindications

Sold by @itachibot

b) Contraindicated in overt AIDS and active hepatitis

c) Complications in donor include hypertension and focal segmental glomerulosclerosis

d) The most common cause of death after renal transplant is infections

Question 20:

Which of the following is incorrect about rejection reactions following a kidney transplant?

a) Acute rejection can occur due to both type 4 and type 2 hypersensitivity reactions

b) Hyperacute rejection is a type 2 hypersensitivity reaction

c) Induction therapy for prophylaxis of rejection is cyclosporine, azathioprine, prednisolone.

d) Alemtuzumab is monoclonal antibody against CD52 used for immunosuppression

Question 21:

A 45-year-old man with polycystic kidney disease underwent renal transplantation 3 months ago. Which of the following infections is he at the highest risk of?

a) Candida

b) Pneumocystis

c) BK virus

d) Cytomegalovirus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 22:

Which of the following lesions does not recur in the transplanted kidney?

a) Diabetic nephropathy

b) Membranous glomerulonephritis

c) Lupus nephritis

d) Alport syndrome

Question 23:

Patients with which of the following conditions have the highest risk of disease recurrence after renal transplant?

Sold by @itachibot

a) Lupus nephritis

b) Membranoproliferative glomerulonephritis

c) Diabetic nephropathy

d) Membranous glomerulonephritis

Question 24:

What is the average life expectancy in patients following a living-donor kidney transplant?

a) 10 years

b) 14 years

c) 18 years

d) 20 years

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	d
2	a
3	a
4	c
5	c
6	a
7	c
8	c
9	a
10	d
11	c
12	d
13	c
14	d
15	a
16	a
17	a
18	d

Sold by @itachibot

19	d
20	c
21	d
22	d
23	c
24	b

Detailed Explanations

Solution to Question 1:

Dialysis is not indicated in patients who initially present with hyperkalemia. It is indicated in the case of refractory hyperkalemia that has failed to respond to other modalities of treatment.

Indications for dialysis can be remembered using the mnemonic AEIOU:

• Acidosis - metabolic acidosis that is refractory to medical treatment.

• Electrolyte abnormality - refractory hyperkalemia and hypercalcemia.

• Intoxication - barbiturates, alcohol, and lithium.

• Overload - hypervolemia and pulmonary edema.

• Uncontrolled hypertension (and)

• Uremia - encephalopathy, pericarditis, asterixis, seizures, intractable nausea and vomiting, and bleeding diathesis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

The most common therapeutic modality for patients with end-stage renal disease (ESRD) is hemodialysis.

The various modalities of renal replacement therapy include the following:

• Dialysis:

• Acute intermittent hemodialysis

• Continuous renal replacement therapy - includes hemodialysis and ultrafiltration techniques

• Peritoneal dialysis

• CAPD - Continuous ambulatory peritoneal dialysis

• CCPD - Continuous cycling peritoneal dialysis

• NIPD - Nocturnal intermittent peritoneal dialysis

• Renal transplant

Sold by @itachibot

Solution to Question 3:

In hemodialysis, the dialyzer is made up of biocompatible membranes derived from polysulfone. Cellulose is bio-incompatible and is no longer used.

Hemodialysis is based on the principle of solute diffusion using an artificial semi-permeable membrane. Smaller molecules (urea) diffuse faster and are cleared more efficiently than larger molecules (creatinine). There are three components to this system:

• Dialyzer - a chamber that contains compartments for blood and dialysate solution, separated by a semipermeable membrane.

• Dialysate - a solution that helps in the removal of metabolic waste products. Bicarbonate is used as a buffer to correct acidosis in CKD patients.

• Blood delivery system - consists of the vascular access and extracorporeal circuit.

The dialyzers used during a session of dialysis are reprocessed and reused for the next dialysis sessions. The reprocessors used for dialyzers are:

• Glutaraldehyde

• Formaldehyde

• Peracetic acid-hydrogen peroxide

The following image depicts hemodialysis.



If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Solution to Question 4:

The most common complication of arteriovenous (AV) grafts is stenosis, not infections.

Sold by @itachibot

In order to perform hemodialysis, vascular or hemodialysis access must first be established. The preferred site is the forearm. The following types of vascular access may be used for hemodialysis:

• AV fistulas:

• Fistula refers to the anastomosis of an artery and an adequately large vein (either side-to-side or end-to-side).

• This causes arterialization of the vein, thereby causing venous wall thickening and increasing venous diameter.

• Fistulas need 4-6 weeks to mature before they can be used.

• The access of choice is radiocephalic AV fistula (Brescia-Cimino fistula).

• Fistulas have higher long-term patency rates than grafts.

• Thrombosis is a common complication.

The following image depicts a radiocephalic fistula.

• AV grafts:

• The artery and vein are connected by a prosthetic graft such as PTFE (polytetrafluoroethylene).

• The advantage is that it can be performed with smaller veins.

• Complications include stenosis (most common), due to intimal hyperplasia at anastomosis between graft and vein, and infections.

• Central venous catheter:

• Tunneled catheters are used for long-term IV access. As they tunnel through the skin before entering the vein, they have a lower risk of infection.

Solution to Question 5:

Sold by @itachibot

During hemodialysis, heparinized blood flows through the dialyzer at a rate of 250–450 mL/min. On the other hand, the dialysate flows in the opposite counter-current direction at 500–800 mL/min.

Solution to Question 6:

The adequacy of dialysis is determined by the urea reduction ratio. It is defined as the fractional reduction in blood urea nitrogen per hemodialysis session. It is calculated as (Co - C) / Co, where:

Co - predialysis serum urea

C - postdialysis serum urea

The target urea reduction ratio is 65–70%.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

Acetate is used as a buffer in the dialysate solution. It is the most common cause of dialysis-induced hypotension because of its vasodilatory and cardio-depressive effects.

For this reason, acetate is less commonly used these days and has been largely replaced by bicarbonate.

Solution to Question 8:

Hypotension is the most common acute complication of hemodialysis, particularly among patients with diabetes mellitus.

Solution to Question 9:

Cooling, not warming, the dialysate before dialysis must be done to prevent hypotension during dialysis.

Hypotension is the most common acute complication of hemodialysis. The mechanism is not fully understood and occurs due to multiple factors. However, the following precautions have been shown to frequently prevent this complication:

• Careful calculation of the dry weight and adjustment of the filtration rate accordingly.

• Ultrafiltration modeling such that more fluid is removed during the beginning than the end of the procedure.

• Avoiding rapid fluid removal and maintaining the rate < 13 mL/kg/hour.

• Avoiding heavy meals during dialysis.

• Cooling of the dialysate during the dialysis.

Sold by @itachibot

Solution to Question 10:

The given scenario is suggestive of the dialysis disequilibrium syndrome or the reverse urea effect. It is an acute complication that occurs especially in patients undergoing dialysis for the first time.

During dialysis, urea in blood vessels is rapidly removed as compared to that in the interstitium. This decreases the osmolality of blood. This results in a fluid shift from the blood vessels into the interstitium and brain parenchyma, causing cerebral edema.

In order to prevent this, the rate of flow of dialysate must be adjusted to avoid the rapid removal of solutes.

Solution to Question 11:

Dialysis must be discontinued immediately in case of a type A reaction. It is an IgE-mediated immediate hypersensitivity reaction that can lead to shock and death.

Bio-incompatible cellulose-containing membranes are commonly associated with hemodialysis-associated anaphylaxis.

The reactions can be of two types:

Type A reaction	Type B reaction
IgE-mediated immediate hyp ersensitivity reaction in respo nse toethylene oxide. (used t o sterilize dialyzers).	Delayed reaction due tocomp lementactivationand cytokine release.
Occur within minutes.	Occur after hours.
Stop dialysisand give subcuta neous adrenaline or steroids.	It isself-limitingand dialysis c an be continued.

Solution to Question 12:

Hypovolemia is an acute complication of hemodialysis. The chronic complications of hemodialysis are:

• Amyloidosis

• Adynamic bone disease

• Dementia (aluminum toxicity)

Solution to Question 13:

If you bought this from someone else, it has been scammed. Sold by @itachibot

Sold by @itachibot

Carpal tunnel syndrome in the setting of chronic dialysis is suggestive of dialysis-associated amyloidosis. It occurs due to the deposition of amyloid ß-2 in tissues.

This condition is seen in patients undergoing long-term hemodialysis or continuous ambulatory peritoneal dialysis (CAPD). Amyloid deposition can lead to the following complications:

• Destructive osteoarthropathies - carpal tunnel syndrome, flexor tenosynovitis

• Subchondral bone cysts and pathologic fractures

• Paraplegia - due to deposits along paravertebral ligaments and intervertebral discs

• Visceral involvement - GI tract, heart, tongue

Management involves providing symptomatic relief by means of analgesics, anti-inflammatory drugs, physical and occupational therapy, and surgical procedures. Renal transplantation is the treatment of choice.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

Renal osteodystrophy does not show improvement with dialysis.

Failure of renal function leads to decreased renal vitamin D production. The resulting hypocalcemia and hyperphosphatemia, in turn, lead to secondary hyperparathyroidism. This can be corrected only by vitamin D and calcium supplementation.

Other manifestations of renal dysfunction and uremia show improvement with regular dialysis sessions.

Solution to Question 15:

Cardiovascular disease due to accelerated atherosclerosis is the major cause of death in patients with ESRD on chronic dialysis.

Cardiovascular mortality is higher in dialysis patients compared to patients after transplantation.

Solution to Question 16:

In peritoneal dialysis, 1.5–3 L of dialysis fluid is infused into the peritoneal cavity.

In this process, the peritoneum itself acts as a semipermeable membrane. A peritoneal catheter is used to infuse the fluid into the cavity. It is allowed to dwell for a set period of time usually 2–4 hours to allow for the removal of metabolic waste products. The dialysis fluid contains osmotically active agents such as glucose or icodextrin.

The following image shows the process of peritoneal dialysis.

Sold by @itachibot



Sold by @itachi... else,
If you purchased this from some...
you may have been scammed.

Solution to Question 17:

This clinical scenario in a patient undergoing peritoneal dialysis is suggestive of peritonitis. It is most commonly caused by gram-positive cocci such as Staphylococcus, which spread from the skin.

Solution to Question 18:

Weight gain is a complication of peritoneal dialysis, not weight loss. The complications of peritoneal dialysis include:

- Peritonitis
- Catheter-associated non-peritonitis infections
- Weight gain
- Metabolic disturbances - hypoproteinemia and hyperglycemia
- Residual uremia - especially among patients with no residual kidney function
- Insulin resistance
- Hypertriglyceridemia

Solution to Question 19:

The most common cause of death in renal transplant recipients is cardiovascular causes. It occurs in > 50% of patients.

Sold by @itachibot

Other complications in recipients include:

• Infections

• Most common - CMV.

• Most specific - BK (polyoma) virus

• Most common fungal - pneumocystis.

• Malignancy - most commonly of skin and lip

Donors can develop long-term complications related to premature failure of the single remaining kidney including:

• Hypertension

• Proteinuria

• Focal segmental glomerulosclerosis (FSGS)

ABO and HLA mismatch are relative, not absolute contraindications due to the availability of desensitization procedures. The absolute contraindications for transplantation are as follows:

Recipient	Donor
Any active infections/ malign ancy	Any active infections (includi ng HIV)/ malignancy
Shortened life expectancy du e to other causes	Uncontrolled diabetes/ hyper tension
Active substance abuse/ unco ntrolled psychiatric condition	Uncontrolled psychiatric con dition or suicidality
Treatment nonadherence	Age < 18 or lack of mental ca pacity
Reversible renal failure	

If you @itachibot someone else, you ned

Solution to Question 20:

Cyclosporine, azathioprine, prednisolone constitute the prophylactic maintenance therapy used after renal transplant.

Immunosuppressive therapy is given to reduce the risk of transplant rejection reactions. The following regimen is commonly followed:

• Induction therapy:

• Given at the time of transplant to reduce the risk of early acute rejection.

• The preferred treatment is anti-lymphocyte serum and alemtuzumab.

• Maintenance therapy:

• Given to all kidney transplant patients except when the donor and recipient are identical twins.

• The most common combination used is :

Sold by @itachibot

cyclosporine, azathioprine, and prednisolone (also known as triple therapy).

• Belatacept is a co-stimulatory blocking antibody approved by the FDA for maintenance therapy.

Rejection reactions can be hyperacute, acute, or chronic. Treatment of rejection is as follows:

• First episode - IV prednisolone for 3 days

• Failure to respond to prednisone or second episode - anti-thymocyte globulin

Solution to Question 21:

The most common opportunistic infection occurring 1-6 months after renal transplant is Cytomegalovirus (CMV).

Pneumocystis carinii is rare these days due to universal prophylaxis for 6-12 months following transplantation.

Opportunistic infections after renal transplant		
< 1 month	1-6 months	> 6 months
Wound infectionsHerpesvirusOral candidiasisUrinary tract infection	Pneumocystis cariniiCytomegalovirusLegionellaListeriaHepatitis B, C	AspergillusNocardiaBK virus Herpes zosterHepatitis B, C

Solution to Question 22:

Alport syndrome does not recur in the transplanted kidney. This is because it occurs due to an inherited defect in type 4 collagen in the glomerular basement membrane. This defect is absent in the transplanted kidney.

Diabetic nephropathy, membranous glomerulonephritis, and lupus nephritis can recur in the transplanted kidney.

Solution to Question 23:

Patients with diabetic nephropathy have the highest risk of disease recurrence after renal transplant. Nearly 52% of transplant recipients have histological evidence of diabetic kidney disease/mesangial sclerosis at 10 years after transplantation.

The recurrence rates of the other conditions are as follows:

• Membranoproliferative glomerulonephritis: 20-48%

• Membranous glomerulonephritis: 10-45%

• Lupus nephritis: 2-11%

Sold by @itachibot

Solution to Question 24:

The average life expectancy of a living-donor graft is around 14 years and that of a deceased-donor graft is around 10 years.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Cystic and Inherited Disorders of the Kidney

Question 1:

Which of the following is true in patients with autosomal dominant polycystic kidney disease?

a) ADPKD progresses to end-stage renal disease in early adulthood

b) Calcium oxalate stones are the most common kidney stones

c) Renal cyst infections are rarely seen

d) Cardiovascular complications are a major cause of mortality

Question 2:

A 48-year-old woman developed a sudden severe headache and vomiting. Her neighbor rushed her to the hospital where she was declared dead on arrival. She lived alone and her family history is unknown. An autopsy revealed a subarachnoid hemorrhage and the following findings in her kidneys. Which of the following is incorrect about this condition?



a) Mutation of PKD-1 is responsible for 85% of cases

b) PKD-2 mutation has higher risk of end-stage renal disease than PKD-1

c) The PKD genes code for the polycystin protein

d) Cysts are formed only in 5% of the tubules of the kidney

If you bought this from someone else, it has been scammed. Sold by @itachibot

Sold by @itachibot

Question 3:

In patients with ADPKD, cysts are least likely to be present in which of the following sites?

a) Arachnoid membranes

b) Lungs

c) Pancreas

d) Seminal vesicles

Question 4:

A 60-year-old man presents with loss of weight and appetite over the last 3 months. He was diagnosed with autosomal dominant polycystic kidney disease 15 years ago. A CT abdomen reveals bilateral, multicentric renal masses along with enlarged retroperitoneal lymph nodes. What is the likely diagnosis?

a) Clear cell RCC

b) Papillary cell RCC

c) Chromophobe RCC

d) Sarcomatoid RCC

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 5:

A 25-year-old woman presented with persistent headaches over the last 1 month. Her father and paternal uncle suffered from kidney disease and both died in their 50's of a 'brain bleed'. Her blood pressure is 160/110 mm Hg and had also been elevated during a previous visit. Which of the following investigations should be avoided in this patient?

a) Ultrasonography of the abdomen

b) Contrast enhanced CT of the abdomen

c) Non-contrast CT scan of the abdomen

d) MR angiography of the cerebral vasculature

Question 6:

How many renal cysts on USG are required to diagnose a 45-year-old patient with autosomal dominant polycystic kidney disease?

a) At least 2 cysts in each kidney

Sold by @itachibot

b) At least 3 cysts in each kidney

c) At least 4 cysts in each kidney

d) Even a single cyst in each kidney

Question 7:

A 56-year-old woman on dialysis is being planned for a renal transplant with her son as an organ donor. He undergoes a USG scan as part of routine donor evaluation. The following findings are seen. Which of the following will you not recommend for the son?



a) Increased fluid intake

b) Angiotensin-converting enzyme inhibitors

c) Surgical decompression of cysts

d) Tolvaptan

Question 8:

A 24-year-old G2P1L0 female patient came for an antenatal check-up at 26 weeks gestation. On examination, her abdominal girth was less for gestational age and the fetal parts were easily palpable. Medical records from her previous pregnancy showed that the deceased infant had been positive for a PKHD1 mutation. Which of the following statements is correct regarding the most likely diagnosis of the fetus?

a) It is commonly an asymptomatic and mild disease

b) In severe cases, it can be diagnosed antenatally on USG

Sold by @itachibot

c) Respiratory insufficiency is uncommon in newborns

d) It is commonly diagnosed at 5-10 years of age

Question 9:

Which of the following is not a typical feature seen in patients with autosomal recessive polycystic kidney disease?

a) Biliary dysgenesis

b) Portal hypertension

c) Caroli's disease

d) Hepatic cysts

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Match the following:

a) 1-d, 2-b, 3-c, 4-a

b) 1-b, 2-d, 3-a, 4-c

c) 1-c, 2-a, 3-d, 4-b

d) 1-a, 2-c, 3-b, 4-d

Question 11:

Which of the following is an autosomal recessive renal cystic disease?

a) Tuberous sclerosis

b) Medullary cystic kidney

c) Nephronophthisis

d) Von Hippel-Lindau disease

Question 12:

What is the most common inherited cause of kidney failure in children?

a) Medullary sponge kidney

b) Nephronophthisis

Sold by @itachibot

c) Medullary cystic kidney disease

d) Autosomal recessive polycystic kidney disease

Question 13:

A child with Joubert syndrome will have which of the following conditions?

a) Juvenile nephronophthisis

b) Medullary cystic kidney disease

c) Medullary sponge kidney

d) Autosomal recessive polycystic kidney disease

Question 14:

The following finding on intravenous pyelography is seen in which of the following conditions?

a) Medullary cystic kidney disease

b) Medullary sponge kidney

c) Senior loken syndrome

d) Infantile nephronophthisis

If yo ld by @itachibot is from someone else, scammed.

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	d
5	b
6	a
7	c
8	b
9	d
10	c
11	c
12	b
13	a
14	b

If you purchased this from somewhere else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Cardiovascular complications are the most common cause of mortality in autosomal dominant polycystic kidney disease (ADPKD). Infections are the second most common cause.

ADPKD is usually clinically silent. Many cases remain asymptomatic until the fifth decade of life and are diagnosed incidentally. PKD-1 mutations cause more severe disease than PKD-2 mutations. The following clinical features may be seen in symptomatic patients:

• Abdominal mass

• Back or flank pain

• Recurrent UTIs - Renal cyst infections by Gram-negative bacteria are common

• Nephrolithiasis - most commonly uric acid stones

• Hematuria

• Cardiac:

• Hypertension - early feature

• Left ventricular hypertrophy

• Mitral valve prolapse, valvular insufficiency

Sold by @itachibot

• Extra-renal:

• Intracranial aneurysms

• Hepatic cysts

As the cysts grow they compress on the surrounding normal tissue which leads to a progressive reduction in the GFR. Eventually, it leads to end-stage renal disease (ESRD) by the fifth decade of life.

Solution to Question 2:

This clinical scenario with a subarachnoid hemorrhage (SAH) and multiple renal cysts is suggestive of autosomal dominant polycystic kidney disease (ADPKD). The PKD-1 mutation has a higher risk of end-stage renal disease (ESRD) than PKD-2.

ADPKD is a disorder which is characterized by progressive bilateral formation of renal cysts. It is caused by the following mutations:

• PKD1 gene on chromosome 16 (85%) - codes for polycystin-1

• PKD2 gene on chromosome 4 (15%) - codes for polycystin-2

Polycystins are transmembrane proteins that are present on the primary cilia of the tubular cells. The mutated proteins in ADPKD alter the normal functioning and triggers cell growth and cyst formation.

Though cysts are formed only in 5% of the tubules of the kidney, they compress surrounding renal tissue due to their enormous growth. This results in loss of renal function.

Solution to Question 3:

In patients with ADPKD, cysts in the lungs are very rare.

There are very few case reports with lung cysts in ADPKD patients, but the association between ADPKD and lung cysts has not been proved. However, there is a three-fold increase in the incidence of bronchiectasis in ADPKD patients.

The common extrarenal manifestations of ADPKD include:

• Hepatic cysts (50-70%) - most common

• Seminal vesicle cysts (40%) - can result in infertility

• Arachnoid membrane cysts (8%) - usually asymptomatic, may increase the risk of subdural hematoma

• Pancreatic cysts (7-36%)

• Splenic cysts

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Solution to Question 4:

This clinical scenario is suggestive of renal cell carcinoma (RCC) with lymph node metastases. The sarcomatoid type of RCC is commonly associated with ADPKD.

Renal cell carcinoma (RCC) is a relatively rare complication of ADPKD. There does not seem to be an increased frequency of RCC in these patients compared to the general population. However, when it occurs, it tends to be bilateral, multicentric and of sarcomatoid type.

Solution to Question 5:

This clinical scenario is suggestive of autosomal dominant polycystic kidney disease (ADPKD). These patients usually have impaired renal function. Hence, contrast-enhanced CT (CECT) would usually be avoided due to the risk of contrast nephrotoxicity.

The following investigations are done in patients with ADPKD:

• USG abdomen - for screening and diagnosis

• Non-contrast CT abdomen - more sensitive and picks up cysts of smaller size

• MR-angiography of brain - pre-symptomatic screening for intracerebral aneurysms

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

Between 40-59 years of age, the presence of ≥2 cysts in each kidney is needed to diagnose ADPKD.

This condition is diagnosed based on the USG findings using the Ravine's criteria:

Age (in years)	Imaging findings
15-39	Total of ≥3 cysts, unilateral o r bilateral
40-59	≥2 cysts in each kidney
≥ 60	≥ 4 cysts in each kidney

Solution to Question 7:

This clinical scenario and USG showing a cystic appearance of the kidney is suggestive of asymptomatic autosomal dominant polycystic kidney disease (ADPKD). Surgical decompression is not a routine procedure performed in these patients. It may be required only in cases of very large cysts causing persistent pain and infected cysts.

The management includes the following:

• Analgesics

Sold by @itachibot

- Increased fluid intake
- Prevents renal stone formation
- Reduces growth of cysts by suppressing ADH release
- ACE inhibitors and ARBs
- Blood pressure control
- Slowing of cyst growth
- Tolvaptan (V2 receptor antagonist)
- Reduce pain and slow the decline in renal function
- Renal transplant
- Definitive management
- Pre-transplant nephrectomy may be required in patients with very large polycystic kidneys

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

A PKHD1 mutation in the deceased infant is suggestive of autosomal recessive polycystic kidney disease (ARPKD). As the woman has signs of oligohydramnios, it is likely that this fetus also has ARPKD. Severe cases of ARPKD can be diagnosed in-utero with a USG. Prenatal testing can also be done.

ARPKD is a hereditary renal disease of childhood caused by a mutation in the PKHD1 gene on chromosome 6. It codes for fibrocystin/ polyductin (FPC), a membrane protein present in the cells of the cortical and medullary collecting ducts.

Affected fetuses show greatly enlarged kidneys and reduced fetal urine production. This leads to oligohydramnios and pulmonary hypoplasia. About 30% of neonates die soon after birth due to respiratory insufficiency. Most deaths occur < 1 month of life. Others develop chronic kidney disease and hypertension in early childhood.

Solution to Question 9:

Hepatic cysts are not a typical feature of autosomal recessive polycystic kidney disease (ARPKD). They occur only in rare cases.

ARPKD is usually diagnosed in the neonatal period. These patients commonly have the following findings:

- Hypertension
- Chronic renal insufficiency
- Congenital hepatic fibrosis
- Biliary dysgenesis with periportal fibrosis is seen
- Can lead to portal hypertension

Sold by @itachibot

• Caroli's disease - dilatation of intrahepatic bile ducts

Note: Caroli's disease is also associated with medullary sponge kidney.

Solution to Question 10:

The gene mutations causing the following cystic disorders are:

• Medullary cystic kidney disease-1 (MCKD-1) - MUC1

• Medullary cystic kidney disease-2 (MCKD-2) - UMOD

• Autosomal recessive polycystic kidney disease (ARPKD) - PKHD1

• Adolescent nephronophthisis - NPHP 1-20

Medullary cystic kidney disease (MCKD) is an autosomal dominant disorder. It presents with a late-onset, slowly progressive renal failure. Renal cysts are not common but can be seen occasionally. Medullary cystic kidney diseases are of two types, MCKD1 and MCKD2.

MCKD1 is due to a defect in the mucin 1 (MUC1) gene. The mutated gene codes for a large protein fragment that has a toxic effect on the tubules. It is not associated with elevated uric acid levels and gout.

MCKD2 is due to a defect in the UMOD gene, which codes for the protein uromodulin. It is associated with hyperuricemia and gout.

Solution to Question 11:

Nephronophthisis is an autosomal recessive renal cystic disease.

It belongs to a category of conditions called ciliopathies. It occurs due to an underlying defect in the primary cilia of the tubular epithelial cells. The cilia play an important role in cell signaling and maintenance of cell growth, proliferation, and secretion.

Other ciliopathies include autosomal dominant polycystic kidney disease, autosomal recessive polycystic kidney disease, and medullary cystic kidney disease.

Solution to Question 12:

The most common inherited cause of kidney failure in childhood is nephronophthisis.

Nephronophthisis (NPHP) refers to a group of related, but genetically distinct, autosomal recessive disorders. Mutations are seen in the NPHP genes 1-20. These genes code for nephrocystins which are expressed on the kidney tubules.

NPHP is classified into infantile, juvenile, and adolescent forms. The juvenile form is the most common form. Common features include tubulointerstitial fibrosis, renal cysts, and chronic kidney disease.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Solution to Question 13:

Joubert's syndrome is defined by the presence of juvenile nephronophthisis (NPHP) and multiple neurologic findings.

A few examples of multiorgan syndromes associated with nephronophthisis are:

• Senior-Loken syndrome: NPHP with retinitis pigmentosa

• Joubert syndrome: NPHP with multiple neurologic findings

• Bardet-Biedl syndrome: NPHP with truncal obesity, cognitive impairment, retinal dystrophy, and polydactyly

Solution to Question 14:

The given image shows a paintbrush or bouquet of flowers appearance on intravenous pyelography (IVP). It occurs in medullary sponge kidney (MSK).

MSK is a sporadic benign condition that is often diagnosed incidentally. It is a developmental defect affecting the formation of collecting tubules and results in cystic dilatation of the medullary and papillary portions of collecting ducts. It is characterized by medullary cysts that are of variable size. It is associated with distal RTA which leads to hypercalciuria. The hypercalciuria leads to a tendency to form calcium oxalate and calcium phosphate stones. The patients can also have recurrent UTIs and hematuria.

CT urography has largely replaced intravenous pyelography as the investigation of choice.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Renal Tubular Disorders

Question 1:

What acid-base disturbance is seen in patients with renal tubular acidosis?

a) Normal anion gap hyperchloremic metabolic acidosis.

b) High anion gap hyperchloremic metabolic acidosis

c) Normal anion gap hypochloremic metabolic acidosis

d) High anion gap hypochloremic metabolic acidosis.

Question 2:

What is the most common type of renal tubular acidosis?

a) Type 1

b) Type 2

c) Type 3

d) Type 4

Question 3:

Which of the following is incorrect about type 4 renal tubular acidosis?

a) The most common cause is diabetes mellitus

b) It can occurs in patients with interstitial renal diseases

c) It is always congenital

d) It is caused by hyporeninemic hypoaldosteronism

Question 4:

A 52-year-old man presented with dizziness and infrequent urination. He was diagnosed with stage 4 chronic kidney disease last month. Investigations reveal metabolic acidosis and hyperkalemia. What is the likely diagnosis?

a) Type 1 RTA

b) Type 2 RTA

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

c) Type 3 RTA

d) Type 4 RTA

Question 5:

Which of the following would not be seen in patients with type 1 renal tubular acidosis?

a) Reduced NH■■ excretion

b) Serum potassium < 3.5 mEq/L

c) Urinary pH is < 5.5 after acid loading

d) Increased bone resorption

Question 6:

A 6-month-old infant is being evaluated for failure to thrive. Lab investigations are as follows. Hypokalemia is found to improve following alkalinization therapy. What type of renal stones is this infant at a higher risk of developing?

a) Uric acid

b) Citrate

c) Cystine

d) Calcium phosphate

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following would you not expect in a patient with Lowe's syndrome?

a) Glaucoma

b) Cataract

c) Hypertonia

d) Intellectual disability

Question 8:

Which of the following is unlikely to be seen in patients with Dent's disease?

a) X-linked inheritance

Sold by @itachibot

b) Inactivating mutation of chloride channel

c) Recurrent kidney stones

d) Distal renal tubular defect

Question 9:

Patients with which of the following are not at risk of developing type 2 renal tubular acidosis?

a) Wilson's disease

b) Galactosemia

c) Hereditary fructose intolerance

d) Medullary sponge kidney

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following is incorrect about renal tubular acidosis type 2?

a) It is associated with hypophosphatemia

b) It can be caused by Lowe's syndrome

c) Renal stones may occur in patients

d) Minimum urinary pH is < 5.5

Question 11:

A 14-month-old baby was brought with complaints of refusal to walk after a minor fall while playing. Imaging revealed a left tibial fracture along with generalized sclerosis and a bone-in-bone appearance. Additionally, his developmental milestones were found to be delayed. Investigations are as follows. What is the likely renal tubular defect present in this patient?

a) Type 1 RTA

b) Type 2 RTA

c) Type 3 RTA

d) Type 4 RTA

Question 12:

Sold by @itachibot

A 30-year-old man is being evaluated for high blood pressure noted on multiple occasions. Investigations reveal metabolic alkalosis, suppressed renin levels, and high urine chloride levels. What is the likely cause?

a) Bartter syndrome

b) Liddle syndrome

c) Gitelman syndrome

d) Fanconi syndrome

Question 13:

A young patient presented with polyuria and polydipsia. His BP is 130/80 mm Hg. Laboratory parameters revealed serum K+ of 3.2 mmol/L, blood pH of 7.46, and hypomagnesemia. Urinary calcium was elevated. He is not on any medications. This condition is likely due to a defect in which of the following?

a) Proximal tubule

b) Distal convoluted tubules

c) Ascending loop of Henle

d) Epithelial sodium channels

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 14:

A 17-year-old boy is being evaluated for fatigue and muscle weakness for 3 months. On examination, Chvostek's sign is positive and BP is 90/60 mm Hg. Investigations show hypokalemia and hypercalcemia with metabolic alkalosis. What is the likely diagnosis?

a) Bartter syndrome

b) Gitelman syndrome

c) Liddle syndrome

d) Fanconi's syndrome

Answer Key

Question No.	Correct Option
1	a

Sold by @itachibot

2	d
3	c
4	d
5	c
6	d
7	c
8	d
9	d
10	c
11	c
12	b
13	c
14	b

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

In renal tubular acidosis (RTA), there is normal anion gap hyperchloremic metabolic acidosis.

RTA refers to a group of disorders in which there is impaired tubular transport of H+ and/or HCO3- despite relatively well-preserved renal function. There are 4 types:

• Type I - impaired H+ secretion (distal), leading to decreased HCO3- production

• Type II - impaired HCO3- reabsorption (proximal)

• Type III - carbonic anhydrase deficiency causing defects in the transport of both H+ and HCO3- (mixed)

• Type IV - hypoaldosteronism (hyperkalemic)

The overall deficiency of HCO3- in all the types is compensated by Cl- secretion. Hence, the serum anion gap remains normal.

Solution to Question 2:

The most common type of renal tubular acidosis (RTA) is the type 4 or hyperkalemic type.

Solution to Question 3:

Type 4 renal tubular acidosis can be a congenital or acquired disorder.

Sold by @itachibot

It is caused by conditions that result in hypoaldosteronism or hyporeninemic hypoaldosteronism including:

• Diabetic nephropathy - most common

• Tubulointerstitial disease

• Addison's disease

• Obstructive uropathy

• Drugs like NSAIDs, ACEI, ARBs can increase the risk

• Psuedohypoaldosteronism type 1 (due to aldosterone resistance)

• Psuedohypoaldosteronism type 2/ gordons syndrome (due to defects in Na-Cl cotransporter)

• Congenital isolated hypoaldosteronism (due to enzyme deficiency)

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Solution to Question 4:

Hyperkalemia is a characteristic feature of type 4 RTA or generalized distal RTA. It commonly occurs in patients with chronic kidney disease (CKD).

In CKD patients, renal dysfunction results in hyperkalemia only when the GFR is severely impaired (stage 5). Hyperkalemia in the earlier stages of CKD is suggestive of type 4 RTA.

In CKD, volume overload leads to low levels of renin and, hence, aldosterone. This leads to Na+ loss and K+ retention in the distal tubule. The resultant hyperkalemia causes impaired NH_3 synthesis in the proximal tubule. Hence, H+ cannot be excreted as NH_4^+ ions. Retained H+ ions lead to reduced HCO_3^- production and acidosis.

Solution to Question 5:

Urinary pH will be > 5.5 after acid loading in RTA type 1.

In this condition, there is a defect in H+ secretion by the distal tubules. Hence, there is a failure of urine acidification. H+ combines with NH_3 to form NH■■ and thus in RTA type 1, NH_4^+ excretion is reduced. In the acid loading test, acidosis is induced by the administration of NH_4Cl (releases H+). In normal individuals, this results in increased excretion of H+, leading to acidic urine (pH < 5.5). However, this fails to occur in RTA type 1.

Impaired secretion of H+ leads to potassium wasting and hypokalemia (via the H+/K+ATPase antiporter). The resulting metabolic acidosis causes bone demineralization and resorption.

Solution to Question 6:

This clinical scenario is suggestive of type 1 RTA. In this condition, calcium phosphate and oxalate stones are formed in the distal convoluting tubules (DCT).

Sold by @itachibot

In this condition, there is a defect in H+ secretion by the distal tubules. This leads to:

• Normal anion gap metabolic acidosis

• Urinary pH > 5.5

• Hypokalemia

• Impaired H+ secretion causes K+ wasting

• It improves following alkalinization therapy with $NaHCO_3$

Acidemia leads to hypercalcemia and then hypercalciuria. There is also hypocitraturia as acidemia induces increased cell citrate utilization. Normally, citrate in the urine binds to calcium and increases its solubility. In the absence of citrate in the urine, calcium begins to precipitate. This leads to nephrocalcinosis and calcium phosphate and oxalate stones.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

Hypertonia is not a feature of Lowe's syndrome or oculo-cerebro-renal syndrome. The following clinical features are seen in this condition:

• Hypotonia

• Cataract

• Glaucoma

• Intellectual disability

• Renal tubular acidosis (RTA) type 2

Solution to Question 8:

Dent's disease is an inherited cause of type 2 RTA. A proximal renal tubular defect is seen in this condition, not a distal tubular defect.

It is an X-linked recessive condition. It is caused by inactivating mutations in CLCN5, a chloride transporter expressed in endosomes of the proximal tubule. The resulting dysfunction of the proximal convoluted tubule and failure of reabsorption of substances leads to:

• Hypercalciuria

• Hypophosphatemia

• Recurrent kidney stones

Solution to Question 9:

Medullary sponge kidney is a cause of RTA type 1, not type 2.

The causes of RTA type 2 include:

Sold by @itachibot

• Amyloidosis

• Wilson's disease

• Galactosemia

• Hereditary fructose intolerance

• Cystinosis

• Lowe syndrome

• Drugs - tenofovir, acetazolamide, topiramate

The causes of RTA type 1 include:

• Familial defects

• Autoimmune conditions - SLE, Sjogren syndrome

• Drugs - amphotericin, lithium, aminoglycosides, ifosfamide

• Medullary sponge kidney

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

Renal stones are not a feature of type 2 or proximal RTA. This is because high levels of citrate in the urine bind with calcium and prevent stone formation.

In this condition, the cells of the proximal convoluted tubule (PCT) fail to reabsorb HCO3-. In the initial stages, this HCO3- loss leads to a urine pH > 5.5. However, as the body's HCO3- stores are depleted, the urinary pH falls to < 5.5 as the secretion of H+ in the DCT remains intact. Hence, the minimum urinary pH is < 5.5.

It can be caused by Lowe's syndrome but a majority of type 2 RTA cases are associated with Fanconi's syndrome. This syndrome is characterized by a generalized defect of reabsorption of various substances at the PCT. This leads to loss of HCO3-, glucose, amino acids, phosphate, citrate, and calcium in the urine. Loss of phosphate leads to vitamin-D resistant hypophosphatemic rickets /osteomalacia.

Note: An exception is Dent's disease, a genetic cause of RTA type 2 in which renal stones are present.

Solution to Question 11:

Pathological fractures and a bone-in-bone appearance point towards a diagnosis of osteopetrosis. The lab values are significant for hypokalemia with normal anion gap metabolic acidosis. These findings are suggestive of renal tubular acidosis (RTA) type 3.

Type 3 or mixed RTA is a rare condition that usually occurs in neonates and infants. It is due to an autosomal recessive, inherited carbonic anhydrase-2 (CA-2) deficiency. This results in features of both distal and proximal RTA with:

• Failure to reabsorb HCO3 at PCT

Sold by @itachibot

• Failure to excrete H+ at DCT

This results in the following findings:

• Normal anion gap metabolic acidosis

• Hypokalemia

• Urinary pH > 5.5

• Positive urine anion gap

As CA-2 is widely expressed, mutations of this gene result in a condition called marble brain disease or Guibaud-Vainsel syndrome. It presents with multiple clinical findings including:

• Cerebral calcifications

• Mental retardation

• Mixed RTA (Type 3 RTA)

• Osteopetrosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

The clinical scenario with hypertension and metabolic alkalosis is suggestive of Liddle's syndrome.

It is an autosomal dominant condition due to a gain of function mutation in the ENaC gene. This gene codes for the epithelial sodium channels present in the kidney. This leads to increased sodium retention and results in a state of pseudo-hyperaldosteronism. The following findings are seen:

• Hypertension

• Hypokalemic metabolic alkalosis

• Normal-low aldosterone

• Low renin

Treatment is with diuretics that directly block the ENaC channels, like amiloride and triamterene. Spironolactone is not effective as it is an aldosterone-receptor antagonist.

Solution to Question 13:

This clinical scenario is suggestive of Bartter syndrome. It occurs due to a defect in the Na+/K+/Cl- channel in the thick ascending loop of Henle.

A defect in the Na+/K+/Cl- channel impairs the transport functions of the thick ascending loop of Henle. It results in the following features:

• Polyuria and polydipsia

• Natriuresis with high aldosterone (due to RAAS activation)

Sold by @itachibot

• Hypokalemia

• Metabolic alkalosis

• Hypomagnesemia

• Hypocalcemia and hypercalciuria

The clinical features of this condition are similar to the effects of loop diuretics. These drugs also act on the Na+/K+/Cl- channel. Hence, response to loop diuretics may be blunted. Treatment is with potassium-sparing diuretics like spironolactone and amiloride. NSAID's like indomethacin are second line drugs however they can reduce renal function.

Option A: Defect in the proximal tubule results in Fanconi's syndrome. These patients have acidosis, with glycosuria, generalized aminoaciduria, and phosphaturia.

Option B: Defect in the Na+/Cl- cotransporter in distal convoluted tubule (DCT) results in Gitelman syndrome. These patients also present with hypokalemic alkalosis and hypomagnesemia. However, they also develop, hypocalciuria, and chondrocalcinosis.

Option D: Increased activation of the epithelial sodium channels (ENaC) channels results in Liddle syndrome. This causes increased reabsorption of Na+ exchanged for K+ causing hypokalemia. Increased Na absorption leads to hypertension.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

Chvostek's sign (spasm of facial muscles on tapping the facial nerve) indicates tetany. In this patient with hypercalcemia, it is likely due to hypomagnesemia. These clinical features are suggestive of Gitelman syndrome.

This condition occurs due to a mutation of Na+/Cl- cotransporter in the distal collecting tubule (DCT). The resulting impairment of the tubular transport leads to the following features:

• Mild natriuresis with normal-low BP

• Increased renin and aldosterone

• Hypokalemia

• Metabolic alkalosis

• Hypercalcemia and hypocalciuria

• Hypomagnesemia

The clinical features of this condition are similar to the effects of thiazide diuretics. These drugs also act on the Na+/Cl- cotransporter. Hence, response to thiazide diuretics may be blunted. Treatment is with potassium-sparing diuretics like spironolactone and amiloride.

Option A: Bartter syndrome affects the Na+/K+/Cl- channels. It causes hypocalcemia and hypercalciuria.

Option C: Liddle syndrome causes increased activation of ENaC channels. It results in hypokalemia and hypertension.

Option D: Fanconi's syndrome occurs due to a defect in proximal tubular function. It results in metabolic acidosis.

Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Vascular Diseases of Kidney

Question 1:

Which of the following is true regarding renal vein thrombosis?

a) Right renal vein is more commonly involved

b) Most common cause in children is nephrotic syndrome

c) Investigation of choice is doppler ultrasound of renal vein

d) Most common cause in elderly is renal cell carcinoma

Question 2:

A 10-year-old child develops hematuria after 4 days of severe diarrhea. Peripheral smear shows fragmented RBCs and thrombocytopenia. Ultrasound reveals the enlargement of both kidneys. What is the likely diagnosis?

a) Renal vein thrombosis

b) Hemolytic uremic syndrome

c) Henoch-Schonlein purpura

d) Post-infectious glomerulonephritis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

What is the most common cause of hemolytic uremic syndrome globally?

a) Shigella dysenteriae type I

b) E. coli O157: H7

c) Streptococcus pneumoniae

d) Autoimmunity

Question 4:

Which of the following is false about hemolytic uremic syndrome?

a) Symptoms and signs of HUS develops as diarrhea begins to improve

b) Only supportive managment of acute renal failure is done in typical HUS

Sold by @itachibot

c) Eculizumab improves microangiopathic hemolytic anemia in atypical HUS

d) Coomb's positive hemolytic anemia is seen in HUS

Question 5:

A 56-year-old woman is brought to the emergency department with sudden onset headache and photophobia. On examination, she is disoriented and febrile. The investigations and peripheral smear are given below. Which of the following statements is incorrect regarding this disease?



a) It is caused by ADAMTS13 deficiency

b) 20% patients will have falsely positive RA factor

c) Arrhythmias are a frequent cause of death

d) Mainstay of treatment is blood transfusion

Question 6:

What is the most■ common cause of clinically significant renal artery stenosis?

a) Hypertensive nephrosclerosis

b) Atherosclerosis

c) Fibromuscular dysplasia

d) Infrarenal aortic dissection

Sold by @itachibot

Question 7:

Which of the following statements is incorrect about fibromuscular dysplasia?

a) String of beads appearance on angiography

b) Predominantly affects females

c) Most commonly affected peripheral artery is subclavian artery

d) Segmental, non-atheromatous dysplastic angiopathy

Question 8:

Which of the following is unlikely in patients with atheroembolic kidney disease?

a) Livedo reticularis

b) Aortic aneurysms

c) Transient eosinophilia

d) Normal complement levels

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 9:

A 50-year-old patient is being evaluated for uncontrolled hypertension despite being on 3 antihypertensive drugs. He has a history of myocardial infarction which was managed with percutaneous intervention last month. On examination, an abdominal bruit is noted. Lab values indicate elevated serum urea and creatinine. Which of the following investigations is the gold standard test to diagnose this condition?

a) MR angiography

b) Renal arteriography

c) CT angiography

d) Duplex Doppler Ultrasound

Question 10:

Which of the following drugs must be avoided in patients with bilateral renal artery stenosis?

a) ACE inhibitors

b) Loop diuretics

c) Beta blockers

Sold by @itachibot

d) Thiazide diuretics

Answer Key

Question No.	Correct Option
1	d
2	a
3	b
4	d
5	d
6	b
7	c
8	d
9	b
10	a

Detailed Explanations

Solution to Question 1:

The most common cause of renal vein thrombosis in the elderly is renal cell carcinoma (RCC).

The most common causes of renal vein thrombosis in different age groups are as follows:

• Children - Dehydration, followed by sepsis

• Adults - Nephrotic syndrome (most commonly membranous glomerulonephritis)

• Elderly - Renal cell carcinoma

The left renal vein is more commonly involved and 2/3rd of cases are bilateral. Doppler USG is used for screening but the investigation of choice is CT angiography.

Solution to Question 2:

This clinical scenario is suggestive of renal vein thrombosis secondary to dehydration.

Hematuria in children following diarrhea can be due to the following:

Henoch-Schonlein purpura would present 1-3 weeks after an upper respiratory infection. It causes hematuria, non-blanching rash over the buttocks, and abdominal pain.

Sold by @itachibot

	Hemolytic uremic syndrome	Renal vein thrombosis
Onset	5-10 days after diarrhea	Depends on severity of dehydration
Renal failure	Present	Present
Hematuria	+	+
Peripheral smear fragmented RBCs	+	+
Platelets	Decreased	Decreased
USG kidney	Normal	Enlarged due to congestion by renal vein thrombosis (bilateral in 67% cases)

Sold by @itachibot
If you purchased this from someone
you may have been scammed.

Solution to Question 3:

The most common cause of hemolytic uremic syndrome (HUS) globally is Escherichia coli O157:H7.

The causes of HUS are as follows:

- Infections: Usually occur 5-10 days after diarrheal diseases due to bacterial exotoxins.
- E.coli O157:H7 - Shiga-like toxin (verotoxin)
- Shigella dysenteriae type I - Shiga toxin
- Streptococcus pneumoniae
- Autoimmunity
- Complement activation (in atypical HUS)

Solution to Question 4:

The hemolytic anemia in hemolytic uremic syndrome (HUS) occurs as a result of mechanical RBC destruction caused by microthrombi. Hence, Coomb's test is negative.

There are two types of HUS, namely typical and atypical.

Typical HUS	Atypical HUS
5-10 daysafter diarrheal illnessE.coliO157:H7orShigella dysenteriaetype I	Mutation in genes encoding complement regulatory protein

Sold by @itachibot

Typical HUS	Atypical HUS
Shiga-liketoxinsproduced cau seendothelialdamage.This lea ds to platelet activation and t hrombi.	Uncontrolledactivationof the complementsystem
No specific treatmentSupport ivemanagement of renal failu re	Plasma exchangeto replace th e deficient proteinsEculizum ab, aC5 inhibitor

Solution to Question 5:

This clinical scenario and image showing schistocytes (fragmented RBCs) in a woman are suggestive of thrombotic thrombocytopenic purpura (TTP). Blood transfusion is not preferred in TTP as it can aggravate the condition.

TTP is a type of thrombotic microangiopathy that occurs due to deficiency or autoantibodies against the ADAMTS13 metalloprotease, which is responsible for cleaving von Willebrand factor multimers. In the absence of this protein, vWF multimers aggregate and form thrombi. It presents with a classical pentad of clinical features:

• Fever

• Thrombocytopenia

• Microangiopathic hemolytic anemia

• Renal failure

• Neurological manifestations

It is treated with plasmapheresis and rituximab.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

The most common cause of clinically significant renal artery stenosis is atherosclerosis.

Renal artery stenosis is a relatively common finding in older patients with chronic essential hypertension. However, it is considered clinically significant only when renovascular hypertension develops. It occurs in the case of:

• > 70% stenosis in both renal arteries

• > 50% stenosis with post stenotic dilatation

The causes of renal artery stenosis are:

• Atherosclerosis (most common) - common in older patients with cardiovascular risk factors

• Fibromuscular dysplasia – more common in young, white females and is usually bilateral

• Less common causes include:

• Thromboemboli

Sold by @itachibot

• External compression by tumors

• Systemic vasculitis

• Antiphospholipid antibody syndrome

• Transplant renal artery stenosis

• Infrarenal aortic aneurysms and dissections

The investigation of choice for renal artery stenosis is renal angiography.

Solution to Question 7:

This clinical scenario and the string of beads appearance on CT angiography is characteristic of fibromuscular dysplasia. The most common peripheral (limb) arteries affected are the iliac arteries.

Fibromuscular dysplasia is a segmental, non-atheromatous, dysplastic type of angiopathy that predominantly affects females. The most common subtype is medial fibroplasia. It is one of the major causes of renovascular hypertension. The most common sites of involvement in decreasing order of frequency are as follows:

• Renal arteries

• Carotid arteries

• Peripheral (limb) arteries - most commonly iliac arteries

On angiography, a string of beads appearance is seen as shown below.

Solution to Question 8:

Atheroembolic kidney disease is associated with decreased complement levels.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Sold by @itachibot

Atheroembolic kidney disease is more frequent in males with a history of diabetes, hypertension, and ischemic heart disease. Most clinical atheroembolic events follow angiographic procedures, often of the coronary vessels.

It is associated with aortic aneurysms and renal artery stenosis. Systemic embolic manifestations such as fever, abdominal pain, and weight loss are also present, although cutaneous manifestations are more common like livedo reticularis and localized toe gangrene.

The lab findings include:

• Rising serum creatinine

• Transient eosinophilia

• Elevated erythrocyte sedimentation rate

• Hypocomplementemia

Definitive diagnosis depends on kidney biopsy which demonstrates microvessel occlusion with cholesterol crystals that leave a cleft in the vessel as shown below.



Solution to Question 9:

This clinical scenario is suggestive of renal artery stenosis. Renal intra-arterial angiography is the gold standard for diagnosis.In a patient with hypertension and increased creatinine, an ultrasound of the abdomen must be performed first to visualize the renal size and morphology to look for a unilateral shrunken kidney and thereby renal artery stenosis.

Option A: MR angiography is less preferred as gadolinium contrast can lead to nephrogenic fibrosis, particularly in patients with reduced kidney function.

Option C: CT scan with image reconstruction provides an excellent image but also carries a risk of contrast nephropathy.

Option D: Renal arterial doppler ultrasound measures renal artery velocities and can predict hemodynamically important lesions. However, it is not the gold standard test. Duplex

Sold by @itachibot

ultrasonography is used as a screening investigation.

Solution to Question 10:

ACE inhibitors must be avoided in patients with bilateral renal artery stenosis (RAS).

In renal artery stenosis, blood flow to the kidney is reduced, thereby reducing blood flow to the afferent arterioles. Hence, constriction of efferent arterioles is needed to maintain hydrostatic pressure within the glomerulus in order to maintain GFR.

ACEIs and ARBs inhibit efferent renal arteriolar vasoconstriction. In bilateral RAS, this causes a decline in the GFR within a few days of starting the drug.

Note: Bilateral renal artery stenosis has been classically considered as an absolute contraindication to the use of ACE inhibitors. However, new studies show that they may be used in some cases with careful monitoring.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Diagnosis of cardiovascular disorders

Question 1:

'a' wave in JVP corresponds to?

a) Right atrial contraction

b) Closure of tricuspid valve

c) Onset of ventricular systole

d) Maximal atrial filling

Question 2:

The c wave seen in a normal JVP waveform represents?

a) Bulging of tricuspid valve into the right atrium

b) Atrial relaxation during atrial diastole

c) Venous filling of the right atrium

d) Emptying of the right atrium into the right ventricle

Question 3:

The following abnormality on a JVP waveform can be seen in which of the given conditions?



©MARROW

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Cardiac tamponade

b) Ventricular tachycardia

c) Tricuspid regurgitation

d) Atrial fibrillation

Question 4:

A 37-year-old male patient comes to the clinic with complaints of weakness, restlessness, and chest pain. On examination, his blood pressure is found to be 90/60 mmHg, his heart sounds are muffled, and his neck veins are distended. What is the typical JVP finding in this patient?

a) Absent a wave

b) Prominent a wave

c) Rapid y descent

d) Absent y descent

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 5:

Deep y descent in JVP is seen in all except:

a) Cardiac tamponade

b) Restrictive cardiomyopathy

c) Constrictive pericarditis

d) Tricuspid regurgitation

Question 6:

The following four patients were examined on a tilt table for postural hypotension. The blood pressures before and immediately after performing head-up tilt are given below. Choose the correct option regarding the patients who suffer from orthostatic hypotension.

a) 2, 3

b) 1, 2, 4

c) 1, 4

d) 2, 3, 4

Question 7:

Sold by @itachibot

Pulsus paradoxus is seen in all of the following except:

a) COPD

b) Tension pneumothorax

c) Hypertrophic cardiomyopathy

d) Cardiac tamponade

Question 8:

Pulsus bisferiens is seen in all of the following conditions except:

a) Advanced aortic regurgitation

b) AS with AR

c) Left ventricular dysfunction

d) Hypertrophic cardiomyopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

Water hammer pulse is seen in which of the following conditions?

a) Aortic stenosis

b) Aortic regurgitation

c) Aortic stenosis and aortic regurgitation

d) Mitral regurgitation

Question 10:

A loud S1 is associated with?

a) Rheumatic mitral stenosis

b) Calcified mitral stenosis

c) Aortic regurgitation

d) All of the above

Question 11:

Sold by @itachibot

A hypertensive patient presented with chest pain and was found to have dilated cardiomyopathy on an echocardiogram. On auscultation, a reversed splitting of second heart sound was heard. Which of the following could most likely be responsible for the same?

a) RBBB

b) ASD

c) LBBB

d) VSD

Question 12:

Disappearance of loud S1 in mitral stenosis occurs in all except:

a) First degree heart block

b) AR

c) Calcified valve

d) Mild MS

Question 13:

While examining a patient, you hear a widely split S2 on auscultation. Which of the following is not a possible diagnosis?

a) Atrial septal defect

b) Mitral regurgitation

c) Pulmonary stenosis

d) Left bundle branch block

Question 14:

Which of the following conditions is associated with a fixed wide splitting of S2?

a) Mitral regurgitation

b) RBBB

c) Atrial septal defect

d) LV pacing

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 15:

Which of the following statements is true?

a) S3 and S4 are high pitched systolic sounds

b) S3 is always pathological

c) S4 is heard during the ventricular-filling phase

d) S3 occurs due to rapid filling of the ventricles during atrial systole

Question 16:

An ECG of a patient who underwent pulmonary artery catheterization showed a right bundle branch block. On auscultation, a widely split S1 was observed. Which of the following is true regarding this auscultatory finding?

a) The split increases with inspiration

b) The split increases with expiration

c) The split remains constant with inspiration and expiration

d) The change in split depends on the degree of bundle branch block

Question 17:

A 23-year-old woman presents with a history of occasional palpitations. She has no history of chest pain or dyspnea. On examination, blood pressure is 110/70 mmHg, the pulse is 82/min and regular. Joint hypermobility and thoracolumbar scoliosis are present. Gorlin's sign is positive. On auscultation, a click is heard after S1, followed by a late systolic murmur. What is the probable diagnosis?

a) Ehlers-Danlos syndrome

b) Marfan's syndrome

c) Loeys-Dietz syndrome

d) Osteogenesis imperfecta

Question 18:

Which of the following is not a diastolic sound?

a) S3

b) S4

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

c) Opening snap

d) Ejection click

Question 19:

On auscultation of a patient, you hear a murmur between S1 and S2. Which of the following conditions is not likely to be associated with this finding?

a) Hypertrophic cardiomyopathy

b) Tricuspid regurgitation

c) Ventricular septal defect

d) Pulmonary regurgitation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 20:

While examining a patient, you hear a mid-diastolic murmur with presystolic accentuation on auscultation of the precordium. Which of the following condition is associated with this finding?

a) Mitral regurgitation

b) MVP

c) Aortic stenosis

d) Mitral stenosis

Question 21:

You are a part of a health camp conducting check-ups for cardiovascular ailments. On auscultation of a patient's precordial area, you hear a systolic murmur. On asking the patient to perform the following maneuver, the intensity of the murmur increased. Which of the following conditions is associated with this finding?

Sold by @itachibot

a) MR

b) TR

c) AS

d) HCM

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 22:

While auscultating a patient's precordium, a murmur was heard that decreased in intensity on performing the following sustained maneuver. Which of the given condition is associated with this finding?



a) Mitral regurgitation

Sold by @itachibot

b) Ventricular septal defect

c) Aortic regurgitation

d) Aortic stenosis

Question 23:

Which of the following murmur increases on standing?

a) HOCM

b) MR

c) MS

d) VSD

Question 24:

A child was being evaluated for an underlying cardiac defect. On auscultation, a murmur was heard which began in systole, enveloped the second heart sound, and continued through some portion of the diastole. Which of the following condition is not associated with this finding?

a) Ventricular septal defect with aortic regurgitation

b) Patent ductus arteriosus

c) Coronary arteriovenous fistula

d) Venous hum

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 25:

Which of the following is true about Gallavardin effect?

a) It is associated with a diastolic murmur

b) It is seen in aortic stenosis

c) It refers to the radiation of murmur to the carotids

d) The transmission makes the murmur low pitched

Question 26:

Sold by @itachibot

A patient presents with easy fatiguability and shortness of breath on moderate physical activity. On auscultation, a holosystolic murmur is heard over the lower left sternal border and radiating to the epigastrium. Which of the following condition is most likely to be associated with this finding?

a) Mitral regurgitation

b) Tricuspid regurgitation

c) Ventricular septal defect

d) Atrial septal defect

Question 27:

Which of the following is true about Still's murmur?

a) It is heard in mitral regurgitation

b) It is best heard in left lateral decubitus position

c) It is a mid-systolic murmur

d) It is of grade 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 28:

Graham Steell murmur is heard in?

a) Aortic regurgitation

b) Pulmonary regurgitation

c) Mitral regurgitation

d) Tricuspid regurgitation

Question 29:

Carey Coombs murmur is heard in?

a) Mitral valvulitis

b) Mitral regurgitation

c) Tricuspid stenosis

d) Tricuspid regurgitation

Sold by @itachibot

Question 30:

Austin-Flint murmur is associated with which of the following conditions?

a) Chronic AR

b) Severe MS

c) Pulmonary hypertension

d) Severe MR

Question 31:

A 60-year-old man presented with an episode of acute onset dizziness and loss of consciousness which lasted for a few seconds followed by a regain of full consciousness. There were no similar episodes in the past. Which of the following is a true statement regarding this scenario?

a) ECG to rule out atrial fibrillation

b) If ECG is normal, CT scan should be done

c) Tilt-table testing

d) Vestibular neuritis is a possible condition that can cause these symptoms

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	d
5	a
6	b
7	c
8	c
9	b
10	a
11	c
12	d

Sold by @itachibot

13	d
14	c
15	c
16	a
17	a
18	d
19	d
20	d
21	d
22	d
23	a
24	a
25	b
26	b
27	c
28	b
29	a
30	a
31	a

If you purchase... you may have... else, ...d.

Detailed Explanations

Solution to Question 1:

The "a" wave in jugular venous pressure (JVP) is a positive wave, which corresponds to right atrial contraction.

The waveforms seen in a normal JVP are:

• a wave corresponds to atrial contraction and occurs just after the P wave (on ECG) and just before the first heart sound.

• c wave represents the bulging of the tricuspid valve into the right atrium during ventricular systole.

• x descent corresponds to the atrial relaxation during atrial diastole.

• v wave represents atrial filling during ventricular systole and peaks at the second heart sound.

• y descent represents passive emptying of the right atrium into the right ventricle after tricuspid valve opening.

Sold by @itachibot



MARROW
Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 2:

The c-wave in a JVP waveform represents the bulging of the tricuspid valve into the right atrium during ventricular systole.

JVP measured in the internal jugular vein is a reflection of the right atrial pressure.

The waveforms seen in a normal JVP are:

• a wave corresponds to atrial contraction and occurs just after the P wave (on ECG) and just before the first heart sound.

• c wave represents the bulging of the tricuspid valve into the right atrium during ventricular systole.

• x descent corresponds to the atrial relaxation during atrial diastole.

• v wave represents atrial filling during ventricular systole and peaks at the second heart sound.

• y descent represents passive emptying of the right atrium into the right ventricle after tricuspid valve opening.

Sold by @itachibot



Sold by @itachi... if you purchased this from some... else, ...may have been scammed.

Solution to Question 3:

The given waveform shows a cannon 'a' wave seen in ventricular tachycardia. It occurs since atria contract against a closed tricuspid valve in ventricular tachycardia.

JVP measured in the internal jugular vein is a reflection of the right atrial pressure.

Note: Giant/prominent a-waves occur during every beat while cannon a-waves result from arrhythmias, therefore, are usually intermittent. The ECG is helpful to look for rhythm disturbances that are associated with cannon a-waves whereas echocardiography can be used to assess structural defects (of right atrium) that are associated with giant a-waves.



Sold by @itachibot

Solution to Question 4:

Hypotension, soft or absent heart sounds, and jugular venous distention (Beck's triad) are the three principal features of cardiac tamponade. The typical JVP finding here is absent y-descent.

JVP shows a prominent x (early systolic) descent and an absent y (early diastolic) descent because of the ventricle's limited or absent early diastolic filling. JVP is almost always elevated in cardiac tamponade and may be associated with venous distention in the forehead and scalp.

There may be reductions in the amplitude of the QRS complexes, and electrical alternans of the P, QRS, or T waves in cardiac tamponade.

Note: Alternation of QRS complexes, usually in a 2:1 ratio, on ECG is called electrical alternans. It is caused by the movement of the heart in the pericardial space. Electrical alternans is also seen in patients with myocardial ischemia, acute pulmonary embolism, and tachyarrhythmias.



Solution to Question 5:

The y-descent in jugular venous pulse (JVP) corresponds to the fall in right atrial pressure after the tricuspid valve opening in the early diastolic phase of the right ventricle. Y-descent is attenuated or absent in cardiac tamponade because of resistance to ventricular filling in the early diastole.

Other options:

Options B and C: Restrictive cardiomyopathy and constrictive pericarditis can have deep y descent due to rapid filling of the right ventricle during early diastole.

Option D: In patients with tricuspid regurgitation, increased blood flow to the right atria through the incompetent tricuspid valve during ventricular systole results in a prominent v wave. When the tricuspid valve opens during ventricular diastole, this increased blood in the right atria enters rapidly into the right ventricle during early diastole resulting in deep y-descent.

Sold by @itachibot

Solution to Question 6:

Orthostatic hypotension is defined as a reduction in systolic blood pressure of at least 20 mmHg or diastolic blood pressure of at least 10 mmHg within 3 min of standing or head-up tilt on a tilt table.

On standing, blood is displaced to the lower extremities, causing a drop in the venous return to the heart. This drop leads to a decrease in cardiac output and stimulation of compensatory mechanisms which trigger a reflex increase in sympathetic (vasoconstrictor) outflow. Orthostatic hypotension is a manifestation of sympathetic (autonomic) failure.

Symptoms include light-headedness, dizziness, and presyncope (near-faintness) in response to sudden postural change.

Solution to Question 7:

Pulsus bisferiens is seen in hypertrophic cardiomyopathy, not pulsus paradoxus.

Pulsus paradoxus refers to the fall in systolic pressure by > 10 mmHg with inspiration. It is an exaggeration of the normal response. It is seen in the following conditions:

- Cardiac tamponade
- Massive pulmonary embolism
- Hemorrhagic shock
- Obstructive lung disease
- Tension pneumothorax

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 8:

Pulsus alternans (characterized by alternating strong and weak beats) is seen in left ventricular dysfunction.

Pulsus bisferiens is a bifid pulse with two systolic peaks. It is seen in the following conditions:

- Advanced aortic regurgitation (AR)
- Aortic stenosis (AS) with AR
- Hypertrophic cardiomyopathy

Sold by @itachibot



MARROW
Sold by @itachi... If you purchased this from some... you may have been scammed. else,

Solution to Question 9:

Water hammer pulse, also known as Corrigan's pulse, is characterized by a sharp rise and a rapid fall-off and is seen in aortic regurgitation (AR) due to the fact that arterial pressure falls rapidly in late systole and early diastole.

In AR, the pulse pressure is widened due to systolic hypertension and decreased diastolic pressure. The diastolic pressure is measured at Korotkoff phase IV. In certain patients with AR or a combination of aortic stenosis (AS) and AR, the carotid arterial pulse may exhibit a bisferiens pattern, characterized by two systolic waves separated by a trough.

The apical impulse is heaving and displaced inferiorly. A diastolic thrill may be palpable along the left sternal border.

A2 is usually absent in AR. An early diastolic murmur can be auscultated in chronic AR. It is best heard in the third intercostal space along the left sternal edge.

Sold by @itachibot



Severe AS (Option A) would show pulsus parvus et tardus, characterized by the carotid pulse rising slowly to a delayed peak.

Chronic mitral regurgitation (Option D) shows a sharp, low volume upstroke due to reduced cardiac output and acute mitral regurgitation demonstrates narrow pulse pressure.

Sold by @itachi... If you purchased this from some-else, you may have been scammed.

Solution to Question 10:

Rheumatic mitral stenosis is associated with loud S1.

The greater the distance the valve travels to close, the louder is the sound. Increased left atrial pressures in rheumatic mitral stenosis force the mobile portion of the mitral valve leaflets far apart such that they are forced closed from a relatively far distance, resulting in a loud S1. In calcific mitral stenosis, leaflet mobility is poor, and they are not separated by a large distance, resulting in a soft S1.

Loud S1 is heard in the following:

- Short PR intervals
- Early stages of mitral stenosis as long as the valves are pliable
- Atrial myxoma
- High output states
- Tachycardia

Soft S1 is heard in the following:

- Mitral regurgitation
- Aortic regurgitation
- Long PR intervals
- Calcified Mitral stenosis

Sold by @itachibot

- Left ventricular contractile dysfunction
- Beta-receptor blockers
- Pericardial effusion, obesity, COPD- due to decreased conduction through the chest wall

Solution to Question 11:

Reversed (or paradoxical) splitting refers to the situation when P2 is heard before A2. This is seen in the following:

- LBBB
- HOCM
- Right ventricular pacing
- Severe AS
- Acute myocardial ischemia

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

Loud S1 is heard in mild mitral stenosis (MS).

The intensity of S1 depends on the distance over which the anterior leaflet of the mitral valve moves, leaflet mobility, LV contractility, and PR interval. It is independent of the severity of MS.

Loud S1 is seen in early rheumatic MS, short PR interval, and hyperdynamic circulatory states.

S1 becomes soft in case of calcified leaflets, long PR interval, associated aortic regurgitation and LV dysfunction.

Solution to Question 13:

A wide split of S2 is not seen in the left bundle branch block (LBBB), however paradoxical splitting is seen.

S2 is normally heard as A2-P2. This split is exaggerated in conditions that cause early aortic valve closure or delayed pulmonary valve closure.

Option A: In atrial septal defect (ASD), left ventricle (LV) ejection time decreases due to the shunting of the blood from the left atrium (LA) to the right atrium leading to early aortic valve closure.

Option B: In mitral regurgitation (MR), due to the regurgitation of blood from the LV to the LA, the LV ejection time decreases leading to early aortic valve closure.

Option C: In pulmonary stenosis (PS), the delayed closure of pulmonary valves leads to the splitting of S2.

Sold by @itachibot

Wide splitting of S2 is also heard in RBBB and LV pacing.

Solution to Question 14:

Fixed wide splitting of S2 is seen in atrial septal defect (ASD).

The ASD creates a left to right shunt that increases the blood flow to the right side of the heart. This causes the pulmonary valve to close later than the aortic valve independent of inspiration/expiration.

Solution to Question 15:

S4 is heard during the ventricular-filling phase.

S3 and S4 are diastolic sounds that are low pitched, and best heard with the bell of the stethoscope at the cardiac apex in the left lateral decubitus position.

The presence of an S3 and S4 heart sound is called a summation gallop and sounds like a galloping horse.

S3 - Ventricular gallop	S4 - Atrial gallop
Occurs during early ventricular diastole	Occurs near the end of ventricular diastole
During the first rapid ventricular filling	During active ventricular filling (filling by atrial contraction)
Due to a sudden rush of blood entering a volume-depleted ventricle (normal ventricular compliance)	Due tovibration of the ventricular wall during atrial contraction because of increased stiffness of ventricular walls (decreased ventricular compliance)
Itcan be physiologicalin children and young adults.	Always pathological
In older patients, it indicates systolic heart failure	Indicates diastolic heart failure

Solution to Question 16:

In a patient with a right bundle branch block (RBBB), there's a widely split S1 which increases with inspiration.

S1 is normally heard as a single sound. Any delay in the closure of the tricuspid valve or early closure of the mitral valve produces a split in S1. During inspiration, venous return to the right

Sold by @itachibot

atrium increases. Therefore, the atria take a longer time to empty into the ventricles delaying the closure of the tricuspid valve. This increases the split in S1 during inspiration. In a patient with an RBBB, the tricuspid valve closure is further delayed leading to a wide split in S1.

Solution to Question 17:

The probable diagnosis of the above scenario is Ehlers-Danlos syndrome (EDS).

EDS is a group of clinically and genetically heterogeneous connective tissue disorders. Typical presenting features include joint hypermobility, skin hyperextensibility, and tissue fragility. EDS type 6 is also characterized by scoliosis. Gorlin's sign refers to the ability of patients of Ehlers-Danlos syndrome to touch the tip of the nose with the tip of their tongue.

Mitral valve prolapse (MVP) is a frequent finding in patients with EDS. An early systolic click which may be followed by a late systolic murmur is the auscultatory finding heard in MVP.

Solution to Question 18:

The ejection click is a systolic sound.

Ejection clicks are high-pitched sounds that occur at the moment of the maximal opening of the aortic or pulmonary valves. They are heard just after the first heart sound.

Ejection clicks occur in:

• Non-calcific aortic or pulmonary stenosis

• Aortic or pulmonary root dilatation

• Congenital bicuspid aortic or pulmonary valves

Nonejection clicks, which occur after the onset of the carotid upstroke, are related to the mitral valve prolapse.

Solution to Question 19:

A murmur occurring between S1 and S2 is a systolic murmur. Pulmonary regurgitation produces a diastolic murmur.

The causes of systolic murmurs are as follows:

• Mid-systolic- aortic stenosis, pulmonic stenosis, ASD, HOCM.

• Holosystolic/pansystolic- mitral regurgitation, tricuspid regurgitation, VSD.

• Late systolic- mitral valve prolapse.

The causes of diastolic murmurs are as follows:

• Early diastolic- aortic regurgitation, pulmonic regurgitation.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

• Mid/late diastolic- mitral stenosis, tricuspid stenosis, Austin-Flint.

The causes of continuous murmurs are as follows:

• Patent ductus arteriosus (machinery or Gibson's murmur).

• Venous hum

• AV fistula

• Ruptured sinus of Valsalva aneurysm

Solution to Question 20:

A mid-diastolic murmur with presystolic accentuation and an opening snap is typically seen in mitral stenosis.

The presystolic accentuation is due to the atrial contraction forcing blood through the stenosed mitral valve. The presystolic accentuation disappears in atrial fibrillation.

Option A: Mitral regurgitation presents with a holosystolic murmur, best heard at the apex with radiation to the left axilla.

Option B: Mitral valve prolapse presents with an early systolic click followed by a late systolic murmur is heard.

Option C: Aortic stenosis presents with a harsh crescendo-decrescendo ejection systolic murmur heard best at the right upper sternal border, with radiation of the murmur to the carotid arteries.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 21:

The above image shows the Valsalva maneuver. The murmur of hypertrophic cardiomyopathy (HCM) increases in intensity while performing the Valsalva maneuver.

The murmurs produced by TR (tricuspid regurgitation), AS (aortic stenosis), and MR (mitral regurgitation) decrease in intensity while performing the Valsalva maneuver.

Note: Valsalva maneuver decreases the intensity of all murmurs except MVP (mitral valve prolapse) and HCM.

Solution to Question 22:

The above image shows a sustained handgrip maneuver. The murmur of aortic stenosis decreases in intensity during sustained handgrip.

Maneuvers like sustained handgrip, simultaneous inflation of blood pressure cuffs above the systolic in both upper limbs, and infusion of vasopressors increase the systemic vascular resistance (afterload). Since increasing afterload will prevent blood from flowing in a normal forward path, it will increase any murmurs that are due to backward flowing blood and decresase the ones due to forward blood flow.

Sold by @itachibot

Murmurs that increase in intensity on sustained handgrip:

• Mitral regurgitation

• Ventricular septal defect

• Aortic regurgitation

Murmurs that decrease in intensity on sustained handgrip:

• Aortic stenosis

• Hypertrophic cardiomyopathy

Solution to Question 23:

The murmur of hypertrophic cardiomyopathy (HOCM) increases with standing, as there is a reduction in preload. In conditions of reduced preload, there is lesser end diastolic volume in the left ventricle, leading to a narrower outflow tract, and a louder murmur is heard.

The murmur heard in HOCM is a mid-systolic murmur, which is loudest along the left sternal border. It increases in conditions of decreased preload and afterload (standing, valsalva, vagal maneuvers) or conditions of increased contractility (inotropic stimulation) and decreases when preload and afterload are increased (squatting, vasopressors).

Most murmurs decrease in intensity with reduced preload and afterload except HOCM and mitral valve prolapse (MVP).

Solution to Question 24:

The murmur described above is a continuous murmur.

A continuous murmur is produced when there is a pressure gradient between the cardiac chambers or the blood vessels throughout the systole and diastole. Continous murmurs are characterised by the flow of blood in the same direction and through the same channel during systole and diastole. These murmurs begin in systole, envelope the second heart sound (S2), and continue through some portion of diastole.

Conditions producing a continuous murmur:

• Patent ductus arteriosus

• Venous hum

• AV fistula

• Ruptured sinus of Valsalva aneurysm

Ventricular septal defect with aortic regurgitation does not produce a continuous murmur. It produces a systolic-diastolic murmur as the blood flows through different channels during systole and diastole. These murmurs do not envelope S2.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Sold by @itachibot

Solution to Question 25:

The Gallavardin effect refers to the phenomenon of the transmission of the mid-systolic murmur of aortic stenosis to the apex where it becomes higher pitched.

Solution to Question 26:

A holosystolic murmur that is best heard over the lower left sternal border and radiating to the epigastrium most likely arises from tricuspid regurgitation.

Solution to Question 27:

Still's murmur refers to the benign grade 2 mid-systolic murmur heard at mid or lower left sternal border in normal children and adolescents. Louder in hyperdynamic states(fever, anxiety).

The anatomic basis for the murmur has not been definitively determined. It is best heard in a supine position.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 28:

Graham Steell murmur is the early to mid-diastolic murmur heard in pulmonary regurgitation.

It is best heard in 2nd left intercostal space and radiates along left sternal border. It can increase with inspiration.

Solution to Question 29:

Carey Coombs murmur is heard in mitral valvulitis.

It is a mid-diastolic murmur heard during an episode of acute rheumatic fever causing mitral valvulitis.

It is produced due to the flow of blood through an edematous mitral valve and dissipates with the resolution of the acute attack.

Solution to Question 30:

Austin-Flint murmur is a low-pitched mid to late diastolic murmur of chronic severe AR.

It results from the turbulence at the mitral inflow area from the mixture of regurgitant and forward blood flow. It is best heard at the apex.

Sold by @itachibot

Austin-Flint murmur is often confused with the murmur of mitral stenosis. This can be differentiated based on the following:

Features	Chronic AR	MS
Opening snap	Absent	Present
Effect of vasodilator administration	Murmur becomes shorter and softer.Mechanism- Fall in the diastolic pressure gradient between the aorta and left ventricle.	Murmur remains constant or increases in intensity.Mechanism- Reflex increase in cardiac output and mitral valve flow.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 31:

The clinical scenario of transient loss of consciousness in this patient is suggestive of syncope. Hence, an ECG must be done to rule out cardiac causes for syncope like - arrhythmias (atrial fibrillation, sinus node dysfunction, supra ventricular tachycardias) or a structural cardiac abnormality.

A CT would only be required if a seizure or transient ischemic attack (TIA) is suspected. However, in this case - the episode only lasted for a few seconds, the patient has no history of similar episodes in the past and there are no lasting neurological deficits after the episode. Therefore, option B is incorrect.

Syncope is a transient loss of consciousness caused by global cerebral hypoperfusion. It can be due to multiple causes, but is broadly divided into - neurally mediated syncope, orthostatic hypotension and cardiac syncope.

The initial workup for a case of syncope includes:

• Detailed history of the episode and past history of similar episodes

• Physical examination mainly to check BP in standing and supine positions to

• rule out orthostatic hypotension.

• A resting ECG to look for cardiac arrhythmia

• Option B: If the ECG is normal, no further tests are needed. If the ECG is suggestive of cardiac arrhythmia, then further tests including continuous ECG monitoring and electrophysiological studies may be done.

Option C: Tilt-table testing is done in patients with recurrent syncopal attacks who are suspected to have vasovagal syncope. Psychiatric evaluation is done in patients with recurrent unexplained syncopal attacks.

Option D: Vestibular neuritis is associated with vertigo, where the patients typically complain of dizziness and a sense of spinning of the head.

Sold by @itachibot

ECG Basics

Question 1:

Which electrophysiological activity is responsible for the T wave in ECG?

a) Ventricular repolarisation

b) Ventricular depolarisation

c) Atrial depolarisation

d) Atrial repolarisation

Question 2:

The PR intervals obtained from the ECG recording of 4 patients are given below. Which of the following patients has a normal PR interval?

a) Patient 1

b) Patient 2

c) Patient 3

d) Patient 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

What is the normal QRS interval?

a) < 0.20 seconds

b) <0.25 seconds

c) < 0.11 seconds

d) < 0.15 seconds

Question 4:

Which lead does the given ECG tracing represent in a normal ECG?

Sold by @itachibot

a) aVL

b) aVR

c) V1

d) V6

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 5:

In a normal ECG obtained from a patient, a biphasic P wave may be seen in which of the following lead(s)?

a) Leads II and V2

b) Leads II and V1

c) Lead V1

d) Lead V5

Question 6:

Which of the following leads would show maximum R wave with the cardiac axis as +90 degrees?

a) Lead aVF

b) Lead aVL

c) Lead I

d) Lead II

Sold by @itachibot

Question 7:

Pseudo P pulmonale is seen in which electrolyte abnormality?

a) Hypokalemia

b) Hyponatremia

c) Hypocalcemia

d) Hypercalcemia

Question 8:

Which of the following ECG findings are seen in a patient with hypokalemia?

a) Prolonged PR interval with ST elevation

b) Decreased PR interval with ST depression

c) Decreased QT interval with tall T waves

d) Prolonged QT interval with flat T waves

Question 9:

A 45-year-old male patient with a long-standing history of rheumatic heart disease comes to the clinic with complaints of dyspnea. Interpret the ECG finding.



a) Rabbit ear pattern

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

b) P mitrale

c) P pulmonale

d) Sawtooth pattern

Question 10:

A 67-year-old man with a history of COPD presented with worsening dyspnea for 2 weeks. On examination, pedal edema was present. An ECG was done and lead II showed the following pattern. Which of the following is responsible for this ECG finding?



a) Right atrial overload

b) Left atrial overload

c) Right ventricular hypertrophy

d) Left ventricular hypertrophy

Question 11:

What is the likely diagnosis of this patient based on the given ECG?

Sold by @itachibot



a) Right ventricular hypertrophy

b) Left ventricular hypertrophy

c) Torsades de pointes

d) Atrial flutter

Sold by @itachi... purchased this from some... else, ...u may have been scammed.

Question 12:

A 45-year-old man comes to the emergency room with complaints of sudden onset chest pain that radiates to the left arm and jaw. He is profusely sweating. An ECG was taken immediately. Interpret his ECG.



a) Right ventricular hypertrophy

Sold by @itachibot

b) Right bundle-branch block

c) Left bundle-branch block

d) Ventricular tachycardia

Question 13:

A 50-year-old patient comes to the emergency room with a history of severe retrosternal pain radiating to the jaw. The patient initially felt some uneasiness while eating lunch after which he experienced pain. His ECG is as shown. Interpret his ECG.



a) Atrial flutter

b) WPW syndrome

c) Myocardial infarction

d) Ventricular tachycardia

Question 14:

A 72-year-old man comes to the emergency room complaining his heart is pounding and that he is going to die. The emergency physician orders an ECG. What is the diagnosis?

Sold by @itachibot



©MARROW

a) Inferior wall MI

b) Extensive anterior wall MI

c) Anterolateral wall MI

d) Anteroseptal MI

Sold by @itachi... purchased this from some... else, If you may have been scammed.

Question 15:

A 52-year-old diabetic man presents with retrosternal chest pain and profuse sweating and lightheadedness. Based on the ECG findings, what is the diagnosis?



a) Extensive anterior wall MI

Sold by @itachibot

b) Anterolateral wall MI

c) Anteroseptal MI

d) Atrial flutter

Question 16:

What is the diagnosis based on the given ECG?



©MARROW

a) Anterior wall MI

b) Lateral wall MI

c) Inferior wall MI

d) Anteroseptal MI

Question 17:

A 60-year-old patient presents with chest pain on the left side radiating to the arm and jaw on the same side. Cardiac troponins are found to be raised. What is the diagnosis based on this ECG?

Sold by @itachibot



a) Anterior wall MI

b) Lateral wall MI

c) Inferior wall MI

d) Posterior wall MI

Question 18:

Which of the following ECG findings are unlikely to be present in a patient with hyperkalemia?

a) Prolonged PR interval

b) Prolonged QT interval

c) Prolonged QRS interval

d) Loss of P waves

Question 19:

A patient was brought to the ER with complaints of multiple episodes of vomiting, increased frequency of urination, and constipation for the past 5 days. He also complained of lethargy and muscle pain. He is a known case of SCC of the lung. What is likely to be seen in the ECG of this patient?

a) ST segment depression

b) Decreased QT interval

c) Increased PR interval

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Sold by @itachibot

d) Tall T waves

Question 20:

ECG of a patient revealed a QT interval of 0.55s. Which of the following will not be a differential diagnosis?

a) Hypothermia

b) Digitalis toxicity

c) Hypocalcemia

d) Romano-Ward syndrome

Question 21:

Osborn waves in ECG are seen in:

a) Hypothyroidism

b) Hypothermia

c) Hypocalcemia

d) Hypokalemia

Question 22:

In a patient with a left-sided massive pneumothorax, all of the following ECG changes are seen except:

a) Reduced R wave amplitude

b) Precordial T wave inversion

c) Decreased QRS amplitude

d) Left axis deviation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a

Sold by @itachibot

2	a
3	c
4	b
5	c
6	a
7	a
8	d
9	b
10	a
11	a
12	c
13	c
14	d
15	a
16	b
17	d
18	b
19	b
20	b
21	b
22	d

If you pu... you may ... else, ...d.

Detailed Explanations

Solution to Question 1:

The T wave in ECG is due to ventricular repolarisation.

Solution to Question 2:

The normal duration of the PR interval is 0.12-0.2 seconds (3 to 5 small boxes).

PR interval is said to be prolonged if it is > 0.2 seconds and short if it is < 0.12 seconds.

It is measured from the beginning of the P wave to the beginning of the QRS complex. It represents the time from the start of atrial depolarization to the start of ventricular depolarization.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

The normal duration of the QRS interval is normally 0.1-0.11 seconds (less than 3 small boxes).

QRS interval represents the duration of ventricular depolarization. When the QRS complex is > 0.11 seconds, it is called a wide QRS complex. It can occur in:

- Bundle branch blocks (RBBB or LBBB)
- Depolarizations originating in the ventricles
- Wolff-Parkinson-White syndrome
- Hyperkalaemia

Solution to Question 4:

As all the waves (P, QRS, T) are negative in the given ECG, it should be lead aVR. All the depolarisations waves (P, QRS, T) move away from lead aVR in the normal sinus.

Depolarisation wave moving towards the positive pole of the lead (or active electrode) produces a positive deflection in the ECG and depolarization moving away from the positive pole of the lead produces a negative deflection in the ECG.

The lead aVR provides information from the right upper side of the heart.

Sold by @itachibot



The given image shows a normal 12 lead ECG:



Solution to Question 5:

In a normal ECG, a biphasic P wave is seen in lead V1 only.

Lead V1 is a chest lead. So, it records potentials in the horizontal plane. The P wave in lead V1 consists of:

- Positive component reflecting right atrial depolarization (due to anteriorly oriented vector)
- Negative component reflecting left atrial depolarization (due to posteriorly oriented vector)

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

With the cardiac axis at +90 degrees, lead aVF would show the maximum R wave.

The cardiac axis represents the mean QRS axis (direction of mean depolarization vector).

The normal adult QRS axis is directed downward and to the left, which is between -30 degrees and +100 degrees.

The position of lead aVF is at +90 degrees. If the cardiac axis is also at +90 degrees, it would be in the same direction as lead aVF and hence aVF would show maximum R wave amplitude.

If the mean axis is:

- More negative than –30 degrees: left axis deviation
- More positive than +90 to +100 degrees: right axis deviation
- Lies between -90 to +180 degrees: extreme axis deviation

Sold by @itachibot



Solution to Question 7:

Pseudo P pulmonale refers to tall peaked P waves seen in hypokalemia.

P-pulmonale refers to P waves taller than >2.5mm in an ECG lead.



Hypocalcemia (Option C) prolongs the QT interval.

Hypercalcemia (Option D) shortens the QT interval.

Solution to Question 8:

Prolonged QT interval with flat T waves is seen in hypokalemia.

Sold by @itachibot

Hypokalemia prolongs ventricular repolarization. The ECG changes seen are as follows:

• Flattening of T wave

• ST-segment depression

• Prolongation of QT interval

• The appearance of U wave



Solution to Question 9:

The P waves in the given ECG are broad and notched. This is the P mitrale pattern which is seen in left atrial enlargement.

This patient has likely developed mitral stenosis (heard as a mid-diastolic murmur) secondary to his rheumatic heart disease. Mitral valve stenosis and atrial inflammation in rheumatic carditis lead to left atrial dilation.

The features of P mitrale are:

• lead II: broad (>2.5mm) notched P waves

• Lead V1: Biphasic wave with a prominent negative terminal component (representing delayed depolarization of the LA)

P pulmonale is seen in right atrial enlargement.

Sold by @itachibot



Option A: The rabbit ear pattern is seen in the right bundle branch block. The ECG of RBBB below shows the rSR' pattern of the QRS complex (rabbit ear) in leads V1-V2 and slurred S wave in lead V6.



Option D: Sawtooth pattern in ECG is seen in atrial flutter (shown below).

Sold by @itachibot



Solution to Question 10:

In the given ECG, a peaked P wave can be seen. This is called P pulmonale and it is seen in cases of right atrial enlargement. The patient is showing features of heart failure possibly as a consequence of dysfunction caused by the overload.

Signs of heart failure in this patient:

• Bilateral pitting edema in lower extremities (systemic congestion)

• Worsening dyspnea due to increased PCWP causing pulmonary hypertension and pulmonary edema (pulmonary congestion)

The features of P pulmonale include:

• The amplitude of the P wave is >2.5mm on Lead II

• Prominent positive wave in the biphasic P wave of lead V1

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 11:

The given ECG shows deep S-waves in V6 and large R-waves in V1. This is suggestive of right ventricular hypertrophy (RVH).

Diagnostic criteria for RVH:

• Dominant R wave in V1 (R/S ratio > 1).

• Dominant S wave in V5 or V6 (R/S ratio < 1).

• Right ventricular strain pattern - ST depression/T wave inversion in the right precordial (V1-4) and inferior leads (II, III, aVF)

• Right axis deviation of +110° or more.

For left ventricular hypertrophy, Sokolow Lyon criteria are commonly used.

It is based on the presence of tall left precordial R waves and deep right precordial S waves (S wave in V1 + [R wave in V5 or V6] >35 mm).

Sold by @itachibot



©MARROW

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

The patient has a myocardial infarction (MI) with ECG changes of a left bundle-branch block (LBBB), which can be identified by the following classical features:

• Wide QRS complex- QRS ≥120 ms

• Predominantly negative (QS) complexes in lead V1 and entirely positive (R) complexes in lead V6.

• M pattern seen in left chest leads (V5, V6)

MI can have the following ECG changes:

• ST-segment elevation- zone of injury

• T wave depression

• Deep Q waves- late sign to develop

• New left bundle branch block (LBBB)

Sold by @itachibot



The ECG findings below are characteristic of LBBB: Mnemonic WILLIAM (W pattern in V1-V2; L-LBBB; M pattern in V3-V6)



The ECG findings below are characteristic of RBBB: Mnemonic MARROW (M pattern in V1-V2; R-RBBB; W pattern in V3-V6)

Sold by @itachibot

Sold by @itachi
If you purchased this from some
you may have been scammed.
else,

Solution to Question 13:

The given ECG shows ST-segment elevation in the anterior chest leads suggestive of myocardial infarction.

Features of MI on ECG:

- ST-segment elevation: Zone of injury
- T wave inversions
- Deep Q waves: late sign to develop
- New left bundle branch block (LBBB)

Option A: Atrial flutter shows a sawtooth wave pattern.

Option B: WPW syndrome shows delta waves.

Option D: Ventricular tachycardia has wide QRS complex and increased heart rate.

Solution to Question 14:

The given ECG shows ST elevation in leads V1-V4 which is suggestive of acute anteroseptal myocardial infarction (MI). ST depression in the leads III and aVF signifies an old inferior wall infarction.

Sold by @itachibot

Solution to Question 15:

The ECG shows ST elevations in leads I, aVL, and V1-V6. This is suggestive of extensive anterior wall MI.

Solution to Question 16:

The ECG is suggestive of lateral wall MI and possible posterior wall MI.

ST-elevation is present in the high lateral leads (I and aVL) and in V5 and V6.

Reciprocal ST depression is seen in the inferior leads (III and aVF) with associated ST depression in V1, V2, and V3 (which could represent posterior wall MI).

This pattern is consistent with an acute infarction localized to the superior portion of the lateral wall of the left ventricle (high lateral STEMI).

Solution to Question 17:

The given ECG is of posterior wall MI.

The ECG findings in posterior wall MI include;

• Reciprocal ST depressions in leads V1 to V3. Posterior wall ischemia is indirectly recognized in this way

• Tall and broad R waves (>30ms) in V2 and V3

• Dominant R waves (R/S ratio > 1) in V2

• Upright terminal portions of the T waves in V2 and V3

Solution to Question 18:

Prolonged QT interval is not a feature of hyperkalemia, it is a feature of hypokalemia.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Sold by @itachibot



MARROW
Sold by @itachi... If you purchased this from someone else, they may have been scammed.

Solution to Question 19:

The above scenario is suggestive of hypercalcemia due to ectopic production of PTH-related peptides. It is mostly associated with squamous-cell lung cancer (SCC). ECG is likely to show a decreased QT interval.

Hypocalcemia shows increased QT interval.

Solution to Question 20:

QT interval of 0.55s suggests QT prolongation. It is not seen in digitalis toxicity. Rather, a QT shortening with characteristic “scooping” of the ST–T-wave complex, called a digitalis effect, is seen.

The ECG changes seen in digitalis toxicity are as follows:

• Prolonged PR interval

• ST-segment depression

• Short QT interval

QT interval represents the time taken for the ventricles to depolarize and repolarize. It is prolonged in the following:

• Hypocalcemia

• Hypokalemia

• Congenital long QT syndromes such as Romano-Ward syndrome

• Hypothermia

• Drugs: Class IA and Class III antiarrhythmics

Sold by @itachibot

• Intracranial bleeds: particularly SAH

Solution to Question 21:

Osborn waves are seen in hypothermia.

Osborn waves can also be seen in severe hypothyroidism (Option A), but even in those cases, hypothermia is the reason for the appearance of Osborn waves. Therefore, hypothermia is the better answer here.

It consists of an abrupt ascent right at the J point (the junctional point between the QRS complex and the ST segment) and then an equally sudden plunge back to the baseline. It is also called the J wave. This is because systemic hypothermia prolongs repolarisation.



Osborn wave can also be seen in:

• Hypercalcemia

• Brain injury

• Subarachnoid hemorrhage

• Vasospastic angina

• Cardiopulmonary arrest from oversedation

• Idiopathic ventricular fibrillation

Solution to Question 22:

Right axis deviation is seen in left-sided pneumothorax.

The ECG changes seen in left-sided pneumothorax are as follows:

Sold by @itachibot

- Reduced R wave amplitude
- Inversion of T wave in precordial leads
- Amplitude changes in QRS complex
- Right axis deviation

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Supraventricular Arrhythmias

Question 1:

A 52-year-old man has come to the emergency complaining of a burning type of chest pain. Comment on his ECG recording shown below.



a) Sinus bradycardia

b) Sinus tachycardia

c) Normal sinus rhythm

d) Irregular rhythm

Question 2:

All of the following are supraventricular tachycardias except:

a) AV nodal reentry tachycardia

b) Sinus tachycardia

c) Junctional tachycardia

d) Premature ventricular beats

Question 3:

Sold by @itachibot

ECG of a patient who presented with palpitations to the ER is shown below. Which of the following is a likely diagnosis?



a) Sinus tachycardia

b) Ventricular tachycardia

c) Junctional tachycardia

d) Atrial flutter

If you purchased this from Sold by @ you may have been scammed. else,

Question 4:

Which is the most common supraventricular tachyarrhythmia?

a) Paroxysmal supraventricular tachycardia

b) Sinus tachycardia

c) Atrial fibrillation

d) Junctional tachycardia

Question 5:

A 49-year-old woman was brought to the emergency with complaints of dizziness, shortness of breath, and palpitations. ECG showed tachyarrhythmia with narrow QRS complex and irregular RR intervals. What is the likely diagnosis?

a) Atrioventricular re-entry tachycardia

b) AV nodal re-entry tachycardia

Sold by @itachibot

c) Atrial flutter

d) Atrial fibrillation

Question 6:

A 65-year-old hypertensive patient presents with chest pain and dyspnea following an alcohol binge. His ECG is shown below. What is the most likely diagnosis?



a) Ventricular tachycardia

b) Focal atrial tachycardia

c) Ventricular fibrillation

d) Atrial fibrillation

Question 7:

A 50-year-old man was diagnosed with STEMI. He was admitted and given thrombolytic therapy. The next day he complained of palpitations and an ECG was taken. Which of the following drugs can be used to treat his condition?

Sold by @itachibot



a) Amiodarone

b) Verapamil

c) Flecainide

d) Propafenone

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 8:

A 60-year-old man who is on treatment for chronic AF presents with nausea, dizziness, and anorexia. He has a regular heart rate of 50/min. The most likely cause is:

a) Thromboembolic stroke

b) Myocardial infarction

c) Non compliance to treatment

d) Digitalis toxicity

Question 9:

ECG of a patient who was brought to the casualty due to severe chest pain followed by syncope is given below. What is the likely diagnosis?

Sold by @itachibot



a) Atrial fibrillation

b) Atrial flutter

c) Ventricular tachycardia

d) Ventricular fibrillation

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 10:

A 50-year-old man was brought to the emergency in an unconscious state. He had a fever for the past two days and is a known case of severe COPD. His ECG is given below. What is the most likely diagnosis?



a) Ventricular tachycardia

Sold by @itachibot

b) Multifocal atrial tachycardia

c) Atrial fibrillation

d) Atrial tachycardia

Question 11:

Paroxysmal supraventricular tachycardias do not include:

a) Atrial fibrillation

b) Atrial tachycardia

c) AV nodal reentry tachycardia

d) AV reentry tachycardia

Question 12:

A 32-year-old man is brought to the casualty after he developed sudden onset dyspnea, chest pain, and palpitations at a party. His blood pressure is 100/62 mmHg. His ECG is shown below. What is the drug of choice to treat his condition?



a) Procainamide

b) Adenosine

c) Verapamil

d) Propranolol

If you bought this from someone else, it has been scammed. Sold by @itachibot

Sold by @itachibot

Question 13:

The ECG of a patient who presented to the ER is as given below. The patient was already given a carotid massage and IV Adenosine but the BP is currently 60/30 mm Hg. What is the next best step of management?



a) Repeat Inj adenosine 6 mg

b) Inj amiodarone 300 mg

c) Synchronised cardioversion

d) Defibrillation

If you purchased this ... you may have been ... else, ... d.

Question 14:

A patient came with sudden onset palpitations. ECG showed a narrow complex, regular tachycardia without P waves, and a rate of 160 BPM. His blood pressure is 96/68 mmHg. What is the next step in management?

a) Carotid sinus massage

b) Adenosine

c) Beta-blocker

d) Defibrillator

Question 15:

Sold by @itachibot

A 65-year-old female patient is brought to the emergency department in a state of unconsciousness. Her BP is 70/50 mm of Hg. Her ECG is shown below. What is the next best step in the management of her condition?



a) IV verapamil

b) IV adenosine

c) Carotid massage

d) DC cardioversion

Sold by
if you purchased this fro
you may have been scam
else,
d.

Question 16:

Which of the following is false about accessory pathways?

a) Bundle of Kent has antegrade conduction

b) Concealed pathways have retrograde conduction

c) Antegrade conduction causes preexcitation

d) Retrograde conduction causes preexcitation

Question 17:

Which of the following is the most common reentrant tachycardia caused by an accessory pathway?

a) Orthodromic AV reentry

b) Antidromic AV reentry

Sold by @itachibot

c) WPW syndrome

d) AV nodal reentry

Question 18:

A patient's ECG shows a short PR interval and delta wave. Additional findings include atrial fibrillation and rapid ventricular rate. He is hemodynamically stable. All of the following drugs can be used in the management except:

a) Amiodarone

b) Ibutilide

c) Procainamide

d) Verapamil

Question 19:

The most common location of the accessory pathway in WPW syndrome is:

a) Left free wall

b) Posteroseptal

c) Right free wall

d) Anteroseptal

Question 20:

All are true about the condition that is diagnosed by the given ECG except:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) It is a congenital condition

b) Can be seen in asymptomatic individuals

c) Prevalence increases with age

d) Caused by an abnormal accessory pathway

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 21:

A 30-year-old woman comes to the clinic with complaints of recurrent palpitation and lightheadedness. An ECG was taken and is as shown below. What is the best step in management?



a) DC cardioversion

Sold by @itachibot

b) Carotid sinus massage

c) Synchronised cardioversion

d) Catheter ablation

Question 22:

Which of the following is true regarding sick sinus syndrome?

a) Most patients are symptomatic in the early stages

b) Most commonly in young patients

c) Frequently seen in patients with atrial fibrillation

d) Treated using beta-blockers

If you bought this from someone else, you've been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	c
2	d
3	b
4	b
5	d
6	d
7	a
8	d
9	b
10	b
11	a
12	b
13	c
14	a
15	d
16	d
17	a
18	d
19	a

Sold by @itachibot

20	c
21	d
22	c

Detailed Explanations

Solution to Question 1:

The above ECG strip shows normal sinus rhythm.

The features in this ECG that suggest this diagnosis are as follows:

• The QRS complexes are narrow (<120 ms or 3 small squares) which implies that the impulse is supraventricular and is conducted via the normal conduction pathway.

• Each QRS complex is preceded by a P wave. This means that all the atrial depolarizations that arise in the sinus node are conducted to the ventricles.

Sold by @itachibot

• The PR interval is constant for all the impulses and is normal (<200 ms) in duration.

• The RR interval is constant which implies a regular rhythm.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... you may have been scammed. else,

Solution to Question 2:

Premature ventricular beats are a type of ventricular tachycardias, not supraventricular tachycardias.

Tachycardia is defined as a heart rate of >100 beats/min. It can be classified based on the site of origin of the cardiac impulse as follows:

• Supraventricular tachycardias (SVTs) originate above the ventricles, that is, in the sinoatrial node, atria, or the atrioventricular node. Supraventricular tachycardias include:

• Tachycardias originating from the atrium:

• Sinus tachycardia- from the SA node

• Focal atrial tachycardia

• Multifocal atrial tachycardia

• Atrial flutter

• Atrial fibrillation

• Tachycardias originating from the AV node:

• Junctional tachycardia- automaticity within AV Node

• AV nodal reentry tachycardia (AVNRT)

• Tachycardia associated with accessory atrioventricular pathways: atrioventricular reentry tachycardia (AVRT)

• Ventricular tachycardias originate from the ectopic foci or the reentry pathways in the ventricles. Ventricular tachycardias include:

• Premature ventricular beats

• Ventricular tachycardia (VT)

Sold by @itachibot

• Monomorphic VT- Very rapid monomorphic VT is also known as ventricular flutter

• Polymorphic VT- associated with Torsades de pointes

• Ventricular fibrillation



Solution to Question 3:

The above ECG shows increased heart rate with wide QRS complexes (> 0.12s). This is suggestive of ventricular tachycardia (VT).

Types of ventricular tachycardia based on QRS morphology:

• Monomorphic- a stable QRS morphology from beat to beat

• Polymorphic- changing or multiform QRS variance from beat to beat. Torsades de pointes is an example of polymorphic ventricular tachycardia.

• Bidirectional- has a beat-to-beat alternation in the QRS frontal plane axis. It is associated with digitalis toxicity or catecholaminergic polymorphic VT.

The most common cause of VT is ischemic heart disease.

Ventricular tachyarrhythmias occur due to impulses that originate within the ventricles. These impulses are conducted through the normal ventricular myocardial cells instead of the fast conducting Purkinje system. Hence, they present with wide QRS complexes (>0.12 sec) on the ECG.

Supraventricular tachyarrhythmias occur due to impulses originating above the ventricles. These impulses reach the ventricles through the proper fast conduction pathway resulting in a narrow QRS complex. SVTs are associated with wide QRS in few exceptional conditions like:

• Conduction block in the left or right bundle branch

• Activation of ventricles from an accessory pathway- antidromic AVRT

The given ECG shows wide stable QRS complexes in monomorphic ventricular tachycardia:

Sold by @itachibot

Sold by @itachibot



Solution to Question 4:

The most common supraventricular arrhythmia is sinus tachycardia. It can either be physiological or pathological. Physiological sinus tachycardia is seen in exercise, fever, anemia, hyperthyroidism, etc.

Paroxysmal supraventricular tachycardias are those supraventricular tachycardias that occur with sudden onset and termination, they are non-sustained arrhythmias (lasts < 30 seconds). They include AVNRT, AVRT, and atrial tachycardias.

The most common sustained (lasts > 30 seconds) arrhythmia is atrial fibrillation.

Junctional tachycardia originates from ectopic foci in the AV node. It is an uncommon tachycardia.

Sold by @itachibot

Solution to Question 5:

ECG showing tachyarrhythmia with a narrow QRS complex suggests a supraventricular tachycardia. An irregular RR interval is seen with the following supraventricular tachycardias:

- Atrial fibrillation
- Multifocal atrial tachycardia
- Atrial flutter/ atrial tachycardia with variable block

The ECG below shows atrial fibrillation. Note the narrow QRS complexes and irregular RR intervals. Disappearacne of P-wave is another feature.





Sold by @itachibot

Solution to Question 6:

The ECG shows narrow QRS complexes with irregular RR intervals and no distinct P waves. This is suggestive of atrial fibrillation.

Atrial fibrillation (AF) is a supraventricular tachycardia caused by ectopic foci or small reentry circuits within the atria. It has an atrial rate of 400-600/min and a ventricular rate of 100-160/min.

Risk factors for developing AF include:

• Age- usually above 60 years of age

• Underlying heart disease

• Hypertension

• Diabetes mellitus

• Obesity

• Sleep apnea

• Post-cardiac surgery

• Inflammatory pericarditis

Acute precipitating factors for AF include:

• Hyperthyroidism

• Acute alcohol intoxication

• Acute illness like myocardial infarction and pulmonary embolism

Sold by @Itachibot
-Ihsed this from someone else,
ve been scammed.



Solution to Question 7:

Sold by @itachibot

The above ECG shows narrow QRS complex, irregular RR intervals, and absent P waves which are suggestive of atrial fibrillation. Among the given options, amiodarone can be administered in patients having atrial fibrillation with coronary or structural heart disease.

Drugs for rhythm control in atrial fibrillation:

In patients without structural heart disease, class I agents are used. Class I agents act by blocking the Na+ channels leading to slower depolarization:

• Flecainide

• Propafenone

• Disopyramide

In patients with coronary or structural heart disease, class III agents are used. Class III agents act by blocking the K+ channels leading to slower repolarization causing increased refractory period:

• Amiodarone

• Dofetilide

• Sotalol

• Ibutilide



Solution to Question 8:

A patient of atrial fibrillation usually has an irregular heart rate of >100/ min but this patient presents with a regular slow heart rate. The most likely cause of slow regular heart rate in chronic AF is digitalis toxicity. Symptoms such as nausea, anorexia are also suggestive of digitalis toxicity.

Digitalis lowers the AV nodal conduction by stimulating the vagal nerve endings leading to increased vagal tone. Hence digitalis toxicity can cause AV nodal block. This leads to a junctional escape rhythm taking over the ventricles causing slow and regular heart rate.

Sold by @itachibot

Digitalis toxicity can also lead to a variety of arrhythmias as it increases the intracellular calcium levels increasing the risk of tachyarrhythmias.

Solution to Question 9:

The given ECG is suggestive of atrial flutter.

The given ECG shows narrow QRS complexes, hence it is supraventricular tachycardia. The RR intervals are regular and the normal P waves are replaced by sawtooth-shaped waves, known as flutter waves.

In atrial flutter, the ventricular rate is determined by the AV conduction ratio. The commonest AV ratio is 2:1. In the above ECG, the rhythm is atrial flutter 4:1 block with an atrial rate of about 270 and a ventricular rate of 68/min.

The image below shows sawtooth shaped waves in atrial flutter.



Solution to Question 10:

The given clinical scenario of chronic pulmonary disease and the ECG showing different p wave morphologies are suggestive of multifocal atrial tachycardia.

Multifocal atrial tachycardia (MAT) occurs due to impulses arising from multiple abnormal foci in the atria. This leads to supraventricular tachycardia with p waves and QRS complexes of different morphologies. It is usually encountered in patients with chronic pulmonary disease and acute illness.

Therapy for MAT is directed at treating the underlying disease and correcting any metabolic abnormalities. Calcium-channel blockers and radiofrequency ablation of the AV node are the other treatment modalities.

Sold by @itachibot

The given ECG shows narrow QRS complexes with irregular RR intervals, hence it can be atrial fibrillation or multifocal atrial tachycardia. The next step is to look at the p waves. Multifocal atrial tachycardia (MAT) is characterized by at least 3 distinct P-wave morphologies (shown by arrows in the image). Whereas, in atrial fibrillation, there would be no distinct p waves seen.



Sold by
If you purchased this fr
you may have been sca
else,

Solution to Question 11:

Paroxysmal supraventricular tachycardias do not include atrial fibrillation.

Paroxysmal supraventricular tachycardias are tachycardias that occur with sudden onset and termination, they are not sustained arrhythmias. They include

- AVNRT
- AVRT
- Atrial tachycardia

Whereas, atrial fibrillation is the most common sustained arrhythmia among supraventricular tachycardias.

Features suggestive of PSVT:

- Narrow complex tachycardia at 150 bpm
- No visible P waves
- Pseudo R' waves due to retrograde atrial activation

The given ECG shows pseudo R' waves:

Sold by @itachibot



Solution to Question 12:

The ECG is suggestive of paroxysmal supraventricular tachycardia (PSVT) as it shows:

• Narrow QRS complexes

• No visible p waves

• Pseudo R' waves in V1 lead

Adenosine is the preferred drug in stable paroxysmal supraventricular tachycardia (PSVT).

Treatment of PSVT:

In hemodynamically unstable patients, QRS-synchronous DC cardioversion should be done.

In hemodynamically stable patients,

1. Mechanical measures: These maneuvers result in an acute increase in the vagal tone which decreases AV conduction and terminates the arrhythmia. They include the following:

• Valsalva maneuver

• Lowering the head between the knees

• Coughing

• Splashing cold water on the face

• Breath-holding

• Carotid sinus massage

2. Pharmacologic therapy:

• Adenosine- it is the first-line agent. It blocks conduction across the AV node and terminates the SVT. It is given as a 6-18 mg bolus

• Calcium channel blockers- verapamil and diltiazem

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Sold by @itachibot

• Beta-blockers- esmolol, propranolol, and metoprolol

• IV Procainamide- it is administered if the tachycardia is being mediated through an accessory pathway. It facilitates the AV conduction and therefore causes an initial increase in rate. Therefore, it is given after administering calcium channel blockers or beta-blockers.

The given ECG shows pseudo R' waves:





Solution to Question 13:

The given ECG showing narrow QRS complex (<0.12 s) tachycardia with absent P wave is suggestive of paroxysmal ventricular tachycardia (PSVT). The next best step in the management of a patient with unstable vitals (systolic blood pressure < 90 mm Hg) is synchronized cardioversion.

Sold by @itachibot

Synchronized cardioversion is the treatment of choice in unstable PSVT patients, in which shocks are delivered with the defibrillator/monitor in synchronized mode with an initial energy level of 100J. Synchronization times the delivery so that it is given at the peak of the QRS complex.

If the shock occurs in the ST segment or the T wave (unsynchronized), it can precipitate a more serious arrhythmia such as ventricular fibrillation.

Synchronized cardioversion	Defibrillation
Used inunstable PSVT, atrial flutter, atrial fibrillation, and ventricular tachycardiawith a pulse	Used in ventricular tachycard iawithout pulseand ventricul ar fibrillation.
Shock is delivered in sync wit h the R wave (in absolute refr actory period)	Shock is delivered at any tim e during the cardiac cycle.
Monophasic: 100-200 JBipha sic: 50-100 J	Monophasic: 360 JBiphasic:1 20-200 J

Other options:

Option A: In the absence of hemodynamic instability, a repeat dose of injection adenosine (maximum of 12mg) could be attempted.

Option B: IV Amiodarone is rarely used in the acute management of PSVT. Instead, it is used as an alternative to calcium channel blockers in chronic suppressive therapy.

Option D: Defibrillation is only done for ventricular fibrillation and pulseless ventricular tachycardia.



Solution to Question 14:

Sold by @itachibot

The given clinical scenario and the electrocardiogram (ECG) showing narrow QRS complex regular tachycardia without P waves are suggestive of paroxysmal supraventricular tachycardia (PSVT). The next step in the management of PSVT in a hemodynamically stable (systolic blood pressure >90 mm Hg) patient is vagotonic maneuvres such as carotid sinus massage or Valsalva maneuver.

Carotid sinus massage can be considered in the absence of carotid bruits and no history of stroke. If vagal maneuvers fail or cannot be performed, intravenous adenosine can be used.

Non-DHP calcium channel blockers and beta blockers are second-line pharmacological agents in the management of PSVT. They can cause hypotension before and after terminating the arrhythmia.

If the patient is hemodynamically unstable (systolic blood pressure <90 mmHg or mean arterial pressure <65 mm Hg), immediate QRS-synchronous direct current cardioversion is necessary.

The treatment algorithm for PSVT is described in the flowchart below:



Solution to Question 15:

The given clinical scenario describes an unconscious patient, who is hemodynamically unstable (systolic BP <90 mmHg or mean arterial pressure <65 mmHg). The ECG shows a narrow QRS complex tachycardia (heart rate of 188 BPM) with regular RR interval and absent P waves, characteristic of paroxysmal supraventricular tachycardia (PSVT). The next step in management is QRS-synchronized DC cardioversion.

Shown below is the algorithm for the management of supraventricular tachycardia:

Sold by @itachibot



Options A, B, and C are all used in stable cases. Vagal maneuvers like carotid massage or Valsalva are tried first followed by IV Adenosine. If these are ineffective, IV Verapamil, Diltiazem, or Beta blockers are used.

Synchronized cardioversion	Defibrillation
Used in sustained VTUsed in SVT (unstable cases), atrial fl utter/fibrillation	Used in VT and VF without a pulse
The shock delivered insync w ith R wave(in absolute refract ory period)	The shock is delivered at any time during cardiac cycle
Characteristics of QRS compl ex:Narrow and regular - 50 t o 100JNarrow and irregular - 120 to 200JWide and regul ar - 100JWide and irregular - No synchronised cardiovers ion	Wide and irregular QRS com plexMonophasic: 360 JBipha sic:1 20-200 J

Solution to Question 16:

Retrograde conduction via accessory pathways does not cause preexcitation.

Accessory pathways (APs) are abnormal conduction pathways connecting the atrium and ventricle. They can conduct impulses in the following ways:

Antegrade conduction:

• Occurs from the atria to the ventricle

Sold by @itachibot

• The impulses from atria reach the ventricles faster via the accessory pathway as compared to the normal AV nodal pathway as they bypass the AV nodal delay.

• This leads to an earlier excitation of the ventricles known as the preexcitation of the ventricles and thus a wide QRS complex.

• Examples- WPW syndrome and antidromic AVRT

Retrograde conduction:

• Occurs from the ventricle to the atria.

• The impulses reach the ventricles via the normal AV nodal pathway but re-enter the atria through the accessory pathway.

• Ventricular depolarization is normal, there is no preexcitation and thus a narrow QRS complex. It causes supraventricular tachycardia as the impulses re-enter the atria via the accessory pathway.

• Examples- concealed accessory pathways (can cause SVTs) and orthodromic AVRT.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 17:

The most common reentrant tachycardia caused by an accessory pathway is orthodromic AV reentry.

Accessory pathways (APs) are abnormal conduction pathways connecting the atrium and ventricle. They can conduct impulses in the following ways:

Antegrade conduction:

• Occurs from the atria to the ventricle

• The impulses from atria reach the ventricles faster via the accessory pathway as compared to the normal AV nodal pathway as they bypass the AV nodal delay.

• This leads to an earlier excitation of the ventricles known as the preexcitation of the ventricles and thus a wide QRS complex.

• Examples- WPW syndrome and antidromic AVRT

Retrograde conduction:

• Occurs from the ventricle to the atria.

• The impulses reach the ventricles via the normal AV nodal pathway but re-enter the atria through the accessory pathway.

• Ventricular depolarization is normal, there is no preexcitation and thus a narrow QRS complex. It causes supraventricular tachycardia as the impulses re-enter the atria via the accessory pathway.

• Examples- concealed accessory pathways (can cause SVTs) and orthodromic AVRT.

AV nodal reentry tachycardia (AVNRT) is not caused by an accessory pathway. It involves tissues with different conduction speeds within the AV node or just adjacent to the AV node, known as the fast pathway and the slow pathway. This is known as dual AV nodal physiology.

Sold by @itachibot

Solution to Question 18:

The given clinical scenario describes a patient with WPW syndrome with atrial fibrillation.

Rate-slowing drugs such as non-dihydropyridine calcium channel blockers (eg, verapamil, diltiazem), beta-blockers, and digoxin are contraindicated in these patients. They increase the refractoriness of the AV node with minimal or no effect on the accessory pathway. This can lead to faster ventricular rates, increasing the risk of ventricular fibrillation.

The ECG below shows a short PR interval (<120 ms) and delta waves in WPW syndrome.



Solution to Question 19:

The most common location of an accessory pathway in WPW syndrome is on the left free wall. It is present between left atrium and free wall of left ventricle.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 20:

The given ECG is suggestive of Wolff-Parkinson-White syndrome (WPW). Its prevalence decreases with age.

The classic ECG features in WPW are as follows:

- A shortened PR interval (typically <120 ms)
- A slurring and the slow rise of the initial upstroke of the QRS complex (delta wave)
- A widened QRS complex (total duration >0.12 seconds)
- ST-segment–T wave (repolarization) changes, generally directed opposite the major delta wave and QRS complex, reflecting altered depolarization

Sold by @itachibot



Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Solution to Question 21:

The ECG shows the presence of delta waves (preexcitation wave), wide QRS, and short PR interval which are classically seen in Wolff Parkinson White (WPW) syndrome. In this patient, catheter ablation is the best for long-term management.

Patients with preexcitation who have symptoms of arrhythmia are at risk for developing AF and sudden death if they have an accessory pathway that allows rapid antegrade conduction (preexcitation & tachyarrhythmias are features of WPW syndrome due to the abnormal accessory pathway).

Complications of catheter ablation are rare but include AV block, cardiac tamponade, thromboembolic events, and hematoma at the access site.

Solution to Question 22:

Sick sinus syndrome is frequently seen in patients with concomitant atrial fibrillation.

Sick sinus syndrome (or sinus node dysfunction) is a group of abnormal heart rhythms. It is a disorder of the sinoatrial node caused by impaired pacemaker function and impulse transmission. This leads to a combination of abnormal rhythms, including atrial bradyarrhythmias, atrial tachyarrhythmias, and sometimes a pattern of alternating bradycardia and tachycardia, known as tachy-brady syndrome. These irregular rhythms can cause symptoms such as palpitations, decreased tissue perfusion, subsequent fatigue, lightheadedness, pre-syncope, and syncope.

The management of sick sinus syndrome includes identifying and correcting reversible factors, as an initial step.

Some examples of reversible causes of Sinus Node Dysfunction include,

• Hypothyroidism

Sold by @itachibot

• Sleep apnea

• Anti-hypertensive medications eg. Beta-blockers

• Anti-arrhythmic medications eg. Amiodarone, Procainamide

Ambulatory ECG monitoring is required to diagnose as the symptoms are nonspecific.

Treatment involves permanent pacing.

Other options:

Option A: The course of sinus node dysfunction is usually a slow progression over decades. In the early stages, most patients are asymptomatic.

Option B: It occurs most commonly in elderly patients due to the gradual loss of SA node function.

Option D: Beta-blockers are contraindicated in this condition.

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Ventricular Arrhythmias and Heart Blocks

Question 1:

A hypertensive patient was brought to the emergency due to complaints of palpitations and breathing difficulty. Comment on his ECG.



a) Multifocal PVC

b) Unifocal PVC

c) Ventricular quadrigeminy

d) Ventricular bigeminy

Question 2:

Identify the marked ECG feature in a patient diagnosed with ventricular tachycardia.

Sold by @itachibot

a) Fusion beat

b) Capture beat

c) Polymorphic QRS

d) Triphasic QRS

Sold by @itachi... purchased this from some... else, may have been scammed.

Question 3:

What is the diagnosis of this patient from the ECG shown below?

a) Normal ECG

b) Ventricular fibrillation

Sold by @itachibot

c) Ventricular tachycardia

d) Misplaced leads

Question 4:

A 50-year-old man is brought to the emergency following an episode of syncope during his morning walk. His blood pressure is 80/60 mm Hg. He has a history of myocardial infarction 5 yrs back. His present ECG is given below. Which is the most appropriate treatment for the patient?



a) Lignocaine

b) Diazoxine

c) DC cardioversion

d) Lidocaine

Question 5:

All of the following are effective in the management of patients with ventricular tachycardia except:

a) Procainamide

b) Amiodarone

c) DC cardioversion

d) Carotid sinus massage

Sold by @itachibot

Question 6:

Accelerated idioventricular rhythm is the most common arrhythmia associated with:

a) Myocardial reperfusion

b) Pericarditis

c) Cocaine intoxication

d) Hypertrophic cardiomyopathy

Question 7:

Automatic implantable cardioverter-defibrillator (AICD) implantation is done for all of the following conditions except:

a) Brugada syndrome

b) Ventricular fibrillation

c) Acute coronary syndrome with low ejection fraction

d) Acute myocardial infarction

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

Sudden cardiac death is defined as unexpected nontraumatic death in stable patients within:

a) 1 hour after onset of symptoms

b) 3 hours after onset of symptoms

c) 120 mins after onset of symptoms

d) 90 mins after onset of symptoms

Question 9:

A 60-year-old man with a history of ischemic heart disease is brought to the emergency in an unconscious state. He has been neglecting symptoms of dizziness and palpitations for the past few days. His ECG is given below. The most possible diagnosis is:

Sold by @itachibot



a) Monomorphic VT

b) Polymorphic VT

c) Ventricular fibrillation

d) Torsades de pointes

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 10:

A patient who is a known case of Crohn's disease was brought to the emergency. He complained of sudden onset cold sweats and palpitations. Which among the following is the likely cause of his ECG finding shown in the given image?



a) Hypomagnesemia

Sold by @itachibot

b) Hypophosphatemia

c) Hyperkalemia

d) Hypercalcemia

Question 11:

The ECG of a pregnant lady having pre-eclampsia is shown below. Her vitals are stable. What is the best step in the management of the condition?



a) IV calcium

b) IV $MgSO_4$

c) DC Shock

d) Synchronised cardioversion

Question 12:

Identify the false statement about the bundle of Kent.

a) It is faster than AV nodal pathway

b) It is slower than AV nodal pathway

c) Leads to short PR interval

d) Leads to prolonged QRS duration

Sold by @itachibot

Question 13:

Which of the following is seen in patients with Brugada syndrome?

a) ST segment elevation in V1-V3

b) ST segment elevation in V4-V6

c) ST segment elevation in 1, V5, AvL

d) ST segment elevation in II , III AvF

Question 14:

Brugada syndrome is associated with:

a) Left bundle branch block

b) Left anterior fascicular block

c) Left posterior fascicular block

d) Right bundle branch block

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Question 15:

Romano-Ward syndrome is:

a) Congenital short PR associated with deafness

b) Congenital long QT without deafness

c) Congenital long QT associated with cataract

d) Congenital short PR without cataract

Question 16:

The ECG of a patient as given below is suggestive of:

Sold by @itachibot

a) 1st degree AV Block

b) Mobitz type I

c) Mobitz type II

d) Complete AV Block

Question 17:

Which of the following conditions is associated with AV block?

a) Hypothyroidism

b) Cushing's syndrome

c) Hyperthyroidism

d) Pheochromocytoma

Question 18:

A concerned wife brought her 43-year-old husband to the clinic after an episode of light-headedness and palpitations. An ECG was taken. What is the diagnosis?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) Mobitz type II block

b) Mobitz type I block

c) Complete Heart block

d) 1st Degree AV block

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 19:

All of the following may be seen in Mobitz type II heart block except:

a) Block in bundle of His

b) Atrial rate more than the ventricular rate

c) Constant PR interval

d) P-QRS dissociation

Question 20:

A 56-year-old woman was brought to the emergency room after she lost consciousness briefly during a speech. An ECG was taken immediately. What is the most likely diagnosis?

Sold by @itachibot

a) Mobitz type II block

b) Mobitz type I block

c) Complete Heart block

d) 1st Degree AV block

Question 21:

Interpret the given ECG:

a) Wenckebach block

b) Mobitz block

Sold by @itachi... purchased this from some... else, may have been scammed.

Sold by @itachibot

c) Complete heart block

d) 1st Degree heart block

Question 22:

You observe that an ECG recording revealed a QRS complex > 0.12 seconds. Which of the following would you not include in the differential diagnosis?

a) Ventricular tachycardia

b) Wolff-Parkinson-White syndrome

c) Left anterior fascicular block

d) Right bundle branch block

Question 23:

A 40-year-old male patient came to ED with complaints of weakness, paresthesia, and breathing difficulty. Relevant investigations were done. The ECG obtained is suggestive of :



a) Hypokalemia

b) Hyperkalemia

c) Hypocalcemia

d) Hypercalcemia

Sold by @itachibot
If you bought this from someone else, it is scammed.

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	a
3	c
4	c
5	d
6	a
7	d
8	a
9	c
10	a
11	b
12	b
13	a
14	d
15	b
16	a
17	a
18	b
19	d
20	a
21	c
22	c
23	a

Detailed Explanations

Solution to Question 1:

The above ECG shows a pattern of premature ventricular contractions, one premature beat after one normal beat. This is seen in ventricular bigeminy.

Premature ventricular contractions/beats (PVC) are single ventricular beats that occur earlier than the next normal supraventricular beat. They originate from abnormal foci in the ventricular tissue:

• Unifocal PVCs arise from a single focus and hence have the same QRS morphology

Sold by @itachibot

• Multifocal PVCs arise from multiple foci and therefore have different QRS morphologies

They are not preceded by a P wave as they do not originate in the atria. They show wide QRS complexes (>0.12 ms) on the ECG because they are conducted via normal myocardial tissue instead of the Purkinje system.

Ventricular bigeminy refers to a pair of normal and premature ventricular contractions, whereas ventricular trigeminy refers to a premature beat that occurs after two normal beats. Quadrigeminy is when a premature beat follows three normal beats.

Few other terms associated with premature ventricular contractions:

• Two successive PVCs are known as a couplet

• Three successive PVCs are known as a triplet.

The given ECG shows ventricular bigeminy with premature beats circled in red and normal ventricular contractions circled in green.



Solution to Question 2:

The ECG feature of ventricular tachycardia seen in the given image is a fusion beat. It occurs when a sinus and ventricular beat coincide to produce a hybrid complex.

Ventricular tachycardia (VT) is defined as ≥ 3 consecutive premature ventricular contractions at a rate of >100/min. It is characterized by:

• Wide QRS complex

• AV dissociation- Atrial activity independent of ventricular activity, seen on the ECG as every p wave not followed by QRS complex

• Fusion beats- occur when a sinus and ventricular beat coincide to produce a hybrid complex

• Capture beats occur when the SA node transiently 'captures' the ventricles, that is, ventricles are depolarized by the impulses from the SA node producing a QRS complex of normal duration.

Sold by @itachibot

The given ECG shows a fusion beat (red arrow), and a capture beat (green arrow) can also be seen:

The given image shows capture beats in VT:



Option C: Polymorphic QRS, that is, continuously changing QRS morphology is seen in polymorphic ventricular tachycardia.

Option D: Triphasic QRS (eg. RSR′ pattern on V1) is seen in supraventricular tachycardias with the wide QRS complex.

Solution to Question 3:

The rate of the given ECG is around 188 BPM (>100 BPM) and it shows wide QRS complexes (≥ 0.12 s) with abnormal morphology (red arrows) and absence of P waves which suggests a diagnosis of ventricular tachycardia.

Sold by @itachibot



Ventricular fibrillation (Option B) would not show discreet QRS complexes, and hence is not the answer in this case.



Solution to Question 4:

The clinical scenario describes a man with history of coronary artery disease developing syncope. The ECG shows wide QRS complex (> 0.12 s) tachycardia with a single QRS morphology suggestive of monomorphic ventricular tachycardia (VT). The most appropriate treatment of VT in a hemodynamically unstable (systolic blood pressure < 90 mm Hg) patient is DC cardioversion.

VT can be casued by structural heart disease, scar-related reentry, cardiomyopathies, or abnormalities in the conduction pathways. VT is said to be sustained if it lasts for at least 30 seconds or requires intervention for termination.

Sold by @itachibot

Rapid VT can lead to hypotension and syncope, especially in patients with ventricular dysfunction. Patients with normal cardiac function may tolerate sustained VT, presenting with palpitations. Monomorphic VT associated with structural heart disease can progress to ventricular fibrillation (VF) and sudden cardiac arrest.

QRS synchronous electrical cardioversion is performed for unstable patients. Intravenous amiodarone is the preferred drug for stable VT associated with heart disease.

After restoring sinus rhythm, further evaluation is necessary to determine the underlying heart disease. Treatment options include antiarrhythmic medications, catheter ablation, or implantable cardioverter-defibrillators (ICDs) for recurrent or high-risk VT. ICDs are commonly used for secondary prevention of sudden death in patients with sustained VT and structural heart disease.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

Carotid sinus massage will be effective only in supraventricular tachycardias but not in ventricular tachycardias. This is because it acts by increasing the AV node block which does not play any role in the pathogenesis of ventricular tachycardia.

Hence, AV node blocking maneuvers (carotid sinus massage, valsalva) and drugs (adenosine) are not used in the management of VT.

Solution to Question 6:

Accelerated idioventricular rhythm is the most common arrhythmia associated with myocardial reperfusion, most commonly in the first 2 days after presentation.

Idioventricular rhythm is defined as ≥ 3 consecutive premature ventricular contractions with a heart rate of <50 bpm. If the heart rate is 60-100 bpm, it is known as accelerated idioventricular rhythm, and if it is more than 100 bpm it is known as ventricular tachycardia. It may be the result of an abnormal focus in the ventricle that generates an impulse faster than the sinus node and therefore assumes control.

Other causes of AIVR include:

- Buerger disease
- Congenital heart disease
- Dilated cardiomyopathy
- Myocarditis
- Drugs: Digoxin toxicity, cocaine toxicity, and various anesthesia agents
- Electrolyte abnormality
- Postresuscitation

Episodes of AIVR have not been found to affect prognosis in acute MI and usually do not require intervention. If AIVR is accompanied by hemodynamic compromise, treatment with atropine or atrial pacing will often suppress the rhythm.

Sold by @itachibot

Solution to Question 7:

An automated implantable cardioverter-defibrillator (AICD) is a device placed in the chest or abdomen to detect arrhythmias and perform cardioversion, defibrillation, and pace the heart.

It is contraindicated in patients experiencing tachyarrhythmias with reversible or transient causes such as acute MI.

It is used in:

• Brugada syndrome- refers to a sodium channelopathy with a high risk of arrhythmias that can lead to sudden cardiac death.

• Ventricular fibrillation and ventricular tachycardia.

• Cardiac resynchronization therapy for prevention of sudden cardiac death in heart failure.

• High-risk situations of sudden cardiac death in hypertrophic cardiomyopathy.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

Sudden cardiac death is defined as unexpected non-traumatic death in clinically well or stable patients who die within 1 hour after onset of symptoms.

The causative rhythm in most cases is ventricular fibrillation, which is usually preceded by ventricular tachycardia except in the setting of acute ischemia or infarction.

Solution to Question 9:

The given ECG is suggestive of ventricular fibrillation.

The features in the above ECG that support the diagnosis are:

• Disordered electrical ventricular activation with no identifiable QRS complexes

• No identifiable P-waves, no PR interval

Possible mechanisms of ventricular fibrillation include spiral wave reentry and multiple circulating reentry wavefronts.

The ECG features suggestive of ventricular tachycardia are as follows:

• Wide QRS complex with abnormal morphology (marked with red arrows in image below)

• Absence of the P waves

• Rate >100/min- in the ECG below, it is ~180 beats/min (1500/no. of small boxes between two R waves)

Sold by @itachibot

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Solution to Question 10:

The given ECG finding shows torsades de pointes, which can occur due to hypomagnesemia (Crohn's disease causes chronic diarrhea which can cause excess GI loss of magnesium).

Torsades de pointes means twisting of points, it implies twisting of the QRS complexes around the isoelectric line of the ECG. It is a form of polymorphic ventricular tachycardia that is seen in patients with a prolonged QT interval.

A prolonged QT interval usually results from prolonged ventricular repolarization (lengthening of T wave). A premature ventricular contraction falling in the elongated T wave can initiate torsades de pointes.

The features of torsades de pointes are:

• Polymorphic QRS complexes that change in amplitude

• Cycle length giving an appearance of oscillations around the baseline

• QT prolongation

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 11:

The given ECG showing a wide QRS complex tachycardia with waxing and waning QRS amplitude points to the diagnosis of torsades des pointes. As the patient is hemodynamically stable, intravenous $MgSO_4$ is the treatment of choice.

Polymorphic ventricular tachycardia (VT) has a continually changing QRS morphology. Polymorphic VT that occurs in individuals with congenital or acquired prolongation of QT interval is referred to as torsades des pointes. Patients present with near-syncope, syncope, or cardiac arrest.

IV $MgSO_4$ is highly effective for both treatment and the prevention of recurrence of long QT-related ventricular ectopic beats that trigger torsades des pointes.

If magnesium alone is ineffective, isoproterenol infusion or pacing, to a rate of 100-120 depolarization/minute usually suppresses VT recurrences.

If the patient is hemodynamically unstable, then immediate defibrillation should be done.

Synchronized cardioversion	Defibrillation
Used inunstable PSVT, atrial flutter, atrial fibrillation, and ventricular tachycardiawith a pulse	Used in ventricular tachycard iawithout a pulse, and ventric ular fibrillation.
Shock is delivered in sync wit h the R wave (in absolute refr actory period)	Shock is delivered at any tim e during the cardiac cycle.
Monophasic - 100 to 200J	Monophasic - 360J
Biphasic - 50 to 100J	Biphasic - 120 to 200J

Sold by @itachibot

Solution to Question 12:

The bundle of Kent is the accessory pathway of conduction between the atria and ventricle in Wolff-Parkinson-White syndrome. It conducts impulses faster than the AV nodal pathway and avoids AV nodal delay, leading to shortened PR interval. QRS is prolonged because of the AV nodal impulse reaching the ventricles after the aberrant impulse.

Solution to Question 13:

The major feature of Brugada syndrome is transient or concealed ST-segment elevation in V1 to V3.

Brugada syndrome is a rare syndrome characterized by ST-segment elevation and negative T waves in more than 1 anterior precordial leads (V1- V3). It is associated with mutations involving cardiac sodium channels. The patients have episodes of polymorphic VT which may lead to syncope or cardiac arrest.

It shows an autosomal dominant inheritance pattern and affects males more than females. The episodes of VT may occur either in sleep or maybe provoked by febrile illness or sodium channel-blocking drugs (Class I antiarrhythmics) like ajmaline, flecainide, and procainamide.

The episodes of VT can be suppressed using quinidine or catheter ablation. ICD can be used in individuals who have been resuscitated from cardiac arrest.

Solution to Question 14:

Brugada syndrome is associated with right bundle branch block. It is an inherited disease that predisposes patients to potentially life threatening arrythmias.

The ECG below is depictive of Brugada syndrome with concave ST segments and negative T waves (to be seen in more than one anterior precordial leads, i.e., V1-V3)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Solution to Question 15:

Romano-Ward syndrome is a congenital long QT syndrome without deafness.

Romano-Ward syndrome is associated with specific genetic mutations affecting membrane potassium and sodium channels. The inheritance pattern may be autosomal recessive or autosomal dominant.

The preferred drugs for managing this condition are the beta-blockers. Implantable cardioverter-defibrillator (ICD) implantation can also be done.

Jervell-Lange-Nielsen syndrome is congenital long QT with deafness.

LOF= Loss of Function

GOF= Gain of Function

DISORDER	Gene and its Function								
Long QT syndrome	KCNQ1K+ channel (LOF)KCNH2K+ channel (LOF)SCN5ANa+ channel (GOF)CAV3Caveolin, Na+ current (GOF)	KCNQ1	K+ channel (LOF)	KCNH2	K+ channel (LOF)	SCN5A	Na+ channel (GOF)	CAV3	Caveolin, Na+ current (GOF)
KCNQ1	K+ channel (LOF)								
KCNH2	K+ channel (LOF)								
SCN5A	Na+ channel (GOF)								
CAV3	Caveolin, Na+ current (GOF)								
Short QT syndrome	KCNQ1K+ channel (GOF)KCNH2K+ channel (GOF)	KCNQ1	K+ channel (GOF)	KCNH2	K+ channel (GOF)				

Sold by @itachibot

DISORDER	Gene and its Function								
KCNQ1	K+ channel (GOF)								
KCNH2	K+ channel (GOF)								
Brugadas yndrome	SCN5ANa+ channel (LOF)CACNB2bCa++ channel (LOF)SCN1bNa+ channel (LOF)	SCN5A	Na+ channel (LOF)	CACNB2b	Ca++ channel (LOF)	SCN1b	Na+ channel (LOF)		
SCN5A	Na+ channel (LOF)								
CACNB2b	Ca++ channel (LOF)								
SCN1b	Na+ channel (LOF)								
CPVTsyndrome	RYR2Diastolic Ca++ release (GOF)CASQ2Diastolic Ca++ release (LOF)	RYR2	Diastolic Ca++ release (GOF)	CASQ2	Diastolic Ca++ release (LOF)				
RYR2	Diastolic Ca++ release (GOF)								
CASQ2	Diastolic Ca++ release (LOF)								

Sold by
If you purchased this fro
u may have been scam

Solution to Question 16:

The above ECG rhythm strip shows 1st degree AV Block.

The following features of this ECG support the diagnosis:

• The P wave precedes each and every QRS complex

• PR interval is prolonged

• The PR interval is constant in every beat

Sold by @itachibot

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 17:

Hypothyroidism is associated with AV block. In hypothyroidism, there is reduced myocardial contractility and reduced pulse rate leading to a reduced stroke volume and bradycardia, AV block.

Causes of AV block:

Autonomic:

• Carotid sinus hypersensitivity

Metabolic:

• Hyperkalemia

• Hypermagnesemia

• Hypothyroidism

• Adrenal insufficiency

Drug related:

• Beta blockers

• Calcium channel blockers

• Adenosine

• Digitalis

• Lithium

• Class 1 and class 3 anti arrhythmics

Infections:

• Endocarditis

Sold by @itachibot

• Lyme disease

• Chaga's disease

• Syphilis

• Tuberculosis

• Diphtheria

• Toxoplasmosis

Inflammatory:

• SLE

• RA

• Scleroderma

Infiltrative:

• Amyloidosis

• Sarcoidosis

• Hemachromatosis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

This patient has symptoms and ECG findings suggestive of Mobitz type I or Wenckebach block.

In the given ECG PR segment progressively increases in each successive cycle, until a P wave of the series fails to produce a QRS complex (dropped QRS). In this example, for every 5 'P' waves, there are only 4 'QRS' complexes; hence, this is a 5:4 Wenckebach block.

Solution to Question 19:

The absence of a constant relationship between the P wave and the QRS complex (P-QRS dissociation) is a feature of complete heart block. It is not seen in Mobitz type II second-degree heart block.

The PR interval remains constant in Mobitz type II second-degree heart block but few p waves are not followed by QRS complex. It occurs due to a block in the bundle of His. The atrial rate is more than the ventricular rate.

Solution to Question 20:

The given ECG shows a constant PR interval in each successive cycle followed by a dropped QRS (P wave fails to produce a QRS complex). This pattern is suggestive of Mobitz type II block.

Sold by @itachibot

Mobitz type II block is characterized by an intermittent failure of conduction of the p wave without any change in the preceding impulses.

The dropped QRS is marked in the image below:



Note: Bradyarrhythmias that cause syncope include those due to severe sinus node dysfunction and AV block(Mobitz type II block, complete AV block). Syncope due to bradycardia or asystole is referred to as Stokes-Adams attack.

Sold by
If you purchased this fro
you may have been scam
else,
d.

Solution to Question 21:

The above ECG shows a complete heart block.

Features in the ECG supporting the diagnosis of complete heart block:

• PP interval is constant

• RR interval is constant

• P waves more in number than QRS complexes

• There is no relation between P waves and the QRS complexes.

For a second-degree heart block, the QRS complex should be preceded by the P wave, whereas in a third-degree heart block, the QRS complex is independent of the P wave.

The ECG features are marked in the image below:

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

QRS > 0.12s points towards a wide QRS complex. The left anterior fascicular block does not cause a wide QRS complex.

Fascicular blocks do not prolong the QRS interval. They only lead to axis deviation.

• Left anterior fascicular block: left axis deviation

• Left posterior fascicular block: right axis deviation

Wide QRS complexes (≥ 0.12 seconds) are seen in the following:

• Ventricular tachycardias

• Ventricular premature contractions

• Wolff-Parkinson-White syndrome

• Bundle branch blocks (complete blocks- QRS interval ≥ 0.12 seconds, incomplete blocks- QRS interval between 0.10–0.12 seconds)

• Hyperkalemia

Solution to Question 23:

The symptoms of weakness, paresthesia, and dyspnea along with the given ECG showing depression of the ST segment, a decrease in the amplitude of the T wave (flattening or inversion), and an increase in the amplitude of U waves (at the end of the T wave) are characteristic of hypokalemia.

In the given ECG, leads V1-V6 exhibit T wave inversion. U waves are well appreciated in leads V2-V5.

Sold by @itachibot



Hypokalemia leads to dynamic ECG changes. The image given shows the sequence of changes observed:



Hypokalemia is defined as plasma K+ concentration of <3.5meq/L. It can also cause premature atrial and ventricular beats, sinus bradycardia, paroxysmal atrial or junctional tachycardia, atrioventricular block, and ventricular tachycardia or fibrillation.

Other options:

Option B: Hyperkalemia causes tall peaked T waves with a shortened QT interval. There is a progressive lengthening of the PR interval and QRS duration with an increase in potassium levels, eventually leading to the disappearance of the P wave; and widening of the QRS complex to a sine wave pattern.

Sold by @itachibot



Option C: Hypocalcemia causes prolongation of the QT interval. There is a lengthening of the ST segment, while the T wave (which correlates with the time for repolarization) remains unaltered.

Option D: Hypercalcemia causes shortening of the QT interval and a decrease in the ST segment duration.



Sold by @itachibot

Valvular Heart Diseases

Question 1:

What is the most common cause of valvular heart disease in developing countries?

a) Rheumatic fever

b) Infective endocarditis

c) Congenital

d) Idiopathic

Question 2:

What is the normal range for the mitral valve area in an adult?

a) 4-6 sq mm

b) 4-6 sq cm

c) 6-8 sq mm

d) 6-8 sq cm

Question 3:

A 45-year-old woman is diagnosed with mitral stenosis. Which of the following features is least likely to be seen in this patient?

a) Pulmonary artery hypertension

b) Tricuspid regurgitation

c) Right ventricular hypertrophy

d) Left ventricular hypertrophy

Question 4:

A 45-year-old male patient comes to the clinic with 3 months history of breathlessness. After Evaluation, the patient is diagnosed with mitral stenosis. Which of the following is least likely to be heard on auscultation?

a) Diastolic murmur

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

b) Mid systolic click

c) Opening snap

d) Pansystolic murmur

Question 5:

A 48-year-old female patient was admitted to the hospital in view of severe dyspnea. History revealed that the dyspnea was exertional initially and later worsened to breathlessness during normal daily activities. She died on the 3rd day following admission. The image of the autopsy of the heart is shown below. Identify the condition.



a) Mitral stenosis

b) Mitral regurgitation

c) Tricuspid stenosis

d) Tricuspid regurgitation

Question 6:

A 62-year-old woman presents with breathlessness for 5 months. She is unable to carry out her daily activities and has difficulty sleeping. On examination, she has a parasternal heave. On auscultation, opening snap (OS) is heard followed by a low pitched rumbling diastolic murmur. Which of the following findings is suggestive of increased severity of the given condition?

a) Loud S1

Sold by @itachibot

b) Soft P2

c) Decreased A2-OS interval

d) Increased A2-OS interval

Question 7:

Which of the following cardiac chambers is enlarged in mitral stenosis on chest X-ray initially?

a) Left atrium

b) Right atrium

c) Left ventricle

d) Right ventricle

Question 8:

A patient with mitral stenosis presented with worsening dyspnea and fatigue which are now present at rest. He also complained of a feeling of heavy pounding in the chest with recurrent chest pain. You decided to evaluate the patient using echocardiography. When would you advise the patient to undergo a valvotomy?

a) Mitral orifice is < 0.8 sq cm

b) Mitral orifice is < 1 sq cm

c) Mitral orifice is < 1.5 sq cm

d) In all patients who are symptomatic

Question 9:

A 26-year-old woman undergoing evaluation for panic disorder was referred to the physician for complaints of palpitations and breathlessness. Detailed history revealed the presence of chest pain, which was substernal and not related to exercise. On auscultation, mid-systolic clicks were audible. What is the most likely diagnosis?

a) Unstable angina

b) Aortic stenosis

c) Mitral valve prolapse

d) Mitral regurgitation

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 10:

Carvallo's sign is associated with the murmur of which valvulopathy?

a) Mitral stenosis

b) Tricuspid regurgitation

c) Pulmonary regurgitation

d) Aortic stenosis

Question 11:

The severity of mitral regurgitation can be assessed by:

a) Intensity of murmur

b) Ejection click

c) Wide split of S2

d) Loud S1

Question 12:

A 65-year-old male patient presented with chest pain, exertional dyspnea and syncope. What is the most likely diagnosis?

a) Tricuspid regurgitation

b) Aortic regurgitation

c) Mitral stenosis

d) Aortic stenosis

Question 13:

On auscultation of a patient, a murmur was heard which is depicted by the following phonocardiogram. Which of the given conditions is most likely to produce this murmur?

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) Mitral stenosis

b) Mitral regurgitation

c) Aortic stenosis

d) Aortic regurgitation

Question 14:

A patient presented with breathlessness on exertion and chest pain. Auscultation revealed an ejection systolic murmur heard best at the base of the heart. Echocardiography showed reduced left ventricular ejection fraction and left ventricular hypertrophy. Which of the following features does not signify a severe form of this disease?

a) Pulsus parvus et tardus

b) Single S2

c) Late peaking of murmur

d) Wide splitting of S2

Question 15:

Which of the following conditions is not associated with aortic regurgitation?

a) Ankylosing spondylitis

b) Marfan syndrome

c) Acute myocardial infarction

Sold by @itachibot
If you purchased this from some
you may have been scammed.
else,

Sold by @itachibot

d) Bicuspid aortic valve

Question 16:

A 55-year-old male patient presents to the hospital with palpitations. On examination, a diastolic murmur is heard in the left third intercostal space and pistol shot sound is heard over the femoral arteries. What is the most likely diagnosis?

a) Aortic regurgitation

b) Aortic stenosis

c) Tricuspid regurgitation

d) Mitral regurgitation

Question 17:

A 58-year-old man presented with exertional dyspnea and angina. On examination, the pulse was characterized by abrupt distention and quick collapse. Auscultation revealed an early diastolic murmur. Which of the following is a feature of this murmur with respect to the likely diagnosis in this case?

a) Crescendo-decrescendo murmur

b) Best heard in inspiration

c) High pitched murmur

d) Severity correlates with intensity of murmur

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 18:

Which of the following is the criteria for severe aortic regurgitation, in transthoracic doppler echocardiography?

a) Effective regurgitant orifice <0.3 sq. cm

b) Regurgitant volume >60mL/ beat

c) Regurgitant fraction more than 60%

d) Vena contracta <0.6 cm

Question 19:

Sold by @itachibot

What is the treatment of choice in symptomatic acute severe aortic regurgitation?

a) Beta blockers

b) Aortic valve replacement

c) Periodic monitoring

d) Diuretics

Question 20:

A patient has enlarged neck veins, hepatomegaly and a holosystolic murmur along the lower left sternal margin. What is the most likely diagnosis?

a) Tricuspid regurgitation

b) Mitral regurgitation

c) Left heart failure

d) Right heart failure

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Question 21:

On auscultation of a patient, a murmur was heard which is depicted by the following phonocardiogram. Which of the given conditions is most likely to produce this murmur?



a) 2,5

b) 2,3

c) 1,4

Sold by @itachibot

d) 3,5

Question 22:

A patient has a right upper limb BP of 180/95 mm Hg and a left upper limb BP of 130/90 mm Hg. He also has an early diastolic murmur in right 2nd intercostal space. Which of the following is least likely to be associated with these findings?

a) Supravalvular aortic stenosis

b) Coarctation of aorta

c) Takayasu arteritis

d) Aortic dissection

Question 23:

Barlow's syndrome refers to which of the following?

a) Carcinoid of heart

b) Mitral valve prolapse

c) Aortic regurgitation

d) Hypertrophic cardiomyopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	b
5	a
6	c
7	a
8	c
9	c
10	b

Sold by @itachibot

11	c
12	d
13	c
14	d
15	c
16	a
17	c
18	b
19	b
20	a
21	c
22	a
23	b

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

The most common cause of valvular heart disease in developing countries is rheumatic fever. This is mainly due to the repeated infections with rheumatogenic streptococci and inadequate coverage of antibiotics. It mainly affects the mitral valve.

In developed countries, degenerative or inflammatory processes are the most common cause of valvular heart diseases.

Solution to Question 2:

The size of the mitral orifice in a normal adult is 4-6 sq cm.

Depending on the area of mitral valve opening (MVA), mitral stenosis can be classified as:

• Progressive MS - MVA > 1.5 sq cm

• Severe MS - MVA ≤ 1.5 sq cm

• Very severe MS - ≤ 1 sq cm

Solution to Question 3:

Left ventricular hypertrophy is not seen in mitral stenosis. On the contrary, right ventricle hypertrophy is seen.

Sold by @itachibot

Mechanism of mitral stenosis (MS):

In a normal mitral valve, there is no obstruction to the flow of blood from the left atrium(LA) to the left ventricle. Therefore a small LA pressure is sufficient to empty the blood into the ventricle.

When the MVA is <~2 sq cm, blood flows from the left atrium to the left ventricle, propelled by a mildly elevated atrial pressure. Hence, patients are usually asymptomatic and may develop dyspnea only when the heart rate increases, as seen in exertion, excitement, fever, etc. All these conditions:

- Increase the rate of blood flow across the mitral orifice, resulting in further elevation of the LA pressure
- Decrease the diastolic filling time, resulting in a reduction in cardiac output

The image below explains the mechanism of mitral stenosis.



Solution to Question 4:

Mid-systolic click is least likely to be heard in a patient with mitral stenosis. It is seen in a patient with mitral valve prolapse.

Important clinical features of Mitral stenosis	
Findings	Cause
Inspection	
Malar flush, pinched and blue facies	Decreased cardiac output and increased pulmonary artery hypertension leads to retention of CO_2 and vasodilation
Precordial bulge	

Sold by @itachibot

Important clinical features of Mitral stenosis	
Palpation	
Parasternal lift	Enlarged right ventricle
Palpable P2	Pulmonary hypertension
Diastolic thrill	Increased intensity of diastolic murmur
Auscultation	
Loud S1	It is seen ininitial stageswhen the mitral valve is flexible.
Opening snap (OS)	It is caused due to the sudden opening of thickened leaflets.It follows A2. As the disease progresses, thedurationbetween A2 and OSdecreases.It is best audible at the apex and with the diaphragm of the stethoscope
Diastolic murmur	Lowpitched rumbling murmur heard with thebellof the stethoscope.It follows OS and best heard atapexwith left lateral recumbent position. When it is loud, may radiate to left axilla.The duration of murmur correlates with the severity of stenosis
Loud P2	Pulmonary artery hypertension
Other Auscultatory Findings	
Pansystolic murmur	Due to functional tricuspid regurgitation in severe pulmonary hypertension
Graham Steell murmur	It is adiastolicmurmur due to pulmonary regurgitationCaused due to the dilation of the pulmonary valve ring in severe pulmonary hypertension
JVP	
Prominent a waves	Atrial systole
ECG	
Tall and peaked P waves	Right atrial enlargement
Right axis deviation	Right ventricular hypertrophy

Sold by @itachibot

Solution to Question 5:

The given image is of mitral stenosis showing fish mouth or buttonhole appearance.

Solution to Question 6:

The above clinical scenario is suggestive of mitral stenosis. The more severe the thickening, the earlier the opening snap occurs in diastole. Hence, the decreased interval between A2 and opening snap (OS) indicates increased severity of the stenosis.

The auscultatory findings which are suggestive of increased severity of murmur are:

• Decreased interval between A2 and OS

• Increased duration of the diastolic murmur

Loud S1 is seen in the initial stages of MS when the mitral leaflets are flexible. When there is further thickening and calcification of the leaflets, they become immobile, and the amplitude of S1 decreases.

Increased severity of the stenosis increases the left atrial pressure. This increase in left atrial pressure can increase pulmonary artery pressure which leads to loud P2 on auscultation.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

Left atrium (LA) is the first cardiac chamber to get enlarged in mitral stenosis (MS). It is due to the pressure overload of LA. It can be appreciated on the lateral and left anterior oblique views.

Other radiographic changes seen in MS are straightening of the upper left border of the cardiac silhouette, posterior displacement of the esophagus by an enlarged LA, and Kerley B lines due to distention of interlobular septae and lymphatics with edema.

Severe MS causing pulmonary hypertension leads to enlargement of the pulmonary artery, right ventricle, and right atrium. Left ventricle remains unaffected in isolated MS.

Solution to Question 8:

Mitral valvotomy is indicated in symptomatic (NYHA Functional Class II-IV) patients with mitral stenosis (MS) with orifice < 1.5 cm2 in normal-sized adults or <1 cm2/m2 body surface area.

Treatment of mitral stenosis:

• Administration of warfarin in patients with MS who develop atrial fibrillation.

• Surgical treatment includes:

• Mitral valvotomy: It is done in symptomatic patients whose mitral orifice is less than 1.5 cm2. This can be either percutaneous mitral valvotomy or open surgical valvotomy.

• Mitral valve replacement: This is done in patients whose mitral valve is severely distorted.

Sold by @itachibot

Solution to Question 9:

The above clinical scenario is suggestive of mitral valve prolapse.

It is caused due to the myxomatous degeneration of the mitral valve. It may be associated with Marfan's syndrome, osteogenesis imperfecta, and Ehlers-Danlos syndrome. It is more common in females of the age group of 15- 30 years.

Most patients are usually asymptomatic. However, some patients may present with symptoms such as palpitations, syncope due to arrhythmias (premature ventricular contractions, ventricular tachycardia, and atrial fibrillation), non-specific chest pain, and dyspnea

Auscultation reveals mid-systolic clicks. If there is associated mitral regurgitation, a systolic murmur might also be heard. The duration of the murmur of MVP decreases on squatting and increases on standing. Whereas the duration of the murmur of MR increases on squatting and decreases on standing.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Carvallo's sign refers to the increase in the intensity of systolic murmur of tricuspid regurgitation with inspiration.

During inspiration, the venous return to the heart increases due to the negative intrathoracic pressure. This increases the leak from the right ventricle to the right atrium making the murmur louder.

In general, the right-sided heart murmurs increase in intensity with inspiration and decrease in intensity with expiration, while the opposite is true for left-sided heart murmurs. Pulmonic ejection sound is the only right-sided acoustic event that decreases in intensity with inspiration.

Solution to Question 11:

The severity of mitral regurgitation (MR) can be assessed by the wide splitting of the S2. Due to severe regurgitation of blood, the pressure in the left ventricle falls rapidly, leading to the early closure of the aortic valve.

Clinical findings that indicate the severity of MR is:

• Reduced cardiac output

• Palpable systolic thrill

• Palpable S3

• Wide splitting of S2

Note: The intensity of murmur does not always correlate with the severity of the murmur. Intensity may be diminished in severe MR caused by LV dysfunction, acute MI and periprosthetic

Sold by @itachibot

valve regugitation.

Solution to Question 12:

The most likely diagnosis is aortic stenosis.

Aortic stenosis is the narrowing of the aortic valve. It may be caused due to the calcification of the normal aortic valve, calcification of congenital bicuspid valve, and rheumatic heart disease.

Three cardinal clinical features of aortic stenosis are:

• Exertional dyspnea - this occurs due to the left ventricular (LV) diastolic dysfunction. Raised end-diastolic LV volume and reduced LV compliance lead to increased LV pressure, leading to elevation of pulmonary capillary wedge pressure.

• Angina pectoris - it occurs due to the combination of the increased O_2 needs of hypertrophied myocardium and reduction of O_2 delivery.

• Syncope - it occurs due to the combination of decreased cardiac output and systemic arterial dilation which lead to a reduction of cerebral perfusion.

Solution to Question 13:

The phonocardiogram given above is of an ejection systolic murmur, which is seen in aortic stenosis. It is also heard in pulmonary stenosis. They are differentiated based on the difference in symptoms, site of murmur, etc.

Clinical findings in aortic stenosis:

• Pulse - slow rising and late peaking pulse which is known as pulsus parvus et tardus. A carotid thrill may also be palpated over these arteries.

• Apical impulse - A double apical impulse may be recognized, and in the later stages, it may be displaced laterally.

Auscultation findings in aortic stenosis:

• Early systolic ejection sound - heard in early stages, disappears when the valve becomes calcified and rigid

• Paradoxical splitting of S_2 - With worsening of stenosis, delayed closure of the aortic valve occurs. This leads to narrowing of S_2 sound, and in severe cases, leads to paradoxical splitting of S_2.

• S_4 and S_3 - heard in severe cases

• Ejection systolic murmur - starts after S_1, increases in intensity to reach the peak, and ends before A_2. It radiates to the neck and base of the heart. It is augmented by squatting and reduced by the strain of the Valsalva maneuver.

The image below shows phonocardiograms of normal and abnormal heart sounds.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



MARROW
Sold by @Itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 14:

The above features point to a diagnosis of aortic stenosis. Wide splitting of S2 is not a feature of severe aortic stenosis. On the contrary, there might be single S2 or paradoxical splitting of S2.

Features of severe aortic stenosis are:

- Pulsus parvus et tardus
- Palpable carotid thrill
- Louder and late peaking of systolic murmur
- Single S2 or paradoxical splitting of S2
- Presence of S3 and S4 heart sounds
- Signs of right and left ventricular failure
- Left ventricular hypertrophy

Solution to Question 15:

Acute myocardial infarction is not associated with aortic regurgitation.

Aortic regurgitation is the backflow of the blood from the aorta to the left ventricle.

It may be caused either due to the defect in the valve or defect in the aortic root (dilation of the aortic annulus).

Conditions that affect valves are:

- Congenital
- Endocarditis

Sold by @itachibot

• Rheumatic fever

• Myxomatous (prolapse)

• Traumatic

• Syphilis

• Ankylosing spondylitis

Conditions that affect aortic root are:

• Aortic dissection

• Cystic medial degeneration

• Marfan syndrome

• Bicuspid aortic valve

• Hypertension

Sold by @itachibot
If you got this from someone else, it has been scammed.

Solution to Question 16:

The presence of a diastolic murmur along with the pistol shot sound over the femoral is most likely to be due to aortic regurgitation.

The image below explains the mechanism of aortic regurgitation.



Solution to Question 17:

The given clinical features are suggestive of aortic regurgitation. An early diastolic murmur is a high-pitched decrescendo diastolic murmur.

Auscultatory findings in AR are:

Sold by @itachibot

• Soft S1 - due to the early closure of the mitral valve

• Diminished or absent A2

• S3 - due to increased left ventricular (LV) end-diastolic volume. The presence of S3 may be a sign of impaired LV function.

• Early diastolic murmur -

• High-pitched, blowing, decrescendo, early diastolic murmur

• Best heard in the left third intercostal space

• Heard with the patient sitting up and leaning forward, with breath held in forced expiration

• Duration of the murmur best correlates with the severity

• Systolic ejection murmur - It is heard due to increased LV stroke volume and ejection rate.

• Austin Flint murmur - a low-pitched, mid to late diastolic murmur. It is caused due to the large volume of regurgitant blood, hitting the anterior leaflet of the mitral valve or LV wall.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

Regurgitant volume >60mL/ beat is the criteria for severe aortic regurgitation, in transthoracic doppler echocardiography (TTE).

Doppler echocardiography can be used to assess the severity of aortic regurgitation. According to this, AR is considered to be severe when:

• Central jet width is more than 65% of left ventricular outflow tract

• Vena contracta width >6mm

• Regurgitant fraction >50%

• Regurgitant volume >60mL/beat

• Effective regurgitant orifice >0.3cm2

Solution to Question 19:

The treatment of choice in symptomatic acute severe AR is aortic valve replacement (AVR).

Treatment of aortic regurgitation:

• In non-severe AR, periodic monitoring can be done. The symptoms of dyspnea and decreased cardiac output can be managed by diuretics and vasodilators.

• In severe AR, surgery (AVR) is indicated in the following conditions:

• The patient is symptomatic.

• The patient is asymptomatic, but the LV ejection fraction is less than 50% or LV end-diastolic diameter is more than 65mm.

• The patient is asymptomatic and undergoing other cardiac surgeries.

Sold by @itachibot

Solution to Question 20:

The most likely diagnosis in the above clinical scenario is tricuspid regurgitation.

Clinical features of tricuspid regurgitation are:

• Distended neck veins with prominent c-v waves

• Hepatomegaly with systolic pulsations

• Positive hepatojugular reflex - the distension of the jugular vein when firm pressure is applied over the liver

• Blowing holosystolic murmur along the lower left sternal margin

Solution to Question 21:

The given phonocardiogram is of a pansystolic/holosystolic murmur. It is heard in mitral regurgitation (MR), tricuspid regurgitation (TR), and ventricular septal defect (VSD).



Solution to Question 22:

A significant difference in blood pressure measured between two arms (>10 mm Hg) may be seen in Takayasu arteritis (subclavian artery disease), supravalvular aortic stenosis, aortic coarctation, and aortic dissection (all four options).

An early diastolic murmur in the right second intercostal space (aortic area) is suggestive of aortic regurgitation. Supravalvular aortic stenosis (Option A), which involves the part of the aorta distal to the aortic valve, is unlikely to cause aortic regurgitation and hence the best answer.

If yo... @itachibot ... from someone else, ... scammed.

Sold by @itachibot

Features	Supravalvular AS	Takayasu arteritis	Aortic dissection	Coarctation of aorta
Aortic regurgitation	Unlikely	Present in 25% of cases	Present in acute situations	Present
BP difference between arms	Present	Present	Present	May be present

So the single best answer would be Supravalvular AS > Coarctation of the aorta.

Solution to Question 23:

Barlow's syndrome refers to mitral valve prolapse.

Mitral valve prolapse syndrome is also called as systolic click-murmur syndrome, Barlow's syndrome or floppy valve syndrome. It is the most common abnormality leading to primary mitral regurgitation.

Note: Infantile scurvy is referred to as Barlow's disease.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Infective Endocarditis and Acute rheumatic fever

Question 1:

In patients with infective endocarditis, the vegetations are most commonly present on:

a) Aortic valve

b) Pulmonary valve

c) Mitral valve

d) Tricuspid valve

Question 2:

Which of the following is not a cardiac manifestation of infective endocarditis?

a) Impaired conduction system

b) Heart failure

c) Perivalvular abscess

d) Roth's spots

Question 3:

Which of the following is not included in the minor criteria of modified Duke criteria for clinical diagnosis of infective endocarditis?

a) History of injection drug use

b) Immunological evidence of glomerulonephritis

c) Positive blood culture but not meeting the major criterion

d) Positive echocardiogram not meeting major criteria

Question 4:

According to the modified Duke criteria, a single positive blood culture is included in the major criteria for which organism?

a) Coagulase negative staphylococcus aureus

b) Coxiella burnetii

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Viridans streptococci

d) HACEK group

Question 5:

A known IV drug abuser is referred to medicine OPD from the de-addiction wing due to prolonged fever and new-onset seizures. His fever has been present for 2 weeks and did not come down with antibiotics. An early diastolic murmur is heard on auscultation. Multiple painless skin lesions are present on the hands. What are these lesions?

a) Osler's nodes

b) Janeway lesions

c) Roth's spots

d) Pink rim rashes

Question 6:

Which of the following is true about the vegetations seen in infective endocarditis?

a) Present mainly along the lines of closure

b) Present on both side of valve leaflets

c) Extend into the chordae

d) Small and warty

Sold by @itachibot

Question 7:

You are asked to collect blood samples for culture from a patient suspected to have infective endocarditis by a typical pathogen. Choose the correct statement for the same.

a) 3 separate culture sets - two aerobic and one anaerobic

b) 3 separate culture sets - two anaerobic and one aerobic

c) 2 separate culture sets - two aerobic

d) 2 separate culture sets - one aerobic and one anaerobic

Question 8:

A patient develops prosthetic valve endocarditis 2 years after valve replacement surgery. Which of the following organism is the most likely cause?

a) Streptococci

b) Staphylococcus aureus

c) Coagulase negative staphylococci

d) HACEK organisms

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 38-year-old patient presented with malaise, fatigue, and rigors for 1 week. Multiple puncture sites were present on the forearm along with sclerosed veins. His temperature on admission was 38.5 °C. Examination revealed a pan-systolic murmur which was loudest at the left sternal edge. Three sets of blood cultures were taken. Which organism is most likely to be found in his blood cultures?

a) Streptococcus sanguis

b) Enterococcus faecium

c) Coxiella burnetii

d) Staphylococcus aureus

Question 10:

What is the most common cause of culture-negative infective endocarditis?

a) Coxiella burnetii

b) HACEK group of organisms

Sold by @itachibot

c) Fungal endocarditis

d) Prior antibiotic use

Question 11:

An 80-year-old woman with a history of moderate aortic stenosis presented with fatigue, weight loss, and night sweats for 2 weeks. Blood tests showed the following. Echocardiography revealed an intracardiac mass on the mitral valve. Blood cultures showed growth of Streptococcus gallolyticus. What additional investigation should be performed?

a) Hysteroscopy

b) Cystoscopy

c) Colonoscopy

d) Bronchoscopy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Transesophageal echocardiography is preferred over transthoracic echocardiography as initial investigation in all of the following patients, except:

a) Patient with suspected prosthetic valve endocarditis

b) Patient with suspected IE presenting with fever

c) Patient with suspected IE with a new onset regurgitant murmur

d) Patient with suspected IE who has previous history of IE

Question 13:

Patients with which of the following conditions have the lowest risk of developing infective endocarditis?

a) Atrial septal defect

b) Ventricular septal defect

c) Patent ductus arteriosus

d) Tetralogy of Fallot

Question 14:

Sold by @itachibot

A 50-year-old woman underwent mitral valve replacement for severe MR, 5 years ago. She is now posted for an elective colonoscopy, for evaluation of altered bowel habits. What is the recommendation regarding antibiotic prophylaxis for this patient?

a) Amoxicillin, 2g orally, taken 30 to 60 minutes before the procedure

b) Clindamycin 600mg orally, taken 30 minutes before the procedure

c) Ampicillin, 2g IM or IV 60 minutes before the procedure

d) No antibiotic prophylaxis required

Question 15:

Antibiotic prophylaxis is recommended for each of the following patients undergoing dental procedures, except:

a) A patient with palliative shunt for cyanotic CHD

b) A patient with previous CABG surgery

c) Patient with valvulopathy post cardiac transplantation

d) A patient with prosthetic heart valve

Question 16:

The blood culture of a patient with suspected infective endocarditis is positive for gram-positive cocci in chains. What is the treatment of choice?

a) Ampicillin with gentamicin for 2 weeks

b) Penicillin with gentamicin for 2 weeks

c) Vancomycin for 2 weeks

d) Ceftriaxone for 2 weeks

Question 17:

A 43-year-old patient presented with spikes of fever and difficulty in breathing. Transesophageal echo revealed vegetations in the heart. The culture was positive for Burkholderia cepacia. Which of the following is the first line of management?

a) Aminoglycosides and colistin

b) Cotrimoxazole with 3rd generation cephalosporin

c) Cefepime and tigecycline

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) Carbepenams and 3rd generation cephalosporins

Question 18:

What is the causative organism of acute rheumatic fever?

a) Group A streptococcus

b) Group B streptococcus

c) Group C streptococcus

d) Group D streptococcus

Question 19:

What is the most common clinical feature seen in patients with acute rheumatic fever?

a) carditis

b) Polyarthritis

c) Chorea

d) Erythema marginatum

Question 20:

Which of the following is a feature of Sydenham's chorea?

a) Always associated with arthritis

b) Affects mainly the head and upper limbs

c) Occurs within 3 weeks of streptococcal infection

d) Only restricted to one half of the body

Question 21:

Which of the following is a feature of subcutaneous nodules palpated in a patient with acute rheumatic fever?

a) Painful

b) Early manifestation

c) Common in hands and feet

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) Fixed to underlying structure

Question 22:

Which of the following is a minor criterion according to modified Jones criteria for diagnosis of rheumatic fever in low risk population?

a) 1,5

b) 2,4

c) 2,3

d) 3,5

Question 23:

Which among the following is not a feature of joint pain seen in patients with rheumatic fever?

a) Migratory arthritis

b) Mainly involves large joints

c) Symmetric involvement

d) Good response to salicylates

Question 24:

A 30-year-old patient presented to the clinic with skin lesions as shown in the image. On examination, body temperature is 38.5 °C, ESR and ASLO titers are elevated. What is true about the skin finding?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Pruritic

b) It is called erythema migrans

c) Disappears after hot shower

d) Generally occurs in patients with carditis

Question 25:

Evidence of preceding streptococcal infection is not essential for making a diagnosis of acute rheumatic fever in patients presenting with:

a) Polyarthritis

b) Chorea

c) Subcutaenous nodules

d) Erythema marginatum

Question 26:

A patient diagnosed with acute rheumatic fever develops a new-onset murmur. His echocardiography is likely to reveal:

a) Mitral stenosis

b) Aortic stenosis

c) Mitral regurgitation

d) Aortic regurgitation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 27:

The prevalence of rheumatic valvular heart disease peaks between:

a) 5-15 years

b) 15-25 years

c) 25-40 years

d) >40 years

Question 28:

The drug used for refractory intractable rheumatic chorea is:

a) Probenecid

b) Diazepam

c) Haloperidol

d) Sodium valproate

Question 29:

A 12-year-old boy is admitted with fever and polyarthritis. His ESR on admission is 80mm/hr. There is no evidence of carditis. A diagnosis of acute rheumatic fever is made and treatment is initiated. How long will you put the patient on secondary prophylaxis?

a) Till 21 years of age

b) Till 17 years of age

c) Till 22 years of age

d) Lifelong

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	d

Sold by @itachibot

3	d
4	b
5	b
6	c
7	d
8	a
9	d
10	d
11	c
12	b
13	a
14	d
15	b
16	b
17	b
18	a
19	a
20	b
21	c
22	d
23	c
24	d
25	b
26	c
27	c
28	d
29	a

If yo
else,
d.

Detailed Explanations

Solution to Question 1:

The mitral valve is the most common valve affected in infective endocarditis.

Infective endocarditis is the infection of the endocardial layer of the heart. It may involve the endocardial layer over the heart valves, mural endocardium, or septum. However, it can also affect intracardiac devices.

Sold by @itachibot



Note: Infection of the arteriovenous shunts and the arterio-arterial shunt is called infective endarteritis and not infective endocarditis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 2:

Roth's spots are not the cardiac manifestation of infective endocarditis (IE). They are the immunological phenomenon seen in patients with infective endocarditis.

Symptoms of infective endocarditis can be broadly divided into cardiac and non-cardiac manifestations. Non-cardiac manifestations of infective endocarditis are further divided into immunologic and vascular phenomena.

• Cardiac manifestations of IE:

• Presence of new regurgitant murmurs

• Congestive heart failure due to valve dysfunction

• Perivalvular abscesses and intracardiac fistulae- caused due to the extension of infection beyond the valves and into the adjacent myocardial tissue

• Conduction system defects- due to the damage of the interventricular septum by the abscess.

• Embolization of coronary arteries

• Non-cardiac manifestations of IE:

• Immunological phenomena (mnemonic ROGeR):

• Osler's nodes- small, tender, red papules that are present on the distal finger and toe pads.

• Roth's spots- white-centered retinal hemorrhages

• Immune-complex mediated glomerulonephritis

• Positivity for rheumatoid factor

• Vascular phenomena:

Sold by @itachibot

• Janeway lesions- these are faint, red, macular, painless lesions, commonly present on the thenar and hypothenar eminences.

• Major arterial emboli

• Septic pulmonary infarct

• Mycotic aneurysm

• Intracranial hemorrhages

• Conjunctival hemorrhages

• Splenic abscess

• Cerebral abscess

The image below shows Roth spot on the retina.

Solution to Question 3:

Positive echocardiogram not meeting major criteria was earlier a minor criterion but is now removed.

Solution to Question 4:

According to modified Duke criteria for the diagnosis of infective endocarditis, single positive blood culture for Coxiella burnetii or phase I IgG antibody titre > 1:800 is included in the major criteria.

Solution to Question 5:

Sold by @itachibot

The above scenario is suggestive of infective endocarditis. The skin lesions shown in the image are Janeway lesions. They are non-tender, erythematous, or hemorrhagic macular or nodular lesions, which are present on the palms or soles.

Another skin finding that can be found in patients with infective endocarditis is Osler's nodes. In contrast to Janeway lesions, these are small, tender, red papules which are present on the distal finger and toe pads.

Solution to Question 6:

Vegetations of infective endocarditis can extend onto chordae and are:

- Large and irregular
- Mainly present on valve cusps

Sold by @itachibot

• Prone to embolization

Solution to Question 7:

Option D. According to the Modified Duke 2023 Criteria for infective endocarditis, when a typical organism is suspected, two separate culture sets are required, one aerobic and one anaerobic.

If an atypical organism is suspected, three or more separate culture sets are used.

According to the new Duke's criteria, there is no specific timing to draw blood for blood culture.

Solution to Question 8:

Patient developing endocarditis 2 years after prosthetic valve surgery suggests late-onset endocarditis. In such cases, the most common causative organism is streptococci.

Solution to Question 9:

The above clinical scenario is suggestive of infective endocarditis. Staphylococcus aureus is most likely to be positive in cultures as it is the most common cause of infective endocarditis in IV drug abusers.

Endocarditis in injection drug users is mostly right-sided and involves the tricuspid valve.

Solution to Question 10:

The most common cause of culture-negative infective endocarditis is the prior use of antibiotics.

Some proportion of patients with endocarditis have negative blood cultures. The important causes are:

• Prior antibiotic use (most common cause)

• IE due to fastidious organisms or unusual pathogens or fungi such as:

• Streptococci (Granulicatella and Abiotrophia species)

• HACEK organisms

• Coxiella burnetii

• Bartonella species

• Brucella

• Tropheryma whipplei

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Sold by @itachibot

Solution to Question 11:

The given clinical features and laboratory investigations are suggestive of infective endocarditis caused due to Streptococcus gallolyticus. In this condition, a colonoscopy should be performed.

Streptococcus gallolyticus originates from the gastrointestinal tract and is associated with polyps and colon cancer. Hence, the presence of this organism in the vegetation of infective endocarditis should warrant a gastrointestinal investigation.

Solution to Question 12:

For a patient with low suspicion of IE, transthoracic echocardiography (TTE) will be done as an initial investigation.

Transesophageal echocardiography (TEE) is preferred as the initial investigation in all high-risk patients and/or in patients where a high clinical suspicion of infective endocarditis (IE) exists.

TEE is also preferred over TTE for evaluating:

- Vegetations < 2mm in diameter
- Prosthetic valve endocarditis (PVE)
- Detection of myocardial abscess
- Detection of valve perforation
- Detection of intracardiac fistulae
- Detection of vegetations in a patient with a cardiovascular implantable electronic device (CIED)

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 13:

Atrial septal defect (ASD) is associated with a very low incidence of IE. In ASD, the velocity of the shunt blood is low due to the decreased pressure gradient between the two atria. This low velocity of the blood does not usually damage the lining endocardium.

Unrepaired VSD is the most frequent congenital heart disease lesion associated with IE, followed by ventricular outflow tract obstructive lesions, such as with tetralogy of Fallot.

Risk factors for infective end ocarditis		
High risk	Moderate risk	Low risk
Prosthetic cardiac valves	Mitral valve prolapse with re gurgitation	Mitral valve prolapse without regurgitation
Previous bacterial endocardit is	Isolated mitral stenosis	Atrial septal defect

Sold by @itachibot

Risk factors for infective end ocarditis		
Cyanotic congenital heart dis eases	Surgically corrected left to rig ht shunt with no residual shu nt	
Mitral regurgitation		
Ventricular septal defect		
Uncorrected left to right shu nt (except ASD)		

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 14:

Antibiotic prophylaxis is not required in patients undergoing genitourinary or gastrointestinal procedures even if they have a high-risk cardiac lesion for developing infecting endocarditis (e.g., prosthetic valve).

Antibiotic prophylaxis is recommended for patients who are undergoing dental procedures, invasive respiratory tract procedures, and procedures on the infected skin.

Solution to Question 15:

Routine antibiotic prophylaxis for infective endocarditis is not recommended for patients with previous coronary artery bypass graft surgery. They do not have a long-term risk of infection.

High-risk cardiac lesions, for which endocarditis prophylaxis is advised before dental procedures are:

• Prosthetic heart valves

• Prior endocarditis

• Unrepaired cyanotic congenital heart disease, including palliative shunts or conduits

• Completely repaired congenital heart defects during the 6 months after repair

• Incompletely repaired congenital heart disease with residual defects adjacent to prosthetic material

• Valvulopathy developing after cardiac transplantation

Antibiotic prophylaxis is not routinely recommended for patients with acquired valvular heart diseases.

Standard regimen: Amoxicillin 2g per orally is given 1-hour prior to surgery. In patients with penicillin allergy, clarithromycin, and azithromycin 500mg per orally can be given.

Solution to Question 16:

Sold by @itachibot

The treatment for infective endocarditis with streptococci consists of penicillin with gentamicin for 2 weeks.

Penicillin G (2-3mU IV Q4h) or Ceftriaxone (2g single dose daily) can be given as a monotherapy for 4 weeks.

Treatment of infective endocarditis	
Streptococci	Penicillin + Gentamicin
Staphylococcus aureus infecting native valve	Vancomycin
Staphylococcus aureus infecting prosthetic valves	Vancomycin + Gentamicin + Rifampin
Enterococci	Ampicillin + Gentamicin
HACEK group	Ceftriaxone or Ampicillin + Sulbactam
Coxiella burnetti	Doxycycline + Hydroxychloroquine
Bartonella	Doxycycline + Gentamicin

Sold by... if you purchased this from... you may have been scammed.

Solution to Question 17:

The first-line agents used in the management of Burkholderia cepacia complex (BCC) infections are trimethoprim-sulfamethoxazole (cotrimoxazole), meropenem, minocycline, and ceftazidime (3rd generation cephalosporin).

Other options:

Option A: Burkholderia species are intrinsically resistant to aminoglycosides and polymyxins.

Option C: Tigecycline is less effective in the case of Burkholderia infections.

Option D: Carbapenems and 3rd generation cephalosporins have the same mechanism of action and compete for the same binding site. Hence, it is not a rational drug combination.

BCC is a group of bacteria known for causing severe respiratory distress and septicemia, particularly in patients with cystic fibrosis (CF). BCC organisms are also found in ICU patients and individuals with chronic granulomatous disease, leading to lung disease. BCC colonization in CF patients prior to lung transplantation increases the risk of death during the early transplant period.

Solution to Question 18:

Acute rheumatic fever is caused due to infection with group A streptococcus (GAS), which is Streptococcus pyogenes. It usually follows a streptococcal sore throat or scarlet fever. Other groups of streptococci do not cause acute rheumatic fever.

Sold by @itachibot

The pathogenesis of the development of ARF is based on the concept of molecular mimicry. The antibodies which are produced against M protein and N-acetyl glucosamine of group A streptococci, bind to and damage the human endothelial cells.

It mainly affects children between 5-14 years of age. Patients have various rheumatologic, cardiac, and neurological manifestations. Modified Jone's criteria is used for the diagnosis of acute rheumatic fever.

The main aim of treatment of rheumatic fever is to relieve symptoms, prevent inflammation and prevent further infections by group A streptoccus by the use of penicillins.

Solution to Question 19:

The most common clinical feature of acute rheumatic fever is carditis followed by polyarthritis.

Solution to Question 20:

Choreiform movements in Sydenham's chorea is mainly seen in the head & upper limbs.

Sydenham chorea is characterized by the presence of involuntary, purposeless, jerky movements of the hands, arms, shoulders, feet, legs, face, and trunk. The movements may be generalized or restricted to one side of the body.

It follows a long latent period (more than 6 months) and therefore can occur alone without any other clinical manifestations.



Solution to Question 21:

Subcutaneous nodules are commonly present in hands, feet, elbows, and occiput.

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Sold by @itachibot

They are firm, painless, and freely movable subcutaneous nodules, present over the bony prominences of hands, feet, elbow, and occiput. They are a delayed manifestation of rheumatic fever and are usually associated with carditis.

Solution to Question 22:

ESR ≥60 mm and polyarthralgia are a part of minor criteria for the low-risk population among the given options.

The major and minor criteria for arthritis differ in low risk and medium to high-risk populations.

• For low-risk populations, polyarthritis is a major criterion and polyarthralgia is a minor criterion.

• In the medium to high-risk populations, polyarthritis, polyarthralgia, monoarthritis are major criteria and monoarthralgia is a minor criterion.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 23:

In rheumatic fever, joint pain is usually asymmetric.

Characteristic features of arthritis of acute rheumatic fever are:

• Migratory- the pain moves from one joint to another over a period of hours

• Involvement of large joints – most commonly knees, ankle, hips & elbows

• Asymmetric

• Highly responsive to salicylates

Solution to Question 24:

The above clinical features are suggestive of rheumatic fever and the skin finding shown in the image is erythema marginatum. It generally occurs in patients with carditis.

Erythema marginatum is a pink, non-pruritic rash that generally occurs on the upper part of the arms or trunk but not on the face. It extends from the centre with an irregular serpiginous border. As the size of the rash increases, the center of the rash returns to normal, while the borders remain pink. Hence it is named erythema marginatum.

The rash may also become more prominent after a hot shower.

Solution to Question 25:

Evidence of preceding streptococcal infection for making diagnosis of acute rheumatic fever (ARF) is not required in the presence of chorea.

Sold by @itachibot

Chorea is frequently a late finding of ARF. When chorea appears, major findings, minor findings and findings of preceding streptococcal infection would have been disappeared.

Solution to Question 26:

Characteristic initial valvular manifestation in acute rheumatic fever (ARF) is mitral regurgitation.

Damage of the valves leads to mitral regurgitation in the early stages. With repeated episodes of rheumatic fever, there is thickening, calcification, and stenosis of the affected valve.

Mitral regurgitation is the most common cardiac lesion in ARF, while mitral stenosis is the most common lesion in rheumatic heart disease.

Other cardiac manifestations of acute rheumatic fever include myocardial inflammation that can also affect electrical conduction pathways, leading to P-R interval prolongation (first-degree atrioventricular block or rarely higher-level block) and softening of the first heart sound.

Solution to Question 27:

The prevalence of rheumatic valvular heart disease peaks between 25-40 years.

Acute rheumatic fever is mainly a disease of children and is common between 5–14 years of age and rare in a person aged more 30 years. In contrast to this, rheumatic valvular heart disease is common between 25-40 years.

Solution to Question 28:

The drug used for refractory intractable rheumatic chorea is sodium valproate.

Sydenham's chorea is characterized by involuntary, purposeless, jerky movements which mainly affect the head and upper limbs. In the tongue, it is visible as the characteristic darting movements.

Medications used to treat Sydenham chorea are:

• Mild cases: managed by providing a calm environment

• Severe cases: dopamine antagonists like haloperidol, pimozide, or antiepileptic medications like sodium valproate (preferred due to no extrapyramidal side effects) and carbamazepine

• Since Sydenham chorea is thought to be an immune-mediated disorder, steroids are effective in the treatment of severe or refractory cases.

Solution to Question 29:

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

This patient has a rheumatic fever without carditis. He should be given prophylaxis till 21 years of age as the duration of secondary prophylaxis is 5 years or until 21 years of age (whichever is longer).

Secondary prophylaxis is given for patients with acute rheumatic fever (ARF). These patients are at a higher risk of developing a further episode of ARF after a group A streptococcal infection. Hence they should receive long-term penicillin prophylaxis to prevent recurrent infections.

The antibiotics that can be given for secondary prophylaxis are:

- Benzathine penicillin G (1.2 million units) given every 4 weeks (Best antibiotic)
- Oral penicillin V (250 mg) given twice daily (less effective than benzathine penicillin G)
- Erythromycin (250 mg) twice daily (For patients allergic to penicillin)

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Hypertensive Vascular Disease

Question 1:

Which of the following is false with respect to lifestyle modifications for managing essential hypertension as recommended by AHA 2019 guidelines?

a) Low potassium diet

b) Sodium intake < 1.5g /day

c) DASH diet

d) Limit alcohol consumption

Question 2:

A 57-year-old male patient presented to the clinic for a routine clinical check-up. On examination, his BP was found to be 136/88 mmHg. During his subsequent visits, the systolic blood pressure recorded was found to be within the range of 130-138 mmHg and diastolic blood pressure was within the range of 86-88mmHg. According to the European guidelines for hypertension 2018, his blood pressure would be classified as _____

a) Normal

b) High normal

c) Grade 1 hypertension

d) Isolated systolic hypertension

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

An 80-year-old male patient complains of headache and dizziness. His ambulatory BP was recorded to be in the range of 144-152/80-86 mmHg. What is the likely diagnosis?

a) Grade 1 hypertension

b) White coat hypertension

c) Secondary hypertension

d) Isolated systolic hypertension

Question 4:

Sold by @itachibot

Which of the following is the gold standard investigation to measure vessel wall stiffness in patients suspected to have hypertension mediated end-organ damage (HMOD)?

a) Ankle brachial index

b) Carotid Angiography

c) Pulse wave velocity

d) Femoral artery doppler

Question 5:

A 60-year-old male patient has grade 2 hypertension. He is also a known case of type 2 diabetes for 9 years. What is the desired goal for blood pressure range in this patient?

a) 120-130/80-89 mmHg

b) 130-140/80-89 mmHg

c) 120-130/70-79 mmHg

d) 130-140/70-79 mmHg

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

While treating a 45 year old hypertensive patient with underlying coronary artery disease, what target blood pressure should you titrate for, as per the ESC 2018 guidelines?

a) 140-150 mmHg

b) 110-120 mmHg

c) 130-140mmHg

d) 120-130 mmHg

Question 7:

A 58-year-old male patient has been diagnosed with hypertension. On examination, there is no existing cardiovascular disease, no hypertension mediated organ damage (HMOD) and the renal function is normal. He should be immediately started with pharmacological management if his blood pressure is more than _____.

a) >130/80 mmHg

b) ≥160/100 mmHg

c) > 150/100 mmHg

Sold by @itachibot

d) ≥140/90 mmHg

Question 8:

A 72-year-old man who is a known hypertensive for 12 years comes for a regular health check-up. On examination, his blood pressure is 172/100 mmHg. His serum creatinine is 1.5mg/dL, urinary albumin to creatinine ratio (UACR) is 40mg/g, and GFR is 49mL/min. What is the preferred antihypertensive treatment in this scenario?

a) Captopril with amlodipine

b) Captopril with atenolol

c) Telmisartan with atenolol

d) Captopril with telmisartan

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A hypertensive patient is on metoprolol. If verapamil is added, there is a risk of:

a) Atrial fibrillation

b) Bradycardia with AV block

c) Tachycardia

d) Torsades de pointes

Question 10:

A hypertensive patient underwent coronary angiography after he presented with chest pain. His angiographic image is as below. What is the preferred drug for the management of hypertension in this patient?

Sold by @itachibot



a) Spironolactone

b) Atenolol

c) Tamsulosin

d) Hydralazine

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 11:

A 30-year-old man presents with sudden onset of a headache, palpitation, and profuse sweating. He had several such episodes in the past. During these episodes, his BP is found to be high. A 24-hour urinary metanephrine is elevated. What is the most likely etiology?

a) Fibromuscular dysplasia

b) Renal artery stenosis

c) Pheochromocytoma

d) Hyperthyroidism

Question 12:

A 35-year-old female patient has hypokalemia, hypertension and metabolic alkalosis. What is the most likely etiology?

a) Bartter syndrome

b) Gitelman's syndrome

c) Liddle's syndrome

Sold by @itachibot

d) Fanconi's syndrome

Question 13:

Which of the following statements are true with respect to hypertensive emergency as per ESC 2018 guidelines?

a) 1,3,4

b) 1,4

c) 1,2,4

d) 1,2,3,4

Question 14:

Which of the following patients would you diagnose as having hypertensive urgency?

a) 1

b) 2

c) 3

d) 4

Question 15:

A 55-year-old female patient who is a known case of hypertension, diabetes, and hyperlipidemia presents to the emergency department with a severe headache and vomiting. The patient is agitated and her blood pressure is 220/120 mmHg. Ophthalmoscopy reveals the presence of papilledema. Which of the following antihypertensive is least useful in the management of this patient?

a) Furosemide

b) Nicardipine

c) Labetolol

d) Enalaprilat

Question 16:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Which of the following is the most important risk factor for the development of hypertension?

a) High salt intake

b) Age

c) Obesity

d) Genetic factors

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	c
5	c
6	d
7	d
8	a
9	b
10	b
11	c
12	c
13	b
14	c
15	a
16	c

Detailed Explanations

Solution to Question 1:

Low potassium increases blood pressure and hence is not recommended in the dietary modifications. On the contrary, potassium supplementation lowers the BP significantly in hypertensives. It is given preferably by dietary modifications unless contraindicated.

Sold by @itachibot

Solution to Question 2:

According to European guidelines for hypertension 2018, systolic blood pressure between 130-139 mmHg and diastolic blood pressure between 85-89 mmHg is classified as high normal.

Solution to Question 3:

SBP ≥140 mm Hg and DBP <90 mmHg is called isolated systolic hypertension.

White-coat hypertension is characterized by:

• The office blood pressure is ≥140/90 mm Hg on ≥3 separate office visits

• ≥2 blood pressure measurements taken outside the office are <140/90 mm Hg, frequently using home blood pressure monitoring; and

• There is no evidence of hypertensive target organ damage.

Solution to Question 4:

Carotid-femoral pulse wave velocity (PWV) is the gold standard for measuring large artery stiffness due to hypertension.

Long-standing hypertension can lead to structural and functional changes in arteries and target organs like the heart, brain, arteries, and kidneys. This is called hypertension mediated organ damage (HMOD).

Organ	Consequences	Investigations
Heart	Left ventricular hypertrophy Decreased diastolic function	ECGEchocardiography
Blood vessels	Stiffening	Pulse wave velocity of carotid or femoral artery (gold standard)Carotid ultrasoundAnkle-brachial pressure index
Kidney	Hyaline arteriosclerosis (benign hypertension)Hyperplastic arteriosclerosis (severe hypertension)	Renal function testsGFR
Retina	Hemorrhages and cotton wool spots	Fundoscopy
Brain	Stroke, impaired cognition	MRI

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 5:

In a patient (18-65 years) with hypertension and coexisting diabetes, the desired goal for systolic blood pressure is between 120-130 mmHg and diastolic blood pressure is between 70-79 mmHg. In patients above 65 years of age, the desired goal of systolic blood pressure is between 130-139 while the desired diastolic range remains the same.

This goal is in accordance with the European Society of Cardiology guidelines (2018).

Solution to Question 6:

The target systolic blood pressure for hypertensive patients in the age group 18-65 years with coronary artery disease is between 120-130 mmHg.

Solution to Question 7:

In this patient with no markers of cardiovascular disease, no hypertension-mediated end-organ damage (HMOD), and normal renal function, immediate pharmacological treatment is recommended when the blood pressure (BP) is ≥140/90 mmHg according to the AHA and ESH on the management of hypertension.

Guidelines for Hypertension Management:

• Adults 18–79 y with SBP ≥140 mm Hg or DBP ≥90 mm Hg

• Adults ≥80 y with SBP ≥160 mm Hg but ≥140 mm Hg can be considered.

• In frail patients, individualize treatment.

• In patients with CVD, especially CAD, SBP ≥130 mm Hg, or DBP ≥80 mm Hg.

Treatment BP targets:

• Adults 18–79 y, primary SBP/DBP goal <140/90 mm Hg; if treatment well tolerated target SBP/DBP <130/80 mm Hg but not SBP <120 mm Hg or DBP <70 mm Hg. For adults with isolated systolic hypertension, target SBP lowering, albeit cautiously.

• In adults ≥80 y, target SBP/DBP <140/90 mm Hg, if well tolerated

AHA Hypertension grading:

BLOOD PRESSURE CATEGORY	SYSTOLIC BP	DIASTOLIC BP
Normal	<120 mm Hg	<80 mmHg
Elevated	120-129 mmHg	<80 mmHg
Stage 1	130-139 mmHg	80-89 mm Hg

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

BLOOD PRESSURE CATEGORY	SYSTOLIC BP	DIASTOLIC BP
Stage 2	≥140 mm Hg	≥90 mmHg
Hypertensive crisis	≥180 mm Hg	≥120 mm Hg

Solution to Question 8:

In the given clinical scenario, the patient is hypertensive along with chronic kidney disease. The preferred initial antihypertensive treatment is captopril with amlodipine.

The preferred treatment in hypertensive patients with chronic kidney disease is a combination of angiotensin-converting enzyme inhibitors (ACEi) or angiotensin II receptor blockers (ARBs) with calcium channel blockers (CCBs) or diuretics (ACEi/ARBs + CCBs/diuretics).

Solution to Question 9:

Metoprolol, when added with verapamil, leads to bradycardia with AV block.

Verapamil and β blockers have additive effect on inhibiting SA node and AV conduction, and the combination should be avoided to prevent severe bradycardia and conduction block.

These agents can also have additive effect on decreasing left ventricular function which could seriously impair cardiac output in a patient with heart failure.

Solution to Question 10:

The above coronary angiogram shows partial occlusion of the left circumflex coronary artery. Beta-blockers like atenolol in combination with ACE inhibitors or ARBs or calcium channel blockers or diuretics are preferred for the management of hypertension in patients with coronary artery disease.

Other indications for beta-blockers are:

• Congestive heart failure due to systolic dysfunction

• Post myocardial infarction

• Sinus tachycardia

• Ventricular arrhythmias

The image below shows partial occlusion of left circumflex coronary artery.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Solution to Question 11:

The most likely etiology is pheochromocytoma. Patients commonly present with a classical triad of palpitation, headache and profuse sweating. The presence of this triad in association with hypertension and elevated 24-hour urinary metanephrine makes pheochromocytoma a likely diagnosis.

Solution to Question 12:

Hypertension with hypokalemia and metabolic alkalosis is seen in Liddle's syndrome.

Liddle's syndrome (pseudo aldosteronism) is an autosomal dominant condition that is caused due to the increased activation of ENaC channels. This causes increased reabsorption of Na+ exchanged for K+ causing hypokalemia. Increased Na absorption leads to hypertension, and there is also a loss of H+ ion with K+, leading to metabolic alkalosis. It is associated with low renin and low aldosterone levels.

Solution to Question 13:

Hypertensive emergencies are situations in which grade 3 hypertension (BP>180/110 mmHg) is associated with acute hypertension-mediated organ damage (HMOD).

Treatment of hypertensive emergency depends on the type of end-organ damage present (as shown in the table below). It is treated with rapidly acting intravenous antihypertensive agents.

In hypertensive encephalopathy, the initial goal of therapy is to immediately reduce mean arterial blood (MAP) pressure by no more than 25%. Rapid reduction of blood pressure may lead to reduced cerebral blood flow and result in cerebral infarction.

Sold by @itachibot

The table below shows the treatment of various hypertensive emergencies with intravenous drug therapy.

Clinical presentation	Timeline and target for BP reduction	First-line treatment	Alternative
Malignant hypertension with or without acute renal failure	Several hours Reduce MAP by 20-25%	Labetalol Nicardipine	Nitroprusside Urapidil
Hypertensive encephalopathy	Immediately reduce MAP by 20-25%	Labetalol, nicardipine	Nitroprusside
Acute coronary event	Immediately reduce SBP to < 140mmHg	Nitroglycerine, labetalol	Urapidil
Acute cardiogenic pulmonary oedema	Immediately reduce SBP to < 140mmHg	Nitroprusside or nitroglycerine (with loop diuretic)	Urapidil (with loop diuretic)
Acute aortic dissection	Immediately reduce SBP to < 120mmHg and heart rate to < 60 bpm	Esmolol and nitroprusside or nitroglycerine or nicardipine	Labetalol or metoprolol
Eclampsia and severe pre-eclampsia/HELLP	Immediately reduce SBP to < 160mmHg and DBP to < 105mmHg	Labetalol or nicardipine and magnesium sulfate	Consider delivery

©MARROW

Sold by @... If you purchased this from ... you may have been scammed. ... else,

Solution to Question 14:

Patient 3 suffers from hypertensive urgency as BP > 180/110 mmHg and end-organ damage is not seen.

Target organ damage is absent in hypertensive urgency but is present in a hypertensive emergency.

Severe hypertension (grade 3) is generally defined as blood pressure ≥180/110 mmHg. Hypertensive emergency, hypertensive urgency, and malignant hypertension are all forms of severe hypertension.

Option A: Patient 1 suffers from hypertension. Since BP <180/110 mmHg it is not a hypertensive crisis (urgency or emergency).

Option B: Patient 2 has a BP > 180/110 mmHg along with symptoms of hypertensive encephalopathy. This is an example of a hypertensive emergency as there is severe hypertension with signs or symptoms of hypertensive-mediated organ damage (HMOD).

Option D: Patient 4 has a BP > 180/110 mm Hg along with papilledema. This is a case of severe hypertension associated with fundoscopic changes (flame hemorrhages, papilloedema) which is suggestive of malignant hypertension. Microangiopathy, DIC, and encephalopathy may also be present. Fibrinoid necrosis of arterioles and small arteries is seen. The prognosis is very poor if untreated.

Posterior reversible encephalopathy syndrome (PRES) or acute hypertensive encephalopathy is a rare complication in patients with an acute rise in blood pressure. Due to the acute rise, there is increased cerebral perfusion and edema of the brain. Clinically it is recognized by alteration in mental status, nausea, headache, visual disturbance, seizure, or development of focal neurological deficits.

Sold by @itachibot

Solution to Question 15:

The above scenario points towards the diagnosis of malignant hypertension. The drug which is least useful in the management is furosemide(diuretic).

Volume depletion is common in patients with malignant hypertension, and the administration of a diuretic together with a hypertensive agent can lead to an extreme drop in blood pressure.

Diuretics are generally not recommended for the rapid reduction of blood pressure in patients with hypertensive crises. They are used only in conditions of volume overload such as acute pulmonary edema.

Solution to Question 16:

The most important risk factor for the development of hypertension is obesity.

Hypertension is defined as repeated office systolic blood pressure of ≥ 140 mm Hg and/or diastolic blood pressure of ≥ 90 mm Hg.

Primary/essential hypertension is the leading cause of hypertension in adults. It is caused due to the interaction of genetic factors with various other factors such as diet, sedentary lifestyle, etc. In some people, hypertension may result from renal, endocrine, or adrenal disorders (secondary hypertension).

Risk factors of hypertension can be divided into modifiable and non-modifiable:

Non-modifiable risk factors are:

• Age- risk increases with age

• Sex- in young and middle-aged adults, the risk is higher in males

• Genetic factors- mutation in the genes which encode the components of the renin-angiotensin-aldosterone system, atrial natriuretic peptide, and alpha adducin

• Ethnicity

Modifiable risk factors are:

• Obesity (most important)

• High alcohol intake

• High salt intake

• Sedentary lifestyle

• Stress

• Dyslipidemia

• Diet low in potassium and calcium

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Ischemic Heart Disease - Presentation and Diagnosis

Question 1:

Which of the following arteries is most commonly involved in patients with coronary artery disease?

a) Right coronary artery

b) Left anterior descending coronary artery

c) Left circumflex coronary artery

d) Left main coronary artery

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 2:

Which area of the myocardium is most vulnerable to damage in patients with ischemic heart disease?

a) Subepicardium

b) Subendocardium

c) Pericardium

d) Midmyocardium

Question 3:

A patient presented to the emergency with symptoms of unstable angina. ECG done in pain-free interval showed biphasic T waves in precordial leads without ST changes. Cardiac troponins were normal. Coronary angiography revealed stenosis of the proximal left anterior descending artery. What is the likely diagnosis?

a) Broken heart syndrome

b) Wellens syndrome

c) Barlow syndrome

d) Brugada syndrome

Question 4:

Sold by @itachibot

In a patient with suspected stable angina, which of the following findings are likely to be present?

a) CK-MB is elevated ; Troponin T is normal

b) Troponin I is elevated ; Myoglobin is normal

c) Myoglobin is elevated ; CK-MB is normal

d) Troponin I is normal ; CK-MB is normal

Question 5:

While evaluating a patient with chest pain in the emergency, you ask him to localize the sensation. He places a hand over the sternum with a clenched fist. What is this sign called?

a) Mayne's sign

b) Levine's sign

c) Frank's sign

d) Freidriech's sign

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A 60-year-old woman presents with chest discomfort, associated with a squeezing sensation in the substernal area. She added that it was precipitated by physical activity and relieved on taking rest. Which of the following is a feature of this condition?

a) Pain radiating to the trapezius

b) Pain lasting more than 20 minutes

c) Pain localized above the mandible

d) Crescendo decrescendo nature of pain

Question 7:

A patient presented with angina and ECG showed ST-segment elevation. The pulse rate was found to be 54 beats/min and blood pressure was 98/56 mm Hg. What is the most likely diagnosis?

a) Anterior wall MI

b) Posterior wall MI

c) Inferior wall MI

Sold by @itachibot

d) Anteroseptal wall MI

Question 8:

A patient was admitted with acute MI. During a daily examination, a transient murmur was heard. What is the most likely cause for this finding?

a) Mitral stenosis

b) Mitral regurgitation

c) Aortic stenosis

d) Aortic regurgitation

Question 9:

Which of the following cardiac biomarkers is the earliest to rise after acute myocardial infarction?

a) Creatine kinase

b) Myoglobin

c) Troponin T

d) Troponin I

Question 10:

A patient is under evaluation for acute MI. Which of the following biomarkers would you prefer to test for?

a) CK-MB

b) Myoglobin

c) Cardiac troponin-T

d) Cardiac troponin-I

Question 11:

Elevated cardiac troponins may not be seen in which of the following patients?

a) A 24-year-old male with myocarditis

b) A 65-year-old woman suffering from Takotsubo cardiomyopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) A patient who underwent cardioversion for VT in the emergency room

d) A 48-year-old man diagnosed with angina

Question 12:

Which of the following biomarkers is preferred for diagnosing reinfarction in a patient post-NSTEMI?

a) Troponins

b) Myoglobin

c) LDH

d) CK-MB

Question 13:

Which of the following is likely to be seen in a patient diagnosed with acute myocardial infarction?

a) Ratio of LDH 1 / LDH 2 < 1 in plasma

b) Ratio of LDH 1 / LDH 2 > 1 in plasma

c) Ratio of LDH 1 / LDH 4 < 1 in plasma

d) Ratio of LDH 2 / LDH 1 > 1 in plasma

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 14:

What is the maximum predicted heart rate in a 50-year-old gentleman on exercise tolerance test?

a) 150 bpm

b) 170 bpm

c) 200 bpm

d) 160 bpm

Question 15:

Which of the following is not a contraindication for performing an exercise tolerance test?

a) Unstable angina

Sold by @itachibot

b) Symptomatic aortic stenosis

c) Active endocarditis

d) AAA with transverse diameter of 5.5 cm

Question 16:

Which of the following drugs cannot be used for a patient undergoing stress echocardiography?

a) Dobutamine

b) Dipyridamole

c) Regadenoson

d) Phenylephrine

Question 17:

Which of the following findings during an exercise tolerance test is not associated with multivessel coronary artery disease?

a) Early onset of ST depression

b) Persistence of ST segment changes late into recovery phase

c) Failure to increase systolic BP by at least 20mmHg

d) ST segment elevation in aVR

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

Which of the following compounds is used for myocardial perfusion imaging?

a) Iodine-123

b) Iridium-192

c) Thallium-201

d) Gallium-67

Question 19:

A patient is being considered for CABG in order to achieve coronary revascularization. Which of the following test will not be useful to assess myocardial viability for predicting the risk of

Sold by @itachibot

perioperative morbidity?

a) Thallium 201 imaging

b) Cardiac MRI

c) PET

d) CT

Question 20:

Which questionnaire is used for the assessment of a patient who presented with symptoms of angina?

a) ROSE questionnaire

b) IPSS questionnaire

c) CAGE questionnaire

d) YTS questionnaire

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

Which of the following ECG findings is first to appear in a patient suspected of acute myocardial infarction?

a) Elevation of S wave

b) Prolonged QT interval

c) Tall T waves

d) Prolonged PR interval

Question 22:

A patient presented with the complaints of chest pain. Which among the following is the best investigation to diagnose MI within 2-4 hours?

a) LDH

b) CKMB

c) Troponin-T

d) BNP

Sold by @itachibot

Question 23:

Which of the following conditions is not associated with an ST-segment elevation?

a) Prinzmetal's angina

b) Pericarditis

c) Ventricular aneurysm

d) Unstable angina

If you bought this from anyone else, it is stolen property. Please report to @itachibot

Answer Key

Question No.	Correct Option
1	b
2	b
3	b
4	d
5	b
6	d
7	c
8	b
9	b
10	d
11	d
12	a
13	b
14	b
15	d
16	d
17	c
18	c
19	d
20	a
21	c
22	c
23	d

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The left anterior descending (anterior interventricular) coronary artery is most commonly involved in coronary artery disease (CAD) or coronary heart disease or ischemic heart disease.

The regions affected are the anterior wall of the left ventricle near the apex and the anterior portion of the ventricular septum.

CAD is caused by plaque buildup in the walls of the coronary arteries that narrow over time and can partially or totally block the blood flow causing myocardial ischemia and subsequently infarction.

The image below shows coronary circulation.



Solution to Question 2:

Subendocardium is the region of the myocardium most vulnerable to ischemic damage.

The myocardium is perfused from the epicardium to the endocardium. This results in the subendocardium being the least perfused region making it most vulnerable to ischemic damage.

A narrow zone of the myocardium immediately beneath the endocardium is spared from necrosis because it can be oxygenated by diffusion from the ventricle.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 3:

The above scenario is suggestive of Wellens syndrome. It is alternatively known as anterior, descending, T-wave syndrome. The ECG pattern of Wellens syndrome is relatively common in patients who exhibit symptoms consistent with unstable angina.

ECG in the pain-free interval is characterized by deeply inverted or biphasic T waves in precordial leads and with minimal or no ST elevations. Cardiac enzymes are normal or mildly elevated.

These patients are at high risk for impending large anterior wall acute myocardial infarction.

The ECG below shows biphasic T waves in leads V2 and V3 and inverted T waves in leads V4 and V5 in Wellens syndrome.



Option A: Broken heart syndrome is another name for Takotsubo cardiomyopathy.

Sold by @itachibot

Option C: Barlow syndrome is associated with mitral valve prolapse.

Option D: Brugada syndrome is a genetic cardiac rhythm disorder.

Solution to Question 4:

The levels of cardiac markers remain unchanged in stable and unstable angina as there is no necrosis of cardiac myocytes.

The clinical presentations of ischemic heart disease can be broadly classified into:

• Chronic coronary syndrome (CCS) - this includes

• Stable angina - this is an episodic angina that occurs with predictable severity, frequency, and threshold of precipitation by stressors, and is relieved by rest and nitrates

• Acute coronary syndrome (ACS) - this includes

• Unstable angina - this is diagnosed in patients with symptoms suggestive of an ACS with no elevation of biomarkers

• Myocardial infarction - it is the necrosis and death of a portion of the myocardium due to prolonged ischemia. The cardiac biomarkers are elevated in this condition

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 5:

When a patient with angina is asked to localize the sensation, he or she typically places a hand over the sternum, sometimes with a clenched fist, to indicate a squeezing, central, substernal discomfort. This is called Levine's sign.

Option A: Mayne's sign- Drop of at least 15 mmHg in the diastolic blood pressure on raising the arm. It occurs in patients with aortic regurgitation (not reliable).

Option C: Frank's sign- Diagonal crease in the ear lobe indicative of cardiovascular diseases and/or diabetes.

Option D: Friedreich's sign- Exaggerated drop in JVP seen in constrictive pericarditis.

Solution to Question 6:

The above scenario is suggestive of stable angina in which the pain is usually crescendo-decrescendo in nature.

Characteristic features of pain in stable angina:

• Squeezing, central, substernal discomfort

• Crescendo-decrescendo in nature

• Typically lasts 2 to 5 min

Sold by @itachibot

• Can radiate occasionally to neck, jaw, epigastrium, shoulders, left arm.

Angina is rarely localized below the umbilicus or above the mandible. Myocardial ischemic discomfort does not radiate to the trapezius muscles, that radiation pattern is more typical of pericarditis.

Features	Stable angina	Unstable angina
Pain	Squeezing, central, substernal discomfort.Crescendo-decrescendo pattern.Precipitated by exercise, cold weather, or emotional stress duration of 2-5 min.	Same as stable angina but may be more severe.Crescendo pattern.Pain occurs at rest and may last up to 20 min.
Pathology	Ischemia due to fixed stenosis	Unstable plaques causing dynamic obstruction
Predictability	Predictable	Unpredictable
Diagnosis	ECG during exercise	ECG during exercise contraindicatedCoronary angiography recommended
ECG	ECG at rest: normalECG in exercise: ST-segment depression and ischemic T-waves	ECG at rest: ST-segment depression and ischemic T-waves



Solution to Question 7:

The above scenario is suggestive of STEMI with bradycardia which is a feature of inferior wall myocardial infarction. Up to one-half with inferior infarction show evidence of parasympathetic hyperactivity (bradycardia and/or hypotension).

Sold by @itachibot

Many patients have a normal pulse rate and blood pressure within the first hour of STEMI.

About one-fourth of patients with anterior infarction have manifestations of sympathetic nervous system hyperactivity (tachycardia and/or hypertension).

Solution to Question 8:

A transient midsystolic or late systolic apical murmur due to mitral regurgitation secondary to dysfunction of the mitral valve apparatus (papillary muscle dysfunction, LV dilation) may be present in STEMI.

Solution to Question 9:

Myoglobin is the earliest to rise in acute MI followed by troponin, CK-MB and then LDH.

Serum myoglobin and heart-type fatty acid-binding protein are smaller molecules that diffuse through interstitial ■uids more rapidly after cell death than the larger CK and troponin molecules. They become abnormal as early as 30 minutes after myocardial injury. Neither molecule is specific to myocardial tissue. Myoglobin levels are not tested in routine practice.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Solution to Question 10:

The most preferred cardiac biomarker for myocardial injury/acute MI amongst the given options is cardiac troponin-I.

Cardiac troponin-I, CK-MB, and myoglobin are significantly more specific for acute-MI than cardiac troponin-T.

The cardiac troponins are particularly valuable when there is clinical suspicion of a small MI that may be below the detection limit for CK-MB measurement. They have replaced CK-MB as the biomarker of choice for diagnosing myocardial infarction.

Cardiac troponins:

• Most specific cardiac biomarker for myocardial injury/acute MI

• Elevations are useful for short & long-term prognoses in MI

• Elevations correlate with the estimation of infarct size and risk of death

• Newer highly sensitive assays for troponins become positive within the first 3 hours after onset of chest pain in MI

Solution to Question 11:

Sold by @itachibot

Troponins are not elevated in patients suffering from angina. Troponins are elevated in the case of myocardial necrosis (seen in MI) whereas angina is associated with myocardial insufficiency/ischemia.

Conditions associated with an elevation in cardiac troponins:

• MI

• Myocarditis

• Takotsubo cardiomyopathy

• Myocardial contusion

• Cardioversion/defibrillation

• LV strain from CHF

• Hypertensive crisis

• RV strain from PE

• Renal disease

• Sepsis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

Troponin(serial) is the preferred biomarker for diagnosing reinfarction.

Serial troponin levels are used to detect reinfarction, an acute MI that occurs within 28 days of an incident or recurrent MI.

In case of suspected reinfarction, an immediate measurement of cardiac troponin followed by a second sample after 3-6 hours. ≥20 percent rise in the second sample suggests reinfarction.

Solution to Question 13:

In myocardial injury, the LDH-1 fraction increases, and the ratio of LDH-1 to LDH-2 becomes >1 in the plasma. This is called flipped LDH pattern.

Normally, the LDH-2 fraction is greater than the LDH-1 fraction in the plasma, and therefore, the ratio of LDH-1 to LDH-2 is normally <1.

Solution to Question 14:

Maximum predicted heart rate on an exercise tolerance test (treadmill test) is given by: HRmax = 220 – Age

So in a 50-year-old male, the maximum predicted heart rate is 220 - 50 = 170 bpm.

Sold by @itachibot

Solution to Question 15:

Treadmill exercise testing (treadmill test) may be safely performed in patients with an abdominal aortic aneurysm.

Absolute contraindications to exercise testing:

• Acute myocardial infarction (within 2 days)

• High-risk unstable angina

• Uncontrolled cardiac arrhythmia with hemodynamic compromise

• Active endocarditis

• Symptomatic severe aortic stenosis

• Decompensated heart failure

• Acute pulmonary embolism or pulmonary infarction

• Acute myocarditis or pericarditis

• Severe pulmonary hypertension

• Aortic dissection

Relative Contraindications to exercise testing:

• Known left main coronary artery stenosis

• Moderate aortic stenosis with uncertain relation to symptoms

• Tachyarrhythmias with uncontrolled ventricular rates

• Acquired complete heart block

• Hypertrophic cardiomyopathy with severe resting gradient

• Mental impairment with limited ability to cooperate

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 16:

Phenylephrine, an alpha-receptor agonist, cannot be used for a patient undergoing stress echocardiography as it causes vasoconstriction.

Dobutamine (preferred agent) and vasodilators (dipyridamole, adenosine, regadenoson) are used for pharmacological stress echocardiography.

Pharmacologic stress testing is used in those who cannot exercise because of pulmonary veno-occlusive disease, musculoskeletal disease, or exertional dyspnea.

Stress echocardiography can be done using either exercise (e.g.: treadmill test or bicycle test) or pharmacologic agents.

Sold by @itachibot

Solution to Question 17:

A failure to increase systolic blood pressure by 10 mm Hg, or a sustained decrease in systolic blood pressure of 10 mm Hg or more, is suggestive of multivessel (or left main) CAD.

Other abnormalities on exercise testing that are suggestive of multivessel CAD include:

• Early-onset of ischemic ST-segment depression

• ST-segment depression of ≥2 mm (0.2mV) involving 5 or more leads, or persisting ≥5 minutes into recovery

• Exercise-induced ST-segment elevation is also consistent with multivessel CAD

• Recent evidence suggests that ST-segment elevation of ≥1 mm in lead aVR may be a significant predictor of left main coronary artery disease, proximal left anterior descending (LAD) artery disease, or at least multivessel CAD

An exercise tolerance/treadmill test involves walking in place on a treadmill while monitoring the electrical activity of your heart. Throughout the test, the speed and incline of the treadmill increase. The results show how well your heart responds to the stress of different levels of exercise.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 18:

Myocardial perfusion imaging is done with either thallium-201 or technetium-99m labeled compounds (e.g., technetium-99m sestamibi).

Solution to Question 19:

Modalities for the determination of myocardial viability include:

• Thallium-201 imaging

• MRI

• Dobutamine echocardiography

• PET scan (investigation of choice)

The investigation of choice for assessing the ventricular function is cardiac MRI.

Solution to Question 20:

WHO Rose questionnaire is used for the assessment of angina.

Option B: International prostate symptom score (IPSS) questionnaire is used for BPH.

Option C: The CAGE questionnaire is used to check for signs of possible alcohol dependency.

Sold by @itachibot

Option D: The youth tobacco survey (YTS) questionnaire is used to study the prevalence of use of tobacco products amongst youth.

Solution to Question 21:

During acute myocardial infarction, the ECG evolves through 3 stages:

• T-wave peaking and T-wave inversion:

• Peaked T-wave aka hyperacute T-wave- first ECG change (occurs within minutes)

• If the blood flow is restored or the oxygen demands of the heart are met, the T waves will revert to normal

• If myocardial cell death has occurred, T wave inversions appear (over days) and persist for months to years.

• A nonspecific finding and is indicative only of ischemia and not diagnostic of infarction

• ST-segment elevation occurs over minutes to hours (after T-wave peaking)

• The appearance of new pathologic Q-waves:

• Occurs over hours to days (after ST-segment elevation) and is indicative of an evolving MI

• The persistence of pathological Q-waves indicates irreversible myocardial cell death



Solution to Question 22:

Measurement of troponin is the best investigation to diagnose MI within 2-4 hours.

Sold by @itachibot



Troponin T/I:

• Highly specific for MI

• Troponin I is more important than troponin T.

• Troponins will remain elevated longer than creatine kinase (CK), for up to 14 days.

• This makes troponins a superior marker for diagnosing MI in the recent past (late diagnosis).

• Reinfarction: It is now clear that cardiac troponin (cTn) increases rapidly, even if it is from an abnormal baseline in patients with reinfarction.

• Sometimes cTn concentration is elevated, but stable or decreasing at the time of suspected reinfarction. In this case, the diagnosis of reinfarction requires a 20% or greater increase in the cTn measurement

Creatine kinase (CK):

• Less sensitive and specific than troponins

• When cTn is available, CK-MB should not be used for the initial diagnosis of acute MI

Lactate dehydrogenase (LDH):

• Normally, LDH2>LDH1; but, in MI, LDH1>LDH2. This is called flipping of the LDH ratio.

• Current recommendations suggest that LDH no longer has a role in the diagnosis of MI.

Myoglobin:

• Earliest enzyme to increase after MI.

• Not specific to cardiac muscle and can be elevated to any form of injury to skeletal muscle.

Heart-type fatty acid-binding protein (H-FABP), which mainly exists in cardiomyocytes, has recently emerged as a potential biomarker, which was found to be sensitive for the early diagnosis of acute myocardial infarction. H-FABP as an early marker and cTnI as a late marker would be the ideal combination to cover the complete diagnostic window for AMI

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Sold by @itachibot

Solution to Question 23:

There is no ST-segment elevation in unstable angina.

Differential diagnosis of ST-segment elevation (Mnemonic-ELEVATION):

• Electrolytes(Hypercalcemia, Hyperkalemia)

• Left bundle branch block

• Early(Benign) repolarization, Embolism(Acute pulmonary)

• Ventricular hypertrophy

• Arrhythmia(Brugada syndrome), Aneurysm of LV, Aortic dissection, Antiarrhythmic drugs(Class 1c)

• Tako-tsubo syndrome, Tumor invading left ventricle, Trauma to ventricles

• Infarction(MI), Inflammation(Myocarditis, Pericarditis)

• Osborn J waves (Hypothermia or hypercalcemia)

• Non-atherosclerotic vasospasm(Prinzmetal's angina)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Ischemic Heart Disease - Complications and Management

Question 1:

A 38-year-old woman presented with chest pain that was unrelated to physical exertion. ECG showed ST-segment elevation. Intracoronary acetylcholine provocation testing revealed a transient spasm of the right coronary artery. What is the drug of choice in this condition?

a) Dopamine

b) Diltiazem

c) Propranolol

d) Digoxin

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 2:

A patient who presents to the ER with chest pain and dyspnea is diagnosed with STEMI. Swift revascularization by thrombolysis is planned for her. What is the dose of streptokinase to be used?

a) 0.5 million units IV over 1 hour

b) 1.5 million units IV over 1 hour

c) 2 million units IV over 1 hour

d) 2.5 million units IV over 1 hour

Question 3:

Which of the given fibrinolytic agents is least preferred for thrombolysis in a patient with MI?

a) Streptokinase

b) Alteplase

c) Tenecteplase

d) Reteplase

Question 4:

Sold by @itachibot

While planning thrombolysis for a patient with MI, use of which of the following agents will reduce the chances of a dosing error?

a) Streptokinase

b) Tissue plasminogen activator

c) Alteplase

d) Tenecteplase

Question 5:

Which of the following is not an absolute contraindication for the use of a fibrinolytic agent in patients with an acute MI?

a) Non-haemorrhagic stroke within 1 year

b) Marked hypertension

c) Prolonged cardiopulmonary resuscitation

d) Suspicion of aortic dissection

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

All of the following drugs should be avoided in a patient with angina except:

a) Thyroxine

b) Sumatriptan

c) Dipyridamole

d) Trimetazidine

Question 7:

Which of the following combinations should be avoided while treating a patient with angina?

a) Nitrates and β-blocker

b) Nitrates and Verapamil

c) Verapamil and β-blocker

d) Amlodipine and β-blocker

Question 8:

Sold by @itachibot

What is the mechanism of action of ranolazine?

a) Inhibits adenosine reuptake

b) Inhibits late inward Na current

c) Potassium channel opener

d) Rho-kinase inhibitor

Question 9:

A patient on clopidogrel is scheduled to undergo an elective operation. What is your advice regarding the drug?

a) Stop clopidogrel 12-24 hours before operation

b) Continue clopidogrel

c) Stop clopidogrel 7-10 days before operation

d) Stop clopidogrel 5-7 days before operation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following drugs would you prescribe to prolong life in a post-MI patient?

a) Beta blockers

b) Calcium channel blockers

c) Nitrates

d) Aspirin

Question 11:

A 60-year-old man post-MI had the following lab values. Which of the following must be his treatment plan?

a) Rosuvastatin + fenofibrate

b) Atorvastatin 80 mg

c) Fenofibrate

d) Rosuvastatin 10 mg

Question 12:

Sold by @itachibot

A patient presents with an intense exertional chest pain radiating to the neck along with shortness of breath for 30mins. ECG reveals ST-segment depression. Cardiac troponins are elevated. Which of the following is incorrect regarding the management of this condition?

a) Aspirin loading dose should be commenced immediately

b) ACC/ AHA guidelines call a P2Y12 inhibitors as a class I recommendation

c) Glycoprotein IIB/IIIA inhibitors are useful adjuncts

d) Fibrinolytic therapy is useful in this condition

Question 13:

A patient is diagnosed with STEMI on presentation. Coronary angiography showed complete occlusion of the infarct-related artery. Immediate primary PCI with a drug-eluting stent is planned. Which of the following drug-eluting stents will have a relatively higher risk of restenosis and late stent thrombosis?

a) Zotarolimus

b) Biolimus

c) Everolimus

d) Sirolimus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 14:

A 40-year-old male patient presents to the emergency department with chest pain for 2 hours. The ECG shows ST-segment depression and cardiac troponins are elevated. The patient has a history of PCI 3 months back. He is given aspirin, clopidogrel, nitrates, and unfractionated heparin and shifted to the coronary care unit. What is the next best course of action?

a) Continue conservative management and monitoring of cardiac enzymes and ECG

b) Continue conservative management and plan for delayed revascularization procedure after the patient is discharged

c) Immediate revascularization with thrombolytics

d) Revascularization with PCI

Question 15:

For an acute STEMI patient with cardiogenic shock, what would be the treatment of choice?

Sold by @itachibot

a) Intra-aortic balloon pump with PCI

b) Revascularization with PCI

c) Thrombolysis

d) Intra-aortic balloon pump

Question 16:

A 40-year-old chronic smoker comes with acute epigastric discomfort for the past one hour. ECG shows ST-segment depression. What is the immediate intervention?

a) Aspirin

b) Thrombolytic therapy

c) IV pantoprazole

d) PCI

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Question 17:

A previously healthy 58-year-old man is admitted to the hospital because of an acute MI. ECG is shown below. Within several hours, he becomes oliguric and hypotensive. On auscultation, no added sounds were present. What is the most likely reason for this patient being in shock?



a) Mitral regurgitation

b) Infarction causing septal defect

c) Right ventricular infarction

Sold by @itachibot

d) Decreased ejection fraction from the left ventricle

Question 18:

A patient presented with sudden circulatory collapse and engorged neck veins 3 days post-MI. A bedside transthoracic echocardiogram revealed the following. Which of the following is false regarding the cause of this patient's findings?



a) It usually results in immediate death

b) Immediate pericardiocentesis should be done

c) It usually involves inferior wall

d) Definitive management is surgical repair

Question 19:

Which of the following statements about Dressler syndrome is false?

a) It presents as pericarditis

b) It is an autoimmune phenomenon

c) First line therapy consists of aspirin

d) It occurs 2-7 days after MI

Question 20:

Sold by @itachibot

Left ventriculography with contrast was performed in a post-MI patient and is shown below. Infarct involving which portion of the myocardium is most likely to cause this condition?

a) Subendocardial

b) Anterior transmural

c) Posterior transmural

d) Inferior wall

If you purchased this from Sold by @ you may have been scamm... else,

Question 21:

A 50-year-old man started experiencing chest pain and was rushed to a primary care facility at 10 AM. A diagnosis of acute severe MI was made. Due to the unavailability of PCI, he was taken to a tertiary center by 10:30 AM. By what time should PCI be performed in this patient at the latest?

a) 11 AM

b) 12 PM

c) 11:30 AM

d) 12:30 PM

Question 22:

Which of the following defines PCI related MI ?

a) Elevation of cTn > 5 times normal in the absence of any other features

Sold by @itachibot

b) Elevation of cTn > 5 times normal in the presence of symptoms of MI, ECG changes, angiographic findings or imaging abnormality

c) Elevation of cTn > 3 times normal in the absence of any other features

d) Elevation of cTn > 3 times normal in the presence of symptoms of MI, ECG changes, angiographic findings or imaging abnormality

Question 23:

A patient comes to the hospital with chest pain for the last 3 hours. Vitals are stable. ECG revealed ST depression and T wave inversion in anterior chest leads. Which of the following would be the most appropriate line of management in this patient?

a) PCI

b) Thrombolysis with alteplase

c) Prophylaxis for arrhythmia

d) Aspirin with heparin

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	d
5	c
6	d
7	c
8	b
9	d
10	a
11	b
12	d
13	d
14	d
15	b

Sold by @itachibot

16	a
17	c
18	c
19	d
20	b
21	b
22	b
23	d

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The above scenario in which rest pain is associated with ST-segment elevation and transient coronary vasospasm is suggestive of Prinzmetal's angina. The drug of choice is a calcium channel blocker.

Prinzmetal variant angina (PVA) is a clinical syndrome in which chest pain occurs without the usual precipitating factors. It usually results from coronary vasoconstriction (vasospasm) and tends to involve the right coronary artery.

It is a supply ischemia and usually causes transmural ischemia (subendocardial in classic angina pectoris).

It often affects women under 50 years of age. It characteristically occurs in the early morning, awakening patients from sleep, and may be associated with arrhythmias or conduction defects.

The clinical diagnosis of PVA is made by the detection of transient ST-segment elevation with rest pain. Coronary angiography demonstrates transient coronary spasm which is the diagnostic hallmark of PVA.

Nitrates and calcium channel blockers both can be used in alleviating symptoms in Prinzmetal's angina. However due to the development of nitrate tolerance in some individuals, they are less preferred and calcium channel blockers such as verapamil, diltiazem are used as first-line agents.

Solution to Question 2:

Streptokinase is given in a dose of 1.5 million units IV given over 1 hour for treatment of MI.

Fibrinolytic therapy should ideally be initiated within 30 min of presentation. The fibrinolytic agents that have been approved for IV use in patients with STEMI are:

• Tissue plasminogen activator (tPA)

• Streptokinase

• Tenecteplase (TNK)

Sold by @itachibot

• Reteplase (rPA)

Adverse effects of streptokinase:

• Transient hypotension

• Allergic reactions

• Hemorrhage

Solution to Question 3:

Streptokinase has the lowest clot selectivity among the fibrinolytic agents.

Fibrinolytic agent with:

• Least fibrin specificity– streptokinase

• Most fibrin specificity – tenecteplase

Solution to Question 4:

Tenecteplase is given as a single bolus dose (based on body weight) which reduces the rate of medication errors (common with IV infusions).

Tenecteplase and reteplase are referred to as bolus fibrinolytics as their administration does not require prolonged intravenous infusion.

Solution to Question 5:

Prolonged cardiopulmonary resuscitation is a relative contraindication for fibrinolytic therapy.

Contraindications for the use of fibrinolytic agents	
Absolute contraindications	Relative contraindications
History of hemorrhagic stroke at any time	Current use of anticoagulants (INR ≥2)
Non-hemorrhagic stroke or other cerebrovascular events within the past year	Recent (<2 weeks) invasive or surgical procedure
Marked hypertension (SBP>180mmHg and/or a DBP>110 mmHg) at any time during the acute presentation	Prolonged (>10 min) cardiopulmonary resuscitation
Suspicion of aortic dissection	Known bleeding diathesis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Contraindications for the use of fibrinolytic agents	
Active internal bleeding (excluding menses)	Pregnancy
	Hemorrhagic ophthalmic condition (e.g., hemorrhagic diabetic retinopathy)
Active peptic ulcer disease	
History of severe hypertension that is currently adequately controlled	

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

Trimetazidine is an antianginal drug.

It is a PFox (partial fatty acid oxidation) inhibitor. It inhibits the beta-oxidation of fatty acids and thus enhances glucose oxidation. This results in decreased oxygen consumption in ATP synthesis as beta-oxidation consumes more oxygen.

Option A: Thyroxine increases the heart rate, which can increase the myocardial oxygen demand and can precipitate angina.

Option B: Sumatriptan and other triptans cause coronary vasospasm. They are therefore contraindicated in patients with CAD.

Option C: Dipyridamole can produce coronary steal phenomenon and can worsen angina.

Solution to Question 7:

Verapamil and β-blockers have additive effects on inhibiting SA node and AV conduction, the combination should be avoided to prevent severe bradycardia and conduction block. These agents can also have additive effects on decreasing left ventricular function which could seriously impair cardiac output in a patient with heart failure.

Diltiazem can be combined with beta-blockers in patients with normal ventricular function and no conduction disturbances.

Amlodipine and β-blockers have complementary actions on coronary blood supply and myocardial oxygen demands. Whereas the former decreases blood pressure and dilates coronary arteries, the latter slows the heart rate and decreases contractility.

Solution to Question 8:

Sold by @itachibot

The predominant mechanism of action of ranolazine is inhibition of the late sodium current in myocardial cells.

Ranolazine is useful for patients with chronic angina despite standard medical therapy. It is administered at a dose of 500-1000 mg twice daily.

Contraindications:

• Hepatic impairment

• QT prolongation

• Drugs inhibiting CYP3A metabolic system- ketoconazole, diltiazem, verapamil, macrolide antibiotics, HIV protease inhibitors, and large quantities of grapefruit juice

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

Clopidogrel should be stopped 5-7 days prior to surgery to avoid the risk of increased surgical bleeding.

Solution to Question 10:

Beta-blockers are the only antianginal agents that have been demonstrated to prolong life in patients with coronary disease (post-myocardial infarction).

Note: B-blockers and spironolactone have been demonstrated to improve survival in individuals with heart failure.

Solution to Question 11:

In this 60 yr old patient with post-myocardial infarction (Atherosclerotic cardiovascular disease patient), high-intensity statins should be given irrespective of cholesterol levels. So atorvastatin 80mg is the treatment plan in the given case scenario.

Examples of high-intensity statins include high-dose atorvastatin 40–80 mg/day and rosuvastatin 20–40 mg/day.

Note: Fenofibrate is not advocated unless TG is more than 500 mg/dl. It can be administered along with a moderate-intensity statin.

Sold by @itachibot



Solution to Question 12:

The above scenario is suggestive of NSTEMI. Antiplatelet and anticoagulation therapies and coronary intervention are mainstays of treatment.

A distinction in acute coronary syndrome between patients with and without ST-segment elevation at presentation is essential to determine the need for reperfusion therapy. Fibrinolytic therapy is harmful in acute coronary syndrome without ST-segment elevation, unlike with ST-segment elevation, where acute reperfusion saves lives.

Solution to Question 13:

Sirolimus and paclitaxel are a part of first-generation drug-eluting stents that carry a relatively higher risk of restenosis and late stent thrombosis than second-generation drug-eluting stents.

Second-generation drug-eluting stents use newer agents such as everolimus, biolimus, and zotarolimus. These second-generation drug-eluting stents are more effective with fewer complications like stent thrombosis than first-generation stents.

An antiproliferative agent is attached to the stent by the use of a thin polymer coating. This antiproliferative drug elutes from the stent over a 1 to 3 month period after implantation.

Solution to Question 14:

The patient described above is categorized as a high-risk patient due to his history of PCI 3 months ago. The next best course of action for this patient is immediate revascularization with PCI.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Revascularization with PCI (percutaneous coronary intervention) is indicated in high-risk patients who have one or more of the following:

• New ST-segment depression

• Diabetes mellitus

• Renal dysfunction - eGFR <60mL/min per 1.73m2

• Ejection fraction <40%

• Early postinfarction angina

• PCI within the past 6 months

• Prior CABG

• Refractory angina

• Symptoms of congestive heart failure

• Hemodynamic instability

• Recurrent angina at rest/low-level activity despite treatment

• Elevated troponin T or troponin I

• Sustained VT or VF

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 15:

The treatment of choice in patients with acute STEMI with cardiogenic shock is revascularization with PCI.

The 2013 ACCF/AHA guidelines for the treatment of cardiogenic shock in patients with acute MI are:

• Emergency revascularization with PCI or CABG is recommended in suitable patients irrespective of the time delay from MI onset.

• In the absence of contraindications, fibrinolytic therapy should be administered to patients with STEMI and cardiogenic shock who are unsuitable candidates for either PCI or CABG.

Routine use of Intra-Aortic Balloon Pump counterpulsation (IABP) is not recommended for cardiogenic shock (based on IABP-SHOCK II trial). IABP can be useful for those patients with cardiogenic shock after STEMI who do not quickly stabilize with pharmacological therapy or have mechanical complications (eg, acute mitral regurgitation or rupture of the ventricular septum).

Alternative left ventricular assist devices for circulatory support may be considered in patients with refractory cardiogenic shock.

Solution to Question 16:

The above scenario is suggestive of a non-ST-elevation acute coronary syndrome (NSTE-ACS). Immediate intervention is the administration of aspirin.

Sold by @itachibot

Aspirin is the first drug given in acute coronary syndromes. An initial dose of 325 mg to be chewed and swallowed is given. If oral administration is not feasible, it should be given as a rectal suppository.

The following protocol must be followed in the management of patients with acute coronary syndrome:

• Stabilization of airway, breathing, and circulation

• Administration of O2

• Treatment of sustained ventricular arrhythmia rapidly according to ACLS protocols

• Give aspirin 325 mg

• History and examination, assess cardiac biomarkers

• Give three sublingual nitroglycerin tablets each 5 min apart. Add IV nitroglycerin for the persistence of symptoms.

• Treat left heart failure if present with an after-load reducing agent or loop diuretic

• Give beta-blockers

• Give morphine for the persistence of anxiety and discomfort related to MI

• In patients with STEMI - Reperfusion with PCI or fibrinolysis (if PCI is unavailable within 120 min) followed by antiplatelet and anticoagulant therapy. In patients with NSTEMI - antiplatelet and anticoagulant therapy are indicated.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 17:

The above ECG shows ST-segment elevation in leads II, III, and aVF suggestive of an inferior-wall MI. In the given scenario, the most likely reason for the patient to have shock is right ventricular infarction.

The diagnosis of right ventricular MI is suspected in patients with inferior-wall MI, clear lung fields (no added sounds on auscultation), and shock. It can also be confirmed by the use of right-sided electrocardiogram (ECG) leads.

In patients with presumed right ventricular shock, intravenous (IV) fluid replacement is indicated if jugular (central) venous pressure is not elevated.

Options A, B, and D: No added sounds on auscultation clearly rule out mitral regurgitation, septal defect (pansystolic murmur is heard), and decreased ejection fraction from the left ventricle (leads to pulmonary congestion and added sounds), as the cause of shock in this patient.

Solution to Question 18:

The echocardiogram shows pericardial effusion with cardiac tamponade (swinging heart sign). The likely diagnosis in the above scenario is myocardial rupture post-MI. It usually occurs 3-5 days postinfarction and commonly involves the anterior wall.

Sold by @itachibot

This condition usually results in immediate death. It is common in older women.

Incomplete or gradual rupture may be sealed off by the pericardium creating a pseudoaneurysm.

Immediate pericardiocentesis should be done to relieve the symptoms of shock due to tamponade. Early surgical repair is the definitive treatment.

Solution to Question 19:

Dressler syndrome commonly occurs 1-12 weeks after MI, not 2-7 days. However, it can recur over months.

Dressler syndrome is a post-myocardial infarction syndrome that presents as pleuritic chest pain, fever, and sometimes a pleuritic rub and effusion. The ECG findings include PR segment depression or ST-segment elevation. It is an autoimmune phenomenon occurring due to immune-complex deposition in the pericardium and lung pleura.

The first-line therapy consists of aspirin.

Note: NSAIDs and corticosteroids impair the infarct healing and predispose the patients to myocardial rupture. Therefore these drugs must be avoided in the early postinfarction period.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 20:

The above image showing left ventriculography with contrast is suggestive of a left ventricular aneurysm. Anterior transmural infarcts are involved in causing aneurysms as a post-MI complication.

Patients with anterior transmural infarcts are at the greatest risk for:

- Free-wall rupture
- Expansion
- Mural thrombi
- Aneurysm

Posterior transmural infarcts are more likely to be complicated by:

- Conduction blocks
- Right ventricular involvement, or both

Solution to Question 21:

The door-to-balloon time for a patient arriving in a hospital that is not equipped to perform a PCI and is then transported to a hospital where PCI can be performed is 120 min.

The door-to-balloon time is the time interval between the first medical contact and PCI.

Sold by @itachibot

It is 90 min or less for a patient arriving in a hospital equipped with PCI (ACC/AHA guidelines and MoHFW, India guidelines).

In patients who cannot receive PCI within 120 min of first medical contact, fibrinolysis is given. Fibrinolysis is only considered if the patient presents within 12 hours of symptom onset.

Note: The European society of cardiology (ESC) recommends 60 min target time for patients presenting in a PCI-centre and 90 min maximum target time for PCI-mediated reperfusion.

Solution to Question 22:

PCI related MI is defined as elevation of cTn > 5 times normal in the presence of symptoms of MI, ECG changes, angiographic findings or imaging abnormality.

The imaging abnormality could be a new loss of viable myocardium or new regional wall motion abnormality. If the baseline cTn values are elevated but stable or falling, a rise in cTn >20 % defined PCI related MI.

CABG related MI is defined as elevation of cTn > 10 times the normal baseline values in the presence of new pathologic Q waves, angiographic findings or imaging abnormality.

Solution to Question 23:

Based on the given clinical scenario of a patient presenting with sudden onset of chest pain, with ECG findings of ST depression in the anterior chest leads and T wave inversion, the most likely diagnosis is a non-ST-segment elevation-acute coronary syndrome (NSTE-ACS). The most appropriate line of management in this patient would be to administer aspirin with heparin.

Percutaneous coronary intervention is indicated after coronary angiography in high-risk patients.

Fibrinolytic therapy is not indicated in patients with NSTE-ACS.

Arrhythmia prophylaxis is generally not routinely recommended in all patients with NSTE-ACS.

Acute coronary syndrome is classified into 2 groups:

• ST-segment Elevation Myocardial Infarction (STEMI)

• NSTE-ACS

NSTE-ACS further includes:

• NSTEMI (myocyte necrosis present and cardiac troponins elevated)

• Unstable Angina (myocyte necrosis absent and cardiac troponins not elevated)

The diagnosis of NSTE-ACS is based on the following:

Severe chest discomfort with at least one of 3 features:

• Occurring at rest or minimal exertion and lasting for more than 10 minutes

• Recent onset

• Crescendo pattern

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

ECG showing:

• ST segment depression

• T wave inversion

Elevated cardiac troponins in case of NSTEMI.

Management of NSTE-ACS:

To provide relief of pain and discomfort, initial treatment includes bed rest, nitrates, beta blockers, and inhaled oxygen (only if arterial O2 saturation <90%).

Antithrombotic therapy consisting of antiplatelet and anticoagulant drugs:

• Dual antiplatelet therapy (DAPT): Aspirin (162-325 mg) AND one ADP receptor inhibitor (Ticagrelor, clopidogrel or prasugrel)

• Anticoagulation: Enoxaparin, unfractionated heparin, fondaparinux or bivalirudin

• An early invasive strategy including diagnostic coronary angiography (within 2–72 hours of presentation) followed by revascularization (percutaneous coronary intervention/ coronary artery bypass grafting) is indicated for stabilized patients who are at high risk of coronary events (Based on TIMI, GRACE or HEART score).

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Cardiomyopathy and Myocarditis

Question 1:

Which of the following is the most common protein mutation seen in hypertrophic cardiomyopathy?

a) Myosin heavy chain

b) Alpha-tropomyosin

c) Cardiac troponin C

d) Titin

Question 2:

A 19-year-old basketball player collapses during a match and regains consciousness several minutes later. He reveals that he has had similar episodes before. His older brother, a soccer player, had died suddenly at the age of 22 years during a soccer game due to a heart ailment. Which of the following is least likely to be found on echocardiography in this patient?

a) Dynamic left ventricular outflow tract obstruction

b) Systolic anterior motion of mitral valve

c) Mitral regurgitation with anteriorly directed regurgitant jet

d) Predominant diastolic dysfunction

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

While evaluating a young man for hypertrophic cardiomyopathy you hear a harsh and late peaking systolic ejection murmur at the lower left sternal border. On performing which of the following maneuvers, the intensity of this murmur is expected to decrease?

a) Valsalva

b) Squatting

c) Standing posture

d) Exercise

Question 4:

Sold by @itachibot

A 20-year-old patient has been diagnosed with hypertrophic cardiomyopathy. Which of the following is not a major risk factor for sudden death in this condition?

a) Spontaneous non sustained ventricular tachycardia

b) Spontaneous sustained ventricular tachycardia

c) Family history of sudden cardiac death

d) Atrial fibrillation

Question 5:

A 17-year-old boy presents with complaints of syncope. ECG is suggestive of LV hypertrophy. A doppler echocardiography revealed abnormal systolic anterior leaflet motion of the mitral valve. Which of the following drugs should be avoided in this patient?

a) Digoxin

b) Beta blockers

c) Verapamil

d) Disopyramide

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which is the most common sustained arrhythmia in hypertrophic cardiomyopathy?

a) Atrial fibrillation

b) Atrial flutter

c) Multifocal atrial tachycardia

d) Ventricular tachycardia

Question 7:

You are counseling the parents of a 15-year-old boy who was recently diagnosed with hypertrophic cardiomyopathy. What is the recommendation regarding his participation in sports?

a) Participation in competitive sports is not recommended

b) Only non-contact competitive sports are allowed

c) Non-contact competitive sports with beta-blocker therapy is allowed

d) High-intensity exercise is acceptable with verapamil therapy

Sold by @itachibot

Question 8:

A 65-year-old Japanese woman was brought to the emergency department with worsening angina and palpitations. Her physical examination was normal. ECG showed giant T wave inversions and coronary angiography was normal. Left ventriculography showed spade-like deformity. What is the most probable diagnosis?

a) Dilated cardiomyopathy

b) Restrictive cardiomyopathy

c) Hypertrophic cardiomyopathy

d) Apical hypertrophic cardiomyopathy

Question 9:

Two brothers were arguing over a property dispute when the elder of the two complained of chest pain and collapsed and was later declared brought dead by the hospital. His family says he was previously healthy and there was no similar disease in the family members. What is the likely diagnosis?

a) Infective cardiomyopathy

b) Takotsubo cardiomyopathy

c) Dilated cardiomyopathy

d) Hypertrophic cardiomyopathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

The observed myocardial stunning pattern does not match the ECG findings of a patient. What is the probable diagnosis?

a) Takotsubo cardiomyopathy

b) Restrictive cardiomyopathy

c) Brigade's cardiomyopathy

d) Pericardial tamponade

Question 11:

Sold by @itachibot

A 17-year-old Chinese boy presented with prolonged fatigue, shortness of breath, and swelling in the legs. His echocardiography was suggestive of dilated cardiomyopathy. Which of the following nutritional deficiencies can explain his condition?

a) Molybdenum

b) Manganese

c) Zinc

d) Selenium

Question 12:

Endomyocardial biopsy of a patient shows vacuolar degeneration and myofibrillar loss. Which of the following drugs may be responsible for this finding?

a) Alcohol

b) Doxorubicin

c) Cyclophosphamide

d) Trastuzumab

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

In which of the following patients, the risk of developing peripartum cardiomyopathy is less?

a) A 26-year-old pregnant woman with an obstetric score of G2P1

b) A 39-year-old pregnant woman

c) A 22-year-old pregnant woman with pre-eclampsia

d) A primigravida with twin pregnancy

Question 14:

A young male patient presents with progressive dyspnea and weakness. He gives a history of fever, myalgia, and sore throat one week ago. To further evaluate him as per Dallas criteria, which of the following tests need to be performed?

a) ECG

b) Echocardiography

c) Endomyocardial biopsy

d) CT-angiography

Sold by @itachibot

Question 15:

Which is the most common infective cause of myocarditis?

a) Borrelia burgdorferi

b) Trypanosoma cruzi

c) Trichinella

d) Toxoplasma gondii

Question 16:

A 13-year-old boy collapsed while playing football. ECG showed V5, V6 inversion and prominent inverted T-waves. 2D echocardiography showed systolic anterior motion of the mitral valve leaflet. Which of the following features will be seen on clinical examination and which of the following are true regarding the management?

a) 1 & 3, a, c, d

b) 2 & 4, b, c, e

c) 1 & 3, a, d

d) 2 & 4, b, e

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	d
5	a
6	a
7	a
8	d
9	b
10	a

Sold by @itachibot

11	d
12	b
13	a
14	c
15	b
16	a

Detailed Explanations

Solution to Question 1:

The most common mutations in HCM are those affecting the myosin heavy chain.

Hypertrophic cardiomyopathy (HCM) is an autosomal dominant disease. In the majority of cases it is due to missense mutations in the following two sarcolemmal protein genes:

• Myosin binding protein C (MYBC)

• Beta myosin heavy chain (MYH7)

Option B: Alpha-tropomyosin mutation is seen in HCM but is not common.

Options C and D: Mutations in cardiac troponin C and titin are associated with dilated cardiomyopathy (DCM).

Solution to Question 2:

The given clinical scenario of exercise intolerance (syncope) in the patient and family history of sudden cardiac death in his elder brother at a young age is suggestive of hypertrophic cardiomyopathy (HCM). Mitral regurgitation with posteriorly directed regurgitant jet is usually seen on echocardiogram in HCM. ECG and echocardiogram are used to confirm the diagnosis of HCM.

HCM is characterized by myofiber disarray in the heart, resulting in a hypertrophic myocardium. The hypertrophy leads to the following pathophysiological changes in HCM:

• Diastolic dysfunction - presenting as exercise intolerance

• Systolic anterior motion (SAM) of the mitral valve

• The hypertrophied muscles result in a narrow outflow tract

• This leads to the production of a high-velocity jet pulling the mitral leaflets towards the septum posteriorly

• SAM leads to mitral regurgitation

• Dynamic left ventricular outflow tract (LVOT) obstruction

Echocardiography features of HCM are as follows:

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

• Left ventricular wall thickness >15 mm

• Hypertrophied heart more confined to the interventricular septum resulting in banana-shaped ventricular cavity

Solution to Question 3:

Squatting increases the preload and afterload thereby decreasing the outflow obstruction and intensity of murmur in HCM.

In HCM left ventricular size is small, thus the approximation of the interventricular septum with the mitral valve increases. The following mechanisms further increase this approximation and increase blood outflow obstruction and intensity of murmur:

• A decrease in left ventricular cavity volume or pressure (decreased preload)

• A decrease in systemic vascular resistance and/or arterial pressure (decreased afterload)

• Increase in left ventricular contractility

The image below shows a cross-section of a heart with hypertrophic cardiomyopathy.



Note: All murmurs increase in intensity on squatting except that of HOCM and MVP.

Parameters	Murmur intensity increased	Murmur intensity decreased
Preload	ValsalvaStanding(↓preload = ↑obstruction)	SquattingPassive leg raising (↑preload = ↓obstruction)
Force of contraction	Exercise(↑force = ↑obstruction)	
Afterload	Vasodilators(↓afterload = ↑obstruction)	Isotonic handgripSquatting(↑ afterload = ↓obstruction)

Sold by @itachibot

Solution to Question 4:

Atrial fibrillation in HCM is not associated with increased HCM-related mortality. It frequently accounts for unexpected hospital admissions, lost productivity, and impaired quality of life.

Among HCM patients, the greatest risk for sudden cardiac death is seen in patients who have sustained ventricular tachycardia (VT). These patients must undergo placement of an implantable cardioverter-defibrillator (ICD) for secondary prevention.

Solution to Question 5:

The given clinical scenario is suggestive of hypertrophic cardiomyopathy. Digoxin should be avoided as it may increase contractility and worsen obstruction.

Drugs that can lead to possible aggravation of outflow tract obstruction and thus contraindicated in HCM are:

- Digoxin
- Inotropes
- Nitrates
- ACE inhibitors
- Dihydropyridine calcium channel blockers (eg- nifedipine, amlodipine)

Beta-blockers and verapamil are first-line agents in the treatment of HCM. Disopyramide and amiodarone are the preferred antiarrhythmic agents.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

Symptomatic atrial fibrillation (AF) is the most common sustained arrhythmia in hypertrophic cardiomyopathy (HCM). It frequently accounts for unexpected hospital admissions, lost productivity, and impaired quality of life.

Treatment of atrial fibrillation in HCM:

- Amiodarone - drug of choice
- Disopyramide
- Radiofrequency ablation for medically refractory cases
- Anticoagulation to prevent embolic events

Note: A sustained arrhythmia is one that lasts for >30 seconds.

Sold by @itachibot

Solution to Question 7:

Participation in competitive sports is not recommended for patients with hypertrophic cardiomyopathy.

The following general measures are advised for patients with HCM:

• Patients are advised against engaging in competitive athletics or vigorous physical activities

• Maintaining low to moderate levels of exercise as a part of a healthy lifestyle has been proven to be beneficial

• Adequate hydration at all times

Note: Non-contact sports are those sports wherein the participants are physically separated with almost no physical contact (ex: tennis, badminton) between them. Contact sports involve some form of physical contact between the participants (ex: wrestling, boxing)

Solution to Question 8:

The presence of giant T wave inversions and spade-like deformity on left ventriculography points to a diagnosis of apical hypertrophic cardiomyopathy.

Apical hypertrophic cardiomyopathy is also called Yamaguchi syndrome. It is characterized by hypertrophy of the myocardium, predominantly in the left ventricular apex. It was first described in the Japanese population and is uncommon in other populations.

The symptoms are milder and the risk of sudden cardiac death is lower than in the traditional septal form of HCM.

The image below shows the ace of spades appearance on left ventriculography:



The image below shows the ace of spades appearance on echocardiography with massive hypertrophy of the apex and relative sparing of the base:

If yc... Sold by @itachibot ...purchased this from someone else... ...may have been scammed.

Sold by @itachibot

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Solution to Question 9:

The given clinical scenario of sudden heart failure in a previously healthy person under a stressful situation is suggestive of Takotsubo cardiomyopathy.

Takotsubo cardiomyopathy is also called broken heart syndrome, apical ballooning syndrome, or stress-induced cardiomyopathy. It is a transient myocardial stunning from microcirculation spams due to a catecholamine surge during stress. It is usually seen in middle-aged women. The patients can present with pulmonary edema, hypotension, and chest pain.

ECHO reveals apical dilatation with basal hypercontraction of the heart, resembling a Takotsubo (pot for catching octopus). The ECG changes can mimic myocardial infarction but changes extend beyond a particular coronary arterial territory. Coronary angiography, therefore, may be needed to rule out coronary occlusion. (Option B)

It is a self-limiting disorder resolving within days to weeks. The prognosis is generally good.

The image below shows left ventriculography during systole showing apical ballooning resembling a Takotsubo pot.

Sold by @itachibot



If you purchased this from some... else, you may have been scammed. Sold by @itachi...

Other options:

Option A: Infective cardiomyopathy: The most common causes of infective myocarditis are viral infections and the protozoan Trypanosoma cruzi, which is associated with Chagas' disease. Acute viral myocarditis often presents with symptoms and signs of heart failure, although it can also manifest as chest pain and show electrocardiogram (ECG) changes resembling pericarditis or acute myocardial infarction. In Chagas' disease, specific features include abnormalities in the conduction system, particularly dysfunction of the sinus node and AV node, as well as the right bundle branch block.

Option C: Dilated cardiomyopathy: It is the most common cardiomyopathy and is predominantly characterised by systolic dysfunction. Though it may present acutely, it progresses chronically. The familial form is due to mutations in TTN gene which encode the sarcomeric protein titin. Other etiologocal agents include alcohol, infective myocarditis, doxorubicin, hemochromatosis and sarcoidosis.

Option D: Hypertrophic cardiomyopathy: Hypertrophic cardiomyopathy (HCM) is the most common genetic cardiovascular disorder and is a prominent cause of sudden death before the age of 35 years. Family history is commonly present, a sarcomere mutation is present in ~50% of patients. Sudden death can be due to polymorphic ventricular tachycardia/fibrillation. Risk factors for sudden death in this disease include young age, nonsustained ventricular tachycardia, failure of blood pressure to increase during exercise, recent (within 6 months) syncope, ventricular wall thickness >3 cm, and possibly the severity of LV outflow obstruction.

Solution to Question 10:

Takotsubo cardiomyopathy is the probable diagnosis when the observed myocardial stunning pattern does not match ECG changes.

Myocardial stunning is a state where certain segments of the myocardium (usually corresponding to an area of major coronary occlusion) show forms of contractile abnormality (dysfunction) on echocardiography.

Sold by @itachibot

In Takotsubo cardiomyopathy, while the ECG changes mimic a myocardial infarction(MI), the left ventricular dysfunction extends well beyond the distribution of a specific coronary artery. Thus, the myocardial stunning pattern might not fully match the ECG. Ventriculography shows apical dilation with basal contraction.

Takotsubo cardiomyopathy occurs typically in older women and is associated with a sudden increase in emotional or physical stress, presenting with hypotension, chest pain with ECG changes of acute MI, and pulmonary edema. Its also known as apical ballooning syndrome or acute stress-induced cardiomyopathy.

Option B: Restrictive cardiomyopathy is characterized by diastolic dysfunction with reduced contractility and ejection fraction, most commonly seen in amyloidosis, and enlarged ventricles on imaging with low-voltage ECG complexes.

Option C: Arrhythmogenic right ventricular cardiomyopathy (ARVC) is a type of familial dilated cardiomyopathy (DCM), having defects in desmosomal proteins, affecting the ventricles showing thin ventricular walls on Echo, epsilon waves, and low voltage ECG complexes.

Option D: Pericardial tamponade is characterized by hypotension, muffled heart sounds and increased jugular venous pressure with muffled heart sounds with total electrical alternans and sinus tachycardia in ECG.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

The patient is probably suffering from Keshan's disease, which is due to selenium deficiency. It is endemic dilated cardiomyopathy that affects children and women of childbearing age in some areas in China.

The nutritional deficiencies that can lead to cardiomyopathy are:

- Thiamine
- Selenium
- Carnitine
- Calcium
- Phosphate
- Magnesium

Solution to Question 12:

Endomyocardial biopsy showing vacuolar degeneration and myofibrillar loss is seen in cardiotoxicity due to anthracyclines (doxorubicin).

Difference between cardiotoxicity due to anthracyclins and trastuzumab:

Anthracyclins	Trastuzumab
Irreversible	Reversible

Sold by @itachibot

Anthracyclins	Trastuzumab
Dose-dependent	Not dose-dependent
Biopsy shows myocyte destruction	Biopsy does not show myocyte destruction

Solution to Question 13:

Low parity is not a risk factor for peripartum cardiomyopathy. Increased parity is a risk factor.

Peripartum cardiomyopathy (PPCM) is a type of dilated cardiomyopathy. It develops during the last month of pregnancy or within the first 5 months following delivery. The risk factors are:

- Increased maternal age
- Increased parity
- Twin pregnancy
- Malnutrition
- Use of tocolytic therapy for premature labor
- Preeclampsia or toxemia of pregnancy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

The above clinical scenario is suggestive of infective myocarditis. The Dallas criteria used for the diagnosis of myocarditis is based on endomyocardial biopsy. It is used to define the disease.

Active myocarditis is defined as an inflammatory infiltrate of the myocardium with necrosis and/or degeneration of adjacent myocytes not typical of the ischemic damage associated with coronary artery disease.

Borderline myocarditis is the term used when the inflammatory infiltrate is too sparse or myocyte injury is not demonstrated.

Solution to Question 15:

The most common infective cause of myocarditis is Trypanosoma cruzi causing Chagas disease.

Infective myocarditis is most commonly associated with viruses (adenovirus, coxsackie B virus, HHV-6) and the protozoan Trypanosoma cruzi.

The most common noninfective cause of myocarditis is sarcoidosis.

Solution to Question 16:

Sold by @itachibot

The given clinical scenario of a young patient with a history of sudden collapse following physical activity along with ECG findings of V5, V6 inversion, and prominent inverted T-waves, and 2D echocardiography showing the systolic anterior motion of the mitral valve leaflet point toward a diagnosis of hypertrophic obstructive cardiomyopathy (HCM).

In HCM, clinical examination may reveal pulsus bisferiens and an ejection systolic murmur that increases on standing.

Beta-blockers and implantable cardioverter-defibrillators (ICD) are used in the management.

HCM is a leading cause of sudden death in young. It is inherited in an autosomal dominant pattern with variable penetrance. HCM is characterized by the thickening of the septal myocardium below the aortic valve leading to obstruction of the left ventricular outflow tract.

Symptoms:

• Exertional dyspnea

• Syncope

• Angina, palpitations, cardiac arrhythmias

• Sudden cardiac death (particularly during or after intense physical activity)

Examination:

• Systolic ejection murmur best heard at the apex or lower left sternal border

• Holosystolic murmur from mitral regurgitation

• Sustained apex beat

• S4 gallop

• Paradoxical split of S2

• Pulsus bisferiens: LV outflow obstruction causes a sudden quick rise of the pulse followed by a slower longer rise.

ECG findings:

• Left axis deviation

• P wave abnormalities

• Prominent abnormal Q waves

• Deeply inverted T waves

Echocardiography findings:

• LV wall thickness ≥15 mm

• Systolic anterior motion of the mitral valve leaflet

• Mitral regurgitation

Treatment:

• Beta-blockers and calcium channel blockers are the first-line treatment

• Implantable cardioverter-defibrillator (ICD) - indicated when the history of cardiac arrest or sustained ventricular tachycardia is present

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Parameters	Murmur intensity increased	Murmur intensity decreased
Preload	ValsalvaStanding(↓preload = ↑obstruction)	SquattingPassive leg raising (↑preload = ↓obstruction)
Force of contraction	Exercise(↑force = ↑obstruction)	
Afterload	Vasodilators(↓afterload = ↑obstruction)	Isotonic handgripSquatting(↑ afterload = ↓obstruction)

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Heart Failure

Question 1:

A 48-year-old man presents with tiredness and palpitations. He also complains of abdominal pain, increased frequency of urination at night, and unexplained weight gain. On examination, the pulse is irregular at 118 beats/min. Bilateral pitting pedal edema is present. ECG is suggestive of right axis deviation. Which of the following feature is unlikely to be present in this patient?

a) Pulsatile liver

b) Raised right atrial pressure

c) Positive hepatojugular reflux

d) Increased PCWP

Question 2:

A patient who is being evaluated for heart failure undergoes echocardiography for assessment of ejection fraction (EF). When will you categorize his condition as heart failure with reduced ejection fraction based on the results?

a) EF < 30%

b) EF < 40%

c) EF < 50%

d) EF < 60%

Question 3:

A 65-year-old male patient came with complaints of episodes of shortness of breath and coughing at night which awakens him from sleep. On examination, distended neck veins were observed and S3 was heard on auscultation. Which of the following biomarkers can be used for the diagnosis in this patient?

a) Brain natriuretic peptide (BNP)

b) Atrial natriuretic peptide (ANP)

c) Endothelin-1 (ET-1)

d) Lipocalin

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 4:

Falsely low values of BNP can be seen in which of the following?

a) Advanced age

b) Female patients

c) Obesity

d) Renal impairment

Question 5:

Which of the following is a newer biomarker to assess the prognosis in a patient with heart failure?

a) Soluble ST-2

b) Lipocalin

c) Endothelin-1 (ET-1)

d) Cystatin C

Question 6:

A patient is diagnosed with heart failure with an ejection fraction < 40%. What is the treatment of choice in this patient after the euvolemic status has been achieved?

a) ARBs + Diuretics

b) ACE inhibitors + Beta-blockers

c) CCBs + ARBs

d) Diuretics + CCBs

Question 7:

The preferred drug for a patient with chronic symptomatic heart failure with reduced ejection fraction is:

a) Bosentan

b) Sacubitril/Valsartan

c) Omapatrilat

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) Moxonidine

Question 8:

Which of the following drugs is a selective myosin activator?

a) Istaroxime

b) Omecamtiv mecarbil

c) Carperitide

d) Milrinone

Question 9:

Which of the following drugs used in heart failure is an inhibitor of I(f) current in the sinoatrial node ?

a) Omapatrilat

b) Ivabradine

c) Nesiritide

d) Levosimendan

Question 10:

In a patient with heart failure, which of the following is not used as an inotropic agent?

a) Levosimendan

b) Milrinone

c) Omecamtiv Mecarbil

d) Nesiritide

Question 11:

Which of the following beta-blockers would you prefer to use in a patient with heart failure with reduced ejection fraction?

a) Atenolol

b) Nebivolol

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Esmolol

d) Bisoprolol

Question 12:

Which of the following drugs does not reduce mortality in a patient with heart failure with reduced ejection fraction?

a) Digoxin

b) Beta-blockers

c) ACE inhibitors

d) Hydralazine + Isosorbide dinitrate

Question 13:

A 45-year-old female patient has NYHA class III heart failure with dyspnea, pedal edema and K+ levels of 5.5 mg% and creatinine 1.1 mg%. Which of the following drugs can be used in this patient?

a) Digoxin

b) Spironolactone

c) Enalapril

d) Furosemide

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

A patient was brought to the emergency with dyspnea, cough, and chest pain. His blood pressure was 180/100 mmHg and SpO2 was 90%. Chest X-ray is shown below. Echocardiography revealed reduced LV ejection fraction (LVEF). Which of the following will you not use in the management of this patient?

Sold by @itachibot



a) Digoxin

b) Morphine

c) Furosemide

d) Positive pressure ventilation

Question 15:

A patient presents with abrupt worsening of renal function leading to acute cardiac disorder. Which type of cardiorenal syndrome would this be?

a) Type I

b) Type II

c) Type III

d) Type V

Question 16:

Who comes under category III non-heart-beating donor?

a) A patient who died during transportation to the hospital

b) A patient who died after failed resuscitation after reaching the hospital

c) A patient who was brought dead to the hospital

d) A patient who is awaiting death in the hospital

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 17:

A 70-year-old man with a known history of DM and HTN on regular medications presents with a history of progressively worsening dyspnoea on exertion. On examination, there is elevated JVP, positive hepatojugular reflux, crackles at the lung bases, hepatomegaly, ascites, and pedal oedema■. What is your provisional diagnosis?

a) Heart failure

b) Hypertrophic cardiomyopathy

c) Mitral regurgitation

d) Portal hypertension

Question 18:

Which of the following is not a radiological finding in a patient with left heart failure?

a) Kerley b lines

b) Focal oligemia

c) Increased venous blood in lung

d) Change in upper lobe circulation

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	b
3	a
4	c
5	a
6	b
7	b
8	b
9	b
10	d
11	d
12	a

Sold by @itachibot

13	d
14	a
15	c
16	d
17	a
18	b

Detailed Explanations

Solution to Question 1:

The above scenario is suggestive of right-sided heart failure. Increased PCWP is not a sign of right-sided heart failure. PCWP is an indirect measurement of the left atrial pressure. Hence, it is a sign of left-sided heart failure.

The signs of heart failure are:

Right heart failure	Left heart failure
Due tocongestionin thesystemic circulation	Due tocongestionin thelungs.
Symptoms and signs:Raised JVPCongestive hepatomegaly Positive hepatojugular reflux AscitesBilateral pitting edemaWeight gain- due to edemaFatigue- due to ↓pumping of blood to lungs leading to ↓oxygenation of the blood	Symptoms and signs:Dyspnea on exertionOrthopneaParoxysmal nocturnal dyspnea (characteristic of LVF)Pulmonary ralesIncreased PCWPFatigue- due to ↓cardiac output

If you bought this from someone else, it was stolen by @itachibot

Solution to Question 2:

Heart failure with reduced ejection fraction is ejection fraction <40%.

According to the European society of cardiology (ESC) heart failure guidelines 2016 -

• Heart failure with reduced ejection fraction (HFrEF)- ejection fraction <40%

• Heart failure with mid-range ejection Fraction (HFmrEF)- ejection fraction 40-50%

• Heart failure with preserved ejection fraction (HFpEF)- ejection fraction >50%

Solution to Question 3:

Sold by @itachibot

The given clinical scenario suggests a diagnosis of heart failure. The brain natriuretic peptide (BNP) can be used to confirm this.

Natriuretic peptides are protein molecules that are secreted by the ventricular myocytes in response to volume or pressure overload. Ventricular musculature releases pre-proBNP which is cleaved into active BNP and inert NT-proBNP.

BNP causes natriuresis, vasodilatation, and diuresis that leads to improved myocardial relaxation. These peptides also counteract the actions of the renin-angiotensin-aldosterone system. Measurement of both BNP and inert NT-proBNP levels can be used in the diagnosis of heart failure and to determine the prognosis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

Falsely low values of BNP can be seen in obese patients.

It is also falsely low in:

• Flash pulmonary edema

• Heart failure due to- mitral regurgitation, mitral stenosis, constrictive pericarditis, restrictive cardiomyopathy

It falsely increases in:

• Advanced age

• Renal impairment

• Female patients

Solution to Question 5:

Soluble ST-2 is a newer biomarker that can be used for determining the prognosis of patients with heart failure. Other newer biomarkers used are galectin-3, GDF 15.

Options B and D: Lipocalin, also known as neutrophil gelatinase-associated lipocalin (NGAL) or siderocalin, cystatin C, and IL-18 are biomarkers of acute kidney injury (AKI).

Option C: Endothelin-1 is a useful biomarker for early detection of bronchiolitis obliterans in lung transplant recipients.

Solution to Question 6:

ACE inhibitors + beta-blockers are the treatment of choice in patients with heart failure with an ejection fraction of <40% after euvolemic status is achieved using diuretics.

The first-line drugs used in congestive heart failure with reduced ejection fraction are ACE inhibitors + diuretics. Once the euvolemic status is achieved, ACE inhibitors + beta-blockers are the drugs of choice.

Sold by @itachibot

Beta-blockers that improve survival in heart failure with reduced ejection fraction are carvedilol, metoprolol, and bisoprolol.

Solution to Question 7:

Sacubitril/Valsartan, an angiotensin receptor neprilysin inhibitor (ARNI), is used in patients with chronic symptomatic HFrEF to reduce morbidity and mortality (class I recommendation).

Sacubitril (neprilysin inhibitor) is approved in combination with valsartan (ARB), in a ratio of 1:1. The combination is approved for use in chronic heart failure.

Sacubitril is a prodrug that is activated to sacubitrilat. It inhibits the enzyme neprilysin (neutral endopeptidase), which is responsible for the degradation of atrial and brain natriuretic peptides (reduction of blood volume). Thus, it increases the levels of atrial and brain natriuretic peptides and lowers blood pressure.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

Omecamtiv mecarbil is the selective myosin activator. It activates cardiac myosin and prolongs systole without increasing oxygen consumption of the heart.

Solution to Question 9:

Ivabradine is an inhibitor of If current in the sinoatrial node. It reduces the heart rate without decreasing the force of contraction.

It is used in patients with chronic HFrEF with a sinus rhythm and a resting heart rate ≥70 bpm who are either on a maximum tolerated beta-blocker dose or have a contraindication for beta-blockers. The concurrent treatment should include ACE inhibitor (or ARB) and aldosterone receptor antagonist.

Option A: Omapatrilat is a dual neprilysin inhibitor+ ACE inhibitor.

Option C: Nesiritide belongs to the intravenous vasodilators class and is a recombinant human brain natriuretic peptide (BNP).

Option D: Levosimendan belongs to the positive inotropes class and is a combined calcium sensitizer and PDE3 inhibitor.

Solution to Question 10:

Nesiritide is not used as an inotropic agent. It belongs to the intravenous vasodilator class and is a recombinant brain natriuretic peptide (BNP)

Sold by @itachibot

Option A: Levosimendan belongs to the positive inotropes class and is a combined calcium sensitizer and PDE3 inhibitor.

Option B: Milrinone belongs to the positive inotrope class and is a PDE-3/4 inhibitor.

Option C: Omecamtiv mecarbil is a selective myosin activator that increases myocardial contractility without increasing the myocardial oxygen demand.

Solution to Question 11:

Bisoprolol is preferred for use in patients having heart failure with reduced ejection fraction (HFrEF).

The only beta-blockers that have been proved to improve the survival of patients with HFrEF or chronic HF are -

- Bisoprolol
- Metoprolol
- Carvedilol

Note: Nebivolol is not approved by FDA for the treatment of heart failure.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

Digoxin does not reduce mortality in patients with HFrEF.

The drugs which reduce mortality in patients with HFrEF are :

- ACE inhibitors
- Angiotensin receptor blockers (ARBs)
- Angiotensin receptor neprilysin inhibitor (ARNI)
- Beta-blockers
- Aldosterone antagonists
- Spironolactone
- Eplerenone
- Hydralazine+Nitrates

Solution to Question 13:

The clinical scenario describes a patient with symptomatic heart failure and K+ levels of 5.5 mg%. She requires a diuretic for symptom control. Furosemide can be given safely as it causes hypokalemia.

Sold by @itachibot

Other options:

Option A: Digoxin is not used as a first-line drug in the management of symptomatic heart failure.

Options B and C: Spironolactone (K+ sparing diuretic) and enalapril (ACE inhibitor) can cause hyperkalemia.

The first step in managing acute decompensated heart failure (ADHF) is addressing the factors that caused the decompensation. They include nonadherence to medications, dietary indiscretion, medication use that exacerbates heart failure, coronary ischemia, arrhythmias, valvular heart disease, infection, and pulmonary thromboembolism.

Volume management is crucial in ADHF. Intravenous loop diuretics are used to relieve congestion. The effectiveness of diuretic therapy can be assessed by changes in weight, improvement in symptoms, and clinical examination findings. Natriuretic peptide levels can also help assess the adequacy of therapy and predict risk.

Cardiorenal syndrome, characterized by the interplay between heart and kidney dysfunction, is a common complication of ADHF. Diuretic therapy should be carefully managed to avoid worsening renal function in patients with elevated right-sided filling pressures. In cases of severe low cardiac output, inotropic therapy or mechanical circulatory support may be necessary to preserve or improve renal function temporarily.

Vasodilators like nitroglycerin, sodium nitroprusside, and nesiritide (recombinant BNP) are used to reduce filling pressures and systemic vascular resistance.

Inotropic therapy with drugs like dopamine and milrinone is used to increase myocardial contractility and improve cardiac output in patients with low-output ADHF. Dopamine has dose-dependent effects on different receptors, while milrinone is a phosphodiesterase-3 inhibitor.

Sold by @itachibot
If you purchased this from someone else
You may have been scammed.

Solution to Question 14:

The above X-ray showing batwing appearance along with the given symptoms and reduced LVEF is suggestive of acute decompensated heart failure with pulmonary edema. Digoxin has no role in the management of this condition.

The management of acute cardiogenic pulmonary edema consists of:

• Oxygen therapy and positive pressure ventilation

• Reduction of preload:

• Diuretics: Furosemide is the diuretic of choice

• Nitrates: sublingual nitroglycerin is the first-line therapy. If pulmonary edema persists, IV nitroglycerin is administered.

• Morphine: It is a transient venodilator that reduces preload, thereby reducing dyspnea and anxiety.

• ACE inhibitors: They reduce both preload and afterload and are recommended for hypertensive patients.

• Nesiritide: It is an IV recombinant BNP which is a potent vasodilator with diuretic properties.

Sold by @itachibot

• Inotropic and inodilators:

• Dopamine, dobutamine-inotropic agents.

• Milrinone- stimulates myocardial contractility and promotes peripheral vasodilatation.

• Intra-aortic balloon counterpulsation- or LV assist devices are used in cardiogenic shock and end-organ failure.

Solution to Question 15:

Type III cardiorenal syndrome is an abrupt worsening of renal function leading to acute cardiac disorder.

Cardiorenal Syndrome (CRS): A pathophysiologic disorder of the heart and kidneys whereby acute or chronic dysfunction in one organ may induce acute or chronic dysfunction in the other organ.

Types of Cardiorenal Syndromes

Type	Precipitating event	Resulting in
CRS Type I(Acute Cardiorenal Syndrome)	Abrupt worsening ofcardiac function:Acute cardiogenicshockDecompensatedcongestive heart failure	Acute kidney injury
CRS Type II(Chronic Cardiorenal Syndrome)	Chronic abnormalities incardiac function:Chronic congestive heart failure	Chronic kidney disease
CRS Type III(Acute Renocardiac Syndrome)	Acute renal dysfunction:Renal ischemia orglomerulonephritis	Acute cardiac disorder
CRS Type IV(Chronic Renocardiac Syndrome)	Chronic kidney disease:Glomerular disease	Decreased cardiacfunctionCardiac hypertrophyHigh risk of adversecardiovascular events
CRS Type V(Secondary Cardiorenal Syndrome)	Systemic conditions:Diabetes Sepsis	Cardiac and renal dysfunction

Solution to Question 16:

Category III donors are patients with non-survivable injuries who have their treatment withdrawn and if such patients wished in life to be organ donors, the transplant team can attend at the time of treatment withdrawal and retrieve organs after cardiac arrest has occurred.

Modified Maastricht classification- Categories of non-heart-beating donors:

Sold by @itachibot

Category	Description	Circumstances
Category I	Includesvictims of accident and suicidewho arefound dead at the sceneand resuscitation is deemed pointless (e.g. fatal cervical spine fracture).	Uncontrolled - transplant team not present or prepared
Category II	victims of sudden cardiac or cerebral catastrophewho are brought to emergency departments while being resuscitated by ambulance personnel or whodied in the department. Other sources include patients suffering an isolated brain injury, anoxia, and stroke, and victims of major trauma who died soon after hospital admission.	Uncontrolled - transplant team not present or prepared
Category III	Patients who are dying, often in an intensive care unit. These are the patients awaiting cardiac arrest where the treating clinicians have decided to withdraw treatment and not commence resuscitation for various reasons.	Controlled - transplant team present and prepared
Category IV	Comprises patients who sufferunexpected cardiac arrestduring or after the determination ofbrain death.	Controlled - transplant team present and prepared
Category V	Unexpected cardiac arrestdur ingintensive care-	Uncontrolled - transplant team not present or prepared

Solution to Question 17:

The given symptoms and signs are suggestive of heart failure (Option A).

Portal hypertension (Option D) can be a result of right-sided heart failure, however, the presence of worsening dyspnea along with the other symptoms implies a primary cardiac condition.

Heart failure is a common endpoint for most chronic cardiac conditions such as hypertension, valvular heart disease, and coronary heart disease.

The symptoms of heart failure are commonly due to volume overload with pulmonary and/or systemic venous pressures.

RIGHT HEART FAILURE	LEFT HEART FAILURE
Symptoms occur due tosystemic congestion	Symptoms occur due topulmonary congestion

Sold by @itachibot

RIGHT HEART FAILURE	LEFT HEART FAILURE
Raised pulsatile JVPHepatic congestionTender hepatomegalyPitting pedal edemaAscitesEarly satiety	Shortness of breathOrthopnea (dyspnea that occurs within 1-2 mins of lying down)Paroxysmal nocturnal dyspnea(shortness of breath that develops after lying down for 30 mins or longer)Pulmonary rales

Solution to Question 18:

Focal oligemia is not a radiological feature of left heart failure. It is seen in pulmonary embolism and is also called Westermark's sign.



It is defined by a hyperlucent area with decreased vascularity due to oligemia of the involved part of the lung. It is caused by a cutoff of circulation due to an embolus, leading to the loss of visibility of a patch of pulmonary vasculature on an X- ray.

Radiological signs of left heart failure include :

Sold by @itachibot



• Cardiomegaly - enlargement of the cardiac silhouette (increased cardiothoracic ratio)

• Upper lobe pulmonary venous congestion (cephalisation of vessels)

• Interstitial edema.

• Kerley B lines - horizontal lines in the periphery of the lower posterior lung fields as shown below.



Other radiological signs of Pulmonary embolism :

• Hampton's hump - a peripheral wedged-shaped density usually located at the pleural base

• Palla's sign - an enlarged right descending pulmonary artery

Sold by @itachibot

Sold by @itachibot

Pericardial Diseases

Question 1:

What is the defined duration for the diagnosis of acute pericarditis?

a) <2 weeks

b) <4 weeks

c) <6 weeks

d) <1 week

Question 2:

A 30-year-old man presented with low-grade fever and severe retrosternal chest pain for 1 week. The pain got exaggerated on lying down and relieved on sitting up. On auscultation of the precordium, a grating sound was audible. Which of the given statements is false regarding this auscultatory finding?

a) Pathognomonic of acute pericarditis

b) A very high pitched sound

c) Best heard with the patient in left lateral position

d) Loudest at the lower left sternal border

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

A young man presents to the medicine OPD with dyspnea, cough, and severe chest pain for 1 week. Chest pain is pleuritic in nature. On examination, the apical impulse is not palpable and breath sounds are normal. Which of the following is likely to be seen in the ECG of this patient?

a) ST-segment elevations with a convex upward configuration

b) PR-segment depressions early in the disease progression

c) Reciprocal ST-segment depressions in all corresponding leads

d) T-wave inversions developing concurrently with ST-segment elevations

Question 4:

Sold by @itachibot

Auscultation of a patient with pleuritic chest pain revealed a patch of dullness with increased vocal fremitus beneath the angle of the left scapula. What is this sign called?

a) Auenbrugger's sign

b) Moschcowitz's sign

c) Dressler's sign

d) Ewart's sign

Question 5:

Which of the following is not a component of Beck's triad?

a) Pulsus paradoxus

b) Hypotension

c) Muffled heart sounds

d) Raised JVP

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which of the following is a differentiating feature between cardiac tamponade and tension pneumothorax?

a) Breath sounds

b) Muffled heart sounds

c) Increased heart rate

d) Raised JVP

Question 7:

An intern is asked to measure the blood pressure of a patient with cardiac tamponade. He/she should ask the patient to:

a) Hold breath

b) Take rapid shallow breaths

c) Take long deep breaths

d) Breathe normally

Sold by @itachibot

Question 8:

An RTA victim is brought with breathing difficulty. On examination, the following is noted. Which of the following JVP findings is most likely to be seen in this patient?

a) Absent x and prominent y descent

b) Prominent x and absent y descent

c) Prominent x and y descents

d) Absent x and y descents

Question 9:

A patient with dizziness and breathing difficulty came to the emergency department. The ECG is given below. What is the most probable diagnosis?

a) Acute pericarditis

b) Constrictive pericarditis

c) Cardiac tamponade

d) Restrictive cardiomyopathy

Question 10:

A 25-year-old road traffic accident victim presents with chest pain and breathlessness. On examination, his BP was 80/60mmHg.On inspection, his neck veins were distended and on auscultation, his heart sounds were muffled. Which of the following clinical features is likely

Sold by @itachibot

to be seen in this patient?

a) Pericardial knock

b) Kussmaul's sign

c) Third heart sound

d) Pulsus paradoxus

Question 11:

A young motorist sustained injuries in a road traffic accident. He was found to have fractures of multiple ribs anteriorly on both sides. Radial and femoral fractures were also present. On examination, the following was noted. What should be the immediate intervention?

a) Rapid blood transfusion

b) Urgent pericardial tap

c) Intercostal tube drainage on both sides

d) Immediate repair of femoral artery

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

You're an intern examining a patient in the ICU. While assessing the JVP, you notice a lack of fall during inspiration. Which of the following conditions would you not consider as a differential diagnosis?

a) Restrictive cardiomyopathy

b) Cardiac tamponade

c) Constrictive pericarditis

d) Right ventricular infarct

Question 13:

A 20-year-old man presents to the emergency department with respiratory distress and hypotension following a trauma. He has subcutaneous emphysema and there is an absence of air entry into the right side of the lungs. What is the next best step?

a) Start IV fluids with large bore cannula

b) Needle decompression into the 5th intercostal space anterior to the midaxillary line

c) Take the patient to the ICU and intubate him

Sold by @itachibot

d) Start positive pressure ventilation

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	d
5	a
6	a
7	d
8	b
9	c
10	d
11	b
12	b
13	b

If you purc... you may ha... else, ...d.

Detailed Explanations

Solution to Question 1:

Clinically, acute pericarditis is of <6 weeks duration.

Pericarditis is the inflammation of the pericardial sac. This inflammation can lead to excessive secretion of the pericardial fluid leading to pericardial effusion. When the accumulation of fluid causes obstruction to the inflow of blood into the ventricles, it is called cardiac tamponade.

Clinically, pericarditis is classified as:

• Acute - < 6 weeks

• Subacute - 6 weeks to 6 months

• Chronic - > 6 months

Solution to Question 2:

Sold by @itachibot

In this patient, Low-grade fever, chest pain that got relieved on leaning forward, and characteristic auscultatory findings are suggestive of acute pericarditis. The grating sound heard on auscultation is a pericardial friction rub.

Features of the pericardial rub:

• Superficial scratchy/leathery/grating sound caused due to friction between the visceral and parietal pericardium

• Best heard at end-expiration with the patient sitting and leaning forward

• Very high-pitched sound best heard with the diaphragm of the stethoscope, at the lower left sternal border

• Consists of three components corresponding to atrial systole, ventricular systole, and early ventricular diastole

• Pathognomonic of acute pericarditis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 3:

The above scenario is suggestive of acute pericarditis. PR-segment depressions are seen early in acute pericarditis.

The ECG changes in acute pericarditis evolve through 4 stages -

• Stage 1 - Generalized ST-segment elevation with upward concavity with reciprocal depressions only in aVR and V1 and PR segment depression.

• Stage 2 - The ST segments return to normal

• Stage 3 - Diffuse T-wave inversion

• Stage 4 - ECG returns to normal in weeks or months

ECG in acute pericarditis vs acute MI:

Cardiac biomarkers are grossly elevated in acute-MI, whereas in acute pericarditis the elevation if present is only mild and indicative of epi-myocarditis causing myocyte necrosis.

The ECG below shows acute pericarditis causing diffuse ST-segment elevations in leads I, II, aVF, and V2 to V6, with reciprocal ST depressions in lead aVR.

	Acute pericarditis	Acute MI
Distribution of ST-segment elevations	More generalized	Limited to anterior or inferior leads because of the localized area of infarct
Morphology of ST-segment elevations	Concave upwards	Convex upwards
Reciprocal ST-segment depressions in the opposite leads	Not present	Typically present

Sold by @itachibot

	Acute pericarditis	Acute MI
T-wave inversions	Only after ST-segment returns to baseline	T wave inversions can be seen even before the ST-segment returns to baseline
PR-segment	Depressed (or elevated in lead aVR) in 75% to 80% of patients	Usually not depressed (can be depressed only if concomitant atrial infarction is present).
Pathological Q-waves	Not seen	May be seen



The image below shows ECG with ST-segment elevation in Leads II, III, and aVF for an inferior acute myocardial infarction (AMI).



Sold by @itachibot

Solution to Question 4:

A patch of dullness with increased vocal fremitus beneath the angle of the left scapula due to the compression of the base of the lung by the pericardial fluid is called Ewart's sign. It is seen in pericardial effusion.

Acute pericarditis is frequently associated with pericardial effusion.

Option A: Auenbrugger's sign is the bulging of the epigastrium in severe pericardial effusion.

Option B: Moschcowitz's sign is the abrupt transition in percussion from the lungs to the right heart border seen in pericardial effusion.

Option C: Dressler's sign is the dullness to percussion of the lower half of the sternum seen in pericardial effusion.

Solution to Question 5:

Pulsus paradoxus is not part of Beck's triad. Pulsus paradoxus is seen in cardiac tamponade. However, it is not a part of Beck's triad.

Beck's triad is seen in cardiac tamponade and it includes:

• Hypotension

• Soft or absent heart sounds

• Jugular venous distension

Cardiac tamponade is the accumulation of fluid in the pericardial space significant enough to cause an obstruction to the inflow of blood to the ventricles. This reduces cardiac output and arterial pressure causing hypotension.

This also causes a greater than the normal reduction in the systolic arterial blood pressure producing pulsus paradoxus. Pulsus paradoxus refers to the fall in systolic blood pressure >10 mmHg with inspiration.

Other causes of pulsus paradoxus include:

• Massive pulmonary embolism

• Tension pneumothorax

• Severe obstructive lung disease

• Hemorrhagic shock

The echocardiography clip below shows the swinging heart sign, which is characteristic of cardiac tamponade.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



Solution to Question 6:

The differentiating feature between cardiac tamponade and tension pneumothorax is breath sounds. The breath sounds are absent in tension pneumothorax while they are present in cardiac tamponade.

The presence of hypotension, distended jugular veins, and muffled heart sounds (Beck's triad) suggest a cardiac tamponade. These features can also be seen in tension pneumothorax. Other clinical features pointing towards tension pneumothorax are absent breath sounds, hyper resonance to percussion, and shift of the mediastinum to the contralateral side.

Heart sounds can be muffled by structures between the stethoscope and the heart, eg, adipose tissue deposition (obesity), fluid accumulation (pericardial effusion, pleural effusion), gas accumulation (pneumothorax, subcutaneous emphysema), soft tissue masses can lead to attenuation of heart sounds.

Solution to Question 7:

The patient with cardiac tamponade is asked to breathe normally to elicit pulsus paradoxus.

In a patient with cardiac tamponade, both ventricles share an incompressible pericardial covering due to the fluid. As the right ventricle enlarges during inspiration, the interventricular septum bulges into the left ventricle, reducing the left ventricular volume.

This results in reduced stroke volume and hence reduced systolic arterial pressure.

As a result, there is a greater than normal (more than 10 mmHg) fall in systolic arterial blood pressure during inspiration, producing pulsus paradoxus.

This cannot be elicited if the patient is holding his/her breath, taking long or shallow breaths.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Pulsus paradoxus is also seen with effusive constrictive pericarditis, restrictive cardiomyopathy and right ventricular myocardial infarction.

Solution to Question 8:

The given clinical scenario with hypotension, muffled heart sounds, and elevated JVP (Beck's triad) suggest a diagnosis of cardiac tamponade. The JVP finding most likely to be seen is prominent x descent and absent y descent.



Note: JVP in constrictive pericarditis is characterized by prominent x and y descents.

Solution to Question 9:

The ECG shows electrical alternans with sinus tachycardia. This along with the patient's symptoms suggests a likely diagnosis of cardiac tamponade.

Electrical alternans is the beat-to-beat alternation in the ECG waveform. In cardiac tamponade, this is caused by the periodic swinging motion of the heart in a large pericardial effusion.

Sold by @itachibot



Feature	Cardiac tamponade	Constrictive pericarditis	Restrictive cardiomyopathy	
Pulsus paradoxus	+++	+	+/- (Rare)	
JVP	Prominent x descent	+++	+	+++
Prominent y descent	-	++	+	
Kussmaul's sign	-	+	+/- (Rare)	
Pericardial knock	-	++	-	
Third heart sound	-	-	+	
Electrical alternans	++	-	-	
Thickened and calcified pericardium	-	+++	-	

Solution to Question 10:

The symptoms and signs suggest a diagnosis of cardiac tamponade. Out of the given features, pulsus paradoxus is seen in cardiac tamponade.

Feature	Cardiac tamponade	Constrictive pericarditis	Restrictive cardiomyopathy	
Pulsus paradoxus	+++	+	+/- (Rare)	
JVP	Prominent x descent	+++	++	+++
Prominent y descent	-	++	+	
Kussmaul's sign	-	+++	+/- (Rare)	
Pericardial knock	-	++	-	
Third heart sound	-	-	+	

Sold by @itachibot

Feature	Cardiac tamponade	Constrictive pericarditis	Restrictive cardiomyopathy	
Electrical alternans	++	-	-	
Thickened and calcified pericardium	-	+++	-	

Solution to Question 11:

The presence of hypotension, distended jugular veins, and absence of heart sounds (Beck's triad) are suggestive of cardiac tamponade. An urgent pericardial tap needs to be done to relieve the tension on the heart chambers.

Although these features could be seen in tension pneumothorax as well, the presence of adequate bilateral air entry excludes this diagnosis.

Pericardiocentesis, also known as pericardial tap, can be performed with sub-xiphoid, parasternal, or apical approaches. The most common approach is a sub-xiphoid approach under echocardiographic guidance.

Option A: Rapid blood transfusion is needed in massive blood loss indicated by hypotension. Although this patient presents with hypotension, it is most likely a result of cardiac tamponade. Although blood transfusion is indicated, it will be effective only when the pericardial tap decompresses the pericardium.

Option C: Intercostal tube drainage is done in tension pneumothorax. The presence of adequate air entry bilaterally excludes this diagnosis.

Option D: The fixation of femur fracture and femoral artery is not a priority here. The inability to palpate pulses distal to the fracture could be a result of hypotension. This needs to be explored only if the patient fails to improve after the pericardial tap and blood transfusion.

If you purchased this from someone else
you may have been scammed
Sold by @itachibot

Solution to Question 12:

A rise or a lack of fall in JVP with inspiration is known as Kussmaul's sign. It is not seen in cardiac tamponade.

Normally, in inspiration due to the negative intrathoracic pressure and increased intrathoracic volume, the capacity of the right heart increases and the JVP falls. In conditions where the myocardium or pericardium is poorly compliant, it does not allow the right heart chambers to accommodate more blood and this fall is not seen. In fact, due to an increase in venous return during inspiration, the JVP might rise. This is called Kussmaul's sign.

This is seen in:

• Restrictive cardiomyopathy

• Constrictive pericarditis

• Massive pulmonary embolism

• Right ventricular infarcts

Sold by @itachibot

• Advanced cases of left ventricular systolic failure

Kussmaul's sign is not seen in patients with cardiac tamponade because even though the increase in pericardial pressure exerts an inward force compressing the entire heart during inspiration, the increase in negative intrathoracic pressure is still transmitted to the right side of the heart. This allows for an increase in blood flow to the right atrium. Therefore, the JVP falls as seen in normal conditions.

Feature	Cardiac tamponade	Constrictive pericarditis	Restrictive cardiomyopathy	
Pulsus paradoxus	+++	+	+/- (Rare)	
JVP	Prominent x descent	+++	++	+++
Prominent y descent	-	++	+	
Kussmaul's sign	-	+++	+/- (Rare)	
Pericardial knock	-	++	-	
Third heart sound	-	-	+	
Electrical alternans	++	-	-	
Thickened and calcified pericardium	-	+++	-	

Sold by @itachibot
If you purchased this from
you may have been scammed

Solution to Question 13:

The given clinical scenario of a young male developing respiratory distress with hypotension, subcutaneous emphysema, and unilaterally absent air entry is suggestive of tension pneumothorax. The immediate next step, in this case, would be needle decompression at the fifth intercostal space anterior to the mid-axillary line (as the patient is an adult).

In children, the second intercostal space in the midclavicular line is chosen as the site for needle decompression.

Needle decompression turns a tension pneumothorax into an open pneumothorax. It has to be followed by tube thoracostomy. An intercostal chest drain (ICD) tube is inserted in the triangle of safety, just anterior to the midaxillary line on the side of the lung injury. The site of insertion is the same in both children and adults.

The triangle of safety is bounded by the lateral border of the pectoralis major anteriorly, the anterior border of the latissimus dorsi posteriorly, and the fifth intercostal space inferiorly, with the apex below the axilla. This provides a safe landmark for the insertion of an ICD with minimal risk of tissue injury.

Sold by @itachibot

Tension pneumothorax is a condition where air accumulates in the pleural space due to a one-way valve air leak, leading to the collapse of the affected lung. This can cause a mediastinal shift, compression of the opposite lung, and decreased venous return, resulting in obstructive shock. It commonly occurs during positive-pressure mechanical ventilation, but can also happen after chest trauma, subclavian/internal jugular venous catheter insertion, or chest wall defects. The diagnosis is clinical, and immediate treatment is necessary without waiting for radiological confirmation.

Tension pneumothorax is characterized by chest pain, tachypnoea, tachycardia, hypotension, tracheal deviation away from the side of injury, absent breath sounds on the affected side, cyanosis, and neck vein distention. Patients can also develop subcutaneous emphysema as a complication secondary to the pneumothorax.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Pulmonary Hypertension

Question 1:

A patient with a history of frequent exacerbations of COPD is undergoing right heart catheterization for evaluation of pulmonary artery pressure. What is the mean pulmonary arterial pressure above which the diagnosis of pulmonary hypertension can be made?

a) >20 mmHg

b) >25 mmHg

c) >32 mmHg

d) > 30 mmHg

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 2:

Which of the following genes is the most commonly implicated in hereditary pulmonary arterial hypertension?

a) Endothelin

b) PAX – 11

c) Bone morphogenetic protein receptor 2

d) Homebox gene

Question 3:

Which among the following is not a feature of pulmonary arterial hypertension?

a) Mean pulmonary arterial pressure >20 mm Hg

b) Pulmonary capillary wedge pressure ≤ 25 mm Hg

c) Left atrial pressure ≤ 15 mm Hg

d) Pulmonary vascular resistance ≥ 3 Wood units

Question 4:

In a patient diagnosed with pulmonary hypertension, which of the following is an unlikely finding on a chest X-ray?

Sold by @itachibot

a) Enlarged central pulmonary artery

b) Pruning of distal pulmonary vasculature

c) Right ventricular enlargement

d) Focal area of oligemia

Question 5:

Echocardiography was done for a patient who presented with severe exertional dyspnea, chest pain, and occasional fainting spells. It revealed a mitral valve area of 1 sq cm, elevated LA pressure, and significant RV hypertrophy with TR. Which is the next best step in the evaluation of this patient?

a) Cardiac MRI

b) CT pulmonary angiography

c) Right heart catheterization

d) Ventilation / perfusion scan

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 6:

A 32-year-old female patient presents with gradually worsening shortness of breath, fatigue, and exertional chest pain. On auscultation, a loud P2 is heard. Pulmonary function tests reveal an isolated reduction in DLCO. Her family history is positive for similar findings in other members and genetic testing has revealed a mutation in the BMPR2 gene. Based on suspicion, which of the following therapies is this patient most likely to benefit from?

a) Bronchodilator

b) Vasodilator

c) Nitric oxide inhibitor

d) Immunosuppressants

Question 7:

Which of the following group of drugs can not be used in the management of a patient with pulmonary arterial hypertension?

a) Phosphodiesterase type-5 inhibitors

b) Prostacyclin derivatives

c) ACE inhibitors

Sold by @itachibot

d) Endothelin receptor antagonists

Answer Key

Question No.	Correct Option
1	a
2	c
3	b
4	d
5	c
6	b
7	c

Sold by @... If you purchased this from ... else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Pulmonary hypertension is a pathological hemodynamic condition characterized by a mean pulmonary arterial pressure (mPAP) >20mmHg assessed by right heart catheterization.

Pulmonary hypertension could be a complication of cardiac or pulmonary disease or a primary disorder of small pulmonary arteries. Right heart catheterization is the gold standard investigation in diagnosing pulmonary hypertension.

Note: The threshold for diagnosing pulmonary hypertension has been revised from >25 mmHg to >20 mmHg to enable earlier detection.

Solution to Question 2:

Bone morphogenetic protein receptor 2 (BMPR2) mutation is the most commonly implicated gene in hereditary pulmonary arterial hypertension (PAH).

The BMPR2 receptor is involved in the regulation of growth, differentiation, and apoptosis of pulmonary artery endothelial and smooth muscle cells. Due to the mutation, there is enhanced proliferation and reduced apoptosis of vascular endothelial and smooth muscle cells leading to pulmonary hypertension.

Solution to Question 3:

Sold by @itachibot

Pulmonary capillary wedge pressure (PCWP) ≤ 15 mm Hg is a feature of pulmonary arterial hypertension (PAH).

The table below shows the hemodynamic classification of pulmonary hypertension. PAH belongs to WHO group 1.

mPAP- Mean pulmonary artery pressure

PCWP-Pulmonary capillary wedge pressure

PVR- Pulmonary vascular resistance

Parameters	Pre-capillaryWHO groups1,3, 4,5	Post-capillaryWHO grou p 2	Combined pre and post-capill aryWHO groups 2,5
mPAP (mmHg)	>20	>20	>20
PCWP (mmHg)	≤15	>15	>15
PVR (wood uni ts)	≥3	<3	≥3

If you purchased this from so... sold by @ita... may have been scammed.

Solution to Question 4:

A focal area of oligemia or Westermark's sign is seen in pulmonary embolism.

The signs seen on chest radiograph in pulmonary hypertension are:

• Enlargement of central pulmonary arteries

• Rapid tapering of vessels known as vascular pruning

• Right ventricular enlargement



The ECG shows features of RV strain in patients with PH. The pulmonary function tests show a reduction in DLCO in PAH.

Sold by @itachibot

Solution to Question 5:

In the given scenario, a mitral valve area of 1 sq cm suggests severe mitral stenosis. This leads to elevated LA pressure and subsequently pulmonary hypertension. Due to pulmonary hypertension, there is compensatory RV hypertrophy (right ventricular-pulmonary hypertension coupling) and secondary TR. The next step should be to confirm pulmonary hypertension. Right heart catheterization is the gold standard test for diagnosing pulmonary hypertension.

Option A: Cardiac MRI is considered the gold standard for assessing RV function. In the case of pulmonary hypertension, it is used for serial follow-ups due to its non-invasive nature.

Options B and D: CT angiography and ventilation/perfusion scan are done when pulmonary hypertension is suspected to be due to pulmonary embolism, coagulation disorders, or past history of thrombosis (chronic thromboembolic pulmonary hypertension) which should be followed by right heart catheterization.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 6:

The given clinical scenario suggests a diagnosis of pulmonary arterial hypertension. Vasodilator therapy would be beneficial in this patient. Endothelin receptor antagonists, PDE-5 inhibitors, and prostacyclins are examples of vasodilators.

Option A: Since pulmonary arterial hypertension is a vascular disorder, bronchodilation is not beneficial.

Option C: Nitric oxide is a vasodilator and inhibitors of nitric oxide would worsen the disease in pulmonary arterial hypertension.

Option D: Immunosuppressants are not useful in pulmonary arterial hypertension.

Solution to Question 7:

ACE inhibitors are not used in pulmonary arterial hypertension.

Sold by @itachibot

Diseases of Aorta and Peripheral Arteries

Question 1:

Aortic aneurysms caused by spondyloarthropathies are most commonly seen in:

a) Arch of aorta

b) Abdominal aorta

c) Ascending thoracic aorta

d) Descending thoracic aorta

Question 2:

What is the most common pathology associated with ascending aortic aneurysm?

a) Medial degeneration

b) Vasculitis

c) Syphilis

d) Atherosclerosis

Question 3:

A 40-year-old man who is a known case of Marfan's syndrome has come for his regular check-up. His chest X-ray is shown below. Following this, he is being taken up for echocardiography to evaluate the given finding further. Which of the following is true regarding the management of this patient?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) Operative repair if diameter is > 5 cm

b) Operative repair if diameter is > 6 cm

c) Operative repair if diameter is > 5.5 cm

d) Operative repair if diameter is > 4 cm

Sold by @itachi... if you purchased this from someone else, you may have been scammed.

Question 4:

A 50-year-old man came with complaints of irregular bowel movements. While undergoing an abdominal ultrasound examination, an abdominal aortic aneurysm of 5.8 cm diameter was noted. There is a history of hyperlipidemia for the past 10 years. What is the next best step in the management of AAA?

a) Medical management with beta blockers

b) Medical management with statins

c) Operative repair of the aneurysm

d) Follow up with serial abdominal ultrasound

Question 5:

A 55-year-old patient came to the OPD with a complaint of abdominal pain. On further investigations, he was found to have an abdominal aortic aneurysm. Which of the following investigations may indicate the etiology of the aneurysm?

a) Abdominal radiography

b) Abdominal ultrasound

Sold by @itachibot

c) MRI

d) Blood cultures

Question 6:

All of the following are predisposing factors for aortic dissection, except:

a) Takayasu's arteritis

b) Ehlers Danlos syndrome

c) Second trimester of pregnancy

d) Coarctation of aorta

Question 7:

A 65-year-old man presents with acute pain in the chest which radiates to the interscapular region of the back. He is a known hypertensive for the past 10 years. His CT is shown below. Which of the following drug is contraindicated in this patient?

a) Hydralazine

b) Esmolol

c) Labetalol

d) Verapamil

Question 8:

Sold by @itachibot

Sold by @itachibot

The image depicting Stanford type A aortic dissection is?



©MARROW

a) 1

b) 2

c) 3

d) 4

Question 9:

A 50-year-old man presents with a history of severe pain and numbness in his calf muscles following morning walks. The symptoms subside on taking rest. On examination, thickened toenails, decreased pulses, and reduced temperature of the legs are noted. He has a history of smoking for the past 20 years. Which of the following is the most appropriate advice for this patient?

a) Antiplatelet therapy with aspirin

b) Antiplatelet therapy and warfarin

c) Intravenous heparin

d) Percutaneous transluminal angioplasty

Question 10:

Which of the following collagen vascular diseases has the strongest association with Raynaud's phenomenon?

Sold by @itachibot

a) Systemic lupus erythematosus

b) Systemic sclerosis

c) Rheumatoid arthritis

d) Dermatomyositis

Question 11:

A 45-year-old woman complains of bluish discoloration of her fingertips on exposure to cold. Which of the following drugs should be avoided in this patient?

a) Clonidine

b) Prazosin

c) Diltiazem

d) Isradipine

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	c
5	d
6	c
7	a
8	d
9	a
10	b
11	a

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

Aortic aneurysms caused by spondyloarthropathies are most commonly seen in the ascending aorta.

An aneurysm is a pathologic dilation of a segment of a blood vessel. An aortic aneurysm occurs when the structural components of the aortic wall, that is, elastin and collagen, are damaged. This may occur either due to degradation or abnormal synthesis of these components.

The causes of aneurysm are as follows:

	Etiology	Most common site
	Degenerative process-oxidative and biomechanical stressAtherosclerosisAgingCigarette smokingHypertension	Descending thoracic aorta, abdominal aorta
	Medial degeneration/Cystic medial necrosis- degenerative changes in the tunica media of the aortaHypertensionGenetic syndromes likeMarfan's syndrome, Loeys-Dietz syndrome, etc.	Ascending aorta
InfectiousCauses	Syphilitic aneurysms-periaortitis and mesoaortitis damage the elastic fibers	Ascending aorta/aortic arch
Tuberculous aneurysms-granulomatous destruction of the media	Thoracic aorta	
Mycotic aneurysms-bacterial or fungal infections of the aorta	Aorta	
Aortitis	Takayasu's arteritis & Giant cell arteritis (vasculitides)	Aortic arch, descending thoracic aorta
Spondyloarthropathies-ankylosing spondylitis, rheumatoid arthritis, etc.	Ascending aorta	
	Traumatic-chest trauma	Descending thoracic aorta

Solution to Question 2:

The most common pathology associated with ascending aortic aneurysm is medial degeneration.

Medial degeneration (cystic medial necrosis) describes the degenerative process of the tunica media of the aorta. There is degeneration of collagen and elastic fibers in the media and loss of medial cells that are replaced by multiple clefts of mucoid material, such as proteoglycans.

It is seen in the following conditions and is associated with the corresponding mutations:

• Marfan's syndrome- fibrillin 1

Sold by @itachibot

• Loeys-Dietz syndrome- TGFBR1, TGFBR2

• Ehlers-Danlos syndrome type IV - type III procollagen

• Familial thoracic aortic aneurysm syndromes- genes associated with actin and myosin eg. ACTA2, MHC11

• Congenital bicuspid aortic valves

• Turner's syndrome

• Hypertension

Solution to Question 3:

The X-ray shows a widening of the mediastinum with trachea displaced to the right and dilatation of the ascending thoracic aorta (red arrow) i.e. ascending thoracic aortic aneurysm. In the case of Marfan's syndrome, ascending aortic aneurysm of 4-5 cm is taken up for operative repair. This is because of a higher rate of expansion of the aneurysm in these patients.

Management of thoracic aortic aneurysm:

• Beta-blockers and angiotensin receptor antagonists reduce the rate of expansion, especially in Marfan's syndrome.

• Operative treatment is indicated in:

• All symptomatic cases

• Ascending aortic aneurysm - ≥ 5.5 cm in diameter or growth rate of >0.5 cm/year

• Descending aortic aneurysm - > 6 cm in diameter or growth rate of >1 cm/year

• Marfan's syndrome - 4-5 cm in diameter

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

Operative repair of the aneurysm is the next best step in management in this case. Operative repair is indicated for both symptomatic or asymptomatic abdominal aortic aneurysms if they are ≥ 5.5 cm in diameter.

Abdominal aortic aneurysms are mostly asymptomatic and are detected as incidental findings on investigations for other complaints. Sometimes, they present with pain and pulsatile abdominal mass. Rupture of aneurysm presents with acute pain and hypotension.

Management of abdominal aortic aneurysms:

• Statins- atherosclerosis is the most common pathology associated with abdominal aortic aneurysms. Hence, statins are prescribed to reduce the risk of cardiovascular events.

• Asymptomatic aneurysms <5.5 cm in diameter- serial abdominal ultrasound to monitor the rate of expansion of the aneurysm.

• Operative repair is indicated for:

Sold by @itachibot

• Aneurysm of any size associated with symptoms

• Rapidly growing aneurysms

• Aneurysms ≥5.5 cm in diameter

Note: Medical management with beta-blockers though helps in BP control, it is not effective in reducing the rate of growth of abdominal aortic aneurysms.

The given image shows an abdominal aortic aneurysm on CT:



Solution to Question 5:

Blood cultures may indicate the etiology of the aneurysm. They are often positive in patients with mycotic aneurysms.

A mycotic aneurysm is a rare condition that develops as a result of staphylococcal, streptococcal, salmonella or other bacterial or fungal infections of the aorta. The term mycotic aneurysm is usually a misnomer as the causative pathogen is usually bacteria rather than fungi.

These aneurysms are usually saccular. These organisms may involve every major artery, but aortic involvement predominates.

Solution to Question 6:

The third trimester of pregnancy is a predisposing factor for aortic dissection, not second.

Predisposition to aortic dissection increases when there are factors associated with:

• Medial degeneration- seen in:

• Syndromes affecting the connective tissue- Marfan's, Ehlers Danlos syndrome

• Vasculitis- Takayasu's arteritis, giant cell arteritis

Sold by @itachibot

• Increased stress on the aortic wall- seen in:

• Hypertension

• Third trimester of pregnancy

• Other conditions with increased incidence include:

• Bicuspid aortic valve

• Coarctation of aorta

• Aortic trauma

• Cocaine use

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

The given clinical scenario and the CT image showing double-lumen of the aorta are suggestive of aortic dissection. Isolated use of a direct vasodilator such as hydralazine is contraindicated in acute dissection.

It is because these agents can increase hydraulic shear and worsen the dissection by causing vasodilation without reducing the left ventricular ejection force that continues to weaken the arterial wall.

Medical therapy for aortic dissection includes beta-blockers and sodium nitroprusside to reduce cardiac contractility and blood pressure. The heart rate is reduced to around 60 bpm and the SBP is lowered to ≤ 120mmHg.

CCBs like verapamil and diltiazem may be used IV if nitroprusside or beta-blockers cannot be employed. Parenteral angiotensin-converting enzyme (ACE) inhibitor such as enalaprilat may be considered.

Solution to Question 8:

Image D depicts Stanford type A aortic dissection.

The classification schemes for aortic dissection are:

• Stanford classification

• Type A - dissection involves the ascending aorta with or without the involvement of other parts of the aorta

• Type B - dissection is limited to the arch and/or descending aorta without the involvement of ascending aorta

• DeBakey classification

• Type I - dissection involves ascending to descending aorta

• Type II - dissection is limited to the ascending aorta or transverse aorta without descending aorta

Sold by @itachibot

• Type III - intimal tear is located in the descending aorta with distal propagation of the dissection



Hence, if the ascending aorta is involved, it is Stanford type A and DeBakey type I or II. If only the aortic arch or the descending aorta is involved, it is Stanford type B and DeBakey type III.

From a management standpoint, the classification of aortic dissections into type A or B is more practical and useful, since DeBakey types I and II are managed in a similar manner.

Solution to Question 9:

The given clinical scenario describes intermittent claudication. It is the most common symptom of peripheral arterial disease (PAD). The most appropriate advice for this patient is antiplatelet therapy with aspirin.

PAD occurs due to atherosclerotic plaques causing stenosis and occlusion of blood flow. The symptoms of inadequate blood flow exacerbate during increased physical activity.

The following advice is given to patients with PAD:

• Lifestyle modification- regular exercise, discontinue smoking

• Statins to treat hypercholesterolemia

• Reducing the risk of cardiovascular events:

• Controlling hypertension- angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, beta-blockers

• Antiplatelet therapy- aspirin, clopidogrel

• Revascularization procedures (catheter-based or surgical)- For patients not improving with medical therapy

• Endovascular interventions- percutaneous transluminal balloon angioplasty (PTA), stent placement, atherectomy

Sold by @itachibot

• Surgical- aortobifemoral bypass using Dacron grafts, axillofemoral bypass, femoro-femoral bypass

Antiplatelet therapy and warfarin is not preferred because of the high risk of bleeding.

Intravenous heparin is used to treat acute limb ischemia which most often occurs due to thromboembolic events. Acute limb ischemia presents with loss of pulses distal to the occlusion, cyanosis, or pallor. These symptoms do not subside on taking rest. Hence, this patient has intermittent claudication and not acute limb ischemia.

Solution to Question 10:

Among the given collagen vascular diseases, systemic sclerosis has the strongest association with Raynaud's phenomenon.

Raynaud's phenomenon occurs in:

• 80–90% of patients with systemic sclerosis (scleroderma).

• 20% of patients with systemic lupus erythematosus (SLE).

• 30% of patients with dermatomyositis or polymyositis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 11:

The given clinical scenario is suggestive of Raynaud's phenomenon. Clonidine is avoided in Raynaud's phenomenon as it worsens the condition.

Clonidine acts on central alpha-2 adrenoreceptor in CNS and reduces sympathetic outflow. This causes decreased blood pressure and perfusion that can exacerbate Raynaud's phenomenon or any PVD.

Drugs that can be used for the treatment of Raynaud's phenomenon:

Primary Raynaud's:

• Dihydropyridine CCBs- nifedipine, isradipine, felodipine, and amlodipine

• Diltiazem

• Prazosin, doxazosin, and terazosin

Secondary Raynaud's:

• Phosphodiesterase type 5 inhibitors- sildenafil and tadalafil

Digital sympathectomy is helpful in some patients who are unresponsive to medical therapy.

Sold by @itachibot

DVT and Pulmonary Embolism

Question 1:

A patient was brought to the emergency with complaints of chest pain, hemoptysis, and severe dyspnea. CT scan showed vessel sign suggestive of pulmonary infarction. Which among the following is typically responsible for this?

a) Small pulmonary embolism

b) Submassive pulmonary embolism

c) Massive pulmonary embolism

d) Saddle embolism

Question 2:

A patient with a suspected pulmonary embolism underwent a chest X-ray. It showed an enlarged right descending pulmonary artery. What is this sign called?

a) Westermark's sign

b) Hampton's hump

c) Palla's sign

d) Black pleura sign

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

Echocardiography in a patient with pulmonary embolism revealed akinesia of the right ventricular free wall with normal motion at the right ventricular apex. What is this sign called?

a) Moses' sign

b) McConnell's sign

c) Homan's sign

d) Lowenberg's sign

Question 4:

Sold by @itachibot

Which of the following biomarker is least likely to be elevated in a patient with pulmonary embolism?

a) Troponins

b) BNP

c) Heart-type fatty acid–binding protein

d) NGAL

Question 5:

Which of the following conditions is likely to show increased D-dimer values?

a) 1,2

b) 1,3,4

c) 2,3

d) 1,2,3,4

Question 6:

A middle-aged patient presents to the emergency department with a sudden onset pain in the right leg. The pain is associated with swelling, redness, and tenderness. Which of the following is false regarding the management of this condition?

a) Recurrence of this condition despite intensive anticoagulation is an indication for IVC filter

b) IVC filters can prevent recurrent PE in patients of right heart failure

c) IVC filters reduce the rate of recurrence of this condition

d) Pulmonary thromboendarterectomy is done in patients with chronic thromboembolic pulmonary hypertension

Question 7:

A resident is evaluating a patient, who presented with acute onset of chest pain and dyspnea, for pulmonary embolism. Which of the following imaging test should be performed to confirm this diagnosis?

a) Doppler echocardiography

b) CT pulmonary angiography

c) Ventilation perfusion scan of lungs

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) MR pulmonary angiography

Question 8:

A 50-year-old patient is diagnosed with deep vein thrombosis in his right limb. Which of the following parameters is a part of Well's criteria that assess the clinical likelihood of pulmonary embolism in this patient?

a) Dyspnea

b) Orthopnea

c) Chest pain

d) Hemoptysis

Question 9:

In a patient with a Well's score of 1.5, which of the following should be the initial investigation?

a) D-dimer levels

b) CT pulmonary angiography

c) V/Q lung scan

d) Doppler ultrasonography

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following is an indication for fibrinolytic therapy in a patient diagnosed with massive pulmonary embolism?

a) Persistent hypotension

b) RV dysfunction

c) RV enlargement on chest CT

d) Benign intracranial neoplasm

Question 11:

A 32-year-old woman with SLE, secondary anti-phospholipid antibody syndrome develops pulmonary embolism. After admission, she is started on anticoagulants. For how long should

Sold by @itachibot

the anticoagulation therapy be continued?

a) 12-24 months

b) 3-5 years

c) 6-12 months

d) Indefinitely

Question 12:

Which of the following drug was recently approved by FDA for the extended prophylaxis of venous thromboembolism in hospitalized medically ill patients?

a) Unfractionated heparin

b) Apixaban

c) Enoxaparin

d) Betrixaban

Question 13:

What is May-Thurner syndrome?

a) Compression of right iliac vein by right iliac artery

b) Compression of right iliac vein by left iliac artery

c) Compression of left iliac vein by right iliac artery

d) Compression of left iliac vein by left iliac artery

Question 14:

You are evaluating a patient with a cardio-respiratory compromise. Chest X-ray shows a focal area of oligemia. The ECG of the patient is shown below. Choose the false statement regarding this condition.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Most common symptom is dyspnea

b) Most common sign is tachycardia

c) Most common ECG abnormality is sinus tachycardia

d) Most preferred preliminary investigation is D-dimer

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 15:

A 45-year-old woman underwent bariatric surgery. On postoperative day 4, she developed pulmonary embolism as confirmed by a CT pulmonary angiography. Her blood pressure was 70/44 mmHg with poor urine output. IV fluids were administered with dobutamine after which the blood pressure increased to 82/54 mmHg. What is the most appropriate next step to manage this patient?

a) Low molecular weight heparin

b) IVC filter

c) Surgical pulmonary embolectomy

d) Thrombolysis with tPA

Question 16:

A 42-year-old lady was admitted with unprovoked pulmonary embolism from left distal leg DVT. On admission, her coagulation studies were normal. She was started on unfractionated heparin with the first dose of warfarin on the same day.

a) Discontinue warfarin and start long term low molecular weight heparin

Sold by @itachibot

b) Reinstitute heparin and continue warfarin

c) Send HIT tests and start argatroban or lepirudin

d) Continue warfarin

Question 17:

A 35-year-old lady is admitted with unprovoked pulmonary embolism. On admission, her PT was 13.6 seconds (normal 9.6-13.4 seconds) and aPTT was 62 seconds (normal 28.2-34.6 seconds). She has no history of bleeding. Thrombophilia work up was done and she was found to have a strong lupus anticoagulant. How will you manage the patient's initial anticoagulation?

a) Start intravenous unfractionated heparin with aPTT goal in the upper therapeutic range

b) Start intravenous unfractionated heparin and monitor with anti Xa tests

c) Start warfarin without heparin bridge because aPTT is already in the therapeutic range

d) Start intravenous argatroban and monitor by aPTT

Question 18:

A 53-year-old woman underwent hip replacement surgery. A week after the surgery, the patient developed swelling of the legs associated with pain on palpation. Her heart rate is 70 beats per min. There is no history of hemoptysis or significant weight loss. There is no previous history of pulmonary embolism. What is the risk of developing pulmonary embolism in the patient based on Well's score?

a) Low

b) High

c) Moderate

d) Cannot comment without d-dimer values

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	c
3	b

Sold by @itachibot

4	d
5	d
6	c
7	b
8	d
9	a
10	a
11	d
12	d
13	c
14	b
15	c
16	b
17	b
18	b

Sold by @itachibot If you purchased this from elsewhere, your may have been scammed.

Detailed Explanations

Solution to Question 1:

Pulmonary infarction as suggested by CT is typically due to a small pulmonary embolism.

Pulmonary infarction occurs when small thrombi lodge in the segmental and subsegmental vessels. These patients most often have pleuritic chest pain as the site where the thrombus lodges are near the innervation of pleural nerves.

Massive pulmonary emboli and saddle emboli that occlude the main pulmonary artery cause hypotension and shock as opposed to infarction. These are caused by larger-sized emboli that lodge more proximally in the pulmonary circulation.

Pulmonary embolism is classified based on the degree of reduction of pulmonary vascular blood flow and associated hemodynamic disturbances:

• Massive: Hypotension lasting for >15 minutes or BP maintained with inotropes.

• Submassive: No hypotension, but features of cardiac injury present.

• Non-massive: No hypotension or features of cardiac injury.

CT findings associated with pulmonary infarction include a feeding vessel or vessel sign as shown below. It depicts a distinct vessel feeding directly to a lesion, usually of hematogenous origin. However, it is not specific for pulmonary infarction.

Sold by @itachibot



Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 2:

Enlarged right descending pulmonary artery on a chest X-ray in pulmonary embolism is called Palla's sign.

The black pleura sign is seen in alveolar pulmonary microlithiasis. It refers to the shadow of the pleura between the extensive alveolar calcification and the adjacent rib cage.

Solution to Question 3:

The sign described is called McConnell's sign.

It demonstrates right ventricular dysfunction as akinesia of the right ventricular free wall with normal motion at the right ventricular apex. It is the best-known indirect sign of pulmonary embolism on transthoracic echocardiography.

Moses' (Bancroft's), Homan's, and Lowenberg's signs are seen in venous thrombosis. They're not specific to pulmonary embolism.

Solution to Question 4:

NGAL (Neutrophil gelatinase-associated lipocalin) is a kidney injury marker and not a cardiac injury marker.

The following are biomarkers of cardiac injury in pulmonary embolism:

• Plasma brain natriuretic peptide (BNP)

• Heart-type fatty acid binding protein

Sold by @itachibot

• Troponins.

Serum troponin and plasma heart-type fatty acid-binding protein levels increase because of right ventricular microinfarction. Myocardial stretch causes the release of brain natriuretic peptide (BNP) or NT-proBNP.

Solution to Question 5:

All of the conditions mentioned above have raised D-dimer levels.

D-dimer is a fibrin degradation product.

Plasma D-dimer levels commonly rise in DVT and pulmonary embolism due to the breakdown of fibrin by plasmin. D-dimer is more sensitive to pulmonary embolism than DVT. The sensitivity of D-dimer levels in the diagnosis of pulmonary embolism is > 95% but it is is not specific for pulmonary embolism.

Normal D-dimer rules out pulmonary embolism in low and intermediate-risk patients. Patients with a high risk of PE should undergo a CT pulmonary angiogram to effectively rule out PE.

Conditions which have raised D-dimer levels include:

• Myocardial infarction

• Pneumonia

• Sepsis

• Cancer

• Postoperative state

• Pregnancy- second and third trimester

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 6:

The above scenario is suggestive of deep vein thrombosis (DVT). IVC filters increase the rate of recurrence of DVT (by providing a nidus for clot formation) even though they usually prevent pulmonary embolism (over the short term).

The two principal indications for insertion of an IVC filter are:

• Active bleeding (prevents the use of anticoagulation)

• Recurrent venous thrombosis despite intensive anticoagulation

Prevention of recurrent PE in patients with right heart failure who are not candidates for fibrinolysis and prophylaxis of extremely high-risk patients are some other indications for filter placement.

Patients impaired by dyspnea due to chronic thromboembolic pulmonary hypertension should be considered for pulmonary thromboendarterectomy.

Sold by @itachibot

Solution to Question 7:

CT pulmonary angiography should be performed as it is the principal imaging test for the diagnosis of pulmonary embolism (PE).

Option A: Most patients with acute PE have normal echocardiograms. However, echocardiography is a very useful diagnostic tool for detecting conditions that may mimic PE, such as acute myocardial infarction, pericardial tamponade, and aortic dissection.

Option C: Ventilation-perfusion (V/Q) scanning has become a second-line diagnostic test for PE, used mostly for patients who cannot tolerate intravenous contrast or in whom the CT results are inconclusive.

Option D: MR pulmonary angiography may detect large proximal PE but is not reliable for smaller segmental and subsegmental PE. It is used when neither CT angiography nor ventilation-perfusion scan can be done.

The images below demonstrate a pulmonary embolus (marked by arrows). The image of the gross specimen shows saddle thromboembolism.



Sold by @itachibot





Solution to Question 8:

Hemoptysis is a parameter in Well's criteria used to assess the likelihood of pulmonary embolism.

Solution to Question 9:

A patient with Well's score of 1.5 has a low probability of pulmonary embolism. The initial investigation should be the measurement of D-dimer levels.

In a patient with a high probability of pulmonary embolism, the initial investigation is CT pulmonary angiography.

Sold by @itachibot



Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

For fibrinolysis (thrombolytic therapy) in massive pulmonary embolism, the only widely accepted indication is persistent hypotension or shock. It is defined as SBP <90mmHg or a decrease in SBP by ≥40 mmHg from baseline for >15 minutes.

Alteplase or recombinant tissue plasminogen activator (tPA), streptokinase, and urokinase are approved for the treatment of PE. tPA is the most commonly used fibrinolytic, but it is not superior to other agents.

Pulmonary embolism is classified based on the degree of reduction of pulmonary vascular blood flow and associated hemodynamic disturbances:

• Massive: Hypotension lasting for >15 minutes or BP maintained with inotropes.

• Submassive: No hypotension, but features of cardiac injury present.

• Non-massive: No hypotension or features of cardiac injury.

Contraindications for thrombolysis:

• Haemorrhagic stroke

• Ischemic stroke in the last 6 months

• CNS tumours

• Major surgery/trauma in the last 3 weeks

• Active bleeding

• Bleeding diathesis.

Solution to Question 11:

Sold by @itachibot

Duration of anticoagulation for DVT and venous thromboembolism secondary to antiphospholipid antibody syndrome (APLA) is lifelong.

Duration of anticoagulation for DVT and venous thromboembolism:

Clinical Scenario	Duration of Anticoagulation
DVT isolated to an upper extremity or calf that has been provoked by surgery, trauma, estrogen, or an indwelling central venous catheter or pacemaker	3 months
Provoked proximal leg DVT or PE	3 to 6 months
For patients with cancer and VTE	Prescribe LMW heparinContinue anticoagulation indefinitely unless the patient is rendered cancer-free
Idiopathic unprovoked VTE	Indefinite
Antiphospholipid antibody syndrome	Indefinite

Sold to ... if you purchased this ... you may have been st... else, ...d.

Solution to Question 12:

Betrixaban is a new oral, once-daily factor Xa inhibitor approved by the United States Food and Drug Administration (FDA) for extended-duration prophylaxis of VTE in acute medically ill patients.

In patients with no obvious risk factors for VTE, early ambulation with or without mechanical methods (pneumatic compression stockings or venous foot pump) should be advised.

In patients with one risk factor for VTE with no risk of bleeding, LMWH is preferred. In presence of renal failure/cost issues, unfractionated heparin is used. In the case of heparin-induced thrombocytopenia, fondaparinux is preferred.

In patients with multiple risk factors, increasing the dose of anticoagulants or using a mechanical device is advised. For patients with risk factors and a high risk of bleeding or contraindication for anticoagulation, mechanical methods are recommended.

Solution to Question 13:

May-Thurner syndrome (MTS) is caused when the left iliac vein is compressed by the right iliac artery against the fifth lumbar vertebra. It is also called iliocaval venous compression syndrome, iliac vein compression syndrome, Cockett's syndrome, and venous spur.

Sold by @itachibot

Most patients have a partial obstruction and are asymptomatic. It can however progress to symptomatic extensive deep vein thrombosis (DVT). It is a cause of recurrent DVT affecting the left leg.

Solution to Question 14:

The above scenario is suggestive of pulmonary embolism. ECG shows the S1Q3T3 sign (i.e, an S wave in lead I, a Q wave in lead III, and an inverted T wave in lead III) that is relatively specific but insensitive. The most common sign of pulmonary embolism is tachypnea.

Most common symptom of pulmonary embolism is dyspnea. Most common ECG abnormality in pulmonary embolism is sinus tachycardia.Most common preliminary investigation used is D-dimer.



Solution to Question 15:

Surgical pulmonary embolectomy is the most appropriate next step in the management of this patient.

This patient is suffering from a massive pulmonary embolism. In such cases, thrombolytic therapy is given once diagnosis is made using CT pulmonary angiography or bedside ECHO.

But, since this patient underwent a large open abdominal operation (bariatric surgery), thrombolysis is contraindicated due to the increased risk of life-threatening bleeding. Therefore, surgical embolectomy must be done. The main pulmonary arteries are incised, and fresh clots are removed.

Contraindications for thrombolysis:

- Haemorrhagic stroke
- Ischemic stroke in the last 6 months

Sold by @itachibot

- CNS tumours
- Major surgery/trauma in the last 3 weeks
- Active bleeding
- Bleeding diathesis.

Solution to Question 16:

This patient had suboptimal anticoagulation management. The best approach would be to reinstitute heparin and continue warfarin.

The patient had unprovoked PE from lower limb DVT. She was started on warfarin bridged with heparin initially.

Warfarin & parenteral anticoagulation need to be overlapped for at least 5 days, independent of the INR achieved during these 5 days in patients with acute venous thromboembolism.

Warfarin inhibits protein C, protein S (which is part of the fibrinolytic pathway) in addition to factors II, VII, IX, X. Protein C has a short half-life (4-6 hours) and thus decreases early. Since protein C is antithrombotic, decreased levels of protein C by warfarin renders the patient hypercoagulable during the first few days of treatment before factor II with a long half-life (36 hours) decreases and protects the patient from thrombosis.

Thus warfarin creates a prothrombotic state in the first 5 days putting the patient at risk of progression of thrombosis unless a parenteral anticoagulant is given overlapping warfarin for these days.

Solution to Question 17:

This patient is most probably suffering from antiphospholipid antibody syndrome (APS) as her lupus anticoagulant is strongly positive. In such patients, the initial anticoagulation is with unfractionated heparin and is monitored by anti-Xa tests.

Due to the presence of lupus anticoagulant, the lab measurement of aPTT will show erroneously prolonged aPTT. Therefore, even though there is an apparently prolonged aPTT, the patient is at increased risk of thrombosis and not bleeding, and hence patients with APS should be anticoagulated.

Anti Xa tests are not influenced by APS. So, the patient can be monitored for the adequacy of anticoagulation with UFH by monitoring anti-Xa.

Option A: The patient's aPTT is already prolonged due to lupus anticoagulant, so we cannot monitor the patient with aPTT after starting unfractionated heparin.

Option C: Warfarin in the first few days of therapy, causes a decrease in Protein C & S levels, causing a prothrombotic state. Hence, a heparin bridge is necessary for APS which is already a pro-thrombotic state.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option D: Argatroban is a direct thrombin inhibitor used in heparin-induced thrombocytopenia(HIT) to prevent thrombosis and also to increase platelet levels. Therapy is monitored with aPTT.

Note: Lupus anticoagulant may function as an inhibitor in vitro by binding to the reagent phospholipid and blocking the phospholipid-dependent reactions of the intrinsic pathway. The net effect is a prolongation of the aPTT.

Solution to Question 18:

The risk of developing pulmonary embolism in this patient is high (Option B) according to the three-tiered Well's clinical risk assessment score.

Calculation of Well's score:

• Clinical signs of DVT = 3; - swelling of legs, pain on palpation (Moses sign)

• No alternative diagnosis = 3

• Heart rate is <100 BPM = 0; - 70 BPM

• Surgery in previous 4 weeks = 1.5; - hip replacement surgery 1 week ago

• No previous pulmonary embolism = 0

• No Hemoptysis = 0

• No malignancy = 0; - no significant weight loss

Total score = 7.5

Risk stratification:

• High risk = > 6.0

• Moderate risk = 2.0 to 6.0

• Low risk = <2.0 (in patients with low risk, to further confirm the risk - pulmonary embolism rule-out criteria (PERC) may be referred)

Note: There is no other alternative diagnosis which is more likely in this condition, where there is a clear clinical presentation of a deep vein thrombosis, with the associated risk factors of a recent surgery and immobilization. Hence, 3 points can be added to the score.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

General Neurology, Brain Death, and GCS

Question 1:

A 60-year-old man with hemiplegia is brought to the OPD for a follow-up visit. He suffered from a stroke 2 months ago. On examination, he was found to have slurred speech and a positive Babinski sign. Which other examination findings are expected to be seen in this patient?

a) Predominant proximal muscles involvement

b) Hypotonia

c) Fasiculations

d) Exaggerated deep tendon reflexes

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 2:

Which of the following statements about hypertonia is false?

a) Spasticity is velocity dependent

b) Rigidity is present throughout the range of motion

c) Spasticity predominantly affects the extensors of upper limb

d) Rigidity affects flexors and extensors equally

Question 3:

Which of the following features would you not expect to see in a patient with pseudobulbar palsy?

a) Dysarthria

b) Dysphagia

c) Depressed jaw jerk

d) Dysphonia

Question 4:

Match the following pairs of reflexes and their root values:

Sold by @itachibot

a) 1-a, 2-b, 3-e, 4-c, 5-d

b) 1-e, 2-a, 3-d, 4-c, 5-b

c) 1-a, 2-e, 3-b, 4-c, 5-d

d) 1-e, 2-c, 3-d, 4-a, 5-b

Question 5:

All of the following drugs have to be stopped before a lumbar puncture, except:

a) Abciximab

b) Eptifibatide

c) Aspirin

d) Rivaroxaban

Question 6:

A pathologist received a blood-tinged CSF sample for evaluation. The sample was subjected to centrifugation and the following finding was observed. What is the least likely differential diagnosis?

a) Traumatic tap

b) Subarachnoid hemorrhage

c) Liver disease

d) Elevated protein levels

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Sold by @itachibot

Question 7:

A 23-year-old woman presents with nausea, fever, and neck rigidity. A lumbar puncture is ordered. All of the following are effective strategies to prevent post-lumbar puncture headache in this patient, except:

a) Use of atraumatic needle

b) Insertion of a needle with bevel pointed upwards

c) Replacement of stylet prior to removal of the needle

d) Use of 20 gauge needle

Question 8:

A 60-year-old man presents to the OPD with sudden onset speech difficulty, tremors, and difficulty in walking. He is noted to have a wide stepping gait. He is a known hypertensive but is not adherent to treatment. Where is the lesion most likely to be seen on neuroimaging?

a) Frontal lobe

b) Cerebellum

c) Basal ganglia

d) Internal capsule

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 50-year-old man is brought to the emergency ward following an acute episode of inability to move his lower limbs. Examination findings are given below. Which of the following investigations is most appropriate for this patient?

a) Brain MRI

b) Spinal MRI

c) Electromyography

d) Nerve conduction studies

Question 10:

An elderly diabetic and hypertensive patient was carried to the emergency room in a comatose state. On examination, the blood pressure was 170/100 mmHg and pulse rate was

Sold by @itachibot

>100/min. Plantar reflex was bilateral extensor. What is the next step to do?

a) Check blood sugar level

b) Give antihypertensive

c) IV mannitol

d) CT brain

Question 11:

Which of the following is the most common cause of autonomic storm?

a) Botulinum toxin

b) Brain injury

c) Spinal cord injury

d) Guillain-Barré syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

What is the minimum Glasgow coma scale-pupils score for a head injury?

a) 0

b) 3

c) 4

d) 1

Question 13:

A 32-year-old man is brought to the emergency department following a road traffic accident while driving a motorcycle. During the evaluation, the patient opens his eyes in response to voice and says inappropriate words. He is unable to move his right arm but extends his left arm when a painful stimulus is applied to his right arm. What is the GCS score of this patient?

a) 7

b) 8

c) 9

d) 10

Sold by @itachibot

Question 14:

You are evaluating a patient post a neurosurgical procedure. The patient opens her eyes in response to a loud voice, localizes pain, and is confused and disoriented. What is the GCS score of this patient?

a) E3M5V4

b) E3M6V5

c) E4M4V5

d) E3M5V5

Question 15:

Calculate the GCS Score in an intubated patient, with findings of eye movements to pain and abnormal flexion?

a) E2VTM4

b) E2VTM3

c) E2V1M3

d) E2VNTM3

Question 16:

A patient is brought to the emergency room after he was brutally beaten during a mugging incident. He is unresponsive even to painful stimuli and is apneic; however, he does have a pulse. Which of the following clinical findings would exclude a diagnosis of brain death?

a) Absence of oculovestibular reflex

b) Constricted pupils

c) Invariant pulse rate

d) Positive deep tendon reflexes

Question 17:

The family of an ICU patient who is in a coma has given consent for organ donation. An apnea test is performed. Lack of respiratory effort at what arterial pCO2 level will aid in confirming brain death?

a) 40 mmHg

Sold by @itachibot
If you purchased this from someone else, you may have been scammed!

Sold by @itachibot

b) 50 mmHg

c) 60 mmHg

d) 70 mmHg

Question 18:

Clinical criteria for brain death include all except:

a) Coma

b) Absent brain stem reflexes

c) Absent spinal cord reflexes

d) Absent motor activity

If you bought this from someone else, you've been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	c
3	c
4	b
5	c
6	a
7	d
8	b
9	b
10	a
11	b
12	d
13	b
14	a
15	d
16	b
17	c
18	c

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The given patient has slurred speech and a positive Babinski sign, which is suggestive of an upper motor neuron lesion (UMN lesion). Exaggerated deep tendon reflexes are a feature of UMN lesions.

The distal muscle groups are affected more than the proximal muscles and the axial muscles are spared (as the anterior corticospinal tracts supplying the axial muscles have bilateral representation) in UMN lesions. Axial muscles are involved when the lesion is severe or bilateral.

Hypertonia in the form of spasticity is seen in the upper motor neuron lesions. Fasciculations and hypotonia are features of lower motor neuron lesions.

If you bought this from someone else, it was scammed. Sold by @itachibot

Solution to Question 2:

Spasticity predominantly affects the antigravity muscles - flexors in the upper limb and extensors in the lower limb.

Spasticity	Rigidity
Seen in lesions of pyramidal t racts	Seen in lesions of the extrapy ramidal tracts such as rubros pinal and vestibulospinal trac ts
Velocity dependent - More no ticeable with fast movements	Present throughout range of motion without change in int ensity
Clasp-knife phenomenon - su dden release after reaching m aximum	Lead pipe or plastic stiffness
Predominantly affects antigra vity muscles	Affects flexors and extensors equally

Solution to Question 3:

Jaw jerk is brisk in pseudobulbar palsy.

Bilateral corticobulbar lesions produce pseudobulbar palsy. This condition is characterized by dysarthria, dysphagia, dysphonia, emotional lability, bilateral facial weakness, and brisk jaw jerk.

Damage to lower motor neurons of the cranial nerves in the brainstem is called bulbar palsy. Depressed jaw jerk is seen in bulbar palsy.

Pseudobulbar palsy is analogous to UMN lesions exhibiting hyperreflexia. Bulbar palsy is analogous to LMN lesions exhibiting hyporeflexia.

Sold by @itachibot

Solution to Question 4:

The correct match is:

Reflex	Root value
Biceps	C5, C6
Cremasteric	L1, L2
Knee	L3, L4
Plantar	L5 - S2
Ankle	S1, S2

Solution to Question 5:

Aspirin and other NSAIDs can be continued in a patient who is scheduled to undergo a lumbar puncture (LP). These drugs have not shown an increased risk of developing spinal hematomas. Hence an LP can be performed in patients receiving these drugs.

Options A and B: Abciximab and eptifibatide are Gp IIb/ IIIa inhibitors. Platelet aggregation remains abnormal following the discontinuation of these drugs. LP, spinal and epidural procedures must be avoided until the platelet function is normal.

Option D: Rivaroxaban is an oral factor Xa inhibitor. LP should be avoided in patients receiving this drug due to the risk of developing epidural or spinal hematomas. Other factor Xa inhibitors, namely, fondaparinux and apixaban must also be held prior to LP, spinal and epidural procedures.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 6:

The given image shows xanthochromatic CSF on centrifugation. This is unlikely with a traumatic tap. A traumatic tap has clear supernatant following CSF centrifugation.

Xanthochromia refers to the yellowish appearance of CSF. In a lumbar puncture, a traumatic tap can be confused with subarachnoid hemorrhage (SAH) as both show blood in the drawn CSF sample. In such situations, centrifugation of the sample distinguishes the two.

• Clear supernatant following CSF centrifugation supports a traumatic tap.

• Xanthochromatic supernatant suggests SAH.

Xanthochromatic supernatant may be present in the following conditions:

• In Liver disease, due to elevated bilirubin levels.

Sold by @itachibot

• When CSF protein concentration is elevated.

• In subarachnoid hemorrhage (SAH), RBCs are present in the CSF long enough for them to undergo lysis. This releases hemoglobin which is converted to bilirubin, thus staining the CSF yellow.

Solution to Question 7:

The use of a 22 gauge needle (or smaller) can reduce the incidence of post-dural headache after lumbar puncture.

Effective strategies that reduce the incidence of PDPH include:

• Use of small-diameter needle (22-gauge or smaller)

• Use of atraumatic needle (Sprotte and others)

• Replacement of stylet prior to removal of the needle

• Insertion of a needle with bevel oriented in a cephalad to caudad direction (when using a stranded needle)

Ineffective strategies:

• Bed rest (up to 4h) following LP

• Supplemental fluids

• Minimizing the volume of spinal fluid removed

• Immediate mobilization following LP

The customary practice of remaining in a recumbent position post-LP is unnecessary. Studies have shown no benefit in terms of reducing the incidence of headaches.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

The given clinical features are suggestive of cerebellar ataxia, hence the lesion is most likely to be seen in the cerebellum. Cerebellar ataxia can be caused due to an acute vascular event such as infarction or hemorrhage.

Wide stepping gait, a broad-based unstable gait is indicative of cerebellar dysfunction. Few other notable gait abnormalities are:

• Short stepped gait- parkinsonism

• Scissoring gait- spasticity i.e. UMN lesion

• High-stepped, slapping gait- posterior column or peripheral nerve disease

• Waddling gait- proximal muscle weakness

A few of the cerebellar signs include:

• Dysdiadokinesia / dysmetria

Sold by @itachibot

• Ataxia

• Nystagmus

• Intention tremor

• Speech - slurred or scanning

• Hypotonia

Solution to Question 9:

The given clinical scenario is suggestive of acute paraparesis. This is caused most commonly by an intraspinal lesion. A negative Babinski reflex and flaccidity are suggestive of a lower motor neuron lesion. Hence, the most appropriate among the given investigations for acute paraparesis with LMN signs is a spinal MRI.

Spinal causes of acute onset paraparesis include:

• Injury to the spinal cord

• Thrombosis of anterior vertebral artery

• Hematomyelia

• Prolapsed intervertebral disc

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Ruling out glycemic extremes is the first step in the evaluation of this patient who is in a coma as most of the causes of coma are metabolic in origin and with appropriate treatment are completely reversible.

In this elderly patient, with no other relevant history, the most likely cause of coma is diabetic ketoacidosis, hyperglycemic hyperosmolar state, or even hypoglycemia (which may present with hypertension).

Other options :

The patient presents with tachycardia which is not a sign of raised intracranial pressure(bradycardia, altered respiratory pattern, raised systolic blood pressure) and hence starting iv mannitol (option 3) to reduce intracranial pressure or doing a CT (option 4) to rule out intracranial space-occupying lesion is not required.

The patient is a known case of hypertension which may cause hypertensive encephalopathy due to cerebral edema and would present with a similar history and signs of raised intracranial pressure giving antihypertensive (option 2) medications would treat the condition but it is not the first step as metabolic causes are more common and have to be addressed first.

Coma is defined as a deep sleep-like state from which the patient cannot be aroused.Coma could be due to:

• Metabolic and toxic encephalopathies, where brain imaging will be normal

Sold by @itachibot

• Meningitis syndromes

• Diseases with prominent focal signs like stroke, or hemorrhage

Urgent intervention is needed to prevent further brain damage. It is done by providing supplemental oxygen, intravenous thiamine (at least 100 mg), and intravenous 50% dextrose in water (25 g) empirically. It is crucial to measure baseline serum glucose levels before administering glucose.

The initial examination includes evaluating the general appearance,vitals, breath sounds, response to stimulation, pupil size and responsiveness, and any abnormal posturing or movements. In trauma cases, the neck should be stabilized until ruling out cervical spine fracture. Airway protection and intravenous access is necessary.

Immediate therapeutic intervention should be done for hypotension, significant hypertension, bradycardia, arrhythmias affecting blood pressure or marked hyperthermia. Signs of meningeal irritation or fever indicate the need for urgent lumbar puncture. Fundus examination for papilledema and a CT scan of the brain should be performed prior to lumbar puncture in comatose patients.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

Brain injury is the most common cause of autonomic storm and typically follows severe head trauma and postresuscitation anoxic-ischemic brain injury.

An autonomic storm is an acute state of sustained sympathetic surge which can cause alterations in BP and heart rate, body temperature, respiration, and sweating.

Other causes include:

• Spinal cord injury

• Drugs and toxins (phenylpropanolamine, cocaine, amphetamines, and TCA's; tetanus; and less often, botulinum toxin)

• Autonomic neuropathy

• Chemodectomas (e.g., pheochromocytoma).

• Guillain-Barré syndrome can produce a moderate autonomic storm.

Solution to Question 12:

The GCS-P score allows a minimum score of 1.

The Glasgow coma scale – pupils score (GCS-P) was described in response to the wish for a single index combining the coma scale with pupillary reactivity as a reflection of brainstem function. Its possible values range from 1 to 15, reflecting an extended range of severity, and may be particularly useful in relating to prognosis.

The GCS-P is calculated by subtracting the pupil reactivity score (PRS) from the Glasgow coma scale (GCS) total score:

Sold by @itachibot

GCS-P = GCS - PRS

The minimum score in GCS is 3.

Solution to Question 13:

The GCS score of the patient described is 8 (E3 V3 M2).

In the case elaborated above, the score allotted is :

• Eye-opening to speech—3

• Inappropriate words—3

• Extensor response of the upper limb in response to painful stimuli—2

Patients scoring a GCS of 3 or 4 have an 85% chance of dying or remaining vegetative.

However, scores >11 indicate only a 5–10% likelihood of death or vegetative state and an 85% chance of moderate disability or good recovery.

Solution to Question 14:

In this patient, GCS is E3M5V4

• Eye-opening to loud voice - 3

• Motor response localizes pain - 5

• Verbal response is confused and disoriented - 4

Solution to Question 15:

The GCS score is E2VNTM3. In this case:

• Eye-opening to pain: E2

• In case of endotracheal intubation or tracheostomy, the verbal component of the score is denoted as 'not testable/NT'. Earlier in intubated patients, the V score was denoted as VT. According to the current update, the V score should be VNT.

• Abnormal flexion: M3

Solution to Question 16:

Constricted pupils would rule out brain death.

Brain death is the irreversible cessation of all cerebral function with the preservation of cardiac activity and maintenance of respiratory and somatic function by artificial means.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

In brain death, the pupils are usually midsized or maybe dilated. However, constricted pupils are not seen.

The diagnosis of brain death should be confirmed with the following clinical findings:

• Absent brain-originating motor response, including response to pain stimulus above the neck or other brain-originating movements (eg, seizures, decerebrate or decorticate posturing)

• Unresponsiveness to any stimuli

• Brainstem damage, as evidenced by dilated or mid-sized pupils without light reaction

• Absent corneal, gag, and oculovestibular reflexes

• Absent sucking or rooting reflexes

• Absent cough with tracheal suctioning

• Apnea, indicating medullary destruction

• The heart rate should be invariant and does not accelerate to atropine

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 17:

To declare brain death, apnea is confirmed if no respiratory effort has been observed in the presence of arterial pCO_2 levels of ≥60 mmHg (apnea test is positive).

Destruction of the medulla leads to complete and irreversible apnea. Demonstration that apnea is due to medullary damage requires that the PCO_2 be high enough to stimulate respiration during a test of spontaneous breathing.

An apnea test is conducted to confirm that there is no respiratory effort in the presence of a sufficiently elevated arterial PCO_2. Apnea testing can be done safely by the use of preoxygenation with 100% oxygen prior to and following removal of the ventilator. CO_2 tension increases ~0.3–0.4 kPa/min (2–3 mmHg/min) during apnea.

Solution to Question 18:

Clinical criteria for brain death does not include absent spinal cord reflexes. Deep tendon reflexes/spinal cord reflexes may be present in brain death as the spinal cord is intact.

Criteria for brain death:

• Coma - Deep, unresponsiveness to noxious stimuli

• Absence of brainstem reflexes - Pupillary, oculocephalic, corneal, gag reflex, etc.

• Apnea - Absence of spontaneous respiration at arterial pCO_2 tension of at least 50–60mm Hg

Sold by @itachibot

Seizures and Epilepsy

Question 1:

A 30-year-old patient is undergoing EEG with his eyes open. What is the expected frequency of the waves being recorded?

a) < 4 Hz

b) 4- 7 Hz

c) 8- 13 Hz

d) >13 Hz

Question 2:

A middle-aged obese man is being evaluated at the sleep clinic. During his EEG recording, he is awake with his eyes closed. Which of the following waves will be predominantly seen?

a) Predominant alpha waves with intermixed beta activity

b) Predominant beta waves with intermixed alpha activity

c) Predominant theta waves with intermixed delta activity

d) Predominant delta waves with intermixed theta activity

Question 3:

Which of the following is not a type of generalized epilepsy?

a) Childhood absence

b) Juvenile absence

c) Juvenile myoclonic

d) Jacksonian march

Question 4:

A 13-year-old boy presents with involuntary, sudden jerking movements of both upper limbs. They occur most frequently in the morning after awakening and are provoked by lack of sleep. This type of seizure is classified as _____.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

a) Unknown onset

b) Focal and generalized onset

c) Focal onset

d) Generalised onset

Question 5:

Which of the following is not seen in focal seizures?

a) Jacksonian march

b) Todd's paralysis

c) Epilepsia partialis continua

d) Ictal cry

Question 6:

A 10-year-old girl is brought to the OPD with complaints of frequent fainting and sudden falls while playing. She sometimes falls off while sitting in the classroom hitting her head to the desk. She regains consciousness immediately and resumes her activity, and has normal behavior after such episodes. The girl is otherwise healthy. What is the likely diagnosis?

a) Typical absence seizures

b) Atonic seizures

c) Syncope

d) Atypical absence seizures

Question 7:

A 6-year-old boy is brought to the hospital. He is found to have a sudden brief lapse of consciousness with preserved postural control. No confusion is reported after regaining consciousness. What is the diagnosis?

a) Atonic seizures

b) Myoclonic seizures

c) Absence seizures

d) Generalized tonic clonic seizures

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 8:

An unconscious patient with a history of seizures is brought to the ED. He then develops muscular flaccidity and excessive salivation. After a few minutes, he begins to regain consciousness and is in a confused state. This phenomenon is associated with which of the following types of seizures?

a) GTCS

b) Absence seizures

c) Atonic seizures

d) Myoclonic seizures

Question 9:

A 15-year-old boy presents to the clinic with involuntary, sudden jerking movements of both upper limbs. They occur most frequently in the morning after awakening and are provoked by a lack of sleep. EEG demonstrates bilaterally synchronous spike-and-slow- wave discharges. Which of the following features is not seen in this type of seizure?

a) Post-ictal paralysis

b) Preserved postural control

c) Lack of postictal confusion

d) Preserved consciousness

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

The following EEG is that of a 6-year-old girl who is easily distracted in class and exhibits poor scholastic performance. Seizures are precipitated by hyperventilation. What is the diagnosis ?

Sold by @itachibot



a) Myoclonic seizures

b) Atonic seizures

c) Absence seizures

d) Clonic seizures

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 11:

What is the first-line agent for treating a patient with generalized tonic-clonic seizures?

a) Phenytoin

b) Carbamazepine

c) Ethosuximide

d) Valproate

Question 12:

A 16-year-old boy developed abnormal jerking movements that started in his left hand and gradually involved the entire arm. Following this, the patient experienced weakness and difficulty in moving the left arm for about an hour. Two similar episodes have also occurred in the past. Which of the following is the first-line drug for this condition?

a) Carbamazepine

b) Valproate

c) Ethosuximide

Sold by @itachibot

d) Topiramate

Question 13:

All of the following can be used as first-line agents in patients with typical absence seizures, except:

a) Levetiracetam

b) Ethosuximide

c) Lamotrigine

d) Valproate

Question 14:

Which of the following is used as the first-line antiepileptic drug in the treatment of both generalized as well as focal seizures?

a) Valproate

b) Carbamazepine

c) Lamotrigine

d) Phenytoin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 15:

Which of the following is the drug of choice for a patient with generalized epilepsy syndrome having mixed seizures?

a) Ethosuximide

b) Carbamazepine

c) Phenytoin

d) Valproate

Question 16:

A 34-year-old man is brought to the emergency department with violent shaking of arms and legs for the past 1 hour. He is a chronic alcoholic whose last intake was 2 days ago. IV lorazepam and phenytoin are administered to this patient. After 5 minutes, the seizures have

Sold by @itachibot

not stopped. What is the next best step in the management of this patient?

a) Phenobarbitone

b) Lorazepam

c) Propofol

d) Thiamine infusion

Question 17:

A 16-year-old girl, who is taking antiepileptics, has had a seizure-free period of 6 months. She has no family history of epilepsy. Her EEG is now normal and she has a normal neurological exam and intelligence. What would your advice be?

a) Stop the treatment and follow up

b) Gradually taper the drug and stop treatment

c) Continue treatment for another 18 months

d) Continue lifelong treatment with antiepileptics

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A 30-year-old man on antiepileptic therapy for GTCS presents with weight gain, alopecia and thrombocytopenia. Which of the following antiepileptic drugs is the most likely cause for this presentation?

a) Phenytoin

b) Carbamazepine

c) Lamotrigine

d) Valproate

Question 19:

Which of the following features would you not expect to see in a patient with West syndrome?

a) Infantile spasms

b) Abnormal EEG

c) Developmental regression

d) Café au lait

Sold by @itachibot

Question 20:

A 4-month-old child was brought with repeated epileptic spasms. The child had reached his normal milestone at the age of 2 months. But the parents noticed that he lost the ability to hold his head steadily and he recently couldn't track light or sound. The EEG findings are shown below. What is the drug of choice for the treatment of this condition?



a) ACTH

b) Vigabatrin

c) Gabapentin

d) Pregabalin

Question 21:

Which of the following drugs is effective in controlling infantile spasms in a patient who has the neurocutaneous marker as shown in image?

Sold by @itachibot

a) Prednisolone

b) ACTH

c) Vigabatrin

d) Gabapentin

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 22:

A 14-year-old boy develops an episode of involuntary jerky movements of the left hand accompanied by a repetitive chewing movement. He had a blank look on his face and was unresponsive during this episode. He also has no memory of the episode. EEG was normal and MRI showed hippocampal sclerosis. What is the most likely diagnosis?

a) Juvenile myoclonic epilepsy

b) Lennox-Gastaut syndrome

c) Gelastic epilepsy

d) Mesial temporal lobe epilepsy

Question 23:

Which of the following statements is false regarding mesial temporal lobe epilepsy ?

a) It results in focal seizures with dyscognitive features

b) The person experiences an aura before the seizure

c) It is associated with a family history of epilepsy

Sold by @itachibot

d) Seizures can be controlled with anticonvulsants such as carbamazepine

Question 24:

Which of the following drugs must be avoided in myoclonic seizures?

a) Valproate

b) Levatriacetam

c) Ethosuxamide

d) Oxcarbazepine

Question 25:

Which of the following is seen in patients with Lennox-Gastaut syndrome?

a) Atypical febrile seizures

b) Single type of seizures

c) EEG showing less than 3Hz Spike and wave discharge

d) Good prognosis with adequate control

Question 26:

Which of the following is false regarding Lafora's disease?

a) Myoclonus is seen

b) Autosomal recessive

c) Diagnosis by DNA sequencing

d) Treatment with glucocerebrosidase infusions

Question 27:

You are at a criminal court to provide medical testimony. The trial ends with the accused being sentenced to life imprisonment. Abruptly, the convict breaks into uncontrollable laughter. Within minutes, he collapses and starts to have a seizure. His lawyer reports that he has a childhood seizure disorder. What is the likely diagnosis?

a) Gelastic seizures

b) Kluver Bucy syndrome

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

c) Witzelsucht

d) Pseudobulbar affect

Question 28:

A 10-year-old girl is brought to the OPD following a generalized tonic-clonic seizure. She had previous episodes of sudden falls at school and also sometimes stares with a blank face. She is also found to be having poor scholastic performance. Her EEG reveals a slow spike and wave pattern. What is the most likely diagnosis?

a) Focal seizures progressing to generalized seizures

b) Myoclonic epilepsy progressing to generalized seizures

c) Lenox-Gastaut syndrome

d) Atypical absence seizures

Sold by @itachibot
If you bought this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	d
2	a
3	d
4	b
5	d
6	b
7	c
8	a
9	a
10	c
11	d
12	a
13	a
14	c
15	d
16	b
17	c

Sold by @itachibot

18	d
19	d
20	a
21	c
22	d
23	d
24	d
25	c
26	d
27	a
28	c

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Detailed Explanations

Solution to Question 1:

Beta waves are seen on the EEG of an adult recorded with eyes open. The frequency of these waves on an EEG is >13 Hz.

Solution to Question 2:

In a normal adult awake patient lying quietly with his eyes closed, alpha waves are predominantly seen intermixed with a variable generalized beta activity.

This alpha rhythm is attenuated when the eyes are opened.



Sold by @itachibot

Solution to Question 3:

Jacksonian march is a feature seen in focal seizures, not generalized epilepsy.

In Jacksonian march, the abnormal motor movements begin in a very restricted region such as the fingers and gradually progress to include a larger portion of the extremity. It represents the spread of the seizure activity over a progressively larger region of the motor cortex.



Solution to Question 4:

The given scenario is suggestive of juvenile myoclonic epilepsy (JME). Myoclonic seizures are the predominant feature of JME and are now classified under both focal onset and generalized onset seizures.

Generalized seizures arise at some point in the brain but immediately and rapidly engage neuronal networks in both cerebral hemispheres.

Focal seizures arise from a neuronal network either discretely localized within one brain region or more broadly distributed but still within a cerebral hemisphere.

Myoclonic seizures are characterized by:

• Preserved consciousness

• Preserved postural control

• Lack of postictal confusion

JME is an epilepsy syndrome that has familial inheritance and begins in adolescence. It most commonly occurs in the morning and is provoked by sleep deprivation. JME is usually associated with GTCS and sometimes with absence seizures. EEG shows a typical 4-6 Hz polyspike and wave pattern. Treatment includes administration of sodium valproate.

Sold by @itachibot

2017 ILAE-International League Against Epilepsy Classification:



Sold by @ita... If you purchased this from so... else, you may have been scammed.

Solution to Question 5:

The ictal cry is seen in generalized, tonic-clonic seizures (GTCS). Jacksonian march, Todd's paralysis, and Epilepsia partialis continua are features of focal seizures.

Jacksonian march is a feature seen in focal seizures wherein the abnormal motor movements begin in a very restricted region such as the fingers and gradually progress to include a larger portion of the extremity. It represents the spread of seizure activity over a progressively larger region of the motor cortex.

Todd's paralysis is a transient focal temporary paralysis usually persisting for a few minutes to hours (maximum up to 48 hours) in the postictal phase.

Epilepsia partialis continua refer to continued focal seizures for hours or days. It is often refractory to medical therapy.

Tonic contraction of the muscles of expiration and the larynx at the onset of a generalized tonic-clonic seizure will produce a loud moan or ictal cry.

Solution to Question 6:

The given history is most consistent with atonic seizures. Atonic seizures are characterized by a sudden loss of postural control and there is no postictal confusion.

Options A and D: Absence seizures usually present as blank stares or unresponsiveness for a few seconds. Postural control is not lost during absence seizures.

Option C: The girl regains consciousness immediately and has no associated complaints like sweating, fatigue, or tachycardia. She also resumes her activity after the episodes. Also, there is no precipitating factor that points towards syncope. Hence it is unlikely to be syncope.

Sold by @itachibot

Solution to Question 7:

Sudden, brief lapse of consciousness with preserved postural control and no post-ictal confusion is seen in absence seizures. Usually, the seizure lasts only a few seconds.

EEG shows generalized symmetric 3 Hz spike and wave pattern (as shown below). Hyperventilation provokes these EEG discharges and seizures.



Option A: Atonic seizures are characterized by a sudden loss of postural control.

Option B: Myoclonic seizures have preserved consciousness. They involve a sudden brief jerking movement involving the entire body or a part of it.

Option D: Generalised tonic-clonic seizures are characterized by post-ictal confusion.

Solution to Question 8:

The scenario is suggestive of post-ictal confusion which is seen in generalized tonic-clonic seizures.

It is not seen in absence seizures, atonic seizures or myoclonic seizures.

Solution to Question 9:

The given clinical scenario is suggestive of myoclonic seizures. Post-ictal paralysis is not seen in these seizures.

Myoclonic seizures are characterized by preserved consciousness (rarely very short loss of consciousness), preserved postural control, Lack of postictal confusion.

Sold by @itachibot

Myoclonic seizures usually coexist with other forms of generalized seizures but are the predominant feature of juvenile myoclonic epilepsy (JME). The EEG shows bilaterally synchronous spike-and-slow wave discharges.

JME or Janz's syndrome is an epilepsy syndrome that has familial inheritance and begins in adolescence. It most commonly occurs in the morning and is provoked by sleep deprivation. JME is usually associated with GTCS and sometimes with absence seizures. EEG shows a typical 4-6 Hz polyspike and wave pattern. Treatment includes administration of sodium valproate.

Solution to Question 10:

EEG shows generalized symmetric 3 Hz spike and wave pattern. This is seen in absence seizures.

Absence seizures are characterized by a sudden brief lapse of consciousness with preserved postural control that lasts only a few seconds. However, there is no post-ictal confusion.

Hyperventilation provokes these EEG discharges and also the seizures.

Ethosuximide is the drug of choice in typical absence seizures.

If yc (altachibot someone else; med.

EEG in Absence Seizure - 3Hz
Spike and Wave Pattern



Solution to Question 11:

The first-line agents in the treatment of generalized tonic-clonic seizures (GTCS) are Valproate, levetiracetam, and lamotrigine.

Solution to Question 12:

The given clinical scenario describes Jacksonian march followed by Todd's palsy. These are characteristics of a focal seizure. Among the given drugs, the first-line drug for focal seizures is

Sold by @itachibot

carbamazepine.

Solution to Question 13:

Levetiracetam is not the first-line agent for the treatment of typical absence seizures.

First-line drugs for:

• Typical absence seizures: Ethosuximide (drug of choice), valproate, and lamotrigine

• Atypical absence seizures: Valproate

Solution to Question 14:

Lamotrigine is a first-line drug in the treatment of both generalized tonic-clonic and focal seizures.

Solution to Question 15:

Valproate is the drug of choice in generalized epilepsy syndrome having mixed seizures. It is particularly effective in absence, myoclonic and atonic seizures.

Valproate is usually first given alone (monotherapy) and if ineffective, another drug such as lamotrigine may be added.

In generalized epilepsy, drugs such as phenytoin, carbamazepine, and oxcarbazepine can increase the number of seizures. They can also provoke new seizure types - myoclonic or absence seizures. Hence, they are avoided.

Ethosuximide is the drug of choice in typical absence seizures and is not used for GTCS.

Solution to Question 16:

With seizures occurring for 1 hour, this patient is suffering from status epilepticus. A second dose of lorazepam can be given to control this patient's seizures if there is no response after 5 minutes of administration of the first dose.

Status epilepticus refers to continuous seizures or repetitive, discrete seizures with impaired consciousness in the inter-ictal period. For practical purposes, it is defined as any one of the following:

• Continuing seizures >5 minutes or

• No regaining of consciousness between 2 seizures.

Management of status epilepticus:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



Solution to Question 17:

In the given clinical scenario of a girl on antiepileptics for 6 months with no further seizures, no family history, normal EEG, neurological examination, and intelligence, antiepileptics should be continued for another 18 months since she's already taken for 6 months (minimum 2 years) to prevent seizure recurrence.

Anti-epileptic drugs withdrawal is considered only when an adult patient has been seizure-free for ≥2 years along with the following criteria:

• Normal CNS examination including intelligence

• Normal EEG and normal MRI

• Single seizure type

• No family history of epilepsy.

It is preferable to reduce the dose of the drug gradually over 2–3 months.

Other options:

Option A: Sudden cessation of drug therapy may cause recurrence of seizure hence not advised.

Option B: The patient has a seizure-free interval of only 6 months and therefore, is not yet eligible for considering medication withdrawal.

Option D: Life-long treatment is not necessary in this case as the patient meets all the above-mentioned criteria for drug withdrawal after 2 years.

Note: Juvenile myoclonic epilepsy (JME), the most common idiopathic generalized epilepsy syndrome is known to have a high risk of seizure recurrence, and hence antiepileptics are continued lifelong in a majority of patients.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 18:

Weight gain, alopecia, and thrombocytopenia are the side effects of valproate use.

Adverse effects of Sodium Valproate (mnemonic: VALPROATE)

• Vomiting

• Alopecia

• Liver toxicity

• Pancreatitis

• Rash

• Obesity

• Anorexia

• Teratogenicity (most common: spina bifida), thrombocytopenia, tremors

• Edema

The first-line agents for treating GTCS are valproate, leviteracetam, and lamotrigine.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 19:

Café au lait is a neurocutaneous marker seen in neurofibromatosis, not a component of West syndrome.

The triad of West syndrome is:

• Infantile spasms (also called Salaam spells)

• Developmental regression

• Hypsarrhythmia (abnormal EEG)

West syndrome is treated with:

• ACTH injections

• Vigabatrin if associated with tuberous sclerosis

Solution to Question 20:

The given scenario is of infantile spasms, developmental regression and the EEG showing hypsarrhythmia is suggestive of West syndrome. The drug of choice for this condition is ACTH.

West syndrome is a rare type of epilepsy syndrome that begins in infancy. It is characterized by a triad of

• Infantile spasm (also called salaam spells) is a convulsive disease characterized by brief, symmetric axial muscle contraction (neck, trunk, and/or extremities).

• Developmental regression

Sold by @itachibot

• Hypsarrhythmia (Himalayan wave pattern) on EEG: Characterized by a high-voltage, slow, chaotic background with multifocal spikes

Treatment of West syndrome includes:

• ACTH

• Vigabatrin if associated with tuberous sclerosis

Solution to Question 21:

The given image shows a Shagreen patch, which is a typical feature of tuberous sclerosis. Vigabatrin is effective in the control of seizures in patients with tuberous sclerosis.

Tuberous sclerosis (Bourneville disease) is an autosomal dominant disease caused by mutations in either the TSC1 or the TSC2 gene. The clinical features seen are as follows:

• Seizures

• Intellectual disability

• Adenoma sebaceum (facial angiofibromas)

• Shagreen patch

• Hypomelanotic macules

• Periungual fibromas

• Renal angiomyolipomas

• Cardiac rhabdomyomas.

A Shagreen patch is an irregularly thickened, slightly elevated, soft, skin-colored plaque, usually in the lumbosacral region.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

The given clinical scenario of focal seizure with impaired consciousness which along with hippocampal sclerosis is suggestive of mesial temporal lobe epilepsy.

The following image shows left hippocampal sclerosis:

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 23:

Seizures of mesial temporal lobe epilepsy are not controlled with carbamazepine. The condition is refractory to anticonvulsant therapy but responds well to surgical intervention.

Solution to Question 24:

Oxcarbazepine is known to worsen myoclonic seizures and is therefore contraindicated in it.

Drugs like carbamazepine, oxcarbazepine and phenytoin can worsen the myoclonic, absence, tonic and atonic seizures and are therefore avoided.

Solution to Question 25:

EEG showing <3Hz spike and wave discharge is a characteristic feature of Lennox-Gastaut syndrome.

Lennox-Gastaut syndrome is a rare type of epilepsy syndrome that begins in childhood. It may occur due to various reasons such as developmental anomalies, mutations, trauma, infections, etc. It is characterized by a typical triad of

• Multiple seizure types

• Intellectual disability

• Typical <3 Hz spike and wave pattern on EEG

Treatment includes administration of anti-epileptic drugs such as lamotrigine and sodium valproate. It is frequently refractory to medical therapy and usually has a poor prognosis.

Sold by @itachibot

Note: EEG with symmetric 3 Hz spike and wave pattern is seen in typical absence seizure.

Solution to Question 26:

Glucocerebrosidase infusions are not the treatment for Lafora's disease. There is no approved treatment for Lafora body disease.

The mainstay of management is anti-epileptic drugs, valproate in particular; however, other AEDs such as topiramate, ethosuximide, and phenobarbital can be used to control seizures. Most recently, perampanel, a new AMPA receptor antagonist AED, is shown to be effective.

Lafora body disease is an autosomal recessive neurodegenerative disease and has an adolescent onset. It is associated with polyglucosan intracellular inclusion bodies. It is characterized by:

- Progressive myoclonus
- Epileptic seizures
- Dementia
- Ataxia.

Diagnosis is by DNA sequencing. Death usually occurs within 10 years of the onset of the disease.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 27:

The given scenario is suggestive of a gelastic seizure.

This condition is also known as gelastic epilepsy, and is a rare type of seizure that involves a sudden burst of energy, usually in the form of laughing or crying. It is seen when the seizure focus is within the left prefrontal cortex. Gelastic seizures have been associated classically with hypothalamic hamartomas.

Option A: Kluver Bucy syndrome is characterized by hypersexuality, placidity, a tendency to explore the environment with the mouth, inability to recognize the emotional significance of visual stimuli, and constantly shifting attention, called hypermetamorphosis.

Option B: Witzelsucht is a tendency to joke and make puns.

Option C: Pseudobulbar affect is defined by involuntary, sudden, and frequent episodes of laughing or crying that are generally out of proportion or inappropriate to the situation.

Solution to Question 28:

The given clinical scenario reveals multiple types of seizures such as atonic seizures (sudden falls), absence seizures (blank stares and unresponsive), and generalized tonic-clonic seizures in this patient. This is a characteristic feature of Lenox-Gastaut syndrome.

Lennox-Gastaut syndrome is a rare type of epilepsy syndrome that begins in childhood. It may occur due to various reasons such as developmental anomalies, mutations, trauma, infections, etc.

Sold by @itachibot

It is characterized by a typical triad of

• Multiple seizure types

• Intellectual disability

• Typical <3 Hz spike and wave pattern on EEG

Treatment includes administration of anti-epileptic drugs such as lamotrigine and sodium valproate. It is frequently refractory to medical therapy and usually has a poor prognosis.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Ischemic Stroke

Question 1:

Ischemia occurs in the brain when the cerebral blood flow falls below _____.

a) 40 mL/100 g/min

b) 20 mL/100 g/min

c) 30 mL/100 g/min

d) 50 mL/100 g/min

Question 2:

A patient presented with acute onset of weakness of the left arm which resolved completely in 10 days. He is known to have mitral stenosis, hypertension and dyslipidemia. What is the most likely diagnosis?

a) Transient ischemic attack

b) Embolic stroke

c) Hemorrhagic stroke

d) Thrombotic stroke

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A patient presented with right-sided hemiparesis that completely resolved in 4-6 hours. What score will you use to determine the risk of a future stroke in this patient?

a) CHADS2-VASc

b) CURB-65

c) ABCD2

d) APACHE II

Question 4:

A 67-year-old patient with hypertension, diabetes, and asthma is found to have an irregularly irregular pulse during a routine physical examination. ECG findings are shown below. How is

Sold by @itachibot

the risk of stroke determined in this patient?



a) CHA2DS2-VASC

b) NIH Stroke Score

c) ABCD2

d) Modified GCS

Question 5:

A 73-year-old woman who was brought to the ER with hemiparesis was found to have a lacunar infarction. What causes this condition?

a) Emboli in small vessels

b) Thrombosis in large vessels

c) Lipohyalinosis of penetrating arteries

d) Hemorrhagic stroke

Question 6:

Which of the following is not caused by a lacunar stroke?

a) Pure motor hemiparesis

b) Pure sensory stroke

c) Ataxic hemiparesis

d) Crural monoplegia with urinary incontinence

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 7:

A patient is diagnosed with an ischaemic stroke. His son says that he holds the comb incorrectly while combing his hair. During the evaluation, he was unable to button his shirt when requested. The lesion is most likely to be in which lobe?

a) Dominant frontal lobe

b) Dominant parietal lobe

c) Non-dominant frontal lobe

d) Non-dominant parietal lobe

Question 8:

Which of the following features is not likely to be seen in a patient with a temporal lobe lesion?

a) Neglect

b) Superior quadrantanopia

c) Auditory hallucinations

d) Poor memory

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 43-year-old man is brought to the hospital due to extremely aggressive behaviour towards his family. His family members also say that he has become forgetful and has developed urinary incontinence. Which of the following lobes is likely to be involved in his disease process?

a) Frontal lobe

b) Dominant parietal lobe

c) Temporal lobe

d) Non-dominant parietal lobe

Question 10:

In the sagittal section of the brain given below if the coloured area is affected, which of the following is not seen?

Sold by @itachibot



©MARROW

a) Urinary incontinence

b) Gait abnormality

c) Perianal anesthesia

d) Fecal incontinence

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 11:

Which type of aphasia is seen in the lesions of the posterior part of the superior temporal gyrus?

a) Fluent

b) Non fluent

c) Anomic

d) Conduction

Question 12:

A 60-year-old patient with atrial fibrillation complains of sudden onset weakness of the right lower limb, with urinary and faecal incontinence. MRI shows an ischaemic stroke involving the left anterior cerebral artery. Which of the following features would you not expect to see in this patient?

a) Ataxia

b) Prosopagnosia

c) Abulia

Sold by @itachibot

d) Paratonic rigidity

Question 13:

Infarction of area of distribution of which of the following arteries is most likely to be seen in a patient with hemiplegia?

a) Vertebral artery

b) Middle cerebral artery

c) Anterior spinal artery

d) Posterior cerebral artery

Question 14:

Which of the following features is not expected to be seen in a patient with a right middle cerebral artery infarct?

a) Aphasia

b) Hemiparesis

c) Facial weakness

d) Dysarthria

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 15:

Visual field examination of a 52-year-old stroke patient reveals the finding shown in the image. During examination, you notice that he is not able to comprehend your instructions and sometimes gives irrelevant replies. What is the likely site of arterial occlusion?

Sold by @itachibot



©MARROW

a) Left MCA stem

b) Right M1 segment of MCA

c) Left M2 segment of MCA – inferior division

d) Right M2 segment of MCA – superior division

Question 16:

A patient is found to have Wernicke's aphasia without motor involvement. Where is the infarct likely to be?

a) ACA territory of the right hemisphere

b) ACA territory of the left hemisphere

c) MCA territory of the right hemisphere

d) MCA territory of the left hemisphere

Question 17:

Which of the following is not a feature of an anterior choroidal artery lesion?

a) Aphasia

b) Contralateral hemiplegia

c) Hemianesthesia

d) Homonymous hemianopia

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Sold by @itachibot

Question 18:

A man is rushed to the ER after losing control of his car and crashing into a tree. Examination findings are given below. What is the most likely cause for his presentation?

a) Spinal cord trauma

b) Morphine poisoning

c) Pontine hemorrhage

d) Ponto-medullary laceration

Question 19:

A patient presents with left-sided facial paralysis and left-sided limb weakness for the past 1 hour. Her blood pressure is 160/100 mm Hg and her CT appears normal. What would be the next step of management?

a) Nothing, since CT was normal

b) Start on aspirin + clopidogrel

c) Intravenous thrombolysis

d) Advice BP control

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Question 20:

Thrombolysis is contraindicated in which of the following patients with ischaemic stroke?

a) Patients 1, 2 and 4

b) Patient 2 only

c) Patients 3 and 5

d) Patient 3 and 4

Question 21:

A patient with acute ischemic stroke requires IV thrombolysis. His symptoms had begun at 6:30 PM and he was brought to the hospital by 7:00 PM. For this treatment to be beneficial for the patient, it should be administered within what time?

a) 7:30 PM

b) 9:00 PM

Sold by @itachibot

c) 10:30 PM

d) 11:00 PM

Question 22:

Which of the following features of stroke need not be treated?

a) Dysphagia

b) Spasticity

c) Fever

d) Neglect

Question 23:

A woman on oral contraceptive pills presents with a headache for 2 days and paraparesis. What could be the most likely cause?

a) Posterior reversible encephalopathy syndrome

b) Cerebral venous thrombosis

c) Complicated migraine

d) Subarachnoid hemorrhage

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 24:

What is the treatment of choice for a patient diagnosed with sagittal sinus thrombosis?

a) Aspirin

b) Heparin

c) Warfarin

d) rtPA

Question 25:

A 40-year-old patient presents with headache and weakness of the right side of the body. He has difficulty in speaking also. Angiography shows a puff of smoke appearance. Which of the following statements regarding this disease is false?

a) It is an occlusive disease involving large intracranial arteries

Sold by @itachibot

b) Vascular inflammation is absent

c) Dawson's fingers are the characteristic imaging finding

d) Rich collateral circulation is present

Question 26:

Which among the following is not a component of CHA2DS2-VASc score?

a) Age > 60 years

b) Hypertension

c) Diabetes

d) Transient ischemic attack

If you bought this from someone else, you've been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	a
5	c
6	d
7	b
8	a
9	a
10	c
11	a
12	b
13	b
14	a
15	c
16	d
17	a
18	c
19	c

Sold by @itachibot

20	c
21	d
22	d
23	b
24	b
25	c
26	a

Detailed Explanations

Solution to Question 1:

A decrease in the cerebral blood flow below 20 mL/100 g/min causes reversible ischemia.

If this state is prolonged for several hours to days, it can lead to infarction. A decrease in CBF to 0 causes the death of the brain tissue within 4–10 min.

Normal cerebral blood flow (CBF) in an adult brain at rest is 50–55 mL/100 g/min.

Solution to Question 2:

This patient with mitral stenosis and acute onset weakness of the left arm is most likely to have had an embolic stroke, which is a type of ischemic stroke. Embolic strokes tend to occur suddenly with maximum neurologic deficit present at onset.

Mitral stenosis is known to cause atrial fibrillation and is a common cause of cardioembolic stroke.

Transient ischemic attack (TIA) is a temporary, focal neurological deficit of sudden onset (stroke-like symptoms) lasting for less than 24 hours. This occurs due to focal ischemia of the brain, retina or cochlea. Most TIAs last for less than 1 hour to a few minutes.

There is a high risk of developing an ischemic stroke after a TIA, especially for the first 3 months following a TIA.

Solution to Question 3:

This patient's clinical features are suggestive of a transient ischemic attack. The risk of stroke following TIAs is determined by the ABCD2 score.

CHADS2-VASc score is used to determine the risk of stroke in a patient with atrial fibrillation.

CURB-65 is a scoring system used to predict mortality in patients with community-acquired pneumonia.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

APACHE II is a scoring system used to predict ICU mortality.

Solution to Question 4:

The given scenario along with the ECG findings are suggestive of atrial fibrillation. The risk of stroke in this condition is determined by CHA2DS2-VASC score.

The ECG show irregular QRS complexes of normal morphology and absent P waves.

Solution to Question 5:

Lacunar infarcts, also called small vessel stroke, refer to infarction following atherothrombotic or lipohyalinotic occlusion of small penetrating arteries in the brain. These penetrating arteries arise from the stem of the middle cerebral artery, circle of Willis, and basilar and vertebral arteries.

The principal risk factors are hypertension and old age.

The image below shows two lacunar infarctions on a CT.

If yo... by @itachibot ...from someone else, ...scammed.

Solution to Question 6:

Crural monoplegia with urinary incontinence is not caused by a lacunar stroke. It is caused by the occlusion of A2 segment of the anterior cerebral artery. Crural monoplegia refers to a monoplegia of the lower limbs.

The types of lacunar syndromes are as follows:

• Pure motor hemiparesis - infarct in the posterior limb of the internal capsule

Sold by @itachibot

• Pure sensory stroke - infarct in the thalamus

• Ataxic hemiparesis - infarct in the pons or internal capsule

• Dysarthria and clumsy hand syndrome - infarction in the pons or internal capsule.



The ACA is divided into two segments:

• The pre-communal (A1) circle of Willis, or stem, which connects the internal carotid artery to the anterior communicating artery

• The post-communal (A2) segment distal to the anterior communicating artery



Solution to Question 7:

Sold by @itachibot

The given scenario where the patient is unable to perform learned skilled movements is suggestive of apraxia. The lesion is likely to be in the dominant parietal lobe as apraxia is a feature of a lesion in this area.

The features of dominant parietal lobe lesions are as follows:

• Apraxia - inability to perform learned skilled movements

• Acalculia - difficulty performing simple mathematical tasks

• Alexia - inability to recognize or read written words or letters

• Agnosia - inability to recognize and identify objects, persons, or sounds

• Dysphasia - language disorder marked by deficiency in the generation of speech and its comprehension

• Contralateral inferior homonymous quadrantanopia

• Gerstmann syndrome (acalculia, dysgraphia, finger agnosia, right-left confusion)

Features of non-dominant parietal lobe lesions include:

• Spatial disorientation

• Constructional apraxia

• Contralateral inferior homonymous quadrantanopia

• Contralateral hemineglect.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

Neglect is not a feature of temporal lobe lesion. It is a feature of a non-dominant parietal lobe lesion.

Features of dominant temporal lobe lesions:

• Complex hallucinations - visual, auditory or olfactory

• Memory impairment

• Complex partial seizures

• Wernicke's aphasia

• Contralateral superior quadrantanopia

Features of non-dominant temporal lobe lesions:

• Poor non-verbal memory

• Loss of musical skills

• Complex hallucinations

• Contralateral superior quadrantanopia.

Sold by @itachibot

Solution to Question 9:

Antisocial behavior, urinary incontinence, and impaired memory in this patient are suggestive of frontal lobe involvement.

Features of frontal lobe lesions:

• Disinhibition

• Lack of initiative

• Antisocial behavior

• Impaired memory

• Incontinence

• Grasp reflexes

• Anosmia.

Solution to Question 10:

Paracentral lobule (PCL) is the continuation of the precentral and postcentral gyri of the superolateral surface into the medial surface of cerebral hemisphere. It controls the motor and sensory innervations of the contralateral lower extremity. It is also responsible for cortical control of micturition and defecation.

The anterior two-thirds of PCL belong to Brodmann area 4, it reflects the primary motor cortex for the muscles of contralateral leg, foot, and perineum,it is functionally important in control of bladder and anal sphincters. Therefore, it's called the cortical center of defecation and micturition.

The posterior third of the PCL contains 3,2 and 1, it reflects the primary somatosensory representation of the leg and foot.

The A2 segment of the anterior cerebral artery supplies the paracentral lobule. Lesions in the paracentral lobule lead to:

• Weakness and loss of sensation in contralateral lower limb

• Urinary incontinence

• Fecal incontinence.

Solution to Question 11:

Lesions of the posterior part of the superior temporal gyrus cause Wernicke's or Fluent aphasia due to the involvement of the Wernicke's area. The speech is full of jargon and neologisms that make little sense (area no 22).

Broca's (Option B) or motor/non-fluent aphasia is seen in lesions of the inferior frontal gyrus which contains the motor speech area. Speech is non-fluent and interrupted, with a paucity of functional words.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Comprehension of written language and the ability to name objects are compromised in anomic aphasia (Option C). This aphasia is caused by damage to the angular gyrus that disrupts visual information processing and transmission to the Wernicke area.

Conduction aphasia (Option D) is seen in lesions of the arcuate fasciculus, the bundle of nerve fibers connecting the Wernicke and Broca's areas. Repetition of words is impaired and paraphasias are observed, though not as much as in Wernicke aphasia.

Solution to Question 12:

Prosopagnosia, also called face blindness, is the inability to recognize faces including one's own image. This occurs due to occipitotemporal lobe damage, seen in posterior cerebral artery (PCA) strokes involving cortical branch occlusion.

It is not seen in anterior cerebral artery strokes.

The anterior cerebral artery (ACA) supplies parts of the frontal lobe, corpus callosum, paracentral lobule, and other deeper structures.

Signs and symptoms of ACA occlusion and structures involved:

- Urinary incontinence, fecal incontinence
- Paratonic rigidity
- Abulia (Akinetic mutism), lack of spontaneity, slowness
- Gait impairment
- Paralysis of contralateral foot and leg
- Paresis of the contralateral arm
- Cortical sensory loss over the contralateral lower limb
- Dyspraxia of left limbs, tactile aphasia in left limbs.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 13:

Hemiplegia is most commonly associated with infarction of the area of distribution of middle cerebral artery.

Middle cerebral artery supplies the motor cortex responsible for sending the output to the motor neurons. Infarction in this territory leads to hemiplegia.

Infarction of area of distribution of vertebral, anterior spinal and posterior cerebral arteries can also lead to hemiplegia.

Sold by @itachibot





Solution to Question 14:

Aphasia is not a feature of right middle cerebral artery infarct (MCA).

Aphasia occurs when there is damage to the dominant hemisphere, i.e., left hemisphere. The dominant hemisphere contains the language network which includes Broca's area, Wernicke's area, their interconnections and connections with frontal, parietal and temporal areas.

The MCA originates from the internal carotid artery. The stem of the MCA is referred to as the M1 segment. It gives rise to cortical and penetrating/lenticulostriate branches.

MCA stem occlusion leads to :

• Contralateral hemiplegia

• Hemianesthesia

Sold by @itachibot

• Homonymous hemianopia

• Global aphasia (dominant hemisphere)

• Anosognosia, constructional apraxia, neglect (non-dominant hemisphere)

Occlusion of the lenticulostriate branches of MCA (lacunar stroke) may affect the genu of the internal capsule. This can lead to dysarthria with contralateral clumsy hand (fine motor weakness).

Solution to Question 15:

The given scenario is suggestive of homonymous right superior quadrantanopia (pie in the sky appearance) and sensory aphasia. These are seen in occlusion of the inferior division of the opposite side middle cerebral artery (MCA).

Parts of the middle cerebral artery:

• The MCA originates from the internal carotid artery. The stem of the MCA is referred to as the M1 segment.

• In the Sylvian fissure, the MCA divides into superior and inferior divisions. These branches are referred to as the M2 segments of the MCA.

Occlusion of the following sites and their effects are given below:

• MCA stem (M1) - leads to contralateral hemiplegia, hemianesthesia and homonymous hemianopia amongst other features

• Superior division of MCA (M2) on the dominant side - leads to motor aphasia (Broca's) and facio-brachial weakness

• Inferior division of MCA (M2) on the dominant side - leads to sensory aphasia (Wernicke's) and contralateral superior quadrantanopia

The following image of coronal section of cerebral hemisphere shows the segments of MCA:



If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Solution to Question 16:

The presence of Wernicke's aphasia without motor involvement suggests an infarct in the MCA territory of the left (dominant) hemisphere.

Wernicke's area is located in the superior temporal gyrus in the dominant hemisphere (mostly the left side). This area is supplied by the inferior division of MCA.

MCA divides into superior and inferior divisions in the Sylvian fissure. These branches are referred to as the M2 segments of MCA.

• The superior division - supplies the frontal and the superior parietal cortex.

• The inferior division - supplies the inferior parietal and temporal cortex.

An infarct in the territory of the inferior division of MCA in the dominant hemisphere causes Wernicke's aphasia without weakness. As the frontal cortex (containing the motor area) is supplied by the superior division and is not affected, weakness is not observed.

An infarct in the inferior division of MCA of the non-dominant hemisphere (mostly the right side) causes hemineglect without weakness.

Solution to Question 17:

Aphasia (dominant hemisphere), constructional apraxia and neglect (non-dominant hemisphere) are seen in MCA stem occlusion. These features are absent in anterior choroidal artery occlusion.

Features common to both MCA and anterior choroidal artery occlusion include:

• Contralateral hemiplegia

• Hemianaesthesia

• Homonymous hemianopia.

Solution to Question 18:

The given scenario is suggestive of pontine haemorrhage. Clinical features of pontine haemorrhage are:

• Deep coma

• Quadriparesis and multiple cranial nerve dysfunction

• Decerebrate rigidity

• Pin-point pupils

• Hyperpnea, hyperpyrexia, severe hypertension, and hyperhidrosis

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Option A: The presence of hyperpyrexia and pin-point pupils rule out the possibility of spinal cord trauma.

Option B: Morphine poisoning is a differential diagnosis for pin-point pupils. But it leads to slower respiratory rate, not tachypnea.

Option D: In ponto-medullary laceration, medullary involvement would cause respiratory depression, leading to apnea. But here tachypnea is noted.

Solution to Question 19:

Sudden onset hemiparesis with apparently normal CT indicates ischemic stroke in this patient. Intravenous thrombolysis(alteplase 0.9 mg/kg) should be immediately administered to patients with ischemic stroke who arrive in the window period i.e., within 3-4.5 hours of symptom onset.

Aspirin + clopidogrel(Option B) is used in secondary prevention of stroke and also to prevent stroke following TIA.

Transient ischemic attacks (TIAs) refer to episodes of stroke symptoms that have a brief duration, the standard definition of duration is <24 h, but most TIAs last <1 hr. TIAs may arise from emboli to the brain or from in situ thrombosis of an intracranial vessel. With a TIA, the occluded blood vessel reopens and neurologic function is restored. If a relevant brain infarction is identified on brain imaging, the clinical entity is now classified as stroke regardless of the duration of symptoms.

The risk of stroke following a transient ischemic attack (TIA) is approximately 10-15% in the first three months, majorly in the first two days. A reliable estimation of this risk can be obtained using the ABCD2 score.

Other options:

Option A: If CT is normal it only rules out hemorrhagic stroke and changes of ischemic stroke usually take time to appear on NCCT.

Option D: BP control is done when BP exceeds 220/120 mmHg, if there is malignant hypertension or concomitant myocardial ischemia, or if blood pressure is >185/110 mmHg and thrombolytic therapy is anticipated.

Solution to Question 20:

A sustained blood pressure > 185/110 mmHg and recent myocardial infarction are contraindications for initiating thrombolysis in ischemic stroke.

Solution to Question 21:

IV thrombolysis in acute ischemic stroke has been proven to be beneficial up to 4.5 hours after the onset of symptoms according to European Cooperative Acute Stroke Study (ECASS) III.

If you purchased this from someone else, you may have been Scammed. Sold by @itachibot

Sold by @itachibot

According to MoHFW (India), thrombolysis is to be carried out within 4.5 hours of symptom onset.

IV tPA should be administered to all eligible acute stroke patients within 3 hours of last known normal and to a more selective group of eligible acute stroke patients (based on ECASS III exclusion criteria) within 4.5 hours of last known normal.

Solution to Question 22:

Neglect is a feature of stroke that need not be treated. Although neglect indicates a poor prognosis, most patients recover without any treatment in 1-6 months.

Dysphagia(Option A) poses a high risk of aspiration and hence can't be left untreated. Spasticity(Option B) can be treated with baclofen, tizanidine. Untreated, it creates a psychosocial impact on the patient. Fever(Option C) is deleterious in stroke patients and must be treated.

Neglect can arise from injury to cortical-limbic-reticular network, that directs attention and integrates the localization and identification of a stimulus. Neglect is associated with poor scores and slower recovery, but severe neglect is rare beyond 6 months. Recovery is rapid in the first 2 weeks, regardless of the side of stroke. By 3 months, patients might have some residual visual neglect. The choice of an intervention depends on the mechanism of neglect, the patient can be treated for an underaroused right hemisphere which has difficulty processing sensory inputs.

Neurogenic dysphagia is an important cause of pulmonary infection in any patient with stroke, motor neuron disease, even transient dysphagia can lead to malnutrition, dehydration, aspiration pneumonia, and airway obstruction with asphyxia, it increases the risk of death and institutionalization. Early placement of nasogastric tube to feed patients after stroke has improved nutrition and reduced deaths.

Spasticity is seen in less than 20% of stroke patients. Patients with bilateral weakness from hypoxic injury are likely to have signs of spasticity. Exaggerated cutaneous and autonomic reflexes and involuntary spasms are disabling problems associated with spasticity. Physical modalities like slow stretching movements and daily passive exercises will reduce symptoms of spasticity and risk of contractures. Baclofen, tizanidine, and clonidine are used in reducing clonus and extensor spasms.

Solution to Question 23:

The presence of headache and paraparesis in this patient with a history of oral contraceptive use is suggestive of cerebral venous thrombosis.

Cerebral venous thrombosis is commonly seen in hypercoagulable states such as:

- Oral contraceptive pill use
- Pregnancy and puerperium
- Inflammatory bowel disease
- Dehydration

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Intracranial infections.

It presents with:

• Headache (most common)

• Seizures

• Intracranial Hypertension

• Focal neurological Deficits

• Altered Mental status

MR venography is used for diagnosis.

The below image shows the empty delta sign seen on CECT or MRI with contrast in patients with sagittal sinus thrombosis.

Solution to Question 24:

IV heparin is the drug of choice in cerebral venous thrombosis. It prevents further thrombosis and reduces venous hypertension and ischemia.

Anticoagulation is often continued indefinitely if thrombophilia is diagnosed.

If there is no underlying hypercoagulable state, treatment with vitamin K antagonists (VKAs) for 3–6 months may be initiated and then converted to aspirin, depending on the degree of resolution of the venous sinus thrombus.

Treatment of CVT:

• Acute treatment - IV Heparin or LMWH

• Maintenance therapy:

• Warfarin for at least 3 months

Sold by @itachibot

• In pregnant women with a history of CVT - anticoagulation with LMWH throughout pregnancy and till 8 weeks postpartum.

Solution to Question 25:

This patient's clinical features are suggestive of Moyamoya disease and Dawson's fingers is not a finding in this condition.

Dawson fingers are a radiographic feature typically seen in multiple sclerosis (MS)

The picture below demonstrates Dawson fingers.



Moyamoya disease is an occlusive disease involving large intracranial arteries. It has a bimodal age distribution and is seen in the first and fourth decades:

• Childhood moyamoya is characterized by ischemic manifestations

• Adult moyamoya is characterized by hemorrhagic manifestations

The first symptom is a stroke or recurrent TIA.

Treatment is by surgical bypass of extracranial carotid arteries to the dura or middle cerebral artery. This may prevent stroke and haemorrhage.

The below image shows the MR angiography findings in a patient with Moyamoya disease.

Sold by @itachibot

Note: As this is a chronic condition, the lenticulostriate arteries develop a rich collateral circulation around the occlusive lesion. This is seen as a 'puff of smoke' (moyamoya in Japanese) on X-ray angiography.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 26:

Age > 60 years is not a component of the CHADS2-VASc score. Age > 75years and age group 65-75 yrs are the two age criteria considered in the score.

The CHA2DS2-VASc score is used to assess the risk of stroke in patients who have atrial fibrillation. Anticoagulation therapy is given when the score is ≥2. Patients with a score of 1 may also be considered for anticoagulation therapy.

Anticoagulation therapy is given for all the patients of AF having mitral stenosis, hypertrophic cardiomyopathy, or a history of stroke.

Sold by @itachibot

Brainstem Syndromes and Hemorrhagic Stroke

Question 1:

A patient has developed confusion and keeps bumping into objects. He can speak fluently. But on examination, he cannot differentiate between fingers and cannot write. He also gets confused with the examiner's right and left sides. What is the most likely diagnosis?

a) Gerstmann syndrome

b) Anton syndrome

c) Millard Gubler syndrome

d) Dejerine Roussy syndrome

Question 2:

A 78-year-old woman presents with right hemisensory loss followed by agonizing, burning pain in the affected regions. The pain responds poorly to analgesics. These features are classically seen in which of the following conditions?

a) Weber's syndrome

b) Parinaud's syndrome

c) Dejerine-Roussy syndrome

d) Millard-Gubler syndrome

Question 3:

A 73-year-old woman was brought to the OPD since she has been tripping over things at home. On entering, she complimented the doctor's tie even though he was not wearing one and continued to insist that she could see normally. On examination, bilateral blindness was present but pupillary light reflex was normal. What is the likely diagnosis?

a) Anton's syndrome

b) Balint's syndrome

c) Parinaud's syndrome

d) Riddoch phenomenon

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Question 4:

A patient with a history of ischemic heart disease now comes to the hospital due to problems with vision. She complains that while turning her head, the image of the previous target still persists for a few seconds before she can see the new target. This phenomenon is seen in which of the following conditions?

a) Anton's syndrome

b) Balint's syndrome

c) Parinaud's syndrome

d) Claude's syndrome

Question 5:

Which of the following features do you not expect to see in a patient with Balint's syndrome?

a) Opsoclonus

b) Optic ataxia

c) Simultanagnosia

d) Ocular apraxia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A patient presents with ipsilateral oculomotor palsy and contralateral hemiplegia. What is the diagnosis?

a) Anton's syndrome

b) Benedikt syndrome

c) Claude's syndrome

d) Weber's syndrome

Question 7:

Which among the following is wrong regarding Weber's syndrome?

a) Contralateral hemiplegia

b) Ipsilateral oculomotor nerve palsy

c) Contralateral Parkinsonism

Sold by @itachibot

d) Ipsilateral paralysis of lower face

Question 8:

A patient presented with loss of upgaze and convergence-retraction nystagmus. What is the likely diagnosis?

a) Nothnagel's syndrome

b) Balint's syndrome

c) Weber's syndrome

d) Parinaud's syndrome

Question 9:

A patient presented with left-sided hemiplegia and examination revealed right-sided abducens nerve palsy and facial nerve palsy. Further investigation revealed a right-sided pontine haemorrhage. What is the most likely diagnosis?

a) Marie-Foix syndrome

b) Raymond's syndrome

c) Millard–Gubler syndrome

d) Foville's syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following differentiates Foville's syndrome from Millard-Gubler syndrome?

a) Contralateral hemiplegia

b) Ipsilateral hemiplegia

c) Ipsilateral facial palsy

d) Lateral gaze palsy

Question 11:

Thrombosis of posterior inferior cerebellar artery causes _____.

a) Lateral medullary syndrome

b) Weber syndrome

Sold by @itachibot

c) Medial medullary syndrome

d) Millard Gubler syndrome

Question 12:

Wallenberg syndrome is most commonly caused by thrombosis of which of the following arteries?

a) Vertebral artery

b) Posterior inferior cerebellar artery

c) Anterior inferior cerebellar artery

d) Basilar artery

Question 13:

Which of the following features do you not expect to see in a patient with Wallenberg syndrome?

a) Ipsilateral ataxia

b) Contralateral hemiplegia

c) Vertigo

d) Impaired sensation on ipsilateral half of face

Question 14:

Which of the following features would you not expect to see in a patient with medial medullary syndrome?

a) Contralateral hemiparesis

b) Ipsilateral paralysis of tongue

c) Contralateral loss of proprioception

d) Horner's sydrome

Question 15:

Which is the most common site of bleed in patients with intracerebral hypertensive haemorrhage?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Thalamus

b) Subcortical white matter

c) Pons

d) Putamen

Question 16:

Raised intracranial tension is not associated with which of the following manifestations?

a) Seizures

b) Blurring of vision

c) Paraparesis

d) Abducent paralysis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

Which of the following is a component of Cushing's triad?

a) Bradycardia

b) Hypotension

c) Loss of consciousness

d) Papilledema

Question 18:

A 62-year-old man presents to the casualty with a severe headache, fever, and vomiting for the past 5 hours. There is no history of trauma. CT head is shown below. What is the most common cause for his presentation?

Sold by @itachibot



a) Saccular aneurysms

b) Hypertension

c) Microaneurysms

d) Atherosclerosis

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 19:

Which of the following structures is most likely to be compressed by an aneurysm of the posterior communicating artery?

a) Oculomotor nerve

b) Optic nerve

c) Trochlear nerve

d) Abducens nerve

Question 20:

At which of the following locations does a Berry aneurysm have the highest risk of rupture?

a) Origin of posterior communicating artery

b) Origin of anterior cerebral artery

c) Middle cerebral artery

d) Anterior communicating artery

Sold by @itachibot

Question 21:

The CT finding of a patient is shown below. What is the most common cause for this condition?

a) Rupture of berry aneurysm

b) Injury to bridging dural veins

c) Rupture of cerebral AV malformation

d) Injury to middle meningeal artery

Question 22:

The CT scan of a patient who was brought to the casualty after a road traffic accident is shown below. Between what structures does blood accumulate in this condition?

If you purchased this from elsewhere, you may have been scammed.

Sold by @itachibot

a) Outer table and inner table of the skull

b) Outer table and dura mater

c) Inner table and dura mater

d) Periosteum and aponeurotic layer of scalp

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 23:

Which of the following statements is false regarding extradural hemorrhage?

a) CT shows crescentic opacity

b) Lucid interval is characteristic

c) Middle meningeal artery bleed is the most common source

d) Most common cause is trauma

Question 24:

A 50-year-old hypertensive man presented with an acute-onset headache for the past 2 hours, neck stiffness, nausea, and vomiting. He claims it is the worst headache of his life. There are no focal neurological deficits and his CT scan shows blood in the Sylvian fissure. What is the most probable diagnosis?

a) Extradural hemorrhage

b) Subarachnoid hemorrhage

c) Subdural haemorrhage

Sold by @itachibot

d) Hypertensive bleed

Question 25:

A 53-year-old man was admitted with a history of a cerebrovascular accident. On the second day, he becomes drowsy with minimal response. CT finding is shown in the image below. What will be your next line of management?

a) Aspirin and clopidogrel

b) Mannitol

c) Decompressive surgery

d) Mechanical thrombectomy

Question 26:

A 62-year-old man presents to the ER with headache, fever, and vomiting for the past 5 hours. A CT image is shown below. Physical examination shows positive Kernig's and Brudzinski's signs. Vitals: Temp - 101 F, BP - 120/88 mmHg, PR - 110 beats/min, pupils reactive. What is the next best step?

Sold by @itachibot



a) Administer IV ceftriaxone

b) Administer IV tenecteplase

c) Craniotomy

d) Surgical clipping

Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	b
5	a
6	d
7	d
8	d
9	c
10	d
11	a
12	a
13	b
14	d

Sold by @itachibot

15	d
16	c
17	a
18	a
19	a
20	a
21	b
22	c
23	a
24	b
25	c
26	d

Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

The most likely diagnosis for this patient is Gerstmann syndrome. It is commonly associated with damage to the left (dominant) inferior parietal lobule.

Gerstmann syndrome consists of:

• Acalculia - impairment of simple arithmetic

• Dysgraphia - impaired writing

• Finger anomia - an inability to name individual fingers

• Right-left confusion - an inability to tell whether a hand, foot, or arm of the patient or examiner is on the right or left side of the body

Option B: Anton syndrome also known as visual anosognosia occurs due to bilateral infarction in the distal PCAs. It produces cortical blindness and the patients are unaware of the blindness or may even deny it.

Option C: Millard-Gubler syndrome occurs due to ventral pontine injury. It causes functional loss of the 6th and 7th cranial nerves and corticospinal tract. This presents as diplopia, ipsilateral facial palsy, ipsilateral loss of corneal reflex, contralateral hemiparesis.

Option D: Dejerine-Roussy syndrome is associated with hemisensory loss and severe burning pain in the affected region. It is also known as thalamic pain syndrome.

Solution to Question 2:

Sold by @itachibot

Hemisensory loss and severe burning pain in the affected regions is classically seen in Dejerine-Roussy syndrome, also known as thalamic pain syndrome.

Dejerine- Roussy syndrome occurs due to occlusion of the thalamic and thalamogeniculate branches arising from the P1 segment resulting in contralateral hemisensory loss and severe burning pain in the affected regions.

The P1 segment of the posterior cerebral artery extends from the basilar bifurcation to the junction with the posterior communicating artery.

This pain responds poorly to analgesics. Anticonvulsants such as carbamazepine or gabapentin or tricyclic antidepressants may be beneficial.

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Solution to Question 3:

The given scenario is suggestive of Anton's syndrome. It occurs due to bilateral infarction in the distal posterior cerebral artery (P2 segment) and is associated with the following features:

• Cortical blindness with preserved pupillary light reaction

• Visual anosognosia - denial of loss of vision

• Confabulation in the setting of obvious visual loss

Option D: Riddoch phenomenon is a condition where cortical blindness is associated with the perception of only moving targets.

Solution to Question 4:

The persistence of visual image even after shifting gaze is called palinopsia. It is seen in Balint's syndrome.

Balint's syndrome results due to low flow in the watershed areas between distal PCA and MCA territories. It is characterized by the following:

• Palinopsia - persistence of a visual image despite gazing at another scene

• Asimultagnosia - difficulty to focus on more than one item in the visual field

• Optic ataxia - Inability to reach accurately under visual guidance

• Ocular apraxia - Inability to direct gaze accurately to a new target

Claude's syndrome is a distinctive brainstem syndrome characterized by ipsilateral third nerve palsy with contralateral hemiataxia. It is due to an intrinsic or extrinsic lesion in the midbrain.

Note: While some authors use the term simultanagnosia, others use asimultanagnosia instead. A few authors also use these two terminologies interchangeably.

Solution to Question 5:

Sold by @itachibot

Opsoclonus is not seen in Balint's syndrome.

The features of Balint's syndrome are:

• Optic ataxia - the inability to guide the arm to reach a certain object in the visual field

• Simultanagnosia - the inability to perceive more than one object at a time. In other words, it is the inability to recognize multiple elements in a visual presentation, one object or some elements of a scene can be appreciated but not the display as a whole.

• Ocular apraxia - the inability to control voluntary purposeful eye movements

• Palinopsia - the persistence of a visual image even after the stimulus has been removed

Note: While some authors use the term simultanagnosia, others use asimultanagnosia instead. A few authors also use these two terminologies interchangeably.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

Ipsilateral oculomotor palsy and contralateral hemiplegia is seen in Weber's syndrome.

Weber's syndrome occurs due to infarction in the distribution of the penetrating branches of the PCA. It affects the cerebral peduncle, especially medially, and damages III cranial nerve and the pyramidal fibers. It results in contralateral hemiplegia and ipsilateral oculomotor palsy.

Claude's syndrome is a distinctive brainstem syndrome characterized by ipsilateral third cranial nerve palsy with contralateral hemiataxia. This is due to an intrinsic or extrinsic lesion in the midbrain.

Solution to Question 7:

Ipsilateral paralysis of the lower face is not seen in Weber's syndrome. However, there is contralateral paralysis of the lower face.

Weber syndrome is caused by infarction in the distribution of the penetrating branches of the posterior cerebral artery (PCA). It affects the cerebral peduncle, especially medially, with damage to the fascicle of cranial nerve III and the pyramidal fibers.

The resultant clinical findings are contralateral hemiplegia of the face, arm, and leg secondary to corticospinal and corticobulbar tract involvement, and ipsilateral oculomotor paresis including a dilated pupil.

Structure damaged	Effect
Cranial nerve III fibres	Ipsilateral ptosisIpsilateral di vergent squintLoss of accom modation reflexLoss of light r eflex
Corticobulbar tract	UMN lesion leading to contra lateral paralysis of the lower f ace

Sold by @itachibot

Structure damaged	Effect
Corticospinal tract	Contralateral hemiplegia



Solution to Question 8:

The presence of convergence-retraction nystagmus and loss of upgaze is seen in Parinaud's syndrome. It is caused by a lesion in the dorsal midbrain.

The following tetrad of symptoms are seen:

- Loss of upgaze
- Light-near dissociation
- Convergence-retraction nystagmus
- Lid retraction

Convergence■retraction nystagmus is irregular, jerky nystagmus in which both eyeballs rhythmically converge and retract into the orbit, particularly on attempting an upward gaze.

Nothnagel syndrome is associated with oculomotor palsy and contralateral limb ataxia. Underlying structures involved are the superior cerebellar peduncle and tectum of the midbrain.

Solution to Question 9:

Ipsilateral abducens and facial nerve palsy with contralateral hemiplegia are seen in Millard–Gubler syndrome.

In this syndrome, there is a ventral pontine injury (lesion in abducens fascicle) that causes lateral rectus palsy.

Sold by @itachibot

CNS lesion	Effects
Millard–Gubler Syndrome	Ipsilateral abducens nerve weaknessIpsilateral facial nerve palsyContralateral hemiplegia
Raymond's Syndrome	Ipsilateral abducens nerve palsyContralateral hemiplegia
Foville's Syndrome	Lateral gaze palsyIpsilateral facial nerve palsyContralateral hemiplegia
Balint's syndrome	Optic ataxiaOcular apraxiaVisuospatial disorientation
Lateral pontine syndrome (Marie-Foix syndrome)	Contralateral hemiparesisContralateral loss of proprioception & vibrationIpsilateral limb ataxiaIpsilateral facial palsyLateral hearing lossVertigo Nystagmus

Sold by @... this from ... else, If you purchased ... may have been scammed.

Solution to Question 10:

Lateral gaze palsy is a feature of Foville's Syndrome which is a result of dorsal pontine injury.

Millard-Gubler syndrome occurs due to ventral pontine injury. Lateral gaze palsy does not occur in this condition because the abducens fascicle is injured rather than the nucleus.

CNS lesion	Effects
Millard–Gubler Syndrome	Ipsilateral abducens nerve weaknessIpsilateral facial nerve palsyContralateral hemiplegia
Raymond's Syndrome	Ipsilateral abducens nerve palsyContralateral hemiplegia
Foville's Syndrome	Lateral gaze palsyIpsilateral facial nerve palsyContralateral hemiplegia
Balint's syndrome	Optic ataxiaOcular apraxiaVisuospatial disorientation
Lateral pontine syndrome (Marie-Foix syndrome)	Contralateral hemiparesisContralateral loss of proprioception & vibrationIpsilateral limb ataxiaIpsilateral facial palsyLateral hearing lossVertigo NystagmusContralateral loss of pain and temperature

Sold by @itachibot

Solution to Question 11:

Thrombosis of the posterior inferior cerebellar artery (PICA) causes lateral medullary syndrome or Wallenberg syndrome.

PICA is a branch of the fourth part of the vertebral artery, which is a branch of the first part of subclavian artery.



Solution to Question 12:

Thrombosis of the vertebral artery is most commonly implicated in Wallenberg syndrome.

Wallenberg syndrome, also called lateral medullary syndrome, is due to occlusion of the ipsilateral vertebral artery in most cases. In the remainder, posterior inferior cerebellar artery occlusion is responsible.

Wallenberg / lateral medullary syndrome affects the following structures:

• Descending tract and nucleus of cranial nerve V - Ipsilateral impaired sensation over half of the face

• IX and X cranial nerves - Ipsilateral paralysis of palate, vocal cord, diminished gag reflex

• Nucleus of tractus solitarius - Ipsilateral loss of taste

• Vestibular nucleus - Nystagmus, diplopia, vertigo, nausea

• Spinocerebellar tract - Ipsilateral ataxia

• Descending sympathetic tract - Ipsilateral Horner's syndrome

• Lateral spinothalamic tract - Contralateral impaired pain and thermal sensation over half of the body.

Sold by @itachibot

Solution to Question 13:

The pyramidal tracts are not affected in Wallenberg syndrome and therefore hemiplegia is not seen.

Wallenberg / lateral medullary syndrome affects the following structures:

• Descending tract and nucleus of CN V - I/L impaired sensation over half of face

• IX and X CN - I/L paralysis of palate, vocal cord, diminished gag reflex

• Nucleus of Tractus Solitarius – I/L loss of taste

• Vestibular nucleus - Nystagmus, diplopia, vertigo, nausea

• Spinocerebellar tract – I/L ataxia

• Descending sympathetic tract – I/L Horner's syndrome

• Lateral spinothalamic tract – C/L impaired pain and thermal sensation over half of the body.

Solution to Question 14:

Horner's syndrome is not a feature of medial medullary syndrome.

Medial medullary syndrome occurs due to the occlusion of the vertebral artery or a branch of the vertebral or basilar artery. The features of medial medullary syndrome are

• CN XII - Ipsilateral paralysis of the tongue

• Pyramidal tract - Contralateral hemiplegia

• Medial lemniscus - Impaired tactile and proprioception over the contralateral half of the body

Horner's syndrome occurs due to a lesion involving the sympathetic fibres. It is seen in lateral medullary syndrome.

Solution to Question 15:

The most common site for intracerebral hypertensive hemorrhage is the putamen.

The most common type of intracranial haemorrhage is an intraparenchymal i.e., intracerebral haemorrhage. It is most often caused by a hypertensive bleed.

Most common sites of intracerebral hypertensive bleed:

• Putamen (basal ganglia)

• Thalamus

• Cerebellum

• Pons.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 16:

Raised ICT is not associated with paraparesis.

Manifestations of raised ICT include:

• Hypertension

• Bradycardia

• Respiratory depression

• Headache, visual blurring due to papilloedema and abducent nerve palsy

• Seizures

Paraparesis is usually caused by a spinal lesion such as thoracic myelopathy and neoplasms causing compression of the spinal cord.

Rare intracranial causes of paraparesis are:

• Parasagittal meningioma

• Occlusion of unpaired anterior cerebral artery

• Cerebral venous thrombosis affecting the sagittal sinus.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

Bradycardia is a component of Cushing's triad.

Cushing's triad is characteristically seen in raised ICT. Its components are as follows:

• Hypertension

• Bradycardia

• Respiratory depression.

Solution to Question 18:

The given scenario is suggestive of non-traumatic subarachnoid haemorrhage. The most common cause for this condition is the rupture of a berry aneurysm, also known as a saccular aneurysm.

CT shows hyperattenuating material filling the subarachnoid space, which is suggestive of SAH. Most commonly, it occurs around the circle of Willis due to the rupture of a berry aneurysm at this location.

Most unruptured intracranial aneurysms are completely asymptomatic. Prodromal symptoms occasionally suggest the site of the enlarging unruptured aneurysm. For example, cranial nerve III palsy with loss of ipsilateral light reflex and focal pain above or behind the eye suggests an expanding aneurysm at the junction of the posterior communicating artery and internal carotid

Sold by @itachibot

artery.

Ruptured aneurysms cause an excruciating thunderclap headache followed by a transient loss of consciousness.

Non-contrast CT scan is the investigation of choice for SAH. LP shows xanthochromatic CSF.

Treatment involves clipping or coiling the aneurysm to eliminate the risk of rebleeding.

Solution to Question 19:

Aneurysm of the posterior communicating artery will most likely compress the oculomotor nerve. This is because the posterior communicating artery is anatomically located very close to the oculomotor nerve.

Patients with an aneurysm of the posterior communicating artery can present with symptoms of oculomotor nerve palsy such as:

• Unilateral ptosis (LPS)

• Outward eye deviation (MR)

• Dilated pupil.

The nerves controlling the pupils are located very superficially and these are always affected due to the mass effect of the aneurysm. Therefore the pupil sparing third nerve palsies are unlikely to be due to an aneurysm.

Pupil-sparing oculomotor nerve palsy is often assumed to be caused by ischemic injuries such as hypertension and diabetes mellitus.

Expanding unruptured aneurysms can occasionally lead to prodromal symptoms which suggest the location of the aneurysm.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 20:

The size and site of aneurysm are important in predicting the risk of rupture. Aneurysms with the following features have a greater risk of rupture:

• Size >7 mm in diameter

• Located at the top of the basilar artery (i.e origin of the posterior cerebral artery)

• Located near the origin of the posterior communicating artery.

The most common location of Berry aneurysm is the anterior communicating artery.

Solution to Question 21:

The given image is suggestive of a subdural hematoma and it most commonly results from injury to the bridging dural veins.

Sold by @itachibot

Subdural hematoma refers to the haemorrhage between the dura and arachnoid mater. The image of the CT scan reveals a hyperdense crescentic lesion. Mass effect with midline displacement can also be seen.

Causes of SDH are:

• Trauma

• Anticoagulants

• Antiplatelets

• Alcohol abuse

• Shaken baby syndrome.

Non-contrast CT is the investigation of choice. Radiographic features of subdural hemorrhage are:

• Typically unilateral

• Crescent-like distribution (concavo-convex) around the periphery of the brain

• Not limited by sutures and can fill dural reflections

• Chronic subdural hemorrhage: Crescent-shaped hypodense lesion on axial CT

• Acute subdural hemorrhage: Hyperdense collections on CT

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

The given CT image is suggestive of extradural haemorrhage (EDH). In this condition, blood accumulates between the inner table of the skull and the dura mater.

The typical presentation is an initial loss of consciousness following trauma, a complete transient recovery (lucid interval), and a rapid progression of neurological deterioration.

This condition is treated by rapid surgical evacuation and ligation or cautery of the damaged vessel.

Solution to Question 23:

In extradural hemorrhage (EDH), CT brain does not show a crescentric opacity. It shows a lentiform opacity.

Solution to Question 24:

This patient's clinical features of sudden-onset headache with neck stiffness, nausea, and vomiting with no neurological deficits is suggestive of a diagnosis of subarachnoid haemorrhage. The presence of blood in the Sylvian fissure confirms this diagnosis.

Sold by @itachibot

Trauma is the most common cause. However, spontaneous sub-arachnoid bleeding occurs most often due to the rupture of a berry aneurysm. Most of them are found in the anterior circulation, mostly on the circle of Willis. The three most common locations are:

• Terminal ICA

• MCA bifurcation

• Top of the basilar artery.

A non-contrast CT scan is the investigation of choice. LP shows xanthochromatic CSF.

Treatment involves clipping or coiling the aneurysm to eliminate the risk of rebleeding.

Solution to Question 25:

The clinical scenario describes a 53 year old patient admitted in view of a cerebrovascular accident developing neurologic deterioration (drowsy with minimal response) on the second day, and CT brain showing significant edema of MCA territory with midline shift. The next line of management is decompressive surgery.

Cerebral edema is a potential complication in 5-10% of patients with strokes, leading to obtundation and brain herniation. The severity of edema is highest on the second or third day after the stroke and can last for around 10 days.

Medical management may not be successful in case of significant edema or infarcts involving the cerebellum.

Decompressive surgery is recommended for patients who are less than 60 years old and experience deterioration ≤ 48 hours, despite receiving medical therapy. Hypodensity on head CT within the first 6 hours of stroke onset, involving ≥ one-third of the MCA territory, early midline shift, and a diffusion-weighted imaging volume on MRI within 6 hours ≥ 80 mL are predictors of severe cerebral edema.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 26:

The scenario is suggestive of subarachnoid hemorrhage (SAH) and the treatment of choice for this condition is surgical clipping.

CT shows hyperattenuating material filling the subarachnoid space, which is suggestive of SAH. Most commonly, it occurs around the circle of Willis due to the rupture of a berry aneurysm at this location.

SAH can present with fever, positive Kernig's sign, and Brudzinski's sign, as well as mimic meningitis or encephalitis.

Treatment with the calcium channel antagonist nimodipine (60 mg PO every 4 h) improves outcome, by preventing ischemic injury.

Option A: Vancomycin + ceftriaxone is the recommended therapy for bacterial meningitis. Meningitis does not show hyper-attenuated subarachnoid space.

Sold by @itachibot

Option B: IV Tenecteplase is used for ischemic stroke without bleeding complications.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Peripheral Neuropathy

Question 1:

Which of the following functions are likely to be impaired in patients with peripheral neuropathy?

a) Sensory or motor function only

b) Sensory or autonomic function only

c) Motor or autonomic function only

d) Sensory, motor or autonomic function

Question 2:

Disorders of autonomic function should be considered in patients with:

a) 2 and 4

b) 3 and 5

c) 1, 2 and 5

d) 1, 3, and 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 50-year-old woman complains of numbness of her hands, generalized weakness, and easy fatiguability. She is a vegetarian. She looks very pale and her peripheral smear reveals macrocytic hypochromic anemia. If untreated, she is at risk of developing which of the following conditions?

a) Proprioceptive loss

b) Stabbing pain

c) Temperature loss

d) Burning pain

Question 4:

Small fiber neuropathies can be seen in patients with all of the following conditions, except:

Sold by @itachibot

a) Sjogren's syndrome

b) HIV

c) Hansen's disease

d) Amyloidosis

Question 5:

A 30-year-old man presents with weakness of both the lower limbs. On examination, there is a sensory loss in the lower limbs and both the proximal and distal leg muscles have reduced power. This presentation is characteristic of which of the following conditions?

a) Acquired immune demyelinating polyneuropathies

b) Amyotrophic lateral sclerosis

c) Mononeuropathy multiplex

d) Diabetic plexopathy

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A patient presented with peripheral neuropathy with asymmetric distal weakness without sensory loss. The differential diagnosis for this condition would include all except:

a) Motor neuron disease

b) Juvenile monomelic amyotrophy

c) Hereditary motor neuropathy

d) Multifocal motor neuropathy

Question 7:

What is the most common type of hereditary neuropathy?

a) Parsonage-Turner syndrome

b) Friedreich's ataxia

c) Charcot-Marie-Tooth disease

d) Familial amyloid polyneuropathy

Question 8:

Sold by @itachibot

All of the following are associated with autonomic neuropathies, except:

a) Diabetes

b) Acquired amyloidosis

c) Hyperthyroidism

d) Porphyria

Question 9:

A young woman presents with complaints of insomnia due to her feet becoming very hot and painful at night. On examination, she has multiple superficial foot ulcers, and she is unable to feel light touch and crude touch. What is your diagnosis?

a) HSAN I

b) Dejerine-Roussy syndrome

c) Riley-Day syndrome

d) Charcot Marie Tooth disease

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following is the most common type of neuropathy seen in diabetes?

a) Cryptogenic sensory polyneuropathy

b) Mononeuropathies

c) Autonomic neuropathy

d) Distal Symmetric Sensory and Sensorimotor Polyneuropathy

Question 11:

Which of the following can be prescribed as a first-line drug for a patient with painful diabetic neuropathy?

a) Carbamazepine

b) Tramadol

c) Duloxetine

d) Venlafaxine

Sold by @itachibot

Question 12:

Sensorimotor neuropathy may be caused by all of the following except:

a) Amiodarone

b) Lead poisoning

c) Arsenic poisoning

d) Isoniazid

Question 13:

Which of the following vitamins will you prescribe for a patient on antitubercular therapy?

a) B9

b) B1

c) B12

d) B6

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 14:

During a regular health checkup, you notice the following nail finding in a 64-year-old man. He lives near a smelting industry on the banks of a river. Which of the following conditions can be seen in this patient?

a) Pure motor neuropathy

Sold by @itachibot

b) Pure sensory neuropathy

c) Predominant sensory neuropathy

d) Sensorimotor neuropathy

Question 15:

Identify the diagnosis.

a) Facial nerve palsy

b) Plexiform neurofibromatosis

c) Masticator space abscess

d) Fibrous dysplasia

If you purch... you may ha... else, ...d.

Answer Key

Question No.	Correct Option
1	d
2	d
3	a
4	a
5	a
6	c

Sold by @itachibot

7	c
8	c
9	a
10	d
11	c
12	b
13	d
14	d
15	b

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Detailed Explanations

Solution to Question 1:

Peripheral neuropathies can impair sensory, motor, or autonomic function, either singly or in combination.

Peripheral nerves are composed of sensory, motor, and autonomic elements.

Peripheral neuropathies are classified into those that primarily affect the

- Cell body - neuronopathy or ganglionopathy
- Myelin - myelinopathy
- Axon - axonopathy

Solution to Question 2:

Disorders of autonomic function (dysautonomia) is usually considered in patients with:

- Unexplained orthostatic hypotension (OH)
- Syncope
- Sleep dysfunction
- Altered sweating (hyperhidrosis or hypohidrosis)
- Impotence
- Constipation or other GI symptoms like bloating, nausea, vomiting of old food, diarrhea.
- Bladder disorders such as urinary frequency, hesitancy, or incontinence.

Solution to Question 3:

Sold by @itachibot

The given clinical scenario of macrocytic anemia and a vegetarian diet is suggestive of vitamin B12 deficiency. This typically leads to large fiber neuropathy which presents with loss of proprioception and vibration.

Smaller-diameter myelinated and unmyelinated fibers transmit pain and temperature sensation. Hence, small fiber neuropathies present as temperature loss or burning or stabbing pain.

Solution to Question 4:

Sjogren's syndrome does not cause small fiber neuropathy. It causes preferential large fiber sensory neuropathy.

Solution to Question 5:

The given clinical scenario suggests symmetric proximal and distal weakness with sensory loss. This pattern is characteristic of acquired immune demyelinating polyneuropathies.

It consists of:

• Acute form - Acute inflammatory demyelinating polyneuropathy [AIDP] also known as Guillain-Barré syndrome [GBS]

• Chronic form - Chronic inflammatory demyelinating polyneuropathy [CIDP]

Option B: Amyotrophic lateral sclerosis (ALS) is a multiple degenerative motor neuron disease. It leads to muscle weakness and disability in the absence of sensory symptoms and signs.

Option C: Mononeuropathy multiplex refers to multiple compressive mononeuropathies. It presents as asymmetric subacute or acute sensory and motor symptoms and signs.

Option D: Polyradiculopathy or plexopathy due to diabetes mellitus presents with asymmetric weakness.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 6:

Hereditary motor neuropathy does not present with asymmetric distal weakness without sensory loss. It presents with symmetric distal weakness without sensory loss.

Solution to Question 7:

Charcot-Marie-Tooth (CMT) disease is the most common type of hereditary neuropathy.

Other examples of rare hereditary neuropathies are Friedreich's ataxia, familial amyloid polyneuropathy and Tangier disease.

Sold by @itachibot

Solution to Question 8:

In hyperthyroidism, excessive thyroxine causes sympathetic autonomic overactivity, not neuropathy.

Hypothyroidism can be associated with peripheral neuropathy, most typically carpal tunnel syndrome. Treatment is the correction of hypothyroidism.

Solution to Question 9:

The given case scenario is suggestive of burning feet syndrome, seen in HSAN 1.

There are five types of hereditary sensory and autonomic neuropathies (HSAN I–V), of which the most important are HSAN I and HSAN III.

HSAN I has an autosomal dominant inheritance. It presents in adolescence or adulthood as a distal small-fiber neuropathy. It causes burning feet syndrome in which patients present with:

• Burning sensation of feet, especially at night

• Sensory loss

• Foot ulcers.

Option B: Dejerine Roussy syndrome, also known as thalamic pain syndrome usually occurs after a stroke. The patient experiences burning pain over affected areas of the body.

Option C: HSAN III or Riley Day syndrome presents in childhood. It presents with labile BP, decreased tearing, hyperhidrosis, reduced sensitivity to pain, and areflexia.

Option D: Charcot Marie Tooth disease is the most common type of hereditary neuropathy. It is a hereditary motor and sensory neuropathy (HMSN) which presents with motor weakness and skeletal deformities.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

Distal symmetric sensory and sensorimotor polyneuropathy (DSPN) is the most common form of diabetic neuropathy and is seen in approximately one-third of patients with DM.

Other neuropathies seen in diabetes mellitus are:

• The most common mononeuropathies are median neuropathy at the wrist and ulnar neuropathy at the elbow

• Diabetic autonomic neuropathy is typically seen in combination with DSPN

• Cranial neuropathy (third nerve palsy is the most common) and diabetic neuropathic cachexia

The Michigan diabetic neuropathy score is a screening tool used to assess diabetic neuropathy.

Solution to Question 11:

Sold by @itachibot

Duloxetine is a selective serotonin-norepinephrine reuptake inhibitor (SSNRI) and is one of the first-line drugs used in the treatment of painful diabetic neuropathy.

Treatment of painful sensory neuropathies:

• First-line:

• Lidoderm 5% patch

• Tricyclic antidepressants (e.g., amitriptyline, nortriptyline)

• Gabapentin, Pregabalin

• Duloxetine

• Second-line:

• Carbamazepine

• Phenytoin

• Venlafaxine

• Tramadol

• Third-line:

• Mexiletine

Optimal glucose control is considered the cornerstone for the treatment of diabetes and its complications. Intensive glucose control has been shown to prevent the development of peripheral neuropathy.

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

Lead poisoning does not cause sensorimotor neuropathy. It causes symptoms and signs of primarily motor neuropathy. The most common presentation of lead poisoning is encephalopathy.

The sensation is generally preserved but the autonomic nervous system can be affected in lead poisoning.

Solution to Question 13:

Isoniazid (INH), which is a first-line anti-tuberculosis drug, inhibits pyridoxal phosphokinase, resulting in pyridoxine deficiency and neuropathy. Prophylactic administration of vitamin B6 (pyridoxine) 10 mg/day can prevent neuropathy from developing.

10 mg daily suffices for prophylaxis but patients with evidence of vitamin B6 deficiency may require up to 50 mg daily.

Current WHO guidelines recommend routine substitution of pyridoxine to patients who are at risk of peripheral neuropathy, e.g., malnutrition, chronic alcohol dependence, diabetics, HIV.

However, pyridoxine excess (acute intoxication or long-term use of large doses) is also found to cause peripheral neuropathy.

Sold by @itachibot

Solution to Question 14:

The image shows Mee's lines, transverse white striae of the fingernails. This along with history is suggestive of chronic arsenic poisoning probably due to a contaminated water source. Arsenic poisoning causes toxic sensorimotor polyneuropathy.

The neuropathy manifests 5–10 days after ingestion of arsenic and progresses for several weeks, mimicking Guillain–Barre's Syndrome.

Solution to Question 15:

The given image showing thickening of tissues in the temporal and orbital regions with coarse overlying skin suggests a diagnosis of plexiform neurofibroma.

Plexiform neurofibroma represents an uncommon variant (30%) of neurofibromatosis 1 (NF1). Plexiform neurofibromas are usually evident at birth and result from a diffuse thickening of nerve trunks and surrounding tissues that are usually located in the orbital or temporal region of the face. The skin overlying it may be coarse and associated with hyperpigmentation.

Plexiform neurofibromas may produce an overgrowth of an extremity and a deformity of the corresponding bone.

NF1 is clinically diagnosed when any 2 of the following 7 features are present:

• Six or more café-au-lait macules >5mm in prepubertal and >15mm in postpubertal

• Axillary or inguinal freckling

• Two or more iris lisch nodules

• Two or more neurofibromas or one plexiform neurofibroma

• A distinctive osseous lesion such as sphenoid dysplasia or cortical thinning of long bones with or without pseudoarthrosis.

• Optic gliomas

• A first-degree relative with NF-1

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Guillain-Barre Syndrome and Other Immune-mediated Neuropathies

Question 1:

Which of the following is the most common antecedent infectious cause of Guillain-Barre syndrome?

a) Cytomegalovirus

b) Mycoplasma pneumoniae

c) Campylobacter jejuni

d) Epstein Barr virus

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 2:

Which of the following autoantibody isotype is involved in the pathogenesis of Guillian-Barre syndrome?

a) IgA

b) IgG

c) IgM

d) IgE

Question 3:

In Guillain-Barre syndrome due to antecedent infection with Campylobacter jejuni, the autoantibodies are most commonly directed against which of the following?

a) GM1

b) GD1a

c) GD1b

d) GalNac-GD1a

Question 4:

Which of the following is the most common type of Guillian-Barre syndrome?

Sold by @itachibot

a) Acute motor axonal neuropathy

b) Acute inflammatory demyelinating polyneuropathy

c) Acute motor sensory axonal neuropathy

d) Miller Fisher syndrome

Question 5:

Which type of Guillain-Barre syndrome most commonly affects patients in the pediatric age group?

a) Acute motor axonal neuropathy

b) Acute inflammatory demyelinating polyneuropathy

c) Acute motor sensory axonal neuropathy

d) Miller Fisher syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Acute inflammatory demyelinating polyneuropathy differs from acute motor axonal neuropathy, in that, in the latter, the complement deposition along with IgG antibodies occurs at:

a) Nodes of Ranvier

b) Neuromuscular junction

c) Oligodendrocytes

d) Schwann cells

Question 7:

Which of the following features would you not expect to see in a patient with Guillian-Barre syndrome?

a) Absent reflexes in lower limb

b) Ascending paralysis

c) Early bladder dysfunction

d) Bulbar paralysis

Question 8:

Sold by @itachibot

Which of the following sensory deficits is severely affected in a patient with Guillain-Barre syndrome?

a) Pain

b) Temperature

c) Proprioception

d) Touch

Question 9:

A patient presents with complaints of lower extremity weakness. He initially noticed difficulty climbing the stairs and now he is unable to stand. He gives a history of bloody diarrhea a few weeks ago. Which of the following parameters are expected to be elevated in the CSF of this patient?

a) Lymphocytes

b) Neutrophils

c) Opening pressure

d) Proteins

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following conditions does not present with autonomic dysfunction?

a) Amyloidosis

b) Guillain-Barré syndrome

c) Botulism

d) Subclavian steal syndrome

Question 11:

A 23-year-old patient presents with rapidly progressing symmetric ascending paralysis since 2 days. Nerve conduction studies are consistent with demyelination. He is found to have an elevated protein level in his CSF. Which of the following autonomic disturbance is likely to be seen in this patient?

a) Postural hypotension

b) Urinary incontinence

Sold by @itachibot

c) Diarrhea

d) Increased sweating

Question 12:

The triad of Miller-Fisher syndrome includes all of the following, except:

a) Ataxia

b) Weakness

c) Areflexia

d) Ophthalmoplegia

Question 13:

A patient presented to the ER with tingling, numbness, and weakness in all four limbs. She says that the weakness started in the lower limbs and progressed to the upper limbs within a day. History is significant for a recent COVID-19 infection and surgery for cervical spondylosis and disc herniation 3 years ago. CT brain was normal and an MRI of the lumbar spine showed foraminal narrowing at L2-L3. CSF analysis showed increased protein levels. Which of the following is recommended for the initial therapy of this patient?

a) Intrathecal steroids

b) Remdesivir

c) IV Immunoglobulin

d) Plasmapheresis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

In a patient with Guillain-Barre syndrome, chronic inflammatory demyelinating polyneuropathy is suspected when _____

a) GBS deteriorates >5 weeks after onset or relapses at least three times

b) GBS deteriorates >5 weeks after onset or relapses at least two times

c) GBS deteriorates >9 weeks after onset or relapses at least three times

d) GBS deteriorates >8 weeks after onset or relapses at least two times

Question 15:

Sold by @itachibot

Which of the following conditions are not associated with chronic inflammatory demyelinating polyneuropathy?

a) Diabetic mellitus

b) Systemic lupus erythematosus

c) Lymphoma

d) Hepatitis A infection

Question 16:

All are true about chronic inflammatory demyelinating polyneuropathy, except:

a) Symptoms present for > 8 weeks duration

b) Relapse at least 3 times in a patient with Guillain-Barré syndrome

c) Symmetrical weakness

d) Small nerve fibers are affected more than large fibers

Question 17:

A patient with chronic inflammatory demyelinating polyneuropathy is undergoing nerve conduction study. Which of the following findings is least likely to be seen in this patient?

a) Low amplitude

b) Decreased conduction velocity

c) Increased distal latency

d) Conduction block

Question 18:

In which of the following situations is nerve biopsy not indicated in patients with chronic inflammatory demyelinating polyneuropathy?

a) When the diagnosis not clear

b) When other causes of polyneuropathy cannot be ruled out

c) When electromyography shows axonal involvement

d) When there is no clinical or electrophysiological evidence of nerve functionality

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 19:

All of the following can be used for the treatment of a patient with chronic inflammatory demyelinating polyneuropathy, except:

a) IV Immunoglobulin

b) Plasmapheresis

c) NSAIDs

d) Glucocorticoids

Question 20:

A 63-year-old man presented with complaints of fatiguability and chronic back pain. On evaluation, he was found to have anemia and hypercalcemia. The X-ray of his skull is given below. Which of the following statements is true with regard to neuropathy seen in this condition?



a) Polyneuropathy is seen in 50% of patients presenting with osteoporotic bone lesions

b) Polyneuropathy is seen in 5% of patients presenting with osteosclerotic features

c) The pattern is sensorimotor

d) Treatment may reverse severe neuropathy

Question 21:

Sold by @itachibot

What is the most common pattern of peripheral neuropathy in patients with polyarteritis nodosa?

a) Pure motor neuropathy

b) Pure sensory neuropathy

c) Symmetric motor-sensory neuropathy

d) Mononeuropathy multiplex

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	b
5	a
6	a
7	c
8	c
9	d
10	d
11	a
12	b
13	c
14	c
15	d
16	d
17	a
18	d
19	c
20	c
21	d

Detailed Explanations

Sold by @itachibot

Solution to Question 1:

Campylobacter jejuni is one of the most common risk factors for GBS.

Infections that may trigger GBS are:

• Campylobacter jejuni

• Cytomegalovirus (CMV)

• Epstein Barr virus

• Mycoplasma

• Varicella zoster virus

• Human immunodeficiency virus (HIV), dengue, or influenza, Zika, hepatitis

Less commonly vaccinations, surgical procedures, and trauma have been reported to trigger the development of GBS.

Solution to Question 2:

IgG isotype of autoantibody is involved in the pathogenesis of GBS.

IgM isotype autoantibodies are known to play a role in some types of chronic immune neuropathies.

Solution to Question 3:

Autoantibodies in GBS due to Campylobacter jejuni infection are directed most commonly against GM1 gangliosidase.

Solution to Question 4:

Acute inflammatory demyelinating polyneuropathy (AIDP) is the most common type of GBS.

There are several types of Guillain–Barre syndrome, which are recognized by electrodiagnostic studies and pathological distinctions. These include:

A. Demyelinating

• Acute inflammatory demyelinating polyneuropathy (AIDP) - most common

B. Axonal

• Acute motor axonal neuropathy (AMAN)

• Acute motor–sensory axonal neuropathy (AMSAN)

C. Regional GBS syndromes

• Miller–Fisher syndrome (MFS)

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• Pure sensory forms

• Ophthalmoplegia with anti-GQ1b antibodies as part of severe motor-sensory GBS

• GBS with severe bulbar and facial paralysis

• Acute pandysautonomia

Solution to Question 5:

The acute motor axonal neuropathy (AMAN) subtype of Guillain-Barre syndrome (GBS) is more common in pediatric age groups.

Type	Age group affected
Acute inflammatory demyeli nating polyneuropathy (AIDP)	Adults
Acute motor axonal neuropat hy (AMAN)	ChildrenYoung adu lts
Acute motor-sensory axonal neuropathy (AMSAN)	Mostly adults
Miller–Fisher syndrome	AdultsChildren

If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

In AMAN, the complement along with IgG is deposited at the nodes of Ranvier of large motor axons.

In AIDP, complement deposits along the outer surface of the Schwann cell.

Solution to Question 7:

Bladder dysfunction is not usually severe and does not appear early in the course of Guillain-Barre Syndrome(GBS). If bladder dysfunction is a prominent feature and comes early in the course, diagnostic possibilities other than GBS should be considered.

GBS is an acute demyelinating polyradiculoneuropathy. It is an autoimmune disorder triggered by infections like:

• Campylobacter jejuni

• Cytomegalovirus (CMV)

• Epstein Barr virus

• Mycoplasma

• Varicella zoster virus

Sold by @itachibot

• Human immunodeficiency virus (HIV), dengue, or influenza

It manifests as a rapidly evolving areflexic motor paralysis, with or without sensory disturbance. The usual pattern is ascending paralysis. Proximal and distal muscles are equally involved. The lower cranial nerves are frequently affected, causing bulbar weakness.

Solution to Question 8:

Loss of proprioception is severe in Guillain-Barre syndrome (GBS).

GBS affects the large sensory fibers, leading to attenuation or loss of deep tendon reflexes and loss of proprioception.

Cutaneous sensory deficits (e.g., loss of pain and temperature sensation) are usually relatively mild.

Solution to Question 9:

The given scenario is suggestive of Guillain-Barre syndrome (GBS). Protein levels are elevated (100-1000mg/dl) in the CSF of a patient with this condition.

The CSF cell counts are usually normal. Albuminocytologic dissociation, which is an elevation in CSF protein without an elevation in white blood cells, is characteristic in GBS.

Causes of albuminocytologic dissociation are:

• Guillain-Barre Syndrome

• Spinal cord tumours

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Subclavian steal syndrome does not present with autonomic dysfunction.

Option A: Polyneuropathy is seen in sporadic and familial amyloidosis. In some cases, autonomic insufficiency presents before distal painful polyneuropathy accompanied by sensory loss

Option B: Guillain-Barré syndrome is associated with autonomic instability. This can present with BP fluctuations, arrhythmias, and GI disturbances.

Option C: Botulinum toxin binds presynaptically to cholinergic nerve terminals. After uptake into the cytoplasm, it blocks ACh release. Thus, patients with botulism present with motor paralysis and autonomic disturbances.

Solution to Question 11:

Sold by @itachibot

The given scenario is suggestive of Guillian-Barre syndrome (GBS). Postural hypotension is a common autonomic disturbance seen in this condition.

Autonomic involvement is common and may occur even in patients whose GBS is otherwise mild.

The usual autonomic manifestations are:

• Loss of vasomotor control with wide fluctuation in blood pressure, postural hypotension

• Cardiac dysrhythmias

Solution to Question 12:

Weakness is not seen in the triad of Miller-Fisher syndrome (MFS).

Miller Fisher syndrome is a variant of Guillain-Barré syndrome(GBS). Like Guillain-Barré syndrome, symptoms may be preceded by a viral illness. It classically presents as a triad of:

• Ataxia

• Areflexia

• Ophthalmoplegia

The condition is usually not associated with weakness, and even if present, is very mild. It is strongly associated with antibodies to the ganglioside GQ1b.

The prognosis in Miller Fisher syndrome is usually good with recovery beginning within 2 to 4 weeks of the symptom onset, and completing by 6 months. Some individuals are left with residual deficits. Relapses are rare.

Treatment for Miller Fisher syndrome is identical to treatment for Guillain-Barré syndrome: intravenous immunoglobulin (IVIg) or plasmapheresis and supportive care.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 13:

The given scenario is suggestive of Guillain-Barre syndrome(GBS) probably as a result of COVID-19. IV Immunoglobulin is the recommended initial therapy for patients with GBS.

High-dose intravenous immune globulin (IVIg) and plasmapheresis are both equally effective in GBS and can be used as the initial therapy. However, IVIg is often the initial therapy chosen because of its ease of administration and good safety record. It is administered as five daily infusions for a total dose of 2 g/kg body weight. Glucocorticoids have not been found to be effective in GBS.

Solution to Question 14:

Chronic inflammatory demyelinating polyneuropathy (CIDP) should be suspected when GBS deteriorates >9 weeks after onset or relapses at least three times.

Sold by @itachibot

Onset is usually gradual over a few months or longer, but in a few cases, the initial attack is indistinguishable from that of GBS.

CIDP is distinguished from GBS by its chronic course. In other respects, this neuropathy shares many features with the common demyelinating form of GBS, including elevated CSF protein levels and the electrodiagnostic studies findings of acquired demyelination.

Solution to Question 15:

Hepatitis A infection is not associated with CIDP.

Diseases associated with CIDP are:

- Hepatitis B
- Hepatitis C
- Lymphoma
- HIV
- Diabetic mellitus
- Systemic lupus erythematosus (SLE)
- Pregnancy
- Paraneoplastic syndrome
- Inflammatory bowel disease
- Amyloidosis

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

Large nerve fibers are affected more than small fibers in CIDP.

Difference between CIDP and Guillain-barré syndrome(GBS):

	GBS	CIDP
Duration	<4 weeks	>8 weeks
Motor symptoms	Affected	Affected
Sensory symptoms	Mild	More significantLargefiber > small fiber
Autonomic nervous system	Affected	Rare
Antecedent event	Present	Rare

Sold by @itachibot

	GBS	CIDP
Deep tendon reflex	Absent	Absent
Steroid	No role	2nd line of treatment

Solution to Question 17:

In CIDP, the amplitude is normal since the axons are intact.

Nerve conduction study (NCS) in axonal injury and demyelinating diseases:

	Axonal injury	Demyelination
Amplitude	Decreased	Normal
Distal latency	Normal	Prolonged
Conduction velocity	Normal	Slow
Conduction block	Absent	Present
Temporal dispersion	Absent	Present
H wave	Normal or absent	Prolonged or absent
F wave	Normal or absent	Prolonged or absent

Solution to Question 18:

Nerve biopsy should not be done if there is no clinical or electrophysiological evidence of nerve functionality of that nerve in CIDP patients. This is because completely degenerated nerves usually convey limited information.

In CIDP, nerve biopsy is indicated when:

• Diagnosis is not clear

• Other causes of polyneuropathy cannot be ruled out

• Electromyography (EMG) shows axonal involvement

Solution to Question 19:

NSAIDs are not used in the treatment of CIDP.

Sold by @itachibot

The treatment of chronic inflammatory demyelinating polyneuropathy (CIDP) includes high-dose IV immunoglobulin, plasmapheresis, and glucocorticoids.

Initial therapy is usually with IV immunoglobulin. If all the first-line agents fail, immunosuppressants are attempted either alone or as an adjunctive treatment.

Solution to Question 20:

The given scenario is suggestive of multiple myeloma and the pattern of polyneuropathy seen in this condition is sensorimotor type.

Multiple myeloma-associated polyneuropathies occur in ~5% of patients with the commonly encountered type of multiple myeloma.

Myeloma with osteosclerotic features, although representing only 3% of all myelomas, is associated with polyneuropathy in one-half of cases.

Polyneuropathy generally does not reverse even with successful suppression of the myeloma.

Solution to Question 21:

Most common pattern of peripheral neuropathy in polyarteritis nodosa (PAN) is mononeuritis multiplex (multifocal motor-sensory neuropathy). It is asymmetric, due to ischemic lesions of nerve trunks and roots.

Some cases present as a distal, symmetric sensorimotor polyneuropathy. Small- to medium-sized arteries of the vasa nervorum, particularly the epineural vessels, are affected in PAN, resulting in a widespread ischemic neuropathy.

Combination therapy with glucocorticoids and rituximab is increasingly recommended as the standard initial treatment, particularly for ANCA-associated vasculitis.

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Sold by @itachibot

Diseases of the Spinal Cord

Question 1:

A patient has sensory loss just below the circumferential level of the umbilicus. This corresponds to a lesion at which spinal cord segment?

a) T10

b) T11

c) T8

d) T12

Question 2:

A patient presents with quadriplegia and difficulty in breathing. This presentation can be seen in spinal cord lesions at what level?

a) C1 – C4

b) C5 – C6

c) C7

d) C8

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

Beevor's sign is noted in a patient who has spinal cord injury. At what level is the injury likely to be?

a) T7 – T8

b) T8 – T9

c) T10 – T12

d) T6 – T7

Question 4:

A patient has a spinal cord lesion at the level of L2 – L4 spinal nerve roots. All of the following may be seen in this patient, except:

Sold by @itachibot

a) Paralysis of hip flexion and adduction

b) Paralysis of knee flexion

c) Weakened knee extension

d) Absence of the patellar reflex

Question 5:

Which of the following features would you expect to see in a patient with conus medullaris syndrome?

a) 1, 2, and 4 only

b) 1, 4, and 5 only

c) 1, 3, and 5 only

d) 2,3, and 4 only

Question 6:

Which of the following conditions will you suspect in a patient who has bladder dysfunction early in the progression of the disease?

a) Cauda equina syndrome

b) Conus medullaris syndrome

c) Guillian–Barre syndrome

d) Amyotrophic lateral sclerosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

A patient with a history of a backache for 10 days, has now presented with sudden onset difficulty in micturition and defaecation. There was no history of a cough or fever previously. What is the diagnosis?

a) Pott's Spine

b) Guillain Barre Syndrome

c) Cauda Equina Syndrome

d) Multiple Sclerosis

Question 8:

Sold by @itachibot

Which of the following features will help you differentiate conus medullaris syndrome from cauda equina syndrome?

a) Radicular pain

b) Lower motor neuron injury

c) Symmetric weakness of lower limbs

d) Urinary retention

Question 9:

Which is the characteristic pattern seen in Brown-Sequard syndrome?

a) Contralateral loss of joint sense and position

b) Contralateral loss of pain sensation

c) Ipsilateral loss of complete sensory functions

d) Contralateral motor functions

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A patient is diagnosed with Brown–Sequard syndrome with a lesion on the right side. Which of the following sensations on the right side will be affected?

a) Pain

b) Crude touch

c) Proprioception

d) Temperature

Question 11:

A patient was brought to the casualty after he was hit by a plastic pipe on the back by a bully. On examination, the patient was found to have marked upper extremity weakness compared to the lower extremities. He also had a loss of pain and temperature sensations over the shoulders, lower neck, and upper trunk. Other sensations were intact in this region. What is the likely diagnosis?

a) Brown-Sequard syndrome

b) Central cord syndrome

c) Anterior spinal artery syndrome

Sold by @itachibot

d) Foramen magnum syndrome

Question 12:

The MRI spine of a patient is shown below. What is the characteristic deficit seen in this condition?

a) Pyramidal pattern of weakness

b) Around the clock pattern of weakness

c) Suspended distribution of sensory deficit

d) Segmental pattern of weakness

Question 13:

A patient presents with quadriparesis. He says that it started in his left arm, then progressed to the left leg, and then later to the right leg and arm. What is the likely diagnosis?

a) Extramedullary tumor

b) Central cord syndrome

c) Anterior cord syndrome

d) Foramen magnum syndrome

Question 14:

Sold by @itachibot

Anterior spinal artery thrombosis is characterized by all of the following, except:

a) Loss of pain and touch

b) Loss of vibration sense

c) Loss of power in the lower limbs

d) Sphincter dysfunction

Question 15:

A patient with neurofibromatosis 1 is found to have multiple tumors above the level of the lumbar spine. Which of the following is an early feature of dysautonomia in this patient?

a) Large bladder volumes

b) Flaccid bladder

c) Overflow incontinence

d) Spastic bladder

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

Which of the following autonomic disorders do not involve the spinal cord?

a) Syringomyelia

b) Syringobulbia

c) Amyotrophic lateral sclerosis

d) Multiple sclerosis

Question 17:

Which of the following features would you not expect to see in a patient with an extramedullary tumor in the spinal cord?

a) Abnormal CSF

b) Sacral sparing

c) Spastic weakness of lower limbs

d) Radicular pain

Question 18:

Sold by @itachibot

Which of the following is not a cause of subacute combined degeneration?

a) Pernicious anemia

b) Partial gastrectomy

c) High consumption of refined rice

d) Diphyllobothrium latum infestation

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	b
5	a
6	b
7	c
8	c
9	b
10	c
11	b
12	c
13	d
14	b
15	d
16	b
17	b
18	c

Detailed Explanations

Solution to Question 1:

The sensory loss just below the circumferential level of the umbilicus corresponds to the lesion at the T10 spinal cord segment. It indicates the involvement of the cord adjacent to T7-T8 vertebral

Sold by @itachibot

bodies.

The spinal cord has 31 segments, but they do not always correspond to the same vertebral body.

The first seven pairs of cervical spinal nerves exit above the same-numbered vertebral bodies. All the subsequent nerves exit below the same-numbered vertebral bodies. This is because of the presence of eight cervical spinal cord segments but only seven cervical vertebrae.

The lower spinal nerves take an increasingly downward course to exit via intervertebral foramina.



Chest and abdomen dermatome



©MARROW

Solution to Question 2:

Quadriplegia and weakness of the diaphragm (manifesting as difficulty in breathing) are associated with spinal cord lesion at the level of C1-C4 spinal segments.

Diaphragm is supplied by the phrenic nerve arising from C3 - C5.

Sold by @itachibot

The uppermost level of weakness and reflex loss with lesions:

• At C5-C6 : Biceps

• At C7: Finger, wrist extensors, and triceps

• At C8 : Finger and wrist flexion

Note: Horner's syndrome (characterized by miosis, ptosis, and facial hypohidrosis) may accompany a cervical cord lesion at any level.

Solution to Question 3:

The Beevor's sign localizes the spinal cord injury at T10-T12.

Beevor's sign is the abnormal upward movement of the umbilicus on abdominal wall contraction in the supine position. This is because a lesion in the T10-T12 level of the spinal cord paralyzes the lower abdominal muscles but the upper abdominal muscles are normally resulting in the upward movement of the umbilicus.



Solution to Question 4:

Lesions at L2 – L4 spinal nerve roots do not cause paralysis of the knee flexors.

The paralysis of the flexion at the knee occurs due to the lesion at L5 – S1.

Solution to Question 5:

The clinical features of the conus medullaris syndrome include:

Sold by @itachibot

• Bilateral saddle anesthesia

• Bladder and bowel dysfunction

• Impotence

• Absence of bulbocavernosus (S2-S4) and anal reflexes (S4-S5)

The conus medullaris is the tapered caudal termination of the spinal cord, comprising all the sacral and single coccygeal segments. Damage at this level of the cord, often occurs due to injuries to the lumbar vertebra, result in conus medullaris syndrome. Muscle strength is largely preserved in this condition.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

Early loss of bladder control is a feature of the conus medullaris syndrome.

It is characterized by urinary retention and atonic anal sphincter, which cause overflow urinary incontinence and fecal incontinence, respectively.

Option A: In the cauda equina syndrome, there is relative sparing of bladder control.

Options C and D: Bladder dysfunction is uncommon in Guillian–Barre syndrome and ALS.

Solution to Question 7:

This is a case of cauda equina syndrome (CES).

CES signifies an injury of multiple lumbosacral nerve roots within the spinal canal distal to the termination of the spinal cord at L1-L2. It is commonly due to a ruptured lumbosacral intervertebral disk, lumbosacral spine fracture, hematoma within the spinal canal, compressive tumor, or other mass lesions.

Low back pain, weakness, and areflexia in the legs, saddle anesthesia, or loss of bladder function may occur.

Treatment options include surgical decompression, sometimes urgently in an attempt to restore or preserve motor or sphincter function, or radiotherapy for metastatic tumors.

Solution to Question 8:

Symmetric weakness of lower limbs helps in a diagnosis of conus medullaris syndrome.

There is an asymmetric motor loss in cauda equina syndrome.

Mass lesions in the lower spinal canal often produce a mixed clinical picture with elements of both cauda equina and conus medullaris syndromes.

Sold by @itachibot

Solution to Question 9:

The Brown–Sequard syndrome presents with contralateral loss of pain sensation.

Lateral hemisection of the spinal cord produces the Brown–Séquard syndrome (also known as the hemicord syndrome). Knife or bullet injuries and demyelination are the most common causes of this condition.

Clinical features include:

• Ipsilateral weakness (corticospinal tract involvement)

• Ipsilateral loss of joint position and vibratory sense (posterior column involvement)

• Contralateral loss of pain and temperature sense (spinothalamic tract involvement) one or two levels below the lesion

• Ipsilateral segmental signs like radicular pain, muscle atrophy, or loss of a deep tendon reflex

Solution to Question 10:

Proprioception is the type of sensation lost on the same side (ipsilateral) in Brown–Sequard syndrome. This is due to a lesion of the posterior column.

The sensations of crude touch, pain and temperature are lost on the opposite side, due to the lesions of anterior column or spinothalamic tract.

Solution to Question 11:

The given scenario is suggestive of cape distribution of sensory deficit is seen in central cord syndrome.

This condition affects the cervical cord producing arm weakness > leg weakness. A dissociated sensory loss i.e. loss of pain and temperature with preservation of light touch, joint position, and vibration sensations is seen over the shoulders, lower neck, and upper trunk (cape distribution).

It results from selective damage to the gray matter nerve cells and the decussating spinothalamic tracts surrounding the central canal. These decussating fibers carry crude touch, pain, and temperature sensations.

Spinal trauma, syringomyelia, and intrinsic cord tumors are the main causes.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot



Option A: In Brown–Sequard Syndrome, ipsilateral weakness and loss of joint position and vibratory sense, with contralateral loss of pain and temperature sense, one or two levels below the lesion is seen.

Option C: Anterior spinal artery syndrome is caused due to an ASA infarct. It typically presents as loss of motor function and pain/temperature sensation, with relative sparing of proprioception and vibratory sense below the level of the lesion.

Option D: In foramen magnum syndrome, compressive lesions near the foramen magnum may produce weakness of the ipsilateral shoulder and arm followed by weakness of the ipsilateral leg, then the contralateral leg, and finally the contralateral arm. This is referred to as 'around the clock' pattern.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 12:

The MRI shows a syrinx located at C6–C7 in the cervical spine which is suggestive of syringomyelia. Suspended distribution of sensory deficit is a feature of this condition.

Syringomyelia is a developmental cavity of the cervical cord that may enlarge and produce progressive myelopathy or may remain asymptomatic.

It presents as a central cord syndrome with unilateral sensory loss in the hands initially. There is a loss of pain and temperature sensation with the sparing of touch and vibration. The sensory deficit has a distribution that is 'suspended' over the nape of the neck, shoulders, and upper arms (cape distribution) or in the hands. Areflexic weakness is seen in the upper limbs.

Extension of the syrinx into the grey matter of the cord causes muscle wasting in the lower neck, shoulders, arms, and hands with asymmetric or absent reflexes in the arms. Extension of the syrinx into the medulla (syringobulbia) causes palatal or vocal cord paralysis, dysarthria, horizontal or vertical nystagmus, episodic dizziness or vertigo, and tongue weakness with atrophy.

Sold by @itachibot

Solution to Question 13:

The given scenario is suggestive of an 'around-the-clock' pattern of weakness, which is seen in foramen magnum syndrome.

Around the clock pattern of weakness is described as a sequential weakness of the ipsilateral shoulder and arm followed by weakness of the ipsilateral leg, then the contralateral leg, and finally the contralateral arm. The weakness may begin in any of the limbs.

The lesions in the foramen magnum area interrupt decussating pyramidal tract fibers destined for the legs, resulting in weakness of the legs. There is typically suboccipital pain spreading to the neck and shoulders in foramen magnum syndrome.

Solution to Question 14:

Anterior spinal artery thrombosis does not lead to loss of vibration sense. Vibration sense is carried by the posterior column.

Anterior spinal artery syndrome or anterior cord syndrome is the result of occlusion or diminished flow in the anterior spinal artery. There is bilateral tissue destruction at several contiguous levels that spares the posterior columns.

All spinal cord functions (motor, sensory, and autonomic) are lost below the level of the lesion, except vibration and position sensation.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 15:

The early feature of dysautonomia that can occur in this patient is spastic bladder.

Dysautonomia due to brain and spinal cord disease above the level of the lumbar spine results first in increased urinary frequency and small bladder volume. Eventually, incontinence occurs. These features are suggestive of an upper motor neuron lesion, i.e., spastic bladder.

On the other hand, dysautonomia due to peripheral nervous system (PNS) disease of autonomic nerve fibers results in large bladder volumes, increased urinary frequency, and overflow incontinence. These are features suggestive of a lower motor neuron lesion, i.e., flaccid bladder.

Postvoidal residual volume helps to distinguish between upper and lower motor neuron bladder dysfunction in the early stages of dysautonomia.

Solution to Question 16:

Syringobulbia is a condition that affects the brainstem, not the spinal cord. It is a neurological disorder characterized by a fluid-filled cavity (syrinx) which affects the brainstem (medulla).

The following autonomic disorders are associated with spinal cord involvement:

• Syringomyelia

Sold by @itachibot

• Multiple sclerosis and neuromyelitis optica

• Amyotrophic lateral sclerosis (ALS)

• Traumatic quadriplegia

• Subacute combined degeneration

• Tetanus

• Stiff-person syndrome

• Spinal cord tumors

Solution to Question 17:

Sacral sparing is not a feature of extramedullary tumors, it is seen in intramedullary tumors. Differences between extramedullary and intramedullary syndromes are:

Extramedullary tumors lie outside the cord and compress the spinal cord or its vascular supply. Intramedullary tumors arise within the substance of the cord.

The spinothalamic tract has a laminated configuration with the sacral fibers outermost, and hence, there is an early sacral sensory loss in extramedullary tumors.

Extramedullary	Intramedullary
Radicular pain	Poorly localized burning pain
Early sacral sensory loss	Sacral sparing
Incontinence due to the superficial location of the corresponding sensory and motor fibers in the spinothalamic and corticospinal tracts	Corticospinal tract signs appear later

Solution to Question 18:

Subacute combined degeneration (SACD) is due to the deficiency of vitamin B12. Pernicious anemia, partial gastrectomy, and Diphyllobothrium latum infestation can lead to this condition. High consumption of refined rice leads to thiamine (vitamin B1) deficiency.

The most severely affected regions in SACD are the posterior columns at the cervical and upper thoracic levels.

Clinical features include:

• Signs of both peripheral nerve and spinal cord involvement

• Paresthesias in the hands or feet > Weakness and unsteadiness of gait

• Loss of vibration or joint position sense in the legs

Sold by @itachibot

• Mental slowing, depression, confusion, delusions, and hallucinations

SACD is treated by a replacement regimen of intramuscular daily injections of 1000 µg of vitamin B12 for the first week, followed by weekly 1000 µg injections for 1 month, and monthly injections thereafter.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Myasthenia gravis and Other Neuromuscular Junction Disorders

Question 1:

Myasthenia gravis is associated with which of the following?

a) Decreased acetylcholine release at the nerve endings

b) Increased acetylcholine release at the nerve endings

c) Increased synaptic transmission at the myoneural junction

d) Decreased synaptic transmission at the myoneural junction

Question 2:

In myasthenia gravis, the anti-AChR antibodies reduce the number of available AChRs at the neuromuscular junction by all of the following mechanisms except:

a) Accelerated turnover of AChRs

b) Damage to the postsynaptic muscle membrane

c) Damage to muscle-specific receptor tyrosine kinase

d) Blockade of the active site of the AChR

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following statements about myasthenia gravis is false?

a) Weakness and fatigability of muscles is present

b) Complete remissions are seen

c) Lids and extraocular muscles are involved

d) Proximal limb weakness is seen

Question 4:

A school teacher presents to the OPD with excessive fatigability and double vision. She says that her symptoms are aggravated towards the end of the day. Her colleagues told her that, lately, she has developed a nasal twang in her voice. Which of the following statements is true

Sold by @itachibot

regarding this condition?

a) Thymoma is present in more than 75% of the patients with this condition

b) In most of the cases the thymus is hypoplastic

c) Myoid cells in thymus serve as the source of autoantigens

d) Thymus is abnormal in 20% of patients with AChR antibody–positive status

Question 5:

A 55-year old woman presented with shortness of breath, cough with expectoration and fever for ten days. Chest x-ray revealed mediastinal widening. CECT chest showed a well-circumscribed heterogeneous solid enhancing mass lesion. FNAC showed features in favor of thymoma. Which of the following is not associated with this condition?

a) Myasthenia gravis

b) Addison's disease

c) Systemic lupus erythematosus

d) Hypergammaglobulinemia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

What is not a feature of myasthenia gravis among the following?

a) Muscle fatigue

b) Absent deep tendon reflexes

c) Normal pupillary reflex

d) Ptosis

Question 7:

A 45-year-old patient comes with complaints of progressive weakness as the day progresses. He is found to have antibodies against nicotinic acetylcholine receptors. Which is the most common ocular finding in this disease?

a) Diplopia

b) Lagophthalmos

c) Exophthalmos

d) Ptosis

Sold by @itachibot

Question 8:

Which of the following features would you not expect to see in patients with ocular myasthenia?

a) Ptosis may switch from one eye to the other over time

b) Manually raising the more ptotic lid causes increased ptosis on the opposite side

c) Extraocular muscle involvement produces binocular diplopia

d) Pupils are affected in 50% of the cases

Question 9:

Which of the following conditions is not associated with congenital myasthenic syndrome?

a) Slow-channel congenital myasthenic syndrome

b) Fast-channel congenital myasthenic syndrome

c) Endplate acetylcholinesterase deficiency

d) Dysferlin deficiency

Question 10:

A 65-year-old man who is a laborer presents with progressive weakness of the legs. He says that the weakness is more in the morning and as the day progresses, it improves for a while usually. He is a known smoker. Examination reveals diminished deep tendon reflexes in the lower limbs. Which of the following statements is true regarding this condition?

a) Depletion of presynaptic acetylcholine vesicles

b) No immunological involvement

c) Paraneoplastic syndrome seen in lung adenocarcinoma

d) Incremental response on repetitive nerve stimulation

Question 11:

For diagnosing patients with suspected myasthenia gravis, which of the following tests has the highest sensitivity?

a) Single-fiber electromyography

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

b) Repetitive nerve stimulation

c) AChR antibodies

d) Edrophonium test

Question 12:

Which of the following is the first line drug of choice in a patient with myasthenia gravis?

a) Azathioprine

b) Prednisone

c) Intravenous immunoglobulin

d) Pyridostigmine

Question 13:

Which of the following statements is true regarding thymectomy in patients with generalized myasthenia gravis?

a) Indicated in most AChR-Ab positive patients with generalized myasthenia gravis

b) Contraindicated in medication-contolled patients

c) Induces rapid remission from disease

d) Indicated exclusively in patients with thymomatous myasthenia gravis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

A female patient with myasthenia gravis was admitted with complaints of sore throat, fever, cough, and weakness. She had a history of repeated hospital admissions due to exacerbation of myasthenia gravis. During some of the episodes, she even required intubation. Blood culture on admission revealed MRSA infection. Which of the following drugs can be safely administered to this patient for treating the infection?

a) Vancomycin

b) Clindamycin

c) Daptomycin

d) Gentamicin

Question 15:

Sold by @itachibot

Patients with myasthenia gravis are resistant to anesthesia with which of the following drugs?

a) Depolarizing agents

b) Non-depolarizing agents

c) Local anesthetics

d) Inhalation anesthetics

Question 16:

All of the following are the drugs that are used for the treatment of patients with myasthenia gravis. Which of the following does not alter the disease process?

a) Pyridostigmine

b) Prednisone

c) Mycophenolate mofetil

d) Azathioprine

Sold by @itachibot
If you purchased this from someone else,
you have been scammed.

Answer Key

Question No.	Correct Option
1	d
2	c
3	b
4	c
5	d
6	b
7	d
8	d
9	d
10	d
11	a
12	d
13	a
14	c
15	a

Sold by @itachibot

16	a

Detailed Explanations

Solution to Question 1:

Myasthenia gravis is associated with a decrease in synaptic transmission at the myoneural junction.

Myasthenia gravis is a neuromuscular disorder that is characterized by weakness and fatigability of skeletal muscles. It is due to autoantibody-mediated destruction of the acetylcholine receptors (AChRs) at the neuromuscular junction. This consequentially leads to a decreased transmission at the neuromuscular (myoneural) junction.

Solution to Question 2:

In myasthenia gravis, damage to muscle-specific receptor tyrosine kinase is not a mechanism by which anti-AChR antibodies reduce the number of available AChRs at the neuromuscular junction.

The anti-AChR antibodies reduce the number of available AChRs at the neuromuscular junction using three distinct mechanisms:

• Accelerated turnover of AChRs

• Damage to the postsynaptic muscle membrane mediated by antibodies and complement

• Blockade of the active site of the AChR

Patients whose sera test negative for both AChR and MuSK antibodies possess antibodies specific to another protein at the neuromuscular junction called low-density lipoprotein receptor-related protein 4 (LRP4). LRP4 plays a functional role in the clustering of AChRs.

Anti-MuSK antibodies are directed against muscle-specific receptor tyrosine kinase (MuSK), a protein involved in AChR clustering at the neuromuscular junction.

Solution to Question 3:

Remissions may occur spontaneously but are usually not complete or permanent.

The cardinal clinical features of myasthenia gravis include:

• Women are affected by myasthenia gravis more frequently than men, in a ratio of ~3:2.

• Weakness and fatigability of muscles - Weakness may increase upon repeated use or later in the day. It may improve with rest or sleep.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

• Diplopia and ptosis - They are common initial clinical complaints. The cranial muscles, particularly the eyelids and the extraocular muscles, are typically involved early in the course of myasthenia gravis.

• Limb weakness - It is often proximal and may be asymmetric.

• Deep tendon reflexes - They are preserved despite muscle weakness.

Solution to Question 4:

The given scenario is suggestive of myasthenia gravis (MG). Myoid cells in the thymus serve as the source of autoantigens in this condition.

The muscle-like cells within the thymus called myoid cells, express AChRs on their surface. These cells are known to serve as a source of autoantigen and trigger autoimmune reactions within the thymus gland. This stimulates the production of autoantibodies, thus causing myasthenia gravis.

The thymus is abnormal in ~75% of patients with AChR antibody-positive MG. In these cases, the thymus is hyperplastic. 10% of the patients show symptoms of thymoma.

Solution to Question 5:

Thymoma is associated with hypogammaglobulinemia, not hypergammaglobulinemia. Thymoma with hypogammaglobulinemia is also called Good's syndrome.

The following conditions are associated with thymomas:

• Myasthenia gravis (30%)

• Pure red cell aplasia (5-8%)

• Hypogammaglobulinemia (~5%)

• Polymyositis

• SLE

• Thyroiditis

• Sjogren's syndrome

• Ulcerative colitis

• Addison's disease

• Pernicious anemia

• Scleroderma

• Panhypopituitarism

• Aplastic anemia

• Limbic encephalitis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Sold by @itachibot

Solution to Question 6:

Absent deep tendon reflex is not a feature of myasthenia gravis (MG).

MG usually occurs in young females. Patients usually present with weakness and fatiguability of the muscles, which worsens on repeated use or late in the day and improves on rest. The cranial muscles, particularly the eyelids and extraocular muscles are involved early in the course of MG. Diplopia and ptosis are common initial complaints. When MG becomes generalized, the proximal muscles of the limbs are affected first. Despite the muscle weakness, the deep tendon reflexes remain normal. Snarling expression on smiling (facial weakness), nasal timbre of speech (weakness of palate and tongue), and difficulty swallowing may be seen. Bulbar weakness and respiratory depression are seen more frequently in MuSK antibody–positive MG.

MG is an autoimmune disorder most commonly caused by antibodies against the acetylcholine receptor (AChR). These antibodies cause damage to the postsynaptic muscle membrane and increased endocytosis of the AChR. This autoimmune response against the AChR seems to be mediated by the thymus. The myoid (muscle-like) cells present in the thymus express AChRs on their surface and are postulated to provide the autoantigen which triggers this autoimmune reaction. Thymic hyperplasia is seen in 65% of the patients, and thymomas are seen in 10% of the patients.

The tests which are used to confirm the diagnosis of MG include:

• Anti-AChR antibody immunoassay: 85% of myasthenic patients have detectable anti-AChR antibodies, but only 50% of patients with weakness limited to ocular muscles. The presence of anti-AChR antibodies is highly suggestive of MG, but a negative test does not exclude the disease. Antibodies to MuSK and LRP4 are found in some patients

• Recurrent nerve stimulation test: MG shows a decremental response > 10%, while Lambert-Eaton myasthenic syndrome (LEMS) shows an incremental response

• Block and jitter response on single-fiber electromyography

• Positive edrophonium test: improvement of ptosis on the administration of anticholinesterase such as edrophonium

• Positive ice-pack test: improvement of ptosis on the application of ice-pack on the affected eye

The first line of medical management in MG is anticholinesterase drugs. Pyridostigmine 30–60 mg three to four times daily is the most commonly used drug.

Patients are evaluated for associated thymic abnormalities using a CT chest. Thymectomy is considered in patients with thymoma and in generalized MG patients with anti-AChR antibodies.

Immunotherapy therapy can be initiated in patients with poor control on pyridostigmine after thymectomy. Glucocorticoids, cyclosporine, or tacrolimus are generally effective within 1-3 months. Azathioprine and mycophenolate mofetil may take several months to show beneficial effects and are used for the long-term management of MG. Rituximab has shown effectiveness in patients with MuSK antibodies. Intravenous eculizumab, which binds to terminal complement component 5, is also effective.

Myasthenic crisis is the life-threatening exacerbation of MG leading to respiratory failure. Treatment for myasthenic crisis should take place in intensive care units. It is important to rule out excessive anticholinesterase medication as a cause of deterioration by temporarily discontinuing these drugs. Plasmapheresis and intravenous immunoglobulin (IVIg) are beneficial

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

in hastening recovery.

Solution to Question 7:

The given clinical scenario is suggestive of myasthenia gravis and ptosis is the most common ocular feature of the disease.

Myasthenia gravis (MG) is an autoimmune disorder of the neuromuscular junction most commonly caused by anti-acetylcholine receptor antibodies. The disorder is characterized by weakness and fatigability of skeletal muscles where the weakness increases during repeated use and may improve following rest.

Ptosis and diplopia are common initial complaints due to early involvement of the lids and extraocular muscles (EOMs). Of them, Ptosis is a more common manifestation than diplopia. Ptosis in myasthenia gravis is insidious, bilateral, and frequently asymmetrical.

Investigations of myasthenia gravis include:

• Ice pack test -This tests for a transient improvement of the ptotic eyelid on placing an ice pack for 2 minutes, as the cold inhibits the breakdown of acetylcholine by acetylcholinesterase.

• Tensilon (Edrophonium) test - This tests for a transient improvement in the weakness on administration of a short-acting anticholinesterase such as edrophonium.

• Antibody testing - for Acetylcholine receptor (AChR) antibodies, MuSK protein antibodies helps support a diagnosis of myasthenia gravis.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 8:

The pupils are always spared in myasthenia gravis (MG), differentiating it from other disorders marked by similar ocular symptoms (e.g. botulism)

If weakness remains restricted to the extraocular muscles for 3 years, it is likely that it will not become generalized. These patients are said to have ocular MG.

Ptosis and/or diplopia are the initial symptoms of MG in up to 85% of patients. The ptosis may increase with sustained upward gaze or by holding up the opposite eyelid with the examiner's finger (curtain sign).

Cogan lid twitch is characteristic of MG. It refers to a brief overshoot twitch of the lid when the patient is made to gaze upwards after a period of downgaze.

RNS studies (repetitive nerve stimulation) and anti-AChR antibodies are often negative. Single-fiber EMG (SFEMG) testing may be required for confirming the diagnosis in seronegative patients.

Solution to Question 9:

Sold by @itachibot

Dysferlin deficiency is not associated with congenital myasthenic syndrome (CMS). CMS is an uncommon cause of neuromuscular junction failure in newborns. In CMS, the immune system is unaffected.

The common syndromes associated with CMS are as follows:

• Slow-channel congenital myasthenic syndrome (SCCMS)

• Fast-channel congenital myasthenic syndrome (FCCMS)

• Primary acetylcholine receptor deficiency

• Endplate acetylcholinesterase deficiency

Solution to Question 10:

The given scenario is suggestive of Lambert-Eaton myasthenic syndrome (LEMS). Incremental response to repetitive nerve stimulation is a feature of this condition.

LEMS is a neuromuscular junction disorder associated with the reduced acetylcholine (Ach) release from the presynaptic nerve terminals, despite normal ACh vesicle numbers.

Antibodies directed against the P/Q voltage-gated calcium channel (VGCC) interfere with the normal calcium flux required for the release of acetylcholine.

Approximately 50% of LEMS cases are associated with malignancy, mainly small cell lung cancer (SCLC).

Most patients with LEMS present with symptoms as slowly progressive proximal muscle weakness. In LEMS, the weakness improves for a while after exertion. Diminished or absent deep tendon reflexes help differentiate between LEMS and myasthenia gravis.

An increase in the compound muscle action potential (CAMP) amplitude >100% after exercise or with high-frequency repetitive nerve stimulation (RNS) is considered to be of diagnostic significance.

Solution to Question 11:

Single-fiber electromyography has the highest sensitivity for the diagnosis of myasthenia gravis.

The tests used in the diagnosis of myasthenia gravis are:

• Testing for autoantibodies against the acetylcholine receptor (AChR-Ab) should be performed on all patients (most specific).

• If the AChR-Abs test negative, an assay for antibodies against muscle-specific receptor tyrosine kinase (MuSK) should be performed.

• Single-fiber electromyography (SFEMG) simultaneously records action potentials of two muscle fibers innervated by the same motor axon. The variability in time of the second action potential relative to the first is called "jitter" and this is increased in myasthenia gravis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Repetitive nerve stimulation (RNS) is considered abnormal if there is a decremental response of >10%.

• The edrophonium ("Tensilon") test should be used for diagnosis only in those patients with obvious ptosis or ophthalmoparesis and is not confirmatory. This test is not used routinely.

Electrophysiological (RNS/SFEMG) tests are used to diagnose patients affected with seronegative myasthenia gravis.

Solution to Question 12:

Pyridostigmine is the first-line drug of choice in the treatment of symptomatic myasthenia gravis.

Treatment for myasthenia gravis involves:

• Symptomatic treatment: Anticholinesterase agents – Pyridostigmine is the initial therapy of choice.

• Chronic immunomodulating therapy: Glucocorticoids (prednisone) and immunosuppressive drugs (azathioprine, mycophenolate mofetil, cyclosporine) implemented in patients significantly symptomatic on pyridostigmine.

• Rapid immunomodulating treatment: IVIG and therapeutic plasma exchange—Used to tide over severe illness (myasthenic crisis) and as 'bridge' therapy before surgery, or starting chronic immunomodulating treatment.

• Surgical approach: Thymectomy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 13:

Thymectomy is indicated in most AChR-Ab positive patients with generalized myasthenia gravis

Thymectomy is recommended for all patients aged <60 years with non-thymomatous, acetylcholine receptor (AChR) antibody-associated generalized myasthenia gravis.

Significant improvement in symptoms is expected within one year after thymectomy. It is indicated in all patients with thymoma if resection is feasible, regardless of the subtype of myasthenia gravis.

Solution to Question 14:

Daptomycin can be safely administered to treat the patient's MRSA infection. This is because it is not known to unmask or worsen the symptoms of myasthenia gravis.

Drugs that may unmask or worsen myasthenia gravis include:

• Anesthetic agents: Neuromuscular-blocking agents

• Antibiotics:

Sold by @itachibot

- Aminoglycosides (eg., gentamicin, neomycin, tobramycin)
- Clindamycin
- Fluoroquinolones (eg., ciprofloxacin, levofloxacin, norfloxacin)
- Ketolides (eg., telithromycin)
- Vancomycin
- Cardiovascular drugs:
- Beta-blockers (eg., atenolol, labetalol, metoprolol, propranolol)
- Procainamide
- Quinidine
- Other drugs:
- Botulinum toxin
- Chloroquine
- Hydroxychloroquine
- Magnesium
- Penicillamine
- Quinine

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 15:

Patients with myasthenia gravis are resistant to neuromuscular blockade with depolarizing NMBAs (e.g., succinylcholine), possibly because they have a decreased number of acetylcholine receptors.

Local anesthetics and inhalation anesthetics are usually well-tolerated in myasthenia gravis with occasional exacerbations.

Solution to Question 16:

Pyridostigmine is an anticholinesterase agent that does not alter the disease process of myasthenia gravis.

Anticholinesterase agents improve muscle contractions by allowing released acetylcholine to accumulate and act on the receptors over a large area and also have the additional direct depolarizing effect on the motor endplate.

They do not alter the disease process but provide symptomatic relief. The underlying disease process continues and ultimately it may not be possible to restore muscle strength with anticholinesterase agents alone. It is due to the reduction of the receptors by the antibodies.

Sold by @itachibot

Drugs like prednisone, mycophenolate mofetil, and azathioprine are immunosuppressive agents that reduce the antibody titers and can modify the disease process.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Multiple Sclerosis & Other Demyelinating Disorders

Question 1:

Which of the following statements is false regarding demyelinating disorders?

a) They are immune-mediated

b) Commonly presents in adulthood

c) Multiple sclerosis is the most common type

d) It is caused by biochemical abnormality of myelin

Question 2:

Which of the following statements is true regarding multiple sclerosis?

a) It is an acute inflammatory condition of neurons

b) It is an autoimmune disease affecting the central nervous system

c) It may also involve the peripheral nervous system

d) The condition is always benign and resolves spontaneously

Question 3:

A patient with weakness of the limbs and reduced vision, after evaluation, was diagnosed to be a case of multiple sclerosis. Which of the following is not a risk factor for the development of this condition?

a) Cigarette smoking

b) Vitamin D deficiency

c) Exposure to Human Herpesvirus 4

d) HLA-B27

Question 4:

The slowing of conduction in multiple sclerosis is due to which of the following?

a) Gliosis

b) Loss of myelin sheath

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Defect at node of Ranvier

d) Leaky Na+ channel

Question 5:

Autoantibodies in multiple sclerosis are directed against which of the following?

a) Myelin basic protein

b) Both Myelin basic protein & Myelin oligodendrocyte glycoprotein

c) Myelin oligodendrocyte glycoprotein

d) None of the above

Question 6:

Which of the following processes lead to the formation of shadow plaques in patients with multiple sclerosis?

a) Demyelination of axons in the central nervous system

b) Demyelination of axons in the peripheral nervous system

c) Remyelination of axons in the central nervous system

d) Remyelination of axons in the peripheral nervous system

Question 7:

A 40-year-old woman with a history of multiple sclerosis comes to the hospital. She describes an electric shock–like sensation radiating down the back into the legs. What is the name of this sign?

a) Uhthoff's phenomenon

b) Lhermitte's sign

c) Stransky's sign

d) Cornell's sign

Question 8:

An MRI image of a patient with multiple sclerosis is shown below. What is this sign called?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) Black hole lesions

b) Molar tooth sign

c) Tram-track sign

d) Dawson's fingers

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 9:

A woman presents with recurrent episodes of blurry vision, tingling and numbness in the legs, and falls. She says that during hot weather, she finds it difficult to walk and often falls. Recently she had flu, due to which the symptoms aggravated. MRI brain was done which showed Dawson's fingers. What is the criteria used for the clinical diagnosis of her condition?

a) Kurtzke expanded disability status scale

b) Revised McDonald criteria

c) Functional status score

d) Hess and Hunt's scale

Question 10:

In patients with multiple sclerosis, which of the following is used to determine neurological impairment?

a) Hess and Hunt's scale

b) Revised McDonald criteria

c) Kurtzke expanded disability status scale

Sold by @itachibot

d) Modified Well's score

Question 11:

A 45-year-old woman with multiple sclerosis presents with an exacerbation of her symptoms. Which of the following drugs should be prescribed to her for initial management?

a) Glatiramer

b) Methylprednisolone

c) IFN-β-1a

d) Prednisolone

Question 12:

Which of the following patients have Marburg's variant of multiple sclerosis?

a) A patient with chronic form of multiple sclerosis with no remissions

b) A patient with acute form of multiple sclerosis with remissions

c) A patient with acute form of multiple sclerosis with no remissions

d) A patient with chronic form of multiple sclerosis with remissions

Question 13:

A patient has recurrent optic neuritis of both eyes with extensive transverse myelitis. Visual acuity in the right eye is 6/60 and visual acuity in the left eye 6/18. The patient showed a partial response to steroids. What is the likely diagnosis?

a) Acute disseminated encephalomyelitis

b) Devic's disease

c) Riddoch syndrome

d) Acute fulminant multiple sclerosis

Question 14:

Which of the following drugs will you prescribe for a patient with an acute attack of neuromyelitis optica?

a) Rituximab

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) Mycophenolate mofetil

c) Glucocorticoids

d) Azathioprine

Question 15:

A child presents to the casualty with high-grade fever and altered sensorium. He is suspected to have acute disseminated encephalomyelitis. This condition is associated with which of the following?

a) 2 and 4 only

b) 1 and 3 only

c) 1, 2, 3 and 4 only

d) 2, 4 and 5 only

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 16:

Which of the following statements is incorrect regarding acute disseminated encephalomyelitis?

a) Cerebellar involvement is seen in post-varicella ADEM

b) Autoantibodies to myelin basic protein can be found

c) Seizures do not occur commonly

d) Initial treatment is with high-dose glucocorticoids.

Question 17:

Tigroid white matter on MR imaging is seen in which of the following conditions?

a) Pantothenate kinase deficiency

b) Pelizaeus-Merzbacher disease

c) Neuroferritinopathy

d) Aceruloplasminemia

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	d
2	b
3	d
4	b
5	b
6	c
7	b
8	d
9	b
10	c
11	b
12	c
13	b
14	c
15	c
16	c
17	b

If you purchased this, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Demyelinating disorders do not include biochemical abnormalities of myelin. Disorders occuring due to biochemical abnormalities of myelin are called dysmyelinating disorders.

Multiple sclerosis is the most common immune-mediated, inflammatory demyelinating disorder. It commonly presents in adulthood.

Solution to Question 2:

Multiple sclerosis is an autoimmune disease of the central nervous system. It is characterized by chronic inflammation, demyelination, gliosis (scarring), and neuronal loss.

The peripheral nervous system is spared in MS.

The course of the disease may be relapsing-remitting or progressive. The clinical course of manifestation can be variable ranging from a benign condition to a rapidly evolving and incapacitating disease.

Sold by @itachibot

Solution to Question 3:

HLA-DRB1 is a risk factor for the development of multiple sclerosis, not HLA-B27.

Risk factors supporting the course of development of multiple sclerosis include:

• Cigarette smoking

• Vitamin D deficiency

• Family history, HLA-DRB1

• Exposure to Epstein-Barr virus (Human Herpesvirus 4) after early childhood

Solution to Question 4:

In multiple sclerosis, the slowing of conduction is due to the loss of the myelin sheath.

Under normal circumstances, nerve conduction in myelinated axons occurs in a saltatory manner i.e from one node of Ranvier to another. But, when demyelination sets in, the nerve impulse are unable to travel as fast due to not being saltatory. This produces either a conduction block or delay.

Lesions due to multiple sclerosis begin with perivenular cuffing by inflammatory mononuclear cells. Myelin-specific autoantibodies cause degeneration of myelin sheaths. As lesions evolve, prominent astrocytic proliferation (gliosis) is observed.

Demyelination is the hallmark in the pathology of multiple sclerosis. But, it is not solely a disease affecting the myelin sheath. Partial or total axonal destruction (neuronal loss) can also occur.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

The autoantibodies in multiple sclerosis are directed against both myelin basic protein (MBP) and myelin oligodendrocyte glycoprotein (MOG).

Solution to Question 6:

Remyelination of axons in the central nervous system leads to the formation of shadow plaques.

Shadow plaques develop due to the partial remyelination of the naked axons by surviving oligodendrocytes or those that differentiate from precursor cells.

Solution to Question 7:

Sold by @itachibot

An electric shock-like sensation radiating down the back into the legs is called Lhermitte's sign. It is associated with multiple sclerosis.

It is induced typically by flexion or other movements of the neck. Lhermitte's sign can also occur simultaneously with other disorders of the cervical spinal cord like cervical spondylosis.

Option A: Uhthoff 's phenomenon refers to the unilateral visual blurring that may occur during a hot shower or with physical exercise. It is an exacerbation of the symptoms of multiple sclerosis due to heat.

Option C: Stransky's sign is a clinical sign in which vigorous abduction followed by the sudden release of the little toe causes an extensor plantar reflex. It is seen in pyramidal tract lesions.

Option D: Cornell's sign is a clinical sign in which scratching along the inner side of the extensor hallucis longus tendon elicits an extensor plantar reflex. It is seen in pyramidal tract lesions.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

The MRI image shows Dawson's fingers.

Dawson's fingers are demyelinating plaques through the corpus callosum, arranged at right angles along medullary veins (callososeptal location). They are a characteristic feature of multiple sclerosis and presents as T2 hyperintensities.

MRI is the imaging of choice in a patient with multiple sclerosis due to its high sensitivity.

Option A: Black hole lesions are multiple sclerosis plaques indicative of the chronic stage when they display T1 hypointense signals. They signify axonal destruction and irreversible damage.

Option B: The molar tooth sign refers to the appearance of the midbrain in an axial section in which the elongated superior cerebellar peduncles gives the midbrain the appearance of a molar or wisdom tooth. It is seen in Joubert syndrome.

Option C: Tram-track sign in the brain refers to the parallel calcification of the cortex in patients with Sturge-Weber syndrome.

Solution to Question 9:

The given scenario and the MRI finding of Dawson's fingers are suggestive of multiple sclerosis. Revised McDonald's criteria is used in the diagnosis of multiple sclerosis. It takes into consideration the clinical attacks and objective lesions. In addition, it also involves the assessment of the lesions based on MRI and other imaging techniques.

The Kurtzke expanded disability status scale (EDSS) and functional status (FS) score are widely used measures of neurologic impairment, which are performed following the confirmed diagnosis of multiple sclerosis.

Solution to Question 10:

Sold by @itachibot

The Expanded disability status scale (EDSS) gives a measure of neurological impairment in multiple sclerosis.

EDSS scores <3.5: The patient walks normally, shows no disability

EDSS scores >5.5: Progressive disease, gait impairment, and occupational disability

EDSS scores =10.0: Death due to multiple sclerosis

Solution to Question 11:

Glucocorticoids are used in the clinical management of either first attacks or acute exacerbations.

Intravenous methylprednisolone administration at a dose of 500–1000 mg/day for 3-5 days, followed by a course of oral prednisolone beginning at a dose of 60–80 mg/day and gradually tapered over 2 weeks is given.

These medications reduce the severity and shorten the duration of attacks.

Disease-modifying therapies for relapsing forms of multiple sclerosis are:

- IFN-β-1a
- IFN-β-1b
- Glatiramer acetate
- Natalizumab
- Fingolimod
- Dimethyl fumarate
- Teriflunomide
- Mitoxantrone
- Alemtuzumab

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 12:

Marburg's variant is an acute form of multiple sclerosis, typically with no remissions.

It is a fulminant demyelinating process that in some cases progresses, leading to death within 1–2 years. Marburg's variant does not seem to be associated with infection or vaccination.

Clinical management by high-dose glucocorticoids, plasma exchange, and cyclophosphamide have been tried, with uncertain benefit.

Solution to Question 13:

The given scenario is suggestive of neuromyelitis optica, which is also known as Devic's disease.

Sold by @itachibot

It is an aggressive inflammatory demyelinating disorder characterized by recurrent attacks of optic neuritis and myelitis. Attacks of Optic neuritis can be bilateral or unilateral and myelitis can be severe and involve three or more contiguous vertebral segments.

Core clinical characteristics are:

• Optic neuritis

• Acute myelitis

• Area postrema syndrome: episodes of otherwise unexplained hiccups or nausea or vomiting

• Acute brainstem syndrome

• Symptomatic narcolepsy or acute diencephalic clinical syndrome with NMOSD*-typical diencephalic MRI lesions

• Symptomatic cerebral syndrome with NMOSD*-typical brain lesions

Option A: Acute Disseminated Encephalomyelitis (ADEM) is a demyelinating condition most frequently associated with an antecedent infection

Option C: Riddoch syndrome is the ability to consciously perceive the stimulus in a blind field. It is seen when there is a lesion in the occipital lobe.

Option D: Acute fulminant multiple sclerosis is an acute episode of multiple sclerosis, known as Marburg's variant

*NMOSD stands for neuromyelitis optica spectrum disorder.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 14:

Acute attacks of neuromyelitis optica are managed by high-dose glucocorticoids or plasma exchange.

Drugs used for prophylaxis against relapses include:

• Mycophenolate mofetil

• Rituximab

• Glucocorticoids + azathioprine

Solution to Question 15:

Acute disseminated encephalomyelitis (ADEM) is associated with measles infection, measles vaccination, rubella infection, and rubella vaccination.

ADEM is a demyelinating disease of the central nervous system that typically presents as a monophasic disorder associated with multifocal neurological symptoms and encephalopathy. It may be :

• Post-infectious (95%) - associated infections are with varicella, measles, rubella, mumps, influenza, EBV, HHV-6, HIV, other viruses, and Mycoplasma pneumoniae.

Sold by @itachibot

• Post-vaccination (5%) - presently it is most often associated with measles (most common), mumps, and rubella vaccination.

Solution to Question 16:

Seizures are common in ADEM.

ADEM results from a cross-reactive immune response to the infectious agent (most commonly measles virus) or vaccine which then triggers an inflammatory demyelinating response.

Autoantibodies to MBP and to other myelin antigens have been detected in the CSF of some patients with ADEM.

In ADEM due to chickenpox, cerebellar involvement is often conspicuous.

Initial treatment is with high-dose glucocorticoids.

Solution to Question 17:

Tigroid white matter on MR imaging is seen in Pelizaeus-Merzbacher disease.

It is a rare, X-linked recessive disorder characterized by central nervous system dysmyelination.

The proteolipid protein gene (PLP1) is affected, leading to the failure of axon myelination by oligodendrocytes in the central nervous system.

Clinical Features:

• Nystagmus

• Developmental delay

• Psychomotor delay

• Seizure

• Stridor

• Feeding difficulties

• Ataxia

• Hypotonia progressing to spasticity.

The degree of dysmyelination is correlated with the severity of the clinical manifestations.

Genetic analysis can confirm the diagnosis. MRI shows a lack of myelination and MR spectroscopy reveals aberrant N-acetyl aspartate (NAA) and choline concentrations that reflect axonal and myelination abnormalities.

Unlike other demyelinating leukodystrophies such as metachromatic leukodystrophy and adrenoleukodystrophy, where myelin is formed but subsequently destroyed, PMD shows a failure of myelin formation.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Headache Disorders

Question 1:

What is the most common type of headache?

a) Medication overuse headache

b) Tension-type headache

c) Migraine without aura

d) Cluster headache

Question 2:

What is the most common cause of secondary headache?

a) Head injury

b) Systemic infection

c) Vascular disorders

d) Brain tumor

Question 3:

Which of the following case scenarios do not depict tension-type headache?

a) A 34-year-old woman presenting with a chronic headache for >15 days per month

b) A 41-year-old man presenting with a bilateral headache lasting for 4-6 hours

c) A 38-year-old man presenting with a tight band-like discomfort around the head

d) A 34-year-old woman presenting with a pulsatile headache associated with nausea

Question 4:

A patient comes with complaints of non-pulsatile headache of tightening quality on both sides of his forehead. He describes it as wearing a tight cap. He has no nausea, photophobia, or phonophobia. All of the following drugs can be used in the treatment of this condition except

a) Aspirin

b) Sumatriptan

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

c) Acetaminophen

d) Naproxen

Question 5:

A 41-year-old man has a 4-year history of severe left-sided excruciating episodic headaches. Most episodes are accompanied by ipsilateral lacrimation, conjunctival injection, and rhinorrhea. There are no cutaneous triggers. It does not respond to indomethacin and attacks get triggered by alcohol. What is the likely diagnosis?

a) Trigeminal neuralgia

b) Cluster headache

c) SUNCT

d) Paroxysmal hemicranias

If you purchased this from someone else, you have been scammed. Sold by @Itachibot

Question 6:

A man presents with complaints of a recurrent left-sided stabbing headache. He has been experiencing this every day, along with a runny nose during the episodes, for the last week. These episodes last for about an hour. Examination shows the following. Which of the following is not used for the prophylaxis of this condition?



a) Verapamil

b) Indomethacin

c) Steroids

Sold by @itachibot

d) Lithium

Question 7:

What is the most common positive visual aura seen by patients during an attack of migraine?

a) Scotoma

b) Zig zag lines

c) Transient loss of vision

d) Key hole vision

Question 8:

A 26-year old woman presents with a unilateral throbbing headache associated with nausea and vomiting. She has had similar episodes in the past where the headache usually lasts for 1-2 days. Bright light increases her discomfort and she prefers sitting in a dark room. Which of the following is associated with this condition?

a) Interleukin 1

b) Calcitonin gene-related peptide

c) Leukotriene B4

d) Interleukin 6

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

In patients with familial hemiplegic migraine, which of the following is least likely to be defective?

a) Voltage-gated calcium channel gene

b) Sodium potassium ATPase gene

c) Voltage gated sodium channel

d) Voltage gated chloride channel

Question 10:

A 23-year-old woman experiences severe throbbing headaches on the right side of her forehead. Soon after the headache begins, she becomes very nauseous and occasionally gets

Sold by @itachibot

dizzy. She has also vomited multiple times during these episodes and is subsequently diagnosed with severe migraine. Which among the following is the most efficacious triptan used to treat this condition?

a) Sumatriptan

b) Rizatriptan

c) Frovotriptan

d) Naratriptan

Question 11:

A young woman has been suffering from 1-2 episodes of migraine every week. These episodes are affecting her quality of life. All of the following drugs can be used in the prophylaxis of migraine in this patient except

a) Propranolol

b) Amitriptyline

c) Sumatriptan

d) Valproate

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Sumatriptan can be given in which of the following patients diagnosed with migraine?

a) A 42-year-old woman who had taken ergot alkaloids in the past one week

b) A 34-year-old woman presenting with a positive urine pregnancy test

c) A 48-year-old man with a history of cerebrovascular disease

d) A 51-year-old man with a history of coronary artery disease

Question 13:

What is the use of MIDAS questionnaire?

a) To assess severity of migraine

b) To assess disability of migraine

c) To help decide treatment of migraine

d) To assess risk of migraine

Sold by @itachibot

Question 14:

Which of the following is used in the preventive treatment of a patient with chronic migraine?

a) Onabotulinum toxin A

b) Sumatriptan

c) Metoclopramide

d) Ergotamine

Question 15:

Which of the following is not a trigeminal autonomic cephalalgia?

a) Cluster headache

b) SUNA

c) Hemicrania continua

d) Nummular headache

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

All are true about patients with trigeminal autonomic cephalgia except:

a) Cluster headache is more common in males

b) SUNCT has highest attack frequency rate

c) Lidocaine is effective in SUNCT

d) Cutaneous trigger is seen in cluster headache

Question 17:

A patient was prescribed indomethacin for his complaint of headache and he felt relief after taking the drug. Which of the following could be a likely diagnosis?

a) Paroxysmal hemicrania

b) Cluster headache

c) SUNCT

d) Trigeminal neuralgia

Sold by @itachibot

Question 18:

A 40-year-old male is diagnosed to have SUNCT. It can be distinguished from trigeminal neuralgia by the presence of all except:

a) Excessive lacrimation

b) Nasal congestion

c) Conjuctival injection

d) Short attacks of stabbing pain

Question 19:

All are true about patients with trigeminal neuralgia except:

a) More common in females

b) Pain along V2 and V3 division of trigeminal nerve

c) Deep seated pain

d) No objective signs of sensory loss

Question 20:

A 42-year-old man presented with an extremely painful headache that came on suddenly without warning. It reached maximum intensity within 1 minute and lasted for 5 minutes. Which of the following would you not include as a differential?

a) Subdural hemorrhage

b) Subarachnoid hemorrhage

c) Acute hydrocephalus

d) Pituitary apoplexy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	d

Sold by @itachibot

4	b
5	b
6	b
7	b
8	b
9	d
10	b
11	c
12	a
13	b
14	a
15	d
16	d
17	a
18	d
19	c
20	a

If you purchased this elsewhere, you may have been scammed.

Detailed Explanations

Solution to Question 1:

The most common type of headache is tension-type headache (TTH).

The pain of TTH can generally be managed with simple analgesics such as acetaminophen, aspirin, or NSAIDs. For chronic TTH, amitriptyline is the only proven treatment.

Causes of primary headache, in decreasing order:

• Tension-type headache

• Migraine

• Idiopathic stabbing

• Exertional

• Cluster

Causes of secondary headache, in decreasing order:

• Systemic infection

• Head injury

• Vascular disorders

• Subarachnoid hemorrhage

Sold by @itachibot

• Brain tumor

Solution to Question 2:

The most common cause of secondary headache is a systemic infection.

Causes of primary headache, in decreasing order:

• Tension-type headache

• Migraine

• Idiopathic stabbing

• Exertional

• Cluster

Causes of secondary headache, in decreasing order:

• Systemic infection

• Head injury

• Vascular disorders

• Subarachnoid hemorrhage

• Brain tumor

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 3:

A pulsatile headache associated with nausea is not a feature of tension-type headache (TTH).

Tension-type headache is a chronic head-pain syndrome characterized by bilateral, tight, band-like discomfort. The pain typically builds slowly, fluctuates in severity, and may persist more or less continuously for many days. The headache may be episodic or chronic (present >15 days per month).

	Tension headache	Migraine	Cluster headache
Location	Bilateral	Unilateral	Unilateral
Duration	>30 min (typically 4-6 hr)	4-72 hour	15 min-3 hour
Feature	Steady pain	Pulsating pain	Repetitive brief headache wit h excruciating periorbital pai n
Associatio ns	No photophobia or phonoph obiaNo aura	Nausea, photophobia, phono phobia presentMay have aura	Lacrimation and rhinorrhea presentMay present with Hor ner syndrome
Treatment	Analgesics, NSAIDs, acetami nophen	NSAIDs, triptans, dihydroerg otamine	Sumatriptan, 100% oxygen

Sold by @itachibot

	Tension headache	Migraine	Cluster headache
Prophylaxis	Amitriptyline	Lifestyle changes(eg, sleep,exercise, diet)β-blockersCalcium channel blockersAmitriptyline, topiramate, valproateCGRP antagonists - erenumab, fremanezumab, galcanezumab, eptinezumab	Verapamil

Solution to Question 4:

A bilateral, non-pulsatile headache of tightening quality without any nausea or photophobia suggests a diagnosis of tension-type headache (TTH). Triptans are not useful in the treatment of this condition. They are useful when the patient has a migraine headache.

	Tension-type headache	Migraine	Cluster headache
Treatment	AnalgesicsNSAIDsAcetaminophen	NSAIDsTriptansDihydroergotamine	100% oxygenSumatriptan
Prophylaxis	Amitriptyline	Lifestyle changes(eg, sleep,exercise, diet)β-blockersCalcium channel blockersAmitriptyline, topiramate, valproateCGRP antagonists - erenumab, fremanezumab, galcanezumab, eptinezumab	Verapamil

Solution to Question 5:

This patient's clinical features are suggestive of cluster headache. Patients with cluster headaches are more often smokers with a higher than average alcohol consumption.

Cluster headache causes severe, unilateral periorbital pain with autonomic features, such as unilateral lacrimation, nasal congestion, and conjunctival injection. Occasionally features of Horner's syndrome may be seen. The pain, though severe, is characteristically brief (30–90 minutes). There are 1-8 attacks/day lasting for 15min-3 hours.

Solution to Question 6:

Repetitive brief, unilateral, stabbing headaches associated with lacrimation and rhinorrhea suggest a diagnosis of cluster headaches. Indomethacin is not used for the prophylaxis of these headaches.

Sold by @itachibot

The drug of choice used in the prophylaxis of cluster headaches is verapamil. Steroids and lithium can also be used.

Solution to Question 7:

The most common positive visual aura seen in attacks of migraine is a zig-zag line in front of the eyes.

The most common negative visual aura is scotoma.

The classic migraine is associated with an aura. The aura can be:

• Visual (most common):

• Zigzag lines

• Tunnel vision

• Scotoma

• Transient visual loss

• Sensory:

• Tingling sensation of hands, which spreads proximally

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

Multiple unilateral throbbing headache associated with nausea, vomiting and photophobia suggests a diagnosis of migraine. Calcitonin gene-related peptide (CGRP) is a vasoactive neuropeptide associated with this condition. It is present at vascular terminations of the trigeminal nerve and within the trigeminal nucleus.

CGRP antagonists approved for migraine prevention are:

• Erenumab

• Fremanezumab

• Galcanezumab

• Eptinezumab

Sold by @itachibot



Note:

The FDA has recently approved Atogepant, a calcitonin gene-related peptide (CGRP) receptor antagonist for the preventive treatment of episodic migraine in adults. The recommended dosage is 10 mg, 30 mg, or 60 mg orally once daily. In severe renal impairment or end-stage renal disease it is 10 mg once daily.

Solution to Question 9:

Familial hemiplegic migraine (FHM) is due to defective voltage-gated calcium channel gene, sodium-potassium ATPase gene, and voltage-gated sodium channel. The voltage-gated chloride channel is not defective in this condition.

Type	Mutation
FHM 1	Voltage-gated calcium channel (CACNAIA)gene
FHM 2	Sodium -potassium ATPase (ATPIA2) gene
FHM 3	Voltage-gated sodium channel (SCNIA)

Solution to Question 10:

The most efficacious triptans for moderate to severe acute migraines are rizatriptan and eletriptan.

Oral dosage forms of triptans are the most convenient to use. Still, they may not be practical in patients experiencing migraine-associated nausea and vomiting, for whom injectable and nasal

Sold by @itachibot

spray formulations are helpful.

Standard treatment comprises a 6-mg subcutaneous dose of sumatriptan. All triptans have similar pharmacologic properties but vary slightly in terms of clinical efficacy. Rizatriptan and eletriptan are the most efficacious of the triptans currently available.

Solution to Question 11:

Sumatriptan is used in the treatment of an acute attack of migraine not in prophylaxis. It can be given by oral, subcutaneous, intravenous or intranasal routes.

Drugs used in the prophylaxis of migraine:

• Beta-blockers- propranolol

• Calcium channel blocker- flunarizine (verapamil, nimodipine has no role)

• Antidepressants- amitriptyline, venlafaxine (SSRIs like fluoxetine are not used)

• Anticonvulsants- valproate, topiramate

• 5-HT2 antagonist- pizotifen

• CGRP antagonists - erenumab, framanezumab, galcanezumab, eptinezumab

Other modalities of preventative treatment:

• Greater occipital nerve block

• Onabotulinum toxin A

• Supraorbital transcutaneous stimulation

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Solution to Question 12:

Taking ergot alkaloids in the past one week is not a contraindication for sumatriptan. However, they shouldn't be given together as doing so could result in excessive blood vessel narrowing.

Ergots constrict arteries and triptans narrow normal heart arteries by up to 20%. For this reason, ergots should not be taken with triptans on the same day.

Sumatriptan is a 5HT1B/1D receptor agonist used in an acute migraine attack. It is contraindicated in the following conditions:

• Cerebrovascular disease

• Coronary artery disease

Option B- Sumatriptan is the safest triptan in pregnancy, but its use is recommended only if the benefits outweigh the risks, and if all first line options do not provide relief.

Solution to Question 13:

Sold by @itachibot

The Migraine Disability Assessment Score (MIDAS) questionnaire was put together to help measure the impact of migraine on patients' lives.

Solution to Question 14:

Onabotulinum toxin A is used in the preventive treatment of patients with chronic migraine.

Sumatriptan, ergotamine, and metoclopramide are used in the treatment of an acute attack of migraine.

Drugs used in the prophylaxis of migraine:

• Beta-blockers- propranolol

• Calcium channel blocker- flunarizine (verapamil, nimodipine has no role)

• Antidepressants- amitriptyline, venlafaxine (SSRIs like fluoxetine are not used)

• Anticonvulsants- valproate, topiramate

• 5-HT2 antagonist- pizotifen

• CGRP antagonists - erenumab, framanezumab, galcanezumab, eptinezumab

Other modalities of preventative treatment:

• Greater occipital nerve block

• Onabotulinum toxin A

• Supraorbital transcutaneous stimulation

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 15:

Nummular headache is not a trigeminal autonomic cephalalgia. It is characterized by small circumscribed areas of continuous pain on the head.

Trigeminal autonomic cephalalgia is characterized by pain along the distribution of trigeminal nerve (unilateral) and cranial autonomic symptoms. This is due to increased parasympathetic flow and decreased sympathetic flow.

Solution to Question 16:

The cutaneous trigger is not seen in cluster headache. It is seen in short-lasting unilateral neuralgiform headache with conjunctival injection and tearing (SUNCT).

Solution to Question 17:

Sold by @itachibot

Paroxysmal hemicrania could be the likely diagnosis since it responds to indomethacin.

It is characterized by frequent unilateral, severe, short-lasting, and frequent episodes of headache. They have marked autonomic features ipsilateral to the pain.

Indomethacin (25–75 mg t.i.d) can completely suppress attacks of paroxysmal hemicrania. It is also effective in:

• Hemicrania continuum

• Primary sexual / exercise headache

• Tension-type headache

Solution to Question 18:

Short attacks of stabbing pain are common to both SUNCT and trigeminal neuralgia.

SUNCT has autonomic symptoms such as lacrimation, nasal congestion, conjunctival injection, and salivation. Trigeminal neuralgia does not have autonomic symptoms.

Solution to Question 19:

Trigeminal neuralgia is associated with superficial pain and not deep-seated pain.

It is more common (~ 60% of cases) in females (Option A) over 50 years and is characterized by unilateral lancinating facial pain. This most commonly involves the second and/or third divisions (Option B) of the trigeminal nerve territory. Bilateral or early-onset pain is associated with multiple sclerosis. Unilateral trigeminal nerve compression can be due to the superior cerebellar artery.

Pain is repetitive, severe, and brief and can be triggered by touch, cold, wind, or eating. Physical signs are usually absent, although the spasms may make the patient wince and sit silently (tic douloureux). No signs of sensory loss are present (Option D)

The drug of choice is carbamazepine.

Note: It is important to note that trigeminal neuralgia and trigeminal autonomic cephalalgia (TAC) are different entities. TAC is a group of primary headaches that include - cluster headaches, paroxysmal hemicrania, short-lasting unilateral neuralgiform headache attacks with conjunctival injection and tearing (SUNCT)/short-lasting unilateral neuralgiform headache attacks with cranial autonomic symptoms (SUNA), and hemicrania continua.

Solution to Question 20:

The given clinical scenario is suggestive of a thunderclap headache. Subdural hemorrhage will not produce this type of headache.

Causes of thunderclap headache:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed!

Sold by @itachibot

- Subarachnoid hemorrhage
- Acute hydrocephalus
- Pituitary apoplexy
- Cerebral venous thrombosis
- Hypertensive crisis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Movement Disorders

Question 1:

Which of the following diseases primarily involves the marked structure?



©MARROW

a) Paralysis agitans

b) Depression

c) Alzheimers

d) Huntington chorea

Question 2:

What is the most common gene mutation seen in patients with familial Parkinson's disease?

a) Leucine-rich repeat kinase 2

b) Parkin

c) Alpha-synuclein gene missense mutation

d) PTEN-induced putative kinase 1

Question 3:

Sold by @itachibot

Which of the following features would you not expect to see in patients with Parkinson's disease?

a) Genitourinary disturbances

b) Orthostatic hypotension

c) Exaggerated deep tendon reflexes

d) Micrographia

Question 4:

A 66-year-old man presents with complaints of resting tremor in his right hand, which improves with movement. He also says that he has difficulty initiating movements and his wife says his movements have been slow. On examination, rigidity is observed to passive arm movement and shuffling gait with a stooped posture is noticed. Which of the following statements is false regarding the pathology of this condition?

a) Substantia nigra pars reticulata is unaffected by the disease process

b) Decreased striatal dopamine is seen in this condition

c) Degeneration of dopaminergic neurons in the substantia nigra pars compacta

d) Lewy bodies which containing degenerated tau protein are seen

Question 5:

A patient with Parkinson's disease says that he cannot even hold a coffee mug properly due to tremors. What is the frequency of the tremors likely to be?

a) 4–6

b) 8–10

c) 10–12

d) 6-10

Question 6:

What is the most common psychiatric disturbance seen in patients with Parkinson's disease?

a) Anxiety

b) Depression

c) Abulia

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Sold by @itachibot

d) Visual hallucination

Question 7:

Which of the following is not an example of secondary parkinsonism?

a) Corticobasal syndrome

b) Dementia pugilistica

c) Post-encephalitic PD

d) Drug-induced Parkinsonism

Question 8:

What is the most common cause of secondary parkinsonism?

a) Infection

b) Vascular

c) Neuroleptics

d) Trauma

Question 9:

A patient presented to the OPD with a shuffling gait, slowed movements and speech, and cogwheeling. He was started on a new drug recently. Which of the following drugs cannot cause this condition?

a) Metoclopramide

b) Bezatropine

c) α-Methyldopa

d) Tetrabenazine

Question 10:

Which of the following is a peripheral decarboxylase inhibitor?

a) Benserazide

b) Benztropine

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Tetrabenazine

d) Trihexyphenidyl

Question 11:

A 68-year-old woman presents to the casualty after she tripped over a stone. She has been having trouble sleeping and abdominal discomfort. Her daughter said that she has become forgetful and her movements are very slow. She is diagnosed with Parkinson's disease. Which of her symptoms will be relieved by treatment with levodopa?

a) Forgetfulness

b) Trouble sleeping

c) Abdominal discomfort

d) Slow movements

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Which of the following is a non-ergot dopamine agonist?

a) Bromocriptine

b) Apomorphine

c) Cabergoline

d) Pergoilde

Question 13:

A 75-year-old patient presents with an inability to move and speak. He has been experiencing increasing stiffness and slowness from the past 1 hour. He has an 8-year history of Parkinson's disease and takes his levodopa-carbidopa every day. The last dose was 4 hours back. What drug should be prescribed for this patient?

a) Rasagiline

b) Benztropine

c) Apomorphine

d) Selegillene

Question 14:

Sold by @itachibot

Which of the following drugs has disease-modifying action in Parkinson's disease?

a) Levodopa

b) Carbidopa

c) Apomorphine

d) Selegiline

Question 15:

Which of the following statements regarding the management of non-motor symptoms in patients with Parkinson's disease is false?

a) Depression should not be treated with antidepressants

b) Restless leg syndrome is treated by dopamine agonists

c) Sialorrhea can be managed by anticholinergics

d) Low doses of fludrocortisone or midodrine provide control of autonomic symptoms

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

All are true about levodopa-induced motor complications except:

a) More likely to occur in males

b) Seen in younger individuals with severe disease

c) Seen with higher doses of levodopa

d) Rhythmic movements that predominantly involve the lower extremities

Question 17:

A 65-year-old woman on treatment for Parkinson's disease is brought to the OPD. Her husband is concerned because she has been shopping a lot and reached the maximum spending limit on 2 credit cards in the past week. She has also impulsively been buying several rabbits and raising them in their home. She admits feeling guilty after her purchases. Which of the following drugs is associated with this condition?

a) Dopamine precursors

b) Dopamine agonists

c) Dopamine depletors

d) Dopamine antagonists

Sold by @itachibot

Question 18:

A patient with Parkinson's disease has been scheduled to undergo deep brain stimulation. What is the target site for this procedure?

a) Subthalamic nucleus and globus pallidus internus

b) Subthalamic nucleus and globus pallidus externus

c) Substantia nigra pars compacta and globus pallidus externus

d) Substantia nigra pars reticulata and globus pallidus internus

Question 19:

Which of the following is an example of Parkinson's plus syndrome?

a) Progressive supranuclear palsy

b) Dementia pugilistica

c) Post-encephalitic PD

d) Drug-induced Parkinsonism

Question 20:

Which among the following features is not expected in a patient with atypical parkinsonism?

a) Late speech and gait impairment

b) Absence of rest tremor

c) Poor or no response to levodopa

d) No motor asymmetry

Question 21:

A 60-year-old man presented with Parkinsonism and cerebellar features. MRI showed the finding as shown in the image. What is the likely diagnosis?

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



a) Drug induced parkinsonism

b) Multiple systems atrophy

c) Dementia pugilistica

d) Progressive supranuclear palsy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 22:

All of the following are the core clinical features for the diagnosis of Lewy body dementia except:

a) Fluctuating cognition

b) Recurrent visual hallucination

c) REM sleep behavior disorder

d) Orthostatic hypotension

Question 23:

A 60-year-old man presented with Parkinsonian features which is unresponsive to levodopa. MRI image is given below. What is the probable diagnosis?

Sold by @itachibot



a) Progressive supranuclear palsy

b) Multiple system atrophy type C

c) Hallevorden spatz disease

d) Parkinson's disease

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 24:

An 8-year-old girl exhibits rapid, semi-purposeful, graceful, dance-like, involuntary movements. What is this finding?

a) Hemiballismus

b) Myoclonus

c) Chorea

d) Athetosis

Question 25:

During rounds, you see that a patient diagnosed with stroke exhibits wild, flinging movements on his right side. Where is the lesion likely to be?

a) Caudate nucleus

b) Putamen

c) Subthalamic nucleus

d) Red nucleus

Sold by @itachibot

Question 26:

What is the most common acute drug-induced movement disorder?

a) Akathisia

b) Dystonia

c) Tardive Dyskinesia

d) Myoclonus

Question 27:

Which of the following is the most common movement disorder?

a) Dystonia

b) Chorea

c) Athetosis

d) Essential tremor

Question 28:

A 65-year-old man presents with a tremor in both hands that worsens when he is trying to eat or write something. He says that it gets worse when he is stressed and reduces when he drinks wine. His father also had similar symptoms. On examination, there are no resting tremors. Which of the following drugs can be used to control this condition?

a) Valproate

b) Levetiracetam

c) Primidone

d) Carbamazepine

Question 29:

A 70-year-old male patient is scheduled to undergo deep brain stimulation for essential tremor. What is the target site for the procedure?

a) Subthalamic nucleus and globus pallidus internus

b) Anterior limb of internal capsule

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Substantia nigra pars compacta

d) Ventral intermediate nucleus of thalamus

Question 30:

A patient with a family history of Huntington's disease has come for genetic counselling. All of the following are true about this condition, except:

a) Autosomal dominant inheritance

b) Trinucleotide repeat of CAG sequence

c) Westphal akinetic variant has good prognosis

d) Early onset cognitive impairment seen

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 31:

St. Vitus' dance is the other name for which of the following conditions?

a) Huntigton's chorea

b) Sydenham's chorea

c) Athetosis

d) Hemibalismus

Question 32:

All of the following features can be seen in patients with restless leg syndrome, except:

a) Symptoms worsen during the evening or night

b) Accompanied by an unpleasant sensation in the legs

c) Symptoms that begin or worsen with activity

d) Partial or complete relief by movement

Answer Key

Question No.	Correct Option
1	a

Sold by @itachibot

2	a
3	c
4	d
5	a
6	b
7	a
8	c
9	b
10	a
11	d
12	b
13	c
14	d
15	a
16	a
17	b
18	a
19	a
20	a
21	b
22	d
23	a
24	c
25	c
26	b
27	d
28	c
29	d
30	c
31	b
32	c

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

The structure marked in this transverse section of the brain at the level of superior colliculus is substantia nigra. Degeneration of substantia nigra is seen in Parkinson's disease, also known as paralysis agitans.

Substantia nigra is pigmented due to the presence of neuromelanin granules, which help in the synthesis of dopamine. In Parkinson's disease, there is an absence/reduced pigmentation of the substantia nigra.

Bradykinesia, rigidity, and resting tremor are classic features of Parkinson's disease. Because of motor disability, it is also referred to as paralysis agitans.

Other options:

Option B: Depression is often a comorbid finding with Parkinson's disease, but is not due to damage to the substantia nigra.

Option C: Alzheimer's disease is the most common cause of dementia. It is due to atrophy of the entorhinal cortex, located in the medial temporal lobe.

Option D: Huntington's chorea is an autosomal dominant disorder of movement and cognition primarily involving the striatum (caudate nucleus and putamen).



Solution to Question 2:

The most common form of monogenic familial PD is PARK8, caused by mutations in the leucine-rich repeat kinase-2 (LRRK2) gene.

Familial forms of parkinsonism (designated PARK1 through PARK13) have been associated with causative mutations in a number of nuclear genes. A few of these are:

• The LRRK2 gene mutation causes autosomal dominant PD (PARK8). In this condition, Middle to late-onset PD is seen, which resembles typical PD and is responsive to levodopa.

• Mutations in alpha-synuclein (PARK 1 and PARK4) cause rare forms of familial PD and constitute the major component of Lewy bodies in sporadic PD.

Sold by @itachibot

• Parkin-associated PD (PARK 2) & PINK1-associated PD (PARK6) are autosomal recessive seen in young-onset PD.

• Mutation in the glucocerebrosidase gene (GBA) gene of Gauchers represents the most important risk factor for the development of Parkinson's disease. Patients with PD who are GBA mutation carriers have a younger age at onset.

Solution to Question 3:

Exaggerated deep tendon reflexes are not a feature of Parkinson's disease. Cardinal motor features in PD include:

• Resting tremors

• Bradykinesia

• Rigidity

• Gait disturbance / postural instability

Mnemonic: TRAP - Tremors + Rigidity + Akinesia + Postural instability

• Other motor features include:

• Micrographia

• Masked facies (hypomimia)

• Reduced eye blinking

• Soft voice (hypophonia)

• Dysphagia

• Non-motor features are:

• Anosmia

• Sensory disturbance (pain)

• Mood Disorders (depression)

• Sleep disturbances (rapid eye movement behavior disorder)

• Autonomic:

• GI disturbances

• Genitourinary disturbances

• Sexual dysfunction

• Orthostatic hypotension

• Cognitive impairment

Parkinson's disease(PD) is the second most common neurodegenerative disorder after Alzheimer's disease. UK Brain Bank criteria is used for the diagnosis of PD.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 4:

The given clinical scenario is suggestive of Parkinson's disease and Lewy bodies seen in Parkinson's disease do not contain tau protein. They contain the protein α-synuclein.

Pathologic hallmarks of Parkinson's disease include:

• Degeneration of dopaminergic neurons in the substantia nigra pars compacta (SNc)

• Reduced striatal dopamine

• Intracytoplasmic proteinaceous inclusions called Lewy bodies, which contain the protein α-synuclein

Solution to Question 5:

The frequency of resting tremors in Parkinson's disease is 4–6 Hz.

More than 70 percent of patients with Parkinson's disease have tremors as the presenting feature. The classic parkinsonian tremor begins as a low-frequency, pill-rolling motion of the fingers, progressing to forearm pronation/supination and elbow flexion/extension. Other features of the tremors are:

• Typically asymmetric

• Occurs at rest

• Becomes less prominent with voluntary movement

Frequency of various types of tremors: Physiological tremor (12 Hz) > Essential tremor (8–10 Hz) > Resting tremor in PD (4–6 Hz).

Solution to Question 6:

The most common psychiatric disturbance seen in Parkinson's disease is depression.

The non-motor symptoms seen in Parkinson's disease include:

• Cognitive dysfunction and subcortical dementia

• Psychosis and hallucinations - Visual hallucinations are the most common psychotic symptom

• Mood disorders:

• Depression - most common psychiatric disturbance

• Anxiety - second most common psychiatric disturbance

• Abulia/apathy

• Sleep disorders - insomnia, daytime sleepiness with sleep attacks, restless leg syndrome

• Autonomic dysfunction

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 7:

Corticobasal syndrome is not an example of secondary parkinsonism. It s a type of Parkinson's plus syndrome.

Causes of secondary Parkinsonism include:

• Repeated head trauma—dementia pugilistica

• Drug-induced

• Tumor

• Post-encephalitic PD

• Toxins: manganese, MPTP, carbon monoxide, cyanide

• Vascular

• Liver failure

Solution to Question 8:

Dopamine-blocking agents such as neuroleptics are the most common cause of secondary parkinsonism.

Causes of secondary Parkinsonism include:

• Repeated head trauma—dementia pugilistica

• Drug-induced

• Tumor

• Post-encephalitic PD

• Toxins: manganese, MPTP, carbon monoxide, cyanide

• Vascular

• Liver failure

Solution to Question 9:

The given scenario is suggestive of secondary parkinsonism. Bezatropine, also known as benztropine, is used in the treatment of parkinsonism. It does not cause secondary parkinsonism.

Benztropine is a centrally acting anticholinergic. It has been used for controlling tremors in drug-induced Parkinson's. Side effects include urinary dysfunction, glaucoma, and cognitive impairment.

Drugs causing secondary parkinsonism:

• Neuroleptics (typical antipsychotics)

• Metoclopramide, chlorpromazine

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Dopamine depleting agents (reserpine, tetrabenazine)

• Calcium channel blocker: Flunnarizine, Cinnarizine

• Lithium carbonate, amiodarone, α-Methyldopa

Solution to Question 10:

Benserazide is a peripheral decarboxylase inhibitor.

Options B & D: Benztropine and trihexyphenidyl are centrally acting anticholinergic agents.

Option C: Tetrabenazine is a dopamine depleting agent. It causes drug-induced Parkinsonism.

Solution to Question 11:

Levodopa is useful in alleviating the bradykinesia (slow movement) seen in Parkinson's disease.

Levodopa in Parkinson's disease	
Levodopa benefits	Classic motor symptoms of Parkinson's disease:BradykinesiaRigidityTremor
Levodopa doesnotbenefit	Non-dopaminergic features:FreezingFallingSleep disordersDementiaAutonomic dysfunction (GI disturbances, orthostatic hypotension etc.)

Solution to Question 12:

Apomorphine is a non-ergot dopamine agonist.

Solution to Question 13:

The clinical features of the patient are suggestive of a severe 'off' episode. The rescue agent for the treatment of severe 'off' episodes is apomorphine.

Levodopa-induced motor complications consist of fluctuations in motor response ('on' episodes when the drug is working and 'off' episodes when parkinsonian features return) and involuntary movements are known as dyskinesias which typically complicate 'on' periods.

Apomorphine is a non-ergot dopamine agonist. It has efficacy comparable to levodopa, but it must be administered parenterally (intravenous or subcutaneous) and has a very short half-life and

Sold by @itachibot

duration of activity (45 min).

Solution to Question 14:

Selegiline is a monoamine oxidase B (MAO-B) inhibitor known to have some disease-modifying action in Parkinson's disease. Studies have shown that selegiline significantly delayed the time until the emergence of disability necessitating the introduction of levodopa in untreated PD patients.

MAO-B inhibitors include:

• Selegiline

• Rasagiline (more potent)

Cheese effect: inhibition of MAO-A isoform prevents the metabolism of tyramine in the gut, leading to a potentially fatal hypertensive reaction called the 'cheese effect'. It can be precipitated by tyramine-rich foods such as cheese, aged meat, and red wine.

Selegiline and rasagiline do not functionally inhibit MAO-A isoform and are not associated with cheese effect in doses typically used in clinical practice.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 15:

Depression is treated with selective serotonin reuptake inhibitors (SSRIs). In cases of severe depression, other antidepressants are added.

Manifestations in Parkinson's	Treatment
Depression	SSRIs
Insomnia	Zolpidem
Restless leg syndrome	GabapentenoidsDopamine agonist
Erectile dysfunction	Sildenafil
Postural hypotension	Rehabilitation with postural exerciseAdding salt to the diet and elevating the head of the bed to prevent overnight sodiumnatriuresisLow doses of fludrocortisone or midodrine
Sialorrhea	Anticholinergic

Solution to Question 16:

Levodopa-induced motor complications are more likely to occur in females.

Sold by @itachibot

The complications include transient, stereotypic, rhythmic movements that predominantly involve the lower extremities. It is seen in younger individuals with severe disease and also with higher doses of levodopa.

Solution to Question 17:

The given scenario is suggestive of compulsive shopping. Pathological gambling, hypersexuality, and compulsive eating and shopping are the adverse effects of dopamine agonists. Levodopa can also cause these effects, but to a lesser extent than dopamine agonists.

Solution to Question 18:

The target site for deep brain stimulation in Parkinson's disease is the subthalamic nucleus (STN) and globus pallidus internus (GPi).

It helps improve tremors and reduce both 'off' time and dyskinesia. It does not improve features that fail to respond to levodopa. It does not prevent the development or progression of non-dopaminergic features such as freezing, falling, and dementia.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 19:

Progressive supranuclear palsy is an example of Parkinson's plus syndrome.

Parkinson's plus syndrome or atypical parkinsonism refers to a group of conditions that present with parkinsonism (rigidity and bradykinesia) but are different from PD due to underlying differences in clinical manifestations and pathology. They may have degeneration of the striatum, globus pallidus, cerebellum, and brainstem as well as the SNc. These include multiple system atrophy (MSA), progressive supranuclear palsy (PSP), and corticobasal syndrome (CBS).

PSP is often misdiagnosed as Parkinson's disease (PD). The substantia nigra is damaged in both diseases and this accounts for the overlapping symptoms which are observed. In PSP, impairment of cognitive functions occurs before the motor or ocular signs and this serves as an important distinguishing feature.

Immunohistochemistry broadly classifies them into two subtypes: synucleopathies and tauopathies.

Solution to Question 20:

Early speech and gait impairment are seen in atypical parkinsonism.

Features of atypical parkinsonism include:

• Early speech and gait impairment

Sold by @itachibot

• Absence of rest tremor

• No motor asymmetry

• Poor or no response to levodopa

• Aggressive clinical course

• Symptoms of severe early autonomic involvement

• Early severe dementia also points towards atypical parkinsonism

Solution to Question 21:

Parkinsonism, cerebellar features, and the 'hot cross bun' sign on MRI suggest multiple system atrophy (MSA).

MSA manifests as a combination of parkinsonian, cerebellar, and autonomic features. Based on predominant clinical features it is further divided into :

• MSA-P:

• Predominantly parkinsonian features are seen

• T2 weighted MRI demonstrates hyperintense putaminal rim

• MSA-C:

• Predominantly cerebellar features are seen

• MRI shows cerebellar and brainstem atrophy causing pontine hot cross bun sign

Solution to Question 22:

Orthostatic hypotension is not a core clinical feature of Lewy body dementia (LBD) as per DSM-5.

LBD also known as dementia with Lewy bodies is the second most type of progressive dementia after Alzheimer's disease. Protein deposits, called Lewy bodies, deposit in nerve cells in the brain region involved in thinking, memory, and movement (motor control)

Core features of LBD as per DSM-5 are fluctuating cognition, attention, alertness, recurrent visual hallucination, and Parkinsonian symptoms. Suggestive features are REM sleep behavior disorder and severe neuroleptic sensitivity.

Orthostatic hypotension, erectile dysfunction, and constipation are other features of LBD.

Treatment of LBD includes rivastigmine and donepezil which may improve cognition, and reduce hallucinosis and delusion. Pimavanserin may be used to treat psychosis.

Solution to Question 23:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

An elderly male with a history of features of Parkinsonism nonresponsive to levodopa should raise suspicion for Parkinson plus syndromes which include progressive supranuclear palsy (PSP).

Progressive supranuclear palsy (PSP), also known as the Steele-Richardson-Olszewski syndrome, is a tauopathy and is considered a neurodegenerative disease with no current efficacious treatment. It is characterized by decreased cognition, abnormal eye movements (supranuclear vertical gaze palsy), postural instability and falls, as well as parkinsonian features and speech disturbances.

The atrophy of the midbrain results in hummingbird sign of the brainstem (in the sagittal plane). The preserved pons form the body of the bird, and the atrophic midbrain, the head, with beak extending anteriorly towards the optic chiasm. Midbrain atrophy is best confirmed with the reduction in the midbrain to pons ratio.

Mickey mouse appearance, which is the reduction of anteroposterior midline midbrain diameter, at the level of the superior colliculi is seen on axial imaging.

The following image shows the hummingbird sign:



According to a recent imaging study data, the insula and frontal cortex seem to be involved in the development of anxiety in adults with Parkinson's disease.

Solution to Question 24:

Rapid, semi-purposeful, graceful, dance-like, involuntary movements are seen in chorea.

Sydenham's chorea (also called St. Vitus's dance) is a type of chorea that is typically seen in childhood (5-15 years) with prior exposure to group A streptococcal infection and is thought to be autoimmune in nature. It is characterized by the acute onset of choreiform movements and behavioral disturbances and is self-limiting.

Sold by @itachibot

Hyperkinetic movement disorders:	
Tremor	Rhythmic oscillation of a body part due to intermittent muscle contractions
Dystonia	Involuntary, sustained or repeated muscle contractions often associated with twisting movements and abnormal posture
Athetosis	Slow, distal, writhing, involuntary movements with a propensity to affect arms and hands
Chorea	Rapid, semi-purposeful, graceful, dance-like, non patterned, involuntary movements involving distal or proximal muscle groups
Myoclonus	Sudden, brief (<100ms), jerk-like arrhythmic muscle twitches
Tics	Brief, repeated, stereotyped muscle contractions that can often be suppressed for a short time

if you purcha... you may hav... else, ...d.

Solution to Question 25:

Unilateral wild, flinging movements are called hemiballismus and is seen due to a lesion in the subthalamic nucleus. The most common cause of hemiballismus is stroke.

Solution to Question 26:

The most common acute drug-induced movement disorder is dystonia.

Drug-induced movement disorders are primarily associated with drugs that block dopamine receptors(neuroleptics) or central dopaminergic transmission. Neuroleptics are the commonest cause of this condition. Drug-induced movement disorders are classified as:

• Acute: Dystonia is the most common acute hyperkinetic drug reaction; generalized in children, focal in adults.

• Subacute: Akathisia is the commonest reaction in this category

• Tardive syndromes: Develop after months to years of initiation of the drug.

Sold by @itachibot

• Tardive dyskinesia is the most common. Choreiform movements affecting the mouth, lips, and tongue are seen. In severe cases, trunks, limbs, and respiratory muscles may also be affected.

• In one third, TD remits within 3 months of stopping the drug and improvement occur over several years.

Solution to Question 27:

The most common movement disorder is essential tremor.

It can present in childhood but dramatically increases in prevalence over the age of 70 years. Essential tremor is characterized by a high-frequency tremor 6-10 Hz that predominantly affects the upper extremities.

The tremor is characteristically improved by alcohol and worsened by stress.

	Essential tremor	Parkinson's disease
Bradykinesia■rigidity■ micrographia	Absent	Present
Tremor	Usually postural	Usually resting
Tone	Normal	Cogwheel rigidity

Solution to Question 28:

The given scenario is suggestive of essential tremor. It can be controlled by primidone.

The drugs used to treat essential tremors are:

• Beta-blockers: propranolol, atenolol

• Anticonvulsants:

• Primidone

• Gabapentine

• Topiramate

In severe and drug-resistant cases, surgical treatment involves targeting the ventral intermediate (VIM) nucleus of the thalamus.

Solution to Question 29:

The target for deep brain stimulation in essential tremor is ventral intermediate (VIM) nucleus of the thalamus.

Sold by @itachibot

Disease	Target sites for deep brain stimulation
Parkinson's disease	Subthalamic nucleus (STN)Globus pallidus internus (GPi)
Dystonia	Globus pallidus internus (GPi)
Depression	Anterior limb of internal capsule and cingulum
Obsessive-Compulsive Disorder (OCD)	Anterior limb of the internal capsule
Essential tremor	Ventral intermediate (VIM) nucleus

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 30:

Westphal akinetic variant has the worst prognosis.

Huntington's disease shows an autosomal dominant inheritance. It is a trinucleotide repeat expansion disease with multiple CAG repeats (>40 repeats) in the coding region of the huntingtin gene on chromosome 4p. It demonstrates genetic anticipation with earlier onset if inherited from the father.

It presents in the second to third decade of life with cognitive impairment as an early feature. The dominant cognitive feature of HD is executive dysfunction. Additionally, dementia is seen later as the disease progresses.

Decreased acetylcholine leads to dementia. Decreased GABA and increased dopamine lead to chorea. Treatment consists of haloperidol, tetrabenazine (Dopamine depletor).

The following image shows MRI showing dilated ventricles (ex vacuo ventriculomegaly)- due to atrophy of caudate and putamen:



Sold by @itachibot

MRI showing atrophy of caudate and putamen



Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Solution to Question 31:

Sydenham's chorea was originally called St. Vitus's dance.

Sydenham's chorea is more common in females. It is typically seen in childhood (5-1 5 years) with prior exposure to group A streptococcal infection and is thought to be autoimmune in nature.

It is characterized by the acute onset of choreiform movements and behavioral disturbances.

It is self limited.Treatment is generally restricted to those with severe chorea.The chorea generally responds to dopamine-blocking agents, valproic acid, and carbamazepine.

Solution to Question 32:

In restless leg syndrome (RLS), symptoms begin or worsen with rest. It is also called as Willis-Ekbom disease.

The four core symptoms required for the diagnosis of RLS are:

• An urge to move the legs

• Usually caused or accompanied by an unpleasant sensation in the legs (creepy-crawly feeling, paresthesia, or burning)

• Symptoms that begin or worsen with rest and partial or complete relief by movement

• Worsening during the evening or night

Restless leg syndrome is a typical sensory-motor disorder that generally affects the legs during rest periods, especially sleep. There is an irresistible urge to shake the legs associated with burning sensation, itching, and cramps of the legs. In idiopathic cases, dopaminergic

Sold by @itachibot

hypofunction is postulated.

In patients with restless legs syndrome, iron stores are initially repleted, and the aggravating factors are eliminated. If RLS still persists, initial therapy with a gabapentinoid (pregabalin, gabapentin, gabapentin enacarbil) is preferred, rather than a dopamine agonist (pramipexole, ropinirole, rotigotine). This is due to the high frequency of intermediate and long-term augmentation with dopamine agonists, as well as their risk of impulse control disorders.

FDA-approved drugs for the treatment of restless leg syndrome:

• Gabapentin

• Ropinirole

• Pramipexole

• Rotigotine

A dopamine agonist is used as an alternative in patients with increased risk for side effects from gabapentinoids, such as those with obesity and its complications, a past or present history of moderate to severe depression, a disorder causing gait instability or respiratory failure, and a history of substance use disorder.

Non-pharmacological treatments that can be used in restless leg syndrome are:

• Avoiding alcohol at bedtime

• Massaging affected legs

• Hot baths

• Applying hot or cold to the affected area

• Engaging in moderate exercises

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

ALS and Other Motor Neuron Diseases

Question 1:

Which of the following features are lower motor neuron lesions associated with?

a) Fasciculations

b) Spasticity

c) Proximal muscle weakness

d) Preserved stretch reflexes

Question 2:

Which of the following is referred to as bulbar paralysis?

a) UMN palsy of cranial nerves 1 to 6

b) LMN palsy of cranial nerves 1 to 6

c) UMN palsy of cranial nerves 9 to 12

d) LMN palsy of cranial nerves 9 to 12

Question 3:

Which type of disorder is primary lateral sclerosis?

a) Isolated UMN disorder

b) Isolated LMN disorder

c) UMN + LMN disorder

d) Isolated bulbar disorder

Question 4:

Which of the following is involved in amyotrophic lateral sclerosis?

a) Anterior horn cell + corticospinal tract

b) Anterior horn cell + spinothalamic tract

c) Posterior horn cell + corticospinal tract

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) Ventral root ganglia + spinothalamic tract

Question 5:

A middle-aged man presents with progressive atrophy and weakness of his hands and forearms. On examination, he is found to have spasticity of the legs and brisk jaw jerk. Neuroimaging revealed increased signal in the corticospinal tracts on T2 weighted MRI. What is the most likely diagnosis?

a) Multiple sclerosis

b) Amyotrophic lateral sclerosis

c) Subacute combined degeneration

d) Pregressive spinal muscular atrophy

Question 6:

A young college student presents with multiple episodes of falls while climbing stairs. He stated he had been feeling clumsy and weak for the past few months. On examination, he was found to have bilateral positive Babinski's sign and atrophy of muscles of lower limbs. Deep tendon reflexes were absent. What is the likely diagnosis?

a) Kennedy disease

b) Lou Gehrig's disease

c) Pelizaeus–Merzbacher disease

d) Charcot-Marie-tooth disease

Question 7:

A 42-year-old man who presented with progressive weakness was found to have a mix of upper and lower motor neuron findings on examination. His father and his cousin had passed away in their 40s due to similar symptoms. A mutation of the gene encoding which of the following enzymes is likely implicated here?

a) Superoxide dismutase

b) NADPH oxidase

c) Myeloperoxidase

d) Glutathione peroxidase

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 8:

All of the following features can be seen in patients with amyotrophic lateral sclerosis, except:

a) Split-hand syndrome

b) Orthostatic hypotension

c) Frontotemporal dementia

d) Dropped head syndrome

Question 9:

Which of the following patients is unlikely to have amyotrophic lateral sclerosis?

a) A patient with difficulty in swallowing

b) A patient who has spastic dysarthria

c) A patient who starts laughing or crying inappropriately

d) A patient with bowel and bladder dysfunction

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

What is the diagnosis of amyotrophic lateral sclerosis predominantly based on?

a) MRI findings

b) CSF analysis findings

c) Muscle biopsy findings

d) Clinical criteria

Question 11:

Which of the following is the only drug known to improve survival in patients with amyotrophic lateral sclerosis?

a) Fingolimod

b) Riluzole

c) Tocilizumab

d) Rotigotine

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	a
5	b
6	b
7	a
8	b
9	d
10	d
11	b

Sold by @... If you purchased this from ... else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Lower motor neuron lesions are associated with fasciculations.

Solution to Question 2:

Bulbar paralysis refers to LMN palsy of cranial nerves 9 to 12. It is seen due to disease processes affecting primarily the motor nuclei of the cranial nerves.

The manifestations of bulbar palsy include paralysis of tongue, muscles of swallowing, and facial muscles. There is difficulty handling secretions and maintaining an airway.

Causes of bulbar palsy include:

• GBS

• Polio

• Syringobulbia

Solution to Question 3:

Primary lateral sclerosis is a progressive isolated upper motor neuron disorder.

The spectrum of motor neuron disease includes:

Sold by @itachibot

• Amyotrophic lateral sclerosis (ALS):

• It is the most common form of motor neuron disease

• Includes upper and lower motor neuron pathology (UMN +LMN)

• Primary lateral sclerosis:

• It is a progressive isolated upper motor neuron disorder (UMN)

• Some patients develop lower motor neuron signs later in their clinical course, at which point the disease is called upper motor neuron-onset ALS

• Progressive muscular atrophy:

• It is a progressive isolated lower motor neuron disorder (LMN)

• Some patients develop upper motor neuron signs later in their clinical course, at which point the disease is called lower motor neuron-onset ALS

• Progressive bulbar palsy:

• It is a progressive upper and lower motor neuron (UMN +LMN) disorder of cranial muscles

• This condition may occasionally stay isolated to the bulbar segment, but more commonly, upper and lower motor neuron signs and symptoms spread to involve other segments. This is then referred to as bulbar-onset ALS

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 4:

Amyotrophic lateral sclerosis (ALS) involves the anterior horn cells and corticospinal tract.

ALS is the most common form of progressive motor neuron disease.

The pathologic hallmark of motor neuron degenerative disorders is the death of lower motor neurons and upper, or corticospinal, motor neurons. LMN consist of anterior horn cells in the spinal cord and their brainstem homologs innervating bulbar muscles. UMN originate in layer five of the motor cortex and descend via the pyramidal tract to synapse with lower motor neurons, either directly or indirectly via interneurons.

Although at its onset, ALS may cause selective loss of function of only upper or lower motor neurons, it ultimately causes progressive loss of both categories of motor neurons.

Solution to Question 5:

The most likely diagnosis is amyotrophic lateral sclerosis (ALS).

The clinical hallmark of ALS is the presence of signs and symptoms of both upper and lower motor neuron (UMN + LMN) dysfunction. The signs of UMN dysfunction include spasticity and brisk jaw jerk, while the signs of LMN dysfunction include atrophy and weakness of limbs.

Conventional MRI is usually normal in ALS and is used to exclude other possible causes although increased signal in the corticospinal tracts on T2-weighted MRI is supportive of the diagnosis of ALS.

Sold by @itachibot

Solution to Question 6:

The given scenario with both UMN (positive Babinski sign) and LMN (absent reflexes and wasting of muscles) signs are suggestive of amyotrophic lateral sclerosis (ALS). It is also known as Lou Gehrig's disease, after the famous New York Yankee baseball player who was affected with the disorder.

ALS was first described by Charcot in the 19th century and is also referred to as Charcot's disease.

Option A: Kennedy disease is a gradually progressive neuromuscular disorder characterised by wasting of proximal and bulbar muscles. It is caused by a mutation in the androgen receptor gene.

Option C: Pelizaeus–Merzbacher disease is an inherited leukodystrophy characterized by problems with coordination, motor skills, and learning.

Option D: Charcot-Marie-tooth disease encompasses a group of hereditary sensory and motor neuropathies.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 7:

The given scenario is suggestive of familial amyotrophic lateral sclerosis. This condition is linked to mutations of the gene encoding copper-zinc (Cu, Zn) superoxide dismutase type 1 (SOD1). Familial ALS accounts for about 10% of all ALS cases.

Option B: Mutations in the NADPH oxidase genes cause chronic granulomatous disease (CGD).

Option C: Hereditary deficiency of myeloperoxidase (MPO) predisposes to immune deficiency.

Option D: Glutathione peroxidase deficiency leads to hemolytic anemia.

Solution to Question 8:

Orthostatic hypotension is not a feature seen in classical ALS. Autonomic symptoms are not classically seen in ALS, although a few cases with autonomic insufficiency have been reported and these are classified separately as ALS-plus syndromes.

Option A: The split-hand syndrome describes a frequent pattern of weakness and atrophy in ALS that predominantly involves the median and ulnar-innervated lateral (thenar) hand intrinsic muscles with relative sparing of the medial (hypothenar) muscles.

Option C: There is a well-established link between ALS and frontotemporal executive dysfunction (dementia) that may precede or follow the onset of upper and/or lower motor neuron dysfunction

Option D: Dropped head syndrome is an important clinical sign in ALS caused by severe neck extensor weakness (LMN), which leads to an inability to maintain level head position.

Sold by @itachibot

Solution to Question 9:

The patient with bowel and bladder dysfunction is unlikely to have ALS as these findings are not seen in ALS. Even in the late stages of the ALS, sensory & bowel & bladder functions are preserved.

Fasciculations, muscle cramps, weakness, atrophy, and bulbar palsy are LMN features seen in ALS.

Spasticity, increased reflexes, flexor spasms, and pseudobulbar palsy are UMN features in ALS.

A frequent bulbar UMN symptom seen in ALS is the pseudobulbar affect. This is manifested as inappropriate laughing, crying, or yawning. The observed affect is often mood incongruent or may be triggered by stimuli that would not have elicited such a response prior to the development of pseudobulbar affect.

Solution to Question 10:

The diagnosis of ALS is based on clinical criteria that include the presence of upper motor neuron and lower motor neuron signs, the progression of the disease, and the absence of an alternative explanation.

There is no single diagnostic test that can confirm or entirely exclude the diagnosis of motor neuron disease in general and amyotrophic lateral sclerosis (ALS) in particular.

The revised El Escorial criteria and, more recently, the newer Awaji criteria are being used for the diagnosis of ALS.

Practically, the diagnosis is made when it is suggested by the history and physical examination, supported by electrodiagnostic studies (EMG & NCS), and not excluded by neuroimaging (MRI) and laboratory studies (CSF, blood, and urine analysis). Increased signal in the corticospinal tracts on T2-weighted MRI, if seen, is supportive of the diagnosis of ALS

Muscle biopsy is not a routine part of the diagnostic evaluation of ALS but should be performed if there is clinical suspicion of inflammatory myopathy.

Genetic testing is not a routine part of the diagnostic evaluation in ALS except in cases where familial ALS (10%) is suspected.

Solution to Question 11:

Among the given options, riluzole is the only known drug to have any impact on survival in ALS. It can slow ALS progression to a modest degree. It is a neuroprotective drug that blocks glutamatergic neurotransmission in the CNS.

Edaravone is a newer drug that is a free radical scavenger. It is thought to reduce oxidative stress, which has been implicated in the pathogenesis of ALS. Edaravone was found to slow functional deterioration in some patients with ALS. It is recommended as an adjunct to riluzole for all patients with ALS.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Ataxic Disorders

Question 1:

A patient walks with a stomping gait. When asked to close his eyes and walk, he is unable to do so. Which of the following tracts is probably affected?

a) Spinocerebellar tract

b) Posterior column tract

c) Vestibulospinal tract

d) Rubrospinal tract

Question 2:

Which of the following features would you not expect to find in a patient with cerebellar ataxia?

a) Resting tremor

b) Dysdiadochokinesis

c) Scanning speech

d) Hypotonia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 70-year-old man presented with gait instability, slurring of speech, and intention tremors, for a few months. He was a former smoker who quit 10 years back. After extensive workup, a paraneoplastic cause was suspected, and chest CT was ordered. It showed a 3cm wide mass near the right hilum. What is the most likely diagnosis?

a) Adenocarcinoma lung

b) Non small cell lung cancer

c) Small cell lung cancer

d) Squamous cell carcinoma lung

Question 4:

Sold by @itachibot

What is the most common inherited ataxia?

a) Machado-Joseph disease

b) Friedreich's ataxia

c) Ataxia - telangiectasia

d) Spinocerebellar ataxia

Question 5:

A 24-year-old man presents with complaints of swaying while walking, frequent falls and slurring of speech. He also gives a history of involuntary nodding movements of the head. On examination, you note the following findings. Which of the following regions of the spinal cord is involved first in this condition?

a) Dorsal root ganglion

b) Spinocerebellar tract

c) Lateral corticospinal tract

d) Cerebellum

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A middle-aged vegan woman presented with tingling and numbness over the limbs and progressive fatigue. Her peripheral smear is given below. Which of the following neurological manifestation would you not expect to see in this condition?



Sold by @itachibot

a) Sensory ataxia

b) Optic atrophy

c) Psychosis

d) Myopathy

Question 7:

A 30-year-old man presented with swaying while walking, tingling in palms and feet, and blurring of vision. On examination, deep tendon reflexes are absent, proprioception and vibration sense are absent over the lower limbs. MRI spine reveals diffuse demyelination. What is the diagnosis?

a) Multiple sclerosis

b) Tabes dorsalis

c) Transverse myelitis

d) Subacute combined degeneration

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 8:

The investigations of a male truck driver are given below. Which of the following is he unlikely to present with?

a) Ataxia

b) Argyll Robertson pupil

c) Bladder dysfunction

d) Hyperreflexia

Question 9:

Which of the following is false regarding ataxia telangiectasia?

a) Presents in 4th to 5th decade of life.

b) Autosomal recessive inheritance is seen

c) Loss of Purkinje cells occurs

d) Increased incidence of lymphomas is seen

Question 10:

Sold by @itachibot

An elderly woman is brought to the OPD with complaints of behavioral change, a history of multiple falls, urinary incontinence, and dementia. What is your diagnosis?

a) Normal pressure hydrocephalus

b) Frontotemporal dementia

c) Parkinson disease

d) Creutzfeldt-Jakob disease

Answer Key

Question No.	Correct Option
1	b
2	a
3	c
4	b
5	a
6	d
7	d
8	d
9	a
10	a

Detailed Explanations

Solution to Question 1:

The given clinical scenario is suggestive of the involvement of the posterior column.

Posterior column carries proprioception which is essential for coordination of motor movements. Loss of this leads to the development of a stomping gait. This is seen in cases of vitamin B12 deficiency leading to subacute combined degeneration, tabes dorsalis or diabetic neuropathy.

Solution to Question 2:

Intention tremor is a feature of cerebellar ataxia, not resting tremor.

Cerebellar ataxia is characterized by the following symptoms:

Sold by @itachibot

• Gait impairment

• Unclear speech or scanning speech

• Blurred vision due to nystagmus

• Dysdiadochokinesis

• Intention tremors - tremor on goal-oriented movements

Bilateral and symmetric involvement suggests the following etiologies:

• Genetic - SCA gene mutations

• Metabolic - Vitamin B1 or B12 deficiency

• Immune - Antigliadin antibody syndrome

• Toxic - Phenytoin, amiodarone, alcohol

Focal, unilateral symptoms with headache, impaired level of consciousness, ipsilateral cranial nerve palsies and contralateral weakness suggest a space-occupying cerebellar lesion.

Sold by @itachibot
If you purchased this from someone else, may have been scammed.

Solution to Question 3:

The given case scenario is suggestive of paraneoplastic cerebellar ataxia, which is most likely to be caused by small cell lung cancer here.

Paraneoplastic syndromes causing cerebellar ataxia are associated with various antibodies:

Sometimes, paraneoplastic ataxias may be the only presenting symptom of the underlying cancer. In paraneoplastic cerebellar ataxias, serum anti-glutamic acid decarboxylase (GAD) antibodies may be elevated, affecting speech and gait.

Cancer	Antibody seen
Breast and ovarian cancers	Anti-Yo antibody
Small-cell lung cancer	Anti-PQ-type voltage-gated calcium channel autoantibody
Hodgkin's disease	Anti-Tr antibodies

Solution to Question 4:

Friedreich's ataxia is the most common form of inherited ataxia (autosomal recessive), comprising one-half of all hereditary ataxias.

Friedrich's ataxia has autosomal recessive inheritance and presents before 25 years of age. The median age of death is 35 years. Women have a significantly better prognosis than men.

Friedrich's ataxia presents with clinical features such as dysarthria, progressive staggering gait, frequent falls and titubation. Other neurological symptoms include nystagmus, loss of fast

Sold by @itachibot

saccadic movements, loss of vibratory and proprioceptive sensation, and ataxia or trunk and limb movements. The lower extremities are more severely involved than the upper ones.

Cardiac involvement occurs in 90% of patients. It presents with cardiomegaly, symmetric hypertrophy, and conduction defects. Musculoskeletal deformities are common and include progressive scoliosis, pes cavus, and pes equinovarus.

Solution to Question 5:

The given case scenario is suggestive of Friedrich's ataxia, in which the dorsal root ganglion cells are involved first.

The primary sites of pathology are the spinal cord, dorsal root ganglion cells, and the peripheral nerves.

Spinal cord tracts are involved, and degeneration occurs predominantly in the spinocerebellar tracts, lateral corticospinal tracts, and posterior columns. Degeneration of cranial nerves (glossopharyngeal, vagus, hypoglossal) and deep cerebellar nuclei can occur. Peripheral nerves are also involved, and there's a loss of large myelinated fibers. Cerebellar atrophy is also seen.

Cardiac changes occur due to myocyte hypertrophy, fibrosis, and degeneration of cardiac ganglia.

Note: Friedrich's ataxia may also be associated with vitamin E deficiency.

Solution to Question 6:

The given case scenario and smear showing macroovalocytes and hypersegmented neutrophils, is suggestive of megaloblastic anemia caused by vitamin B12 deficiency. Myopathy is not a feature of vitamin B12 deficiency.

Vitamin B12 deficiency may produce the following neurological manifestations:

- Loss of vibration sense and proprioception
- Paresthesia - upper limb paresthesias occur before lower limbs are affected.
- Unsteady gait - due to sensory ataxia
- Peripheral neuropathy
- Optic neuropathy
- Psychiatric manifestations such as:
- Irritated behavior
- Severe dementia
- Frank psychosis

Solution to Question 7:

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

The given case scenario is suggestive of subacute combined degeneration (SACD) due to vitamin B12 deficiency.

SACD presents with the following features:

• Loss of vibration sense and proprioception

• Paresthesia - upper limb paresthesias occur before lower limbs are affected.

• Unsteady gait - due to sensory ataxia

• Loss of deep tendon reflexes

• Peripheral neuropathy

• Optic neuropathy - can present as blurring of vision

• Psychiatric manifestations such as:

• Irritated behavior

• Severe dementia

• Frank psychosis

MRI spine shows myelopathy and areas of demyelination. Myelopathy of SACD tends to be diffuse rather than focal.

Treatment is replacement therapy is with 1000 µg of intramuscular vitamin B12 repeated at regular intervals or by oral vitamin B12.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

The given case scenario is suggestive of tabes dorsalis. It presents with loss of reflexes in lower limbs, not hyperreflexia.

Tabes dorsalis (or syphilitic myelopathy) is a slow degeneration and demyelination of the neural tracts. It is seen in tertiary syphilis. It primarily involves the dorsal columns of the spinal cord and dorsal roots.

It presents with the following clinical features:

• Loss of reflexes in lower limbs

• Romberg's sign

• Bilateral Argyll Robertson pupils - accommodation reflex present, pupillary reflex absent

• Loss of position sense and vibration

• Stabbing/pricking pain in the legs - this is fleeting or repetitive

• Ataxia of the legs and gait disturbances

• Paresthesias

• Bladder disturbances

• Acute abdominal pain with vomiting - visceral crisis

Note: Diabetic polyradiculopathy may simulate tabes.

Sold by @itachibot

Solution to Question 9:

Ataxia-telangiectasia is an autosomal recessive condition presenting in the first decade of life.

It is associated with a high incidence of recurrent pulmonary infections, lymphomas, Hodgkin's disease, breast cancer, and acute T cell leukemias.

Histology shows loss of Purkinje cells, granules, and basket cells in the cerebellar cortex. There is a loss of neurons in the deep cerebellar nuclei, inferior olives of the medulla, and anterior horn and dorsal root of the spinal cord.

Clinical features of ataxia-telangiectasia include:

- Progressive telangiectatic lesions
- Oculomotor apraxia
- Truncal and limb ataxia
- Dysarthria
- Extensor plantar reflexes
- Myoclonic jerks
- Areflexia
- Distal sensory deficits
- Nystagmus
- Cellular and humoral immunodeficiency (absent/ill-developed thymus)

If you purchased this from someone else, you may have been scammed. Sold by @Itachibot

Solution to Question 10:

The given case scenario describes an elderly woman with urinary incontinence, dementia (cognitive decline), and a history of multiple falls (suggestive of a gait abnormality or ataxia). These form the classical triad of normal pressure hydrocephalus (NPH).

Both frontotemporal dementia (Option B) and Parkinson's disease (Option C) are common causes of dementia associated with falls, but urinary incontinence is not commonly appreciated. NPH is, therefore, the most likely diagnosis.

Normal pressure hydrocephalus refers to a disproportionate enlargement of the ventricles compared to the degree of cortical atrophy. It is a communicating hydrocephalus caused by delayed resorption of cerebrospinal fluid. Out of the classical triad of symptoms, most patients present with one or two. Hence, neuroimaging showing dilated ventricles is essential, as well as an L- dopa challenge test to rule out parkinson's disease is recommended.

A characteristic MRI of NPH is given below, showing dilated lateral ventricles which are out of proportion to sulcal deepening:

Sold by @itachibot



Sold by @itachi... else,
If you purchased this from some...
you may have been scammed.

Sold by @itachibot

Cranial Nerve Disorders

Question 1:

Which of the following cranial nerves are affected in cavernous sinus syndrome?

a) III, IV and V

b) IV, V and VI

c) VI and VII

d) III, IV, V and VI

Question 2:

Which of the following statements is true regarding trigeminal neuralgia?

a) The region of ophthalmic division of trigeminal nerve is always spared

b) The pain usually last for a few hours

c) The condition is also known as tic douloureux

d) It is more common in men

Question 3:

The high frequency of trigeminal neuralgia in patients with multiple sclerosis is mainly attributed to the involvement of _____.

a) Oligodendrocyte-derived myelin covering the beginning of the nerve

b) Oligodendrocyte-derived myelin covering the terminal part of the nerve

c) Schwann cell–derived myelin sheath covering the beginning of the nerve

d) Schwann cell–derived myelin sheath covering the terminal part of the nerve

Question 4:

In patients with trigeminal neuralgia, the nerve is most commonly compressed by _____.

a) Superior cerebral artery

b) Inferior cerebral artery

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Superior cerebellar artery

d) Inferior cerebellar artery

Question 5:

A food vlogger presents with complaints of episodes of piercing electric-shock like pain on the left side of her face. The episodes are triggered by chewing food and last for a few minutes. She is further distressed as it interferes with her work. She has tried over the counter analgesics to no avail. What is the drug of choice for her?

a) Gabapentin

b) Lamotrigine

c) Carbamazepine

d) Amitriptyline

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 6:

Which of the following is false about Bell's palsy?

a) Hyperacusis may be present

b) Pregnancy is a risk factor

c) MRI shows enhancement of geniculate ganglion

d) Maximal weakness within 72 hours

Question 7:

Which of the following viruses is most commonly implicated in Bell's palsy?

a) Cytomegalovirus

b) Herpes simplex virus

c) Epstein-Barr virus

d) Varicella zoster virus

Question 8:

Which of the following is true about Bell's phenomenon?

a) Eyeballs roll upwards on attempting to open the eyes

Sold by @itachibot

b) Eyeballs roll downwards on attempting to close the eyes

c) It is a normal phenomenon evident in facial paralysis

d) It is an abnormal phenomenon evident in facial paralysis

Question 9:

On returning from work, a man noticed that his mother had developed mild right-sided facial drooping. Believing it to be due to a stroke, he immediately took her to the hospital. After extensive evaluation, the doctor ruled out stroke and diagnosed her with Bell's palsy. Which of the following would not be a part of the management of her condition?

a) Paper tape to depress the upper eyelid during sleep

b) Prednisone 60-80 mg daily during the first 5 days

c) Massage of the weakened muscles

d) Acyclovir 1g twice daily for 10 days

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Question 10:

A 35-year-old patient presents with facial weakness and pain for the past 2 days. The following signs are seen on examination. What is the cause?



a) UMN facial nerve palsy

b) LMN facial nerve palsy

c) Hemifacial spasm

d) Facial synkinesis

Sold by @itachibot

Question 11:

Identify the correct type of facial nerve palsy in the following patients based on their clinical features:

a) 1 - Right LMN, 2 - Left UMN

b) 1 - Right UMN, 2 - Right LMN

c) 1 - Left UMN, 2 - Right LMN

d) 1 - Left LMN, 2 - Right UMN

Question 12:

During the neurological examination of a patient, you note that he has bilateral lower motor neuron palsy of the facial nerve. Which of the following is least likely to cause this?

a) Guillain-Barre syndrome

b) Sarcoidosis

c) Lymphoma

d) Diabetes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

You are evaluating a woman with unilateral facial weakness. However, you notice forehead wrinkling is intact. On examination, which of the following findings would you not expect?

a) Absent Bell's phenomenon

b) Normal emotional response

c) Normal corneal reflex

d) Normal taste sensation

Question 14:

Which of the following statements is true about facial synkinesis?

a) Refers to simultaneous bilateral facial nerve palsy

b) Refers to facial nerve palsy associated with trigeminal neuralgia

c) Occurs due to anomalous regeneration of the facial nerve fibers

Sold by @itachibot

d) Loss of emotional response is seen in this condition

Question 15:

Which of the following is false regarding hemifacial spasm?

a) The contractions are painful and irregular

b) It is a complication of Bell's palsy

c) Carbamazepine is used for treatment

d) Local injection of botulinum toxin is effective

Question 16:

Which of the following are features of Melkerrson-Rosenthal syndrome?

a) 1, 5

b) 2, 4, 5

c) 2, 3, 4

d) 1, 3, 4

Question 17:

A 70-year-old man presented with complaints of sudden stabs of pain in the throat on yawning. Each episode only lasted for a few seconds and was exacerbated on trying to clear the throat. Which of the following findings can you see on examination?

a) Tongue deviated to one side

b) Loss of taste sensation

c) Bradycardia

d) Decreased movement at TMJ

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	d

Sold by @itachibot

2	c
3	a
4	c
5	c
6	d
7	b
8	c
9	d
10	b
11	c
12	d
13	b
14	c
15	a
16	d
17	c

Sold to ... If you purchased this ... else, you may have been sc... d.

Detailed Explanations

Solution to Question 1:

The cranial nerves III, IV, V, and VI are affected by cavernous sinus syndrome.

The cavernous sinus syndrome is a condition caused by any pathology involving the cavernous sinus. The oculomotor (III), trochlear (IV) and trigeminal (V) nerves pass in relation to the lateral wall of the cavernous sinus. The abducent (VI) nerve passes through the cavernous sinus.

Patients with this condition usually present with the following features:

• Orbital or facial pain

• Orbital swelling and chemosis due to occlusion of the ophthalmic veins

• Fever

• Oculomotor neuropathy (III, IV, and VI)

• Trigeminal neuropathy (V)

Causes of cavernous sinus syndrome include:

• Cavernous sinus thrombosis - most common

• Carotid artery aneurysm

• Carotid-cavernous fistula

• Tumors such as meningioma.

Sold by @itachibot



If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 2:

Tic douloureux is another term for trigeminal neuralgia.

Trigeminal neuralgia is characterized by frequent paroxysms of severe pain in the lips, gums, cheek, or chin. It is sometimes seen in the distribution of the ophthalmic division of the trigeminal nerve.

Middle-aged and elderly females are primarily affected. The pain usually lasts for a short time (seconds to minutes) but be may intense enough that the patient winces, hence the term 'tic'.

A characteristic feature is the presence of trigger zones, typically on the face, lips, or tongue. Patients usually report excruciating pain during activities such as washing the face or brushing the teeth. However, no signs of sensory loss can be demonstrated.

Solution to Question 3:

The high frequency of trigeminal neuralgia in patients with multiple sclerosis occurs due to the involvement of oligodendrocyte-derived myelin covering the beginning of the nerve.

Multiple sclerosis is a disorder of oligodendrocyte myelin. It should be ruled out when trigeminal neuralgia is bilateral or develops in a young adult. In such cases, the cause is a demyelinating plaque near the root entry zone of the fifth cranial nerve in the pons. In these patients, a facial sensory loss is also often found in association.

Solution to Question 4:

Sold by @itachibot

The superior cerebellar artery most often compresses the trigeminal nerve as it emerges from the brainstem. This causes lancinating pain in the distribution of the nerve which is known as trigeminal neuralgia.

Though trigeminal neuralgia can be diagnosed by the clinical features, MRI is recommended to define the underlying cause. In cases refractory to medical management, it also helps to plan for surgical decompression.

Solution to Question 5:

The given clinical scenario is suggestive of trigeminal neuralgia and the drug of choice is carbamazepine.

Oxcarbazepine is an alternative to carbamazepine that has less bone marrow toxicity and is equally efficacious.

Other drugs used are lamotrigine, gabapentin and baclofen.

Solution to Question 6:

Maximal weakness occurs within 48 hours as a rule in Bells' palsy.

Bell's palsy is an idiopathic, peripheral facial paralysis or paresis of acute onset due to a lower motor neuron (LMN) lesion of the facial nerve.

Clinical presentation of Bell's palsy:

• Pain behind the ear (may precede paralysis and last for 1-2 days)

• Bell's phenomenon

• Asymmetrical face

• Hyperacusis (due to stapedial paralysis)

• Loss of taste (involvement of chorda tympani).

MRI may reveal swelling and uniform enhancement of the geniculate ganglion and facial nerve.

Approximately 80% of patients recover within a few weeks or months. The presence of incomplete paralysis in the first week is the most favorable prognostic sign. Recurrence may be seen in some cases.

Solution to Question 7:

Herpes simplex virus is most commonly implicated in causing Bell's palsy.

Varicella-zoster virus is the second most common cause.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 8:

Bell's phenomenon refers to a normal phenomenon that becomes evident in facial paralysis. The eyeballs roll up and outwards on attempting to close the eyes.

It is also known as the palpebral oculogyric reflex. It is a normal protective reflex resulting in the elevation of globes while blinking to protect the eyes from injury. It also serves to provide a wiper blade-like cleansing function of the cornea by the eyelid.

However, it occurs behind the closed eyelids in healthy individuals. It becomes noticeable in Bell's palsy as the eyelids fail to close completely.

The image below shows a patient with evident Bell's phenomenon.

Solution to Question 9:

Acyclovir may be considered as add-on therapy for patients with severe Bell's palsy as 400mg five times daily for 10 days. Monotherapy is not recommended.

Treatment of Bell's palsy involves prednisone 60-80 mg daily during the first 5 days and then tapered over the next 5 days. Randomized trials found no added benefit of the antiviral agents valacyclovir and acyclovir when compared to glucocorticoids alone.

The use of paper tape to depress the upper eyelid during sleep prevents corneal drying.

For patients with permanent paralysis, cosmetic surgical procedures have been used to restore a relatively symmetric appearance to the face.

Solution to Question 10:

The features are suggestive of lower motor neuron (LMN) type of facial palsy.

Sold by @itachibot

In the image, the absence of wrinkling over the right side of the forehead and incomplete closure of the right eyelid is characteristic of LMN type of facial palsy.

In upper motor neuron (UMN) lesion, the upper part of the face is spared due to bilateral UMN innervation.

Features of facial nerve palsy:

Feature	UMN	LMN
Side affected	Contralateral	Ipsilateral
Facial muscles involved	Lower	Both upper and lower
Forehead wrinkling	Present	Absent
Emotional incontinence	Present	Absent
Corneal reflex	Normal	Absent
Taste	Intact	Affected
Atrophy of facial muscles	Absent	Present



Note: Deviation of the angle of mouth should not be confused with drooping of the angle of mouth. Deviation of the angle of the mouth occurs on the normal side due to the action of unaffected facial muscles, while drooping occurs on the affected side.

Solution to Question 11:

Patient 1 has left UMN facial nerve palsy and patient 2 has right LMN facial nerve palsy.

In an upper motor neuron (UMN) lesion, wrinkling on the forehead and eyelid closure remains intact. In this lesion, the contralateral side of the face is affected. The first-order neurons from

Sold by @itachibot

the left cerebral hemisphere cross over (left to right) at the cortical level and relay in the right facial nerve nucleus in the pons. From here the second-order neurons (right facial nerve) descend and supply the right side of the face.

Hence, a left UMN lesion causes paralysis of the lower half of the face on the right side. In facial palsy, the angle of the mouth deviates towards the normal side. In this patient, since the mouth deviates towards the left, the right facial nerve is affected.

In lower motor neuron (LMN) lesions, there is an absence of wrinkling on the forehead and incomplete closure of the eyelid on the ipsilateral side of the face.



Feature	UMN	LMN
Side affected	Contralateral	Ipsilateral
Facial muscles involved	Lower	Both upper and lower
Forehead wrinkling	Present	Absent
Emotional incontinence	Present	Absent
Corneal reflex	Normal	Absent
Taste	Intact	Affected
Atrophy	Absent	Present

Solution to Question 12:

Diabetes is associated with unilateral LMN palsy.

Common causes of facial nerve palsy:

Sold by @itachibot

	UMN	LMN
Unilateral	Usually vascularCerebral tumourMultiple sclerosis	Bell's palsyParotid tumourHead injuriesSkull base tumoursDiabetesHypertension
Bilateral	Vascular(infarct)Motor neuron disease	Guillain-Barré syndromeSarcoidosisLeukemiaLymphomaLeprosy

Solution to Question 13:

The given features are suggestive of an upper motor neuron (UMN) lesion of the facial nerve in which the emotional response is affected. Patients with UMN lesions of the facial nerve have emotional incontinence i.e. involuntary, uncontrollable laughter, or crying with a paradoxical deficit of facial expression.

Normally cerebellum modulates motor outputs in response to sensory inputs. UMN facial palsy (type of pseudobulbar palsy) is associated with disruption of cortico-ponto-cerebellar pathways leading to loss of cerebellar modulation resulting in:

• Emotional dysmetria (difficulty in fine-tuning emotional responses)

• Emotional disinhibition (excessive emotional output)

Difference between UMN and LMN facial palsy:

Feature	UMN	LMN
Side affected	Contralateral	Ipsilateral
Facial muscles involved	Lower	Both upper and lower
Forehead wrinkling	Present	Absent
Emotional incontinence	Present	Absent
Corneal reflex	Normal	Absent
Taste	Intact	Affected
Atrophy	Absent	Present

Solution to Question 14:

Facial synkinesis occurs due to anomalous regeneration of facial nerve fibers.

Following facial nerve injury , the recovering nerve fibers may aberrantly regenerate and supply other muscles. This results in involuntary muscular movements in the face during voluntary facial movements.

If you bought this from someone else, it may have been scammed. Sold by @itachibot

Sold by @itachibot

Some examples include:

• Retraction of the mouth on closure of the eyelids:

• Seen when fibers originally connected with the orbicularis oculi aberrantly innervate the orbicularis oris

• 'Crocodile tears' also known as Bogorad syndrome:

• It refers to anomalous tearing occurring with activity of the facial muscles, such as eating.

• Due to aberrant innervation of the lacrimal gland.

Note: Frey's syndrome is an aberrant connection between the auriculotemporal nerve and the great auricular nerve, resulting in gustatory sweating.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 15:

Hemifacial spasms are painless irregular contractions occurring on one side of the face.

They commonly occur as sequelae to Bell's palsy. It can also be seen in irritative lesions of the facial nerve such as acoustic neuroma and basilar aneurysm.

Treatment is with carbamazepine or gabapentin. In resistant cases, baclofen may be tried. Local injection of botulinum toxin and surgical decompression are alternative options.

Solution to Question 16:

Melkersson-Rosenthal syndrome is a rare disease characterised by the following triad:

• Recurrent facial nerve palsy

• Facial edema (particularly labial)

• Plication of tongue or fissured tongue

Sold by @itachibot

Facial paralysis along with vesicular rash in the external auditory canal and pinna is seen in Ramsay-Hunt syndrome. It occurs due to herpes zoster infection and is also called herpes zoster oticus.

Congenital facial weakness combined with abnormal ocular abduction is seen in Moebius syndrome. Patients may present with bilateral concurrent facial nerve paralysis.

Solution to Question 17:

The clinical features are suggestive of glossopharyngeal neuralgia which is associated with cardiac symptoms such as bradycardia, asystole, hypotension, and fainting.

It involves the glossopharyngeal (IX) and sometimes portions of the vagus (X) nerve. The condition resembles trigeminal neuralgia but is less common. There is no demonstrable sensory or motor deficit.

It can result from vascular compression, multiple sclerosis or tumors.

The drug of choice is carbamazepine. If unsuccessful, surgical procedures are indicated:

• Microvascular decompression (if vascular compression is evident)

• Rhizotomy of glossopharyngeal and vagal fibers in the jugular bulb

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Infections of the Nervous System

Question 1:

A 25-day-old baby is brought to the hospital with complaints of poor feeding and lethargy. On examination, he is found to have bulging fontanelles and hypothermia. Which of the following is the most common cause of his condition?

a) Group B streptococci

b) L. monocytogenes

c) E. coli

d) N. meningitidis

Question 2:

Which of the following is the most likely cause of meningitis in a 30-year-old patient?

a) Streptococcus pneumoniae

b) Staphylococcus aureus

c) Neisseria meningitidis

d) Haemophilus influenzae

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 66-year-old patient presents with fever, neck rigidity, and headache for the past 3 days. He has a history of metastatic non-small cell lung cancer for which he had received intrathecal chemotherapy via an Ommaya reservoir 10 days ago. On examination, Kernig and Brudzinski's signs are positive. Which organism is most likely implicated here?

a) Streptococcus pneumonia

b) Coagulase negative staphylococci

c) Staphylococcus aureus

d) Gram negative bacilli

Question 4:

Sold by @itachibot

A 45-year-old man presents with fever, headache, photophobia and neck rigidity since the last 2 days. He had undergone a heart transplant one year back and is on immunomodulatory drugs. During examination, the following sign is elicited. What is the therapy of choice for this patient?

a) Cefepime + Vancomycin + Ampicillin

b) Ceftriaxone + Vancomycin

c) Isoniazid + Rifampicin + Prazinamide+ Ethambutol

d) Liposomal amphotericin B + Fluorocytosine

Question 5:

A 21-year-old female complains of fever, headache and vomiting from the past 3 days. She has been living in the army recruitment camp for the last three weeks. On examination, she has neck stiffness and a maculopapular rash was noted over the trunk and extremities. Which of the following is false about the disease she's likely suffering from?

a) Carriers of this disease can be treated with Rifampicin

b) It can lead to adrenal hemorrhage

c) Terminal complement deficiency increases the risk of this disease

d) Metronidazole is the drug of choice

Question 6:

If you purchased this from somewhere else, you may have been scammed.

Sold by @itachibot

A patient came with complaints of high-grade fever and altered sensorium. He was diagnosed to be suffering from meningococcal meningitis. Which of the following is the most appropriate empirical treatment option?

a) Penicillin G

b) Ceftriaxone

c) Cefotetan

d) Cefoxitin

Question 7:

Which drug is used for mass chemoprophylaxis for meningococcal meningitis?

a) Ciprofloxacin

b) Chloramphenicol

c) Tetracycline

d) Penicillin

Question 8:

A 45-year-old patient with a history of fever for 1 month presents with headache and ataxia. Imaging reveals dilated ventricles and significant basal exudates. Which of the following would most likely be seen in the CSF?

a) Lymphocytosis, low glucose, high protein

b) Lymphocytosis, normal glucose, high protein

c) Lymphocytosis, low glucose, normal protein

d) Lymphocytosis, high glucose, high protein

Question 9:

A patient comes to ED with fever and headache. On examination he has neck stiffness. CSF analysis was done and it shows the following findings. Most likely diagnosis is:

a) TB

b) Neisseria Gonorrhoea

c) Cryptococcus

d) Coxsackie

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 10:

A college student presents with fever, headache, and confusion. Lumbar puncture was positive for HSV DNA, detected by PCR. An MRI brain is done to localize the lesions. Which area on the brain is most likely to show lesions?

a) Parietal lobe

b) Frontal lobe

c) Temporal lobe

d) Cerebellar cortex

Question 11:

A 35-year-old man presents with fever, headache, seizures and aphasia. MRI shows hyperintensity in the left frontotemporal region. Which of the following would likely be seen in the CSF analysis?

a) Glucose < 20 mg/dl

b) 0-5 WBC/microliter

c) 100 RBC/ microliter

d) Protein < 40mg/dl

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

A 60-year-old man with progressive dementia of recent onset presents with intermittent irregular jerky movements. The EEG is given below. What is the most likely diagnosis?

Sold by @itachibot

a) HSV encephalitis

b) Hepatic encephalopathy

c) Cerebral toxoplasmosis

d) Cruetzfeldt-Jakob disease

Sold by @itachi... purchased this from some... else, If yo... may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	a
5	d
6	b
7	a
8	a
9	d
10	c
11	c
12	d

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The clinical features are suggestive of neonatal meningitis and Escherichia coli is the most common cause of this condition.

The causes of neonatal meningitis in decreasing order of frequency are:

• Escherichia coli

• Group B Streptococci

• Listeria monocytogenes

Empirical treatment of neonatal meningitis is with ampicillin + 3rd generation cephalosporins.

Solution to Question 2:

The most common cause of meningitis in patients aged 2-50 years is Streptococcus pneumoniae.

The common causes of meningitis in this age group include:

• Streptococcus pneumoniae

• Neisseria meningitidis

• Group B streptococci

• Hemophilus influenzae

• Listeria monocytogenes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

The clinical features are suggestive of acute bacterial meningitis. This condition is most likely caused by Staphylococcus aureus in a patient with an Ommaya reservoir.

An Ommaya reservoir is an intraventricular catheter system used for the delivery of drugs into the cerebrospinal fluid. S. aureus and coagulase-negative staphylococci (CoNS) are important causes of meningitis that occurs following invasive neuro-surgical procedures, or as a complication of the use of Ommaya reservoirs. S aureus is a more common cause than CoNS.

Solution to Question 4:

The sign being elicited here is Kernig's sign and this along with the given clinical features are suggestive of acute bacterial meningitis. Empirical treatment with cefepime, vancomycin and ampicillin is recommended for patients on immunomodulatory drugs as they are at risk for Listeria infection.

Sold by @itachibot

Risk factors for Listeria meningitis include:

• Age>50

• Diabetes

• Chronic illness

• Malignancy

• Immunosuppressive therapy

Solution to Question 5:

The clinical features are suggestive of meningococcal meningitis and the drug of choice is a third-generation cephalosporin such as ceftriaxone. Chloramphenicol is used in patients with penicillin allergy.

It is caused by Neisseria meningitidis, a gram-negative, oxidase-positive, catalase-positive diplococci. Outbreaks usually occur in mass gatherings such as in army camps or hostel dorms.

The risk of infection is increased in patients with terminal complement deficiency, hypogammaglobulinemia and hyposplenism. This is because opsonization by complement is a major mechanism by which the bacteria is cleared from the blood.

Characteristic clinical findings include:

• Maculopapular or purpuric rash

• Shock due to capillary leak

• Bilateral adrenal hemorrhage (Waterhouse–Friderichsen syndrome)

Rifampicin, ciprofloxacin and cephalosporins can be used to treat carriers.

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Solution to Question 6:

Ceftriaxone or cefotaxime is the drug of choice for empirical treatment of meningococcal meningitis. However, Penicillin G remains the antibiotic of choice for susceptible strains.

Empirical antimicrobial therapy is initiated in patients with suspected bacterial meningitis before the results of CSF Gram stain and culture are known.

S. pneumoniae and N. meningitidis are the most common etiologic organisms of community-acquired bacterial meningitis. Due to the emergence of penicillin-resistant and cephalosporin-resistant S. pneumoniae, empirical therapy of community-acquired suspected bacterial meningitis should include a combination of:

• Dexamethasone

• Third- or fourth-generation cephalosporin (e.g., ceftriaxone, cefotaxime, or cefepime)

• Vancomycin

• Acyclovir - HSV encephalitis is the main differential

Sold by @itachibot

• Doxycycline - during tick season.

Ceftriaxone or cefotaxime provides better coverage for susceptible S. pneumoniae, H. influenzae and group B streptococci and good coverage for N. meningitidis. Cefepime is a broad-spectrum fourth-generation cephalosporin with activity similar to that of cefotaxime or ceftriaxone against S. pneumoniae and N. meningitidis with additional greater activity against Enterobacter species and Pseudomonas aeruginosa.

Solution to Question 7:

Ciprofloxacin is used for chemoprophylaxis of meningococcal meningitis for close contacts. Ceftriaxone and rifampicin are also used.

Antibiotic chemoprophylaxis among close contacts of a patient with invasive meningococcal disease is recommended to prevent secondary cases.

• Rifampin - was earlier used widely, but is not optimal due to the high risk of adverse events, emerging resistance, and the need for compliance for four doses.

• Ciprofloxacin - widely used as a single oral dose, but it cannot be given in pregnancy and those below 18 years of age.

• Ceftriaxone - can be given in pregnancy and all ages as a single IM or IV dose. However, reduced susceptibility of isolates to ceftriaxone has occasionally been reported.

Recommended chemoprophylaxis for high-risk close contacts for meningococcal meningitis:

Patient group	Drug	Age	Dose	Route and duration
Non-pregnant individuals	Rifampicin	<1 month	5 mg/kg	Oral every 12 hrs for 2 days
>1 month	10 mg/kg	Oral every 12 hrs for 2 days		
Ciprofloxacin	>18 years	500 mg	Oral single dose	
Children and pregnant women	Ceftriaxone	<15 years	125mg	IM single dose
> 15 years	250 mg	IM single dose		

Solution to Question 8:

The features and imaging findings are suggestive of tuberculous meningitis. CSF analysis would likely show lymphocytosis, low glucose and high protein.

If the CSF is allowed to stand, a 'spider's web clot' or 'cobweb' coagulam may form due to the high level of protein.

Sold by @itachibot
Bought this from someone else.
Have been scammed.

Sold by @itachibot

Solution to Question 9:

The given symptoms cluster of fever, vomiting, and neck stiffness is suggestive of meningitis. A CSF analysis showing increased opening pressure, lymphocytic pleocytosis, mildly increased protein, and normal glucose is suggestive of viral meningitis. Among the given options, Coxsackie (Option D) is the only viral cause of meningitis.

Other options:

Option A: Tuberculous meningitis shows lymphocytic pleocytosis with low glucose.

Option B: Bacterial meningitis shows neutrophilic pleocytosis with low glucose.

Option C: Fungal meningitis shows lymphocytic/neutrophilic pleocytosis with low glucose.

Solution to Question 10:

The clinical hallmark of HSV encephalitis has been the acute onset of fever and focal neurologic symptoms and signs, especially in the temporal lobe.

Investigation of choice includes HSV-DNA demonstration in CSF by PCR.

The image below is a coronal T2-weighted MR image showing a high signal intensity in the temporal lobes.



Solution to Question 11:

The clinical features are suggestive of herpes simplex encephalitis and the CSF analysis would likely show 100 RBC/microliter. HSV encephalitis is associated with hemorrhagic necrosis of inferomedial temporal and frontal lobe resulting in significant number of RBCs in the CSF.

Sold by @itachibot

It is caused by both HSV 1 and HSV 2. Patients present with fever, headache, seizures, personality, and behavioral changes.

CSF analysis shows lymphocytic pleocytosis, moderately elevated protein levels and normal glucose concentration. EEG shows periodic lateralized spike-wave discharges.

The drug of choice for the condition is acyclovir.

Solution to Question 12:

The clinical features and EEG point towards a diagnosis of Creutzfeldt-Jakob disease (CJD). The EEG shows periodic biphasic or triphasic sharp-wave complexes seen as spikes.

CJD is a spongiform encephalopathy caused by a prion. It presents as rapidly progressive dementia and is fatal within 1 year of onset of symptoms.

Patients present with myoclonus, akinetic mutism, cerebellar or visual disturbances and pyramidal/extrapyramidal dysfunction (hypokinesia). Insomnia, apathy and other behavioral abnormalities may be seen.

Four types of Creutzfeldt-Jakob disease have been described:

• Sporadic (sCJD): Accounts for 85-90% of cases

• MRI shows cortical ribbon sign

• Variant (vCJD): mad cow disease, Kuru

• MRI shows hockey stick sign

• Familial (fCJD): 10% of cases (these individuals carry a PRPc mutation)

• Iatrogenic

CSF analysis shows a positive assay for 14-3-3 protein. Management is mainly supportive.

Note: Hepatic encephalopathy will show a triphasic wave pattern (negative-positive-negative wave). They are also seen in Hashimoto encephalopathy and CO_2 narcosis.

EEG in Creutzfeldt-Jakob disease (CJD)



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Hypoproliferative Anemia

Question 1:

Which of the following is not a hypoproliferative anemia?

a) Sideroblastic anemia

b) Anemia of chronic disease

c) Hemolytic anemia

d) Megaloblastic anemia

Question 2:

A 25-year-old woman came to the OPD one year after the birth of her last child. She had been treated for iron deficiency anemia while pregnant. On examination, she is pale and investigations show a Hb of 5g/dl and a reticulocyte count of 9%. What is her corrected reticulocyte count? (Take normal Hb as 15g/dl)

a) 6

b) 4.5

c) 3

d) 1

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

What does the red cell distribution width deal with?

a) Hypochromia

b) Anisocytosis

c) Poikilocytosis

d) Anisochromia

Question 4:

Which of the following patients is unlikely to have the following finding on a peripheral smear study?

Sold by @itachibot



a) A 55-year-old alcoholic man with ascites and jaundice

b) A 19-year-old girl who underwent emergency splenectomy after a road accident

c) A 25-year-old man with hemoglobin C disease

d) A 70-year-old woman undergoing dialysis for the past 3 years

Sold by @itachi... If you purchased this from some... else, may have been scammed.

Question 5:

A first-year pathology resident prepares the following smear from a patient's blood sample. Which of the following is it suggestive of?



a) Thalassemia

Sold by @itachibot

b) Autoimmune hemolysis

c) Iron deficiency anaemia

d) Lead poisoning

Question 6:

The following finding is not typically seen in which of the following conditions?



a) Myelodysplasia

b) Myelofibrosis

c) Thalassemia

d) Sideroblastic anemia

Question 7:

The following blood picture is seen in patients with which of the following conditions?

Sold by @itachibot



a) 1, 2, 5

b) 2, 4, 5

c) 1, 3, 5

d) 1, 3, 4

Question 8:

A 30-year-old man with a BMI of 41 kg/m2 is being evaluated for bariatric surgery. Which of the following procedures does not reduce the surface area for iron absorption?

a) Duodenal switch

b) Sleeve gastrectomy

c) Roux-en-Y gastric bypass

d) Biliopancreatic diversion

Question 9:

You have prescribed oral iron supplements for an anemic young woman. Which of the following foods should she avoid taking along with these supplements?

a) Grapefruit juice

b) Fresh kiwi fruit

c) Broccoli

d) Black tea

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 10:

The levels of which of the following will be detected first in a patient with iron deficiency?

a) Marrow iron stores

b) Serum iron

c) Serum ferritin

d) Transferrin saturation

Question 11:

Which of the following is not seen in the stage of negative iron balance in iron-deficiency anemia?

a) Decrease in marrow iron stores

b) Decrease in TIBC

c) Normal RBC morphology

d) Normal serum iron

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 12:

Patients with which of the following do not need to be evaluated for iron deficiency?

a) Unusual craving to eat clay and paper

b) Insomnia due to a compulsive urge to keep moving the legs

c) Incidental finding of esophageal webs on upper GI endoscopy

d) Tingling and numbness of lower extremities

Question 13:

Which of the following PBF findings is least likely to be seen in patients with iron deficiency anemia?

Sold by @itachibot



a) 1

b) 2

c) 3

d) 4

Question 14:

A 28-year-old woman is brought to the ER after she faints at home. Her husband says that she is very body-conscious and does intermittent fasting. She has been on a vegan diet for the past 2 months. Lab values are given below. What is the most likely diagnosis?

a) Anemia of chronic disease

b) B12 deficiency anemia

c) Iron deficiency anemia

d) Folate deficiency anemia

Question 15:

A 62-year-old woman is found to have a Hb of 7.2 g/dl and high ferritin levels. She has been undergoing chemotherapy for breast cancer since last year. Which of the following is responsible for this presentation?

a) Ferritin

b) DMT1

c) Hemojuvelin

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Sold by @itachibot

d) Hepcidin

Question 16:

Normocytic normochromic anemia is seen in patients with which of the following?

a) Liver disease

b) Kidney disease

c) Lead toxicity

d) Thalassemia

Question 17:

A 38-year-old woman presented to the OPD with progressive tiredness over the last 2 months. She also reported having heavy menstrual bleeding during her previous 2 cycles. She is a known case of rheumatoid arthritis. Lab investigations are given below. What is the likely diagnosis?

a) Iron deficiency anemia

b) Anemia of chronic disease

c) Sideroblastic anemia

d) Anemia due to blood loss

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A 60-year-old man with stage 4 chronic kidney disease on hemodialysis is found to have a hemoglobin of 6.3 mg/dl. Which of the following will be additionally seen in this patient?

a) Increased serum ferritin

b) Increased TIBC

c) Increased serum iron

d) Decreased reticulocytes

Question 19:

What is the predominant inheritance pattern of sideroblastic anemia?

a) Autosomal dominant

Sold by @itachibot

b) Polygenic

c) X linked

d) Autosomal recessive

Question 20:

Which of the following would not be seen in patients with sideroblastic anemia?

a) Erythroid hyperplasia

b) Low serum iron

c) High transferrin saturation

d) Dimorphic blood picture

Question 21:

The bone marrow biopsy sample of a patient with chronic anemia is stained with Prussian blue, as shown below. The stained deposits are present in which of the following?



a) Lysosome

b) Ribosome

c) Mitochondria

d) Endoplasmic reticulum

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Sold by @itachibot

Question 22:

Which of the following drugs is known to cause sideroblastic anemia?

a) Cyclophosphamide

b) Chloramphenicol

c) Ethionamide

d) Rifampicin

If you bought this from anyone else, it is stolen.

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	d
5	b
6	d
7	c
8	b
9	d
10	c
11	b
12	d
13	d
14	c
15	d
16	b
17	b
18	d
19	c
20	b
21	c
22	b

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Hemolytic anemia is not a hypoproliferative anemia.

Anemia can be defined as hemoglobin of < 13 g/dl in men and < 12 g/dl in women. Based on the mechanism of reduction of hemoglobin, anemia can be classified as:

• Hypoproliferative anemias - due to impaired production of RBCs

• Hemolytic anemias and hemoglobinopathies - due to increased destruction of RBCs

• Anemia due to blood loss

Hypoproliferative anemias are characterized by an abnormally low reticulocyte response (reticulocyte index <2–2.5) to reduced RBC production. They include:

• Iron-deficiency anemia

• Anemia of chronic disease (inflammation, renal disease, or malignancies)

• Thalassemia

• Sideroblastic anemia

• Aplastic anemia

• Megaloblastic macrocytic anemias (folate and vitamin B12 deficiency)

• Non-megaloblastic macrocytic anemias (alcohol use)

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 2:

The corrected reticulocyte count (CRC) of this patient is 3.

CRC = Reticulocyte % x (Actual Hb / Normal Hb)

Plugging in the values given in the question,

CRC = 9% x 5/15 = 3%

Alternatively, CRC can be calculated using hematocrit (PCV) as follows:

CRC = Reticulocyte % x (Actual PCV / Normal PCV)

For these calculations, take normal Hb as 15g/dL and normal PCV as 45%.

The patient's hemoglobin level (5 g/dL) is significantly lower than the normal hemoglobin level (15 g/dL). To obtain an accurate assessment of reticulocyte production relative to the reduced number of circulating red blood cells, a correction factor has to be applied. CRC accounts for the anemia and provides a more precise estimate of the reticulocyte count.

When polychromatophilic macrocytes (prematurely released reticulocytes/shift cells) are observed in the peripheral blood smear, a second correction is applied to account for the longer life of reticulocytes to yield reticulocyte production index (RPI).

Sold by @itachibot

In the classification of anemia, the patient's reticulocyte count is compared to the expected reticulocyte response. Reticulocytes can be identified by staining their ribosomal RNA with a supravital dye, resulting in the formation of blue or black punctate spots. They can be quantified manually or by using fluorescent dyes. The presence of reticulocytes indicates active red cell production.

The premature release of reticulocytes is typically associated with increased EPO stimulation. In hemolytic anemia, red cell production can increase six to seven times. If the RPI < 2 in a patient with anemia, a defect in erythroid marrow proliferation or maturation has to be suspected.

Solution to Question 3:

Red cell distribution width (RDW) measures the range of red blood cell volumes and provides an assessment of their variability (anisocytosis).

The normal range of RDW is 11-14%. In the presence of uneven cell size (anisocytosis), the RDW increases to 15-18%.

RDW has clinical utility in two main scenarios:

• In cases of microcytic anemia (low MCV), it helps differentiate between iron deficiency and thalassemia. Thalassemia is characterized by uniformly small red blood cells with a normal RDW, while iron deficiency leads to greater size variability and an increased RDW.

• A high RDW can indicate dimorphic anemia caused by chronic atrophic gastritis. Vitamin B12 malabsorption results in macrocytic anemia and blood loss leads to iron deficiency (microcytic anemia).

Solution to Question 4:

The black arrows in the given image denote target cells or codocytes. They are RBCs with a dense center (hyperchromic bull's eye) surrounded by an area of pallor. They are not a feature of renal disease (Patient D).

Target cells are seen in the following conditions:

• Thalassemia

• Iron deficiency anemia

• Liver disease

• Post-splenectomy

• Hemoglobin C disease

• As artifacts in improperly prepared slides

Solution to Question 5:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

The small, dense RBCs with absent central pallor seen in the given peripheral smear are called spherocytes. They appear due to decreased membrane surface area and are seen in the following conditions:

- Autoimmune hemolytic anemia
- Hereditary spherocytosis
- Clostridial sepsis
- Snake bites

Solution to Question 6:

The given peripheral smear shows teardrop cells or dacrocytes. They are not seen in sideroblastic anemia.

Teardrop cells are seen in the following conditions:

- Myelofibrosis
- Myelodysplastic syndromes
- Hemolytic anemia
- Thalassemia
- Severe iron deficiency

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

The given image shows Howell-Jolly bodies. They are remnants of RBC nuclei that occur in conditions with either defective production (dysplasia) or destruction of RBCs. They are observed in the following conditions:

- Sickle cell disease (postsplenectomy)
- Hyposplenism
- Megaloblastic anemia
- Dyshematopoesis
- Hemolytic anemia

Thalassemia presents with microcytic, hypochromic RBCs, and target cells.

Lead poisoning presents with basophilic stippling or punctate basophilia on peripheral smears. These are RNA fragments seen as numerous basophilic granules dispersed in the cytoplasm of RBCs, as shown below.

Sold by @itachibot



Sold by @itachi
d this from some
else,
Keep scammed.

Solution to Question 8:

Iron absorption takes place in the duodenum. In sleeve gastrectomy, a portion of the stomach is resected but the duodenum remains intact. Hence, the surface area of iron absorption is not affected.



Dietary iron is efficiently absorbed in its heme form. Non-heme iron is less efficiently absorbed mainly because of interference by phosphates, tannins, oxalates, and other food constituents. Acidic pH enhances iron absorption by promoting the conversion of ferric to ferrous iron by the enzyme ferrireductase.

Sold by @itachibot



The following are malabsorptive procedures that reduce the duodenal surface area and hence, reduce iron absorption:

• Biliopancreatic diversion

• Roux-en-Y gastric bypass

• Duodenal switch

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

Tea contains tannins, which inhibit iron absorption. Hence, it should not be taken with iron supplements. Iron absorption is also reduced by:

• Calcium supplements and antacids

• Milk and dairy products

• Proton pump inhibitors

Grapefruit and kiwi are acid-containing citrus fruits that create an acidic pH and hence, increase iron absorption.

Broccoli contains ascorbic acid that also enhances iron absorption.

Solution to Question 10:

The earliest sign of iron deficiency is a decrease in serum ferritin levels. This is called the state of negative iron balance.

Iron deficiency anemia develops in 3 stages:

• Stage of negative iron balance - decrease in iron stores with normal serum iron and RBC morphology

Sold by @itachibot

• Stage of iron-deficient erythropoiesis - decrease in serum iron levels with normal RBC morphology

• Stage of iron deficiency anemia - fall in hemoglobin with microcytic hypochromic RBCs

Solution to Question 11:

TIBC (total iron-binding capacity) does not decrease during the stage of a negative iron balance of iron deficiency anemia.

The progression to iron deficiency develops in 3 stages:

• Stage of negative iron balance - Decrease in iron stores with normal serum iron and RBC morphology

• Stage of iron-deficient erythropoiesis - Decrease in serum iron levels with normal RBC morphology

• Stage of iron deficiency anemia - Microcytic hypochromic RBCs

	Negative iron balance	Iron-deficient erythropoiesis	Iron deficiency anemia
Marrow iron stores	Decreased	Decreased	Decreased
S. ferritin	Decreased	Decreased	Decreased
TIBC	Increased	Increased	Increased
S. Iron	Normal	Decreased	Decreased
Transferrin saturation	Normal	Decreased	Decreased
Red cell protoporphyrin	Normal	Increased	Increased
Red cell morphology	Normal	Normal	Microcytic hypochromic

Solution to Question 12:

Tingling and numbness of the lower extremities are suggestive of peripheral neuropathy. It is a feature of vitamin B12 deficiency, not iron deficiency.

Iron-deficiency anemia is associated with the following findings:

• Smooth tongue - due to loss of papillae

• Cheilosis - fissuring at corners of mouth

• Brittle nails

• Koilonychia - spooning of nails

• Pica - craving for non-edible foods like ice chips, hair, paper, clay

Sold by @itachibot

• Plummer–Vinson syndrome - dysphagia due to the formation of esophageal webs

• Restless legs syndrome - compulsive urge to keep moving the legs which results in insomnia

Solution to Question 13:

Image 4 shows dense RBCs with loss of central pallor or spherocytes. They are seen in autoimmune hemolytic anemia (AIHA) and hereditary spherocytosis, not iron deficiency anemia.



The peripheral blood smear in iron deficiency anemia shows:

• Microcytic hypochromic cells

• Target cells

• Pencil-shaped or cigar-shaped cells

• Poikilocytosis- variations in the shape of RBCs

• Anisocytosis- variations in the size of RBCs.

Solution to Question 14:

This clinical scenario is suggestive of iron deficiency anemia due to the following findings:

• Low MCV (N: 87-96 fL)

• Low MCH (N: 27-33 pg)

• Elevated RDW

• High TIBC

Option A: Anemia of chronic disease is associated with low TIBC.

Sold by @itachibot

Options B and D: Anemia due to B12 and folate deficiency presents with a high MCV (macrocytosis).

Solution to Question 15:

This clinical scenario is suggestive of anemia of chronic disease. The chief mediator of this condition is hepcidin.

Anemia of chronic disease is associated with chronic inflammatory states such as inflammatory bowel disease, rheumatoid arthritis, chronic infections, and malignancies.

Hepcidin is a positive acute-phase reactant (ARP) that increases in chronic inflammatory states. It binds with ferroportin which results in both reduced iron intake in the gut and reduced iron transfer from macrophages to erythroid progenitor cells in the bone marrow. Hence, hepcidin is a negative regulator of iron balance.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

Normocytic normochromic anemia is seen in chronic kidney disease (CKD).

It is also seen in the following conditions:

- RBC membrane disorders - hereditary spherocytosis, elliptocytosis
- RBC enzyme disorders - G6PD deficiency, Pyruvate kinase deficiency
- Parvovirus B19 infection
- Anemia of chronic disease - can also present as microcytic anemia

Solution to Question 17:

This clinical scenario is suggestive of anemia of chronic disease due to the following findings:

- Low MCV (N: 87-96 fl)
- Low TIBC (N: 300-360µg/dL)
- Elevated ferritin level (N: 50-200 µg/L)
- Normal transferrin saturation (N: 25-50%)

Solution to Question 18:

Anemia of CKD is a form of hypoproliferative anemia presenting with decreased reticulocytes.

It mainly occurs due to a failure of erythropoietin production by the diseased kidney and a reduction in the survival of RBCs. Also, patients with CKD maintained on chronic hemodialysis

Sold by @itachibot

may develop iron deficiency from blood loss through the dialysis procedure.

The following features are seen:

• Normocytic and normochromic RBCs

• Decreased reticulocytes

• Serum iron, transferrin saturation, TIBC are usually normal or low

• Decreased serum ferritin

The degree of anemia usually correlates with the severity of the renal failure.

Note: Anaemia in CKD is not the same as anaemia of chronic disease.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 19:

Inherited forms of sideroblastic anemia are usually X-linked. Rarely, autosomal recessive forms may also occur.

X-linked sideroblastic anemia results from the deficient activity of the erythroid form of ALA-synthase (ALA-synthase 2) and is associated with ineffective erythropoiesis

Solution to Question 20:

High serum iron levels are seen in patients with sideroblastic anemia.

Sideroblastic anemia presents with the following features:

• Serum iron level - high/normal

• Transferrin saturation - high

• Serum ferritin levels - high

• TIBC - low/normal

• Peripheral smear - dimorphic population (both microcytes and macrocytes).

• Bone marrow study - marked erythroid hyperplasia.

The iron profile of sideroblastic anemia is opposite to that of iron deficiency anemia.

Solution to Question 21:

The given image shows ringed sideroblasts, which are erythroid precursors in the marrow. In these cells, iron deposits in the mitochondria form a ring around the nucleus and are stained by Prussian blue.

The mature red blood cell counterparts of ringed sideroblasts in the circulation are called siderocytes. These cells have iron-containing basophilic granules called Pappenheimer bodies, as

Sold by @itachibot

shown below.



Ringed sideroblasts and Pappenheimer bodies are pathognomonic of sideroblastic anemia.

Sold by @ita... If you purchased this from so... else, you may have been scammed.

Solution to Question 22:

Chloramphenicol is known to cause sideroblastic anemia.

The agents known to cause sideroblastic anemia are:

- Isoniazid
- Chloramphenicol
- Linezolid
- Excessive alcohol

Sold by @itachibot

Macrocytic Anemia

Question 1:

The peripheral blood smear of a patient is given below. All of the following can cause this except:



a) Vitamin B12 deficiency

b) Chronic alcoholism

c) Folate deficiency

d) Orotic aciduria

Question 2:

Patients with which of the following conditions are unlikely to develop cobalamin deficiency?

a) Tropical sprue

b) Crohn's disease

c) Diphyllobothrium latum infection

d) Ancylostoma duodenale infection

Question 3:

Sold by @itachibot

Which of the following would not be seen in a patient with cyanocobalamin deficiency?

a) Glossitis

b) Conjugated hyperbilirubinemia

c) Skin hyperpigmentation

d) Mild fever

Question 4:

A 56-year-old male patient is being evaluated for peripheral neuropathy and excessive tiredness. He has uncontrolled diabetes and consumes alcohol nearly every day. Lab investigations reveal the following findings. What is the likely diagnosis?

a) Folate deficiency

b) Alcohol-induced macrocytosis

c) Vitamin B12 deficiency

d) Thiamine deficiency

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

Which of the following drugs is not associated with megaloblastic anemia?

a) Nitrofurantoin

b) Phenytoin

c) Tetracycline

d) Temafloxacin

Question 6:

A 45-year-old woman is rushed to the ER following a seizure. She is a known case of epilepsy but is non-compliant with her medications. Her lab values are given below. All of the following drugs can cause this presentation except:

a) Lamotrigine

b) Phenytoin

c) Primidone

d) Amobarbital

Sold by @itachibot

Question 7:

A dose of oral 58-Co-labeled cobalamin was administered to a 65-year-old woman. The 24-hour urinary excretion of radiolabeled cobalamin was found to be < 10%. The test was repeated after a 5-day course of antibiotics and showed urinary excretion > 10%. What is the likely diagnosis?

a) Pernicious anemia

b) Lack of pancreatic enzymes

c) Bacterial overgrowth

d) Malnutrition

Question 8:

A child diagnosed with megaloblastic anemia failed to respond to treatment with vitamin C, vitamin B12, and folic acid. He was started on uridine monophosphate which improved his symptoms. What is the likely diagnosis?

a) Thiamine deficiency

b) Liver disease

c) Pernicious anemia

d) Orotic aciduria

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A 28-year-old woman presents with fatigue and a red, painful tongue. On examination, there is reduced vibration sense and proprioception in the feet. She is under treatment for Hashimoto's thyroiditis. Peripheral smear reveals macrocytes and hypersegmented neutrophils. Which of the following is unlikely to be seen in this patient?

a) Intrinsic factor blocking antibodies

b) Antibodies against gastric parietal cells

c) Intrinsic factor binding antibodies

d) Antibodies against gastrin

Question 10:

Sold by @itachibot

A pathologist receives a gastric biopsy sample of a patient with pernicious anemia. Which of the following findings is he unlikely to find?

a) Intestinal metaplasia

b) Mixed inflammatory cell infiltrate

c) Hypertrophy of glandular elements

d) Increased mucus cells

Question 11:

Patients with pernicious anemia are at a higher risk of which of the following?

a) Diffuse gastric adenocarcinoma

b) Gastric carcinoid tumors

c) Primary gastric lymphoma

d) Gastrointestinal stromal tumor

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Which of the following is the most common cause of folate deficiency?

a) Malabsorption

b) Nutritional

c) Use of anti-folate drugs

d) Deficiency of enzyme DHF reductase

Question 13:

Dietary folates are absorbed from which of the following sites?

a) Duodenum

b) Proximal ileum

c) Terminal ileum

d) Ascending colon

Question 14:

Sold by @itachibot

A 49-year-old woman is brought with increased forgetfulness and clumsiness for the past 1 month. Since the death of her husband 3 months ago, she has been tired and eating poorly. She began taking over-the-counter iron and folic acid tablets 2 months ago. Examination reveals positive Romberg's sign with eyes closed and loss of vibration in lower extremities. What is the likely diagnosis?

a) B12 deficiency

b) Severe folate deficiency

c) Thiamine deficiency

d) Pyridoxine deficiency

Question 15:

A 19-year-old girl with a seizure disorder is started on phenytoin therapy. Which of the following must be supplemented in this patient?

a) Pyridoxine

b) Thiamine

c) Cobalamin

d) Folic acid

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 16:

A 23-year-old woman comes to you for pre-conceptional counseling. Her previous pregnancy resulted in fetal death due to a neural tube defect. What dose of folic acid would you recommend?

a) 400 µg pre-conceptionally and in first trimester

b) 400 µg pre-conceptionally and throughout pregnancy

c) 4-5 mg pre-conceptionally and in first trimester

d) 4-5 mg pre-conceptionally and throughout pregnancy

Question 17:

Which of the following would not be seen in a patient with Imerslund-Grasbeck syndrome?

a) Congenital cobalamin malabsorption

b) AMN gene mutation

Sold by @itachibot

c) Autosomal dominant inheritance

d) Non-specific proteinuria

Answer Key

Question No.	Correct Option
1	b
2	d
3	b
4	a
5	d
6	a
7	c
8	d
9	d
10	c
11	b
12	b
13	a
14	a
15	d
16	c
17	c

Detailed Explanations

Solution to Question 1:

The given image shows hypersegmented neutrophils (> 5 nuclear lobes), which are seen in megaloblastic anemia. Chronic alcoholism causes non-megaloblastic macrocytic anemia.

Megaloblastic anemia is a form of macrocytic anemia with impaired nucleic acid metabolism and defective erythropoiesis. The causes include vitamin B12 deficiency, folate deficiency, and orotic aciduria. Nuclear maturation lags behind cytoplasmic maturation giving the appearance of hypersegmented neutrophils. Peripheral smear findings include:

• Oval macrocytes

Sold by @itachibot

• Anisocytosis

• Hypersegmented neutrophils

Other features of ineffective erythropoiesis seen in megaloblastic anemias include:

• Howell-Jolly bodies are nuclear remnants in red cells. They are usually single, round, dark purple-red in color and peripheral in location, as shown below.



• Basophilic stippling refers to blue granules of various sizes dispersed throughout the cytoplasm of the red cell. They represent ribosomal precipitates.

• Cabot rings are red cell inclusions appearing as fine, purple filamentous loops or figure-of-eight forms, as shown below. They are presumed to be remnants of mitotic spindles.



Solution to Question 2:

Sold by @itachibot

Ancylostoma duodenale (hookworm) infection does not affect the absorption of cobalamin or vitamin B12. Small amounts of blood loss during the attachment of hookworms to the intestinal mucosa can cause iron-deficiency anemia.

Vitamin B12-intrinsic factor (IF) complex is absorbed in the ileum. Tropical sprue, Diphyllobothrium latum (fish tapeworm) infection and Crohn's disease affect the ileum leading to defective absorption of vitamin B12.

Solution to Question 3:

Unconjugated hyperbilirubinemia is seen in cyanocobalamin or vitamin B12 deficiency, not conjugated hyperbilirubinemia.

Vitamin B12 deficiency causes megaloblastic anemia due to ineffective erythropoiesis. This leads to the formation of nucleated RBCs in the marrow, which undergo early lysis. This leads to increased unconjugated bilirubin in the plasma.

The clinical features of cyanocobalamin and folate deficiency include:

System	Clinical features
Oral	GlossitisAngular cheilosis
Gastrointestinal	AnorexiaWeight lossDiarrhea Constipation
Skin	Hyperpigmentation(reversible)
Hematological	ThrombocytopeniaLeucopeniaIneffective erythropoiesis and unconjugatedjaundice
Cardiovascular	Atherosclerosis leading to PVD, CAD (due to elevated homocysteine)
Neurological (only B12)	Peripheral neuropathySubacute combined degeneration (degeneration of the posterior and pyramidal tracts of the spinal cord)
Pregnancy (only folate)	Recurrent fetal lossNeural tube defects
Others	Mildfeverin severe casesPsychiatric disturbances

Solution to Question 4:

This clinical scenario is suggestive of folate deficiency.

Sold by @itachibot

It is associated with the following clinical and laboratory findings:

• Raised MCV (normal: 80-100 fL) - macrocytosis

• Normal MCHC (normal: 33-36 g/dl)

• Increased MCH (mean corpuscular hemoglobin)

• Elevated RDW (red cell distribution width)

• Decreased reticulocyte count - due to destruction of abnormal erythroid precursors

• Elevated homocysteine

• Normal MMA (methyl malonic acid)

In vitamin B12 deficiency, elevated homocysteine and elevated MMA levels are seen.

Peripheral neuropathy is a feature of B12 deficiency, not folate deficiency. In this patient, it may be due to diabetic neuropathy.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

Temafloxacin is not associated with megaloblastic anemia.

Megaloblastic anemia is associated with the following drugs:

• Anticonvulsant drugs - phenytoin, primidone, barbiturates

• Sulfasalazine

• Nitrofurantoin

• Tetracycline

• Folate antagonists - trimethoprim, methotrexate, pyrimethamine, pentamidine, cotrimoxazole

Temafloxacin is a fluoroquinolone with severe adverse effects like hemolytic anemia and allergic reactions. It was withdrawn from the market and is no longer used.

Solution to Question 6:

This patient likely has folate deficiency secondary to the use of an anti-folate anti-convulsant drug. Lamotrigine has no effect on folate metabolism and does not cause its deficiency.

The following anticonvulsants have anti-folate properties:

• Phenytoin

• Barbiturates (amobarbital)

• Primidone

Folate deficiency causes megaloblastic macrocytic anemia which results in the following findings in RBCs:

• Anemia (low Hb = 10g/dl)

Sold by @itachibot

• Macrocytic (high MCV = 110 fl)

• Hyperchromic (high MCH = 40 pg)

Solution to Question 7:

This clinical scenario refers to Schilling's test done for the determination of the cause of B12 deficiency. Improvement of urinary B12 excretion > 10% after antibiotic administration is suggestive of small intestinal bacterial overgrowth (SIBO).

Solution to Question 8:

Megaloblastic anemia that is responsive to treatment with uridine is suggestive of orotic aciduria.

Orotic aciduria is a rare inborn error of metabolism in children. It occurs due to a defect in the pyrimidine synthesis pathway causing a failure of conversion of orotic acid to uridine. It manifests in early childhood with anemia and failure to thrive.

Sold by @itachibot
if you purchased this from someone else,
you may have been scammed

Solution to Question 9:

This clinical scenario in a patient with a history of autoimmune thyroiditis is suspicious for pernicious anemia. It is an autoimmune cause of B12 deficiency. Antibodies against gastrin are not seen in these patients. The following autoantibodies are seen:

• Type I or blocking antibodies - prevent the combination of IF and cobalamin

• Type II or binding antibodies - prevent attachment of IF to the ileal mucosa

• Anti-parietal cell antibodies - target the gastric proton pump located in these cells

This leads to the deficiency of the intrinsic factor of Castle (IF) and hydrochloric acid produced by the parietal cells. The resulting atrophic gastritis causes high gastrin and low pepsinogen levels. Lack of IF further causes vitamin B12 deficiency and megaloblastic anemia.

Solution to Question 10:

Gastric biopsy in pernicious anemia shows loss of glandular elements, not hypertrophy.

This is an autoimmune condition that occurs in genetically susceptible individuals. One of the common triggers of the disease process is Helicobacter pylori infection. Gastric biopsy findings in pernicious anemia include the following:

• Atrophy of all layers of the body and fundus

• Loss of glandular elements

• Absence of parietal and chief cells and replacement by mucus cells

Sold by @itachibot

• Mixed inflammatory cell infiltrate (excess of CD4 cells against gastric H+/K+-ATPase)

• Intestinal metaplasia

• Antral mucosa is usually well preserved.

Solution to Question 11:

Patients with pernicious anemia have an increased risk of intestinal-type of gastric adenocarcinoma and gastric carcinoid tumors. These patients must undergo regular endoscopic screening for these conditions.

The presumed mechanism is prolonged achlorhydria due to parietal cell loss. This leads to hypergastrinemia and cellular hyperplasia.

Solution to Question 12:

Reduced nutritional intake is the most common cause of folate deficiency.

Solution to Question 13:

Dietary folates are absorbed from the duodenum and jejunum in the form of monoglutamates.

Vitamin B12 absorption takes place in the terminal ileum.

Solution to Question 14:

This clinical scenario is suggestive of dorsal column involvement of the spinal cord due to underlying vitamin B12 deficiency.

Vitamin B12 deficiency causes subacute combined degeneration of the spinal cord (SACD). It affects the following structures:

• Dorsal column - sensory loss, paresthesias, incontinence, sensory ataxia (positive Romberg sign with eyes closed)

• Lateral column - spasticity and paraplegias in later stages

Folate deficiency is typically not associated with neurological manifestations. In patients with combined deficiency of B12 and folate, B12 deficiency must always be corrected first. This is because the administration of folate supplements may improve hematologic abnormalities but worsens neurological manifestations.

Treatment of cobalamin deficiency is as follows:

• Replenishment of body stores: Six 1000-µg IM injections of hydroxocobalamin given at 3- to 7-day intervals.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Maintenance therapy: 1000 µg hydroxocobalamin IM once every 3 months

Treatment of folate deficiency is as follows:

• 5–15 mg oral folic acid daily for a minimum of 4 months.

Option C: Thiamine deficiency causes Wernicke's syndrome with ophthalmoplegia, ataxia, and nystagmus. As this is due to cerebellar damage, Romberg's sign would be positive even with eyes open.

Option D: Pyridoxine deficiency causes pellagra like syndrome with features of dermatitis, diarrhea, dementia.

Solution to Question 15:

Usage of phenytoin leads to folate deficiency and megaloblastic anemia. Hence, folic acid must be supplemented.

Solution to Question 16:

The recommended prophylactic dose of folic acid in women who have had a previous fetus with a neural tube defect is 4-5 mg daily, pre-conceptionally, and during the first trimester of pregnancy. Following the first trimester, these patients are then switched to 400 µg of folic acid for the remainder of the pregnancy.

This high dose of folic acid (4-5 mg) should be initiated at least one to three months before pregnancy and should be continued for 12 weeks after pregnancy. The patient can be prescribed 1 mg tablet of folic acid and advised to take them 4- 5 times a day or they can be recommended 5mg tablets/capsules once daily.

Prophylactic folic acid with 400 µg daily, pre-conceptionally, and throughout pregnancy is indicated in all pregnant women to prevent megaloblastic anemia and reduce the incidence of neural tube defects.

Solution to Question 17:

Imerslund-Grasbeck syndrome (IGS) is an autosomal recessive disorder.

IGS occurs due to mutations in either the cubulin (CUBN) or amnionless (AMN) genes. They form the cubam receptor responsible for the absorption of the Cbl-IF complex. Hence, this condition leads to congenital cobalamin malabsorption. It is the most common cause of megaloblastic anemia due to B12 deficiency in infancy in Western countries.

Affected children present with lethargy, failure to thrive, and neurological symptoms like hypotonia, developmental delay, and movement disorders. Non-specific proteinuria with normal renal function is seen in 90% of patients.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Hemolytic Anemia

Question 1:

Which of the following will be seen in a patient with intravascular hemolytic anemia?

a) 1

b) 2

c) 3

d) 4

Question 2:

Which of the following is not due to an extracorpuscular defect in RBCs?

a) Autoimmune hemolytic anemia

b) Paroxysmal nocturnal hemoglobinuria

c) Familial hemolytic uremic syndrome

d) Microangiopathic hemolytic anemia

Question 3:

Which of the following is true about paroxysmal nocturnal hemoglobinuria?

a) Inherited defect in PIG –A

b) Extravascular hemolysis

c) Deficiency of CD 55 and CD 59

d) Microcytic anaemia

Question 4:

A patient with hemolytic anemia develops features of pancytopenia and venous thrombosis. What is the likely diagnosis?

a) Shwachman-Diamond syndrome

b) Pyruvate kinase deficiency

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) G6PD anemia

d) Paroxysmal nocturnal hemoglobinuria

Question 5:

A 31-year-old woman presents with a fever and cough. She also reports excessive tiredness and occasional dark-colored urine for the past 6 months. Chest X-ray reveals lower lobe pneumonia. Labs reveal the following. What is the investigation of choice for this condition?

a) Acidified serum / HAM test

b) Eosin-5′-maleimide-binding test

c) Flow cytometry

d) Acidified glycerol lysis test

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A 46-year-old woman presents with abdominal pain and yellowish discoloration of her skin. She gives a history of dark-colored urine which is more pronounced in the early mornings. Imaging reveals hepatic venous thrombosis. Flow cytometry reveals the absence of GPI-anchored proteins. What is the drug of choice?

a) Rituximab

b) Certolizumab

c) Eculizumab

d) Alemtezumab

Question 7:

A 10-year-old boy is brought for evaluation of fatigue. On examination, he is jaundiced and mild splenomegaly is noted. He was adopted and his family history is unknown. Peripheral smear findings are shown below. What is the most specific investigation for this condition?

Sold by @itachibot



a) Eosin-5-maleimide test

b) Acidified glycerol lysis test

c) Osmotic fragility test

d) NESTROFT test

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 8:

Which of the following is true regarding the treatment of hereditary spherocytosis?

a) Cholecystectomy is only indicated for patients with clinical cholelithiasis at the time of splenectomy

b) Cholecystectomy cannot be performed via laparoscopic route for patients with HS and pigment gall stones

c) Splenectomy is indicated in patients with mild and asymptomatic disease only after puberty

d) Lifelong penicillin prophylaxis after splenectomy is mandatory for all adult patients

Question 9:

The peripheral smear of a patient is given below. What is the most common mutation associated with the hereditary form of this condition?

Sold by @itachibot



a) Spectrin

b) Band 4.2

c) Band 4.1

d) Ankyrin

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 10:

A young couple comes to you for genetic counseling. The husband is a known case of G6PD deficiency and is concerned about passing it on to his future children. What is the chance that his children will inherit the disease?

a) Only male children will be affected

b) 1 in 2 chance

c) Females can inherit the mutation

d) 1 in 4 chance

Question 11:

A 35-year-old male with chronic bacterial prostatitis was treated with cotrimoxazole. Within a few days, he developed malaise, jaundice, abdominal pain, and dark urine. The blood picture is given below. What is the likely diagnosis?

Sold by @itachibot



a) Hemolytic uremic syndrome

b) Autoimmune hemolytic anemia

c) Paroxysmal nocturnal hemoglobinuria

d) Glucose 6 phosphate dehydrogenase deficiency

Sold by @itachi... If you purchased this from some... else, may have been scammed.

Question 12:

The peripheral smear of a patient is given below. It is characteristic of which of the following conditions?



a) Glucose 6 Phosphate dehydrogenase deficiency

Sold by @itachibot

b) Pyruvate kinase deficiency

c) 5' nucleotidase deficiency

d) Cytochrome b5 reductase deficiency

Question 13:

A patient with SLE is found to have warm autoimmune hemolytic anemia. Name the antibody and the site of hemolysis in this condition?

a) IgM and Extavascular hemolysis

b) IgG and extravascular hemolysis

c) IgM and intravascular hemolysis

d) IgG and intravascular hemolysis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

Patients with which of the following are least likely to develop autoimmune hemolytic anemia?

a) Infectious mononucleosis

b) Non-Hodgkin's lymphoma

c) Ulcerative colitis

d) Chronic myeloid leukemia

Question 15:

Which of the following are associated with a positive Coombs test?

a) 1, 2, 4

b) 2, 3, 5

c) 2, 4, 5

d) 1, 3, 4

Question 16:

A 20-year-old woman presented with fatigue and intermittent dark-colored urine. Further evaluation revealed positive Donath Landsteiner antibodies. What is the likely diagnosis?

Sold by @itachibot

a) Warm autoimmune hemolytic anemia

b) Cold autoimmune hemolytic anemia

c) Paroxysmal cold hemoglobinuria

d) Paroxysmal nocturnal hemoglobinuria

Question 17:

Microangiopathic hemolytic anemia is seen in all of the following except:

a) Antiphospholipid antibody syndrome

b) Vasculitis

c) Prosthetic heart valves

d) Eclampsia

Question 18:

In patients with atypical hemolytic uremic syndrome, which of the following mutations is associated with the best prognosis?

a) Complement factor H

b) Complement factor I

c) Complement factor B

d) Membrane cofactor protein

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

You have decided to prescribe an oral iron chelator for a 10-year-old girl with thalassemia. Which of the following will you not prefer?

a) Deferiprone

b) Deferasirox

c) Deferoxamine

d) Both A and B

Question 20:

Which of the following is the best test to screen patients for thalassemia?

Sold by @itachibot

a) Cellulose acetate hemoglobin electrophoresis

b) Naked eye single tube red-cell osmotic fragility test

c) Eosin 5' maleimide flow cytometry

d) Automated high performance liquid chromatography

Question 21:

During a master health check-up, a 25-year-old woman was found to have the following lab values. Which of the following will help in establishing a diagnosis in this patient?

a) Free erythorcyte protoporphyrin levels

b) Hemoglobin electrophoresis

c) Schilling test

d) Bone marrow biopsy

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 22:

Which of the following clinical manifestations is not associated with sickle cell trait?

a) Hyposthenuria

b) Painless hematuria

c) Papillary necrosis

d) Acute pain crisis

Question 23:

Patients with sickle cell anemia most commonly present with which of the following?

a) Acute pain crisis

b) Splenic sequestration crisis

c) Acute chest syndrome

d) Isosthenuria

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	b
2	b
3	c
4	d
5	c
6	c
7	a
8	a
9	a
10	c
11	d
12	c
13	b
14	d
15	d
16	c
17	c
18	d
19	c
20	b
21	b
22	d
23	a

Detailed Explanations

Solution to Question 1:

In intravascular hemolysis, there is elevated MCV, LDH, and urinary urobilinogen, with low haptoglobin.

Laboratory abnormalities seen in intravascular hemolysis:

Additionally, hemoglobinuria and hemosiderinemia may be seen on urine examination.

Lab parameters	Derangement	Mechanism
MCV	High	Compensatory reticulocytosis

Sold by @itachibot

Lab parameters	Derangement	Mechanism
S. LDH	High	Released from RBCs
S. Unconjugated bilirubin	High	Hemoglobin from RBCs is converted to bilirubin
Urinary urobilinogen	High	Unconjugated bilirubin is excreted as urobilinogen in urine
S. Haptoglobin	Low	Free hemoglobin in circulation binds to haptoglobin causing reduced haptoglobin levels

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

Paroxysmal nocturnal hemoglobinuria (PNH) is a hemolytic anemia due to an intracorpuscular defect.

It is caused due to an acquired somatic mutation leading to deficiency of membrane protective factors CD59 and CD55. This makes the RBC extremely sensitive to complement activation and causes intravascular hemolysis.

The other conditions occur due to extracorpuscular defects.

Option A: In autoimmune hemolytic anemia, RBCs are coated with autoantibodies. This triggers phagocytosis in the reticuloendothelial system leading to extravascular hemolysis.

Option C: Familial hemolytic uremic syndrome (atypical HUS) is an inherited abnormality with excessive complement activation leading to episodic and excessive production of membrane attack complex destroying normal RBCs.

Option D: Microangiopathic hemolytic anemia is non-immune hemolysis resulting from intravascular red blood cell fragmentation due to abnormalities in the microvasculature.

Solution to Question 3:

Deficiency of CD55 and CD59 is seen in paroxysmal nocturnal hemoglobinuria (PNH).

It is due to an acquired mutation in the X-linked phosphatidylinositol class A PIG-A gene (Option A) causing a deficiency of GPI anchor proteins. GPI is a glycolipid that anchors membrane protective factors such as CD59 and CD55 to the membrane. The deficiency of CD55 and CD59 makes the RBC sensitive to complement-mediated intravascular hemolysis(Option B). The anemia is usually normo-macrocytic (Option D).

Hemolysis tends to occur at night due to a slight decrease in blood pH during sleep, which triggers complement activity (nocturnal hemoglobinuria). PNH is characterized by hemolytic anemia, pancytopenia, and venous thrombosis.

Sold by @itachibot

PNH is diagnosed using flow cytometry. The drugs of choice for treatment are Eculizumab or Ravulizumab which work against complement C5. Supportive management of PNH includes folic acid and iron supplementation. Bone marrow transplantation is another option for the treatment of this condition.

Solution to Question 4:

Paroxysmal nocturnal hemoglobinuria (PNH) is a type of hemolytic anemia associated with pancytopenia and venous thrombosis.

In PNH, there is complement-mediated intravascular hemolysis. Bone marrow dysfunction also occurs, resulting in pancytopenia. Transformation to aplastic anemia or acute myeloid leukemia (AML) may rarely occur.

PNH is also associated with venous thrombosis (cerebral, renal, hepatic). This is the most common cause of death in PNH.

Option A: Shwachman-Diamond syndrome is inherited aplastic anemia with neutropenia with pancreatic insufficiency and malabsorption.

Options B and C: Pyruvate kinase and G6PD deficiency are inherited RBC enzyme disorders causing extravascular hemolytic anemias. They do not affect other cell lines and are not associated with venous thrombosis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 5:

This clinical scenario is suggestive of intravascular hemolysis due to paroxysmal nocturnal hemoglobinuria (PNH). The gold standard for diagnosis is flow cytometry.

Flow cytometry is performed with fluorescently labeled monoclonal antibodies that bind to GPI-anchored proteins (CD59, CD55). Most tests also use FLuorescent AERolysin (FLAER), a reagent that directly binds the GPI anchor. Absence of CD59 and CD55 are diagnostic of PNH.

Other investigations in PNH include:

• Ham test (acidified serum lysis) - lysis of RBCs in a mildly acidic pH due to complement fixation.

• Bone marrow biopsy:

• Hypercellularity with erythroid hyperplasia is seen in classical presentations.

• Patients who develop pancytopenia will have hypocellular bone marrow.

• Imaging - to evaluate venous thrombosis.

Option B and D: Eosin-5′-maleimide-binding and acidified glycerol lysis tests are done in hereditary spherocytosis.

Sold by @itachibot

Solution to Question 6:

This clinical scenario is suggestive of paroxysmal nocturnal hemoglobinuria (PNH). C5 complement inhibitors like eculizumab and ravalizumab are the drugs of choice in PNH.

Management of PNH involves the following:

• C5 complement inhibitors prevent the formation of membrane attack complex and complement-mediated lysis.

• Bone marrow transplant is useful for patients with severe pancytopenia

• Supportive measures:

• Transfusion of filtered RBCs

• Folic acid, and iron supplementation

• Anticoagulants in case of venous thrombosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 7:

The peripheral smear shows hyperchromic RBCs with loss of central pallor or spherocytes. These findings are suggestive of hereditary spherocytosis. The eosin-5-maleimide test is the most specific investigation for this condition.

It is a flow cytometric test based on the interaction between the dye eosin-5-maleimide and band 3 protein.

Other investigations that can be performed in this condition are:

Acidified glycerol lysis test: Spherocytes from patients with hereditary spherocytosis lyse more quickly than normal red blood cells when incubated in a phosphate buffered hypotonic solution with added glycerol. It has greater sensitivity and specificity than the osmotic fragility test.

Osmotic fragility test: Detects hemolysis by measuring the fraction of total hemoglobin released from red cells when suspended in progressively more dilute salt concentrations.

Note: NESTROFT (Naked eye single tube red-cell osmotic fragility test) is a highly sensitive test used as a screening test for thalassemia.

Solution to Question 8:

Cholecystectomy at the time of splenectomy is indicated in patients with hereditary spherocytosis (HS) only if they have symptomatic cholelithiasis at the time of surgery. It is usually carried out via a laparoscopic approach.

Penicillin prophylaxis after splenectomy is controversial. Routine penicillin prophylaxis is not mandatory for adults. It is recommended up to age of five and for at least one year following splenectomy.

Guidelines for the timing of splenectomy in HS:

Sold by @itachibot

- Mild cases - avoid splenectomy.
- Moderate cases - delay until puberty
- Severe cases - at 4–6 years of age

Solution to Question 9:

The given peripheral smear shows oval, elongated, rod, or cigar-shaped RBCs suggestive of elliptocytosis. The hereditary form is most commonly associated with the spectrin genes.

The following proteins are implicated in hereditary elliptocytosis (HE):

- α-Spectrin (SPTA1 gene) - 65%
- β-Spectrin (SPTB gene) - 30%
- Band 4.1

The treatment of choice is splenectomy, preferably after 5 years of age.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 10:

G6PD deficiency is an X-linked recessive disorder. Male children of an affected man are not affected as they receive only the paternal Y chromosome. However, the mutation can be passed on to female children, who are typically carriers of the disease.

G6PD deficiency is the most common enzymatic disorder of RBCs. G6PD enzyme is involved in the generation of NADPH and protects RBCs from oxidative injury. Hence, patients with G6PD deficiency have episodes of acute hemolysis in the setting of oxidant injury from medications, acute infections, and certain foods (fava beans/broad beans).

Solution to Question 11:

The given clinical features are suggestive of an acute hemolytic reaction following the administration of cotrimoxazole. This is suggestive of glucose-6-phosphate dehydrogenase (G6PD) deficiency.

In G6PD deficiency uncontrolled oxidative stress causes hemoglobin to denature and form Heinz bodies. Removal of Heinz bodies by macrophages in the spleen results in the formation of characteristic bite cells or degmacytes or blister cells (as shown in the image).

A hemolytic attack starts with abrupt onset of malaise, weakness, and abdominal or lumbar pain. After an interval of several hours to 2–3 days, the patient develops jaundice and often dark urine, due to hemoglobinuria.

Drugs that carry the risk of clinical hemolysis in persons with G6PD deficiency:

Note: Cotrimoxazole contains sulfamethoxazole.

Sold by @itachibot

Drug groups	Examples
Antimalarials	PrimaquineChloroquineDapsone/chlorproguanil
Sulphonamides/sulphones	SulfamethoxazoleSulfasalazineSulfisoxazole
Antibacterial/antibiotics	CotrimoxazoleCiprofloxacinChloramphenicolNalidixic acid NorfloxacinNitrofurantoin
Others	NaphthaleneMethylene blue Rasburicase

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 12:

The above peripheral smear shows basophilic stippling. It is characteristic of 5' nucleotidase deficiency.

Pyrimidine 5'-nucleotidase (P5'N-1) deficiency is the third most common enzyme abnormality causing hereditary hemolytic anemia. It is an autosomal recessive trait.

The hallmark of this enzyme deficiency is the presence of pronounced basophilic stippling and accumulation of pyrimidine nucleotides (RNA remnants) within erythrocytes.

Cabot rings may also be seen. They are thin, threadlike, red to violet rings or figure-of-8 shaped inclusions in red blood cells. They are remnants of the mitotic spindle and are seen in conditions causing dyserythropoiesis.

Basophilic stippling is also seen in the following conditions:

- Lead toxicity
- Sideroblastic anemia
- Severe thalassemia
- Conditions with impaired cell division

Solution to Question 13:

The antibody involved in warm autoimmune hemolytic anemia (AIHA) is IgG, which causes extravascular hemolysis.

IgG is called a warm antibody, as it binds to target proteins at 37 oC. On the other hand, cold antibodies or IgM are reactive at 4 to 30oC.

In AIHA, autoantibodies against RBC antigens are produced which coat the RBC membrane. The Fc portion of the antibody is recognized by the Fc receptor on macrophages causing phagocytosis. RBC destruction takes place in extravascular sites where macrophages are abundant.

Sold by @itachibot

Solution to Question 14:

Autoimmune hemolytic anemia (AIHA) is commonly associated with CLL (chronic lymphocytic leukemia). It is not a typical feature of CML (chronic myeloid leukemia).

AIHA can develop secondary to several autoimmune and infectious causes, as well as drugs. Drug-induced hemolytic anemia can occur by the following mechanisms:

Mechanism	Drugs
Adsorption:Drugs coat the RBC membrane. Antibodies are directed against the membrane–drug complex	CefotetanCeftriaxonePiperacillin
Innocent bystander:Drugs bind to antibodies in the plasma. This immune complex activates the complement system and destroys nearby RBCs (innocent bystanders)	SulfonamidesQuinineQuinidinePhenothiazines
Some drugs trigger the production ofautoantibodies	Methyldopa

Solution to Question 15:

Paroxysmal nocturnal hemoglobinuria (PNH) and hereditary spherocytosis are not Coomb's positive conditions. The following are Coombs-positive hemolytic anemias:

- Autoimmune hemolytic anemia (AIHA)
- Warm AIHA
- Cold agglutinin disease
- Paroxysmal cold hemoglobinuria (PCH)
- Drug-induced hemolytic anemia

Sold by @itachibot



In PCH, the direct Coomb's test will be positive for complement only. This condition is characterized by the presence of the Donath-Landsteiner antibody. It is a unique antibody with anti-P specificity at 4°C. Hence, it binds to the RBC at 4°C. However, it causes complement-mediated lysis at 37°C.

Interpretation of direct Coomb's test results:

Condition	Agglutination wit h	
anti- IgG	anti- C3	
Warm AIHA	+	+
Drug-induced	+	+
PCH	-	+
Cold agglutinin disease (IgM)	-	+
PNH	-	-

Solution to Question 16:

Donath Landsteiner antibody is specific for paroxysmal cold hemoglobinuria (PCH). PCH is a rare form of autoimmune hemolytic anemia.

The mechanism of this antibody includes:

• Binding to RBCs at 4°C

• Complement-mediated lysis at 37°C

This condition presents with intravascular hemolysis and hemoglobinuria. Blood transfusion is required in severe anemia. It is a self-limiting condition.

Sold by @itachibot

The differences between PCH and PNH are given below:

Condition	Paroxysmal nocturnal hemoglobinuria (PNH)	Paroxysmal cold hemoglobinuria (PCH)
Onset	Chronic with acute exacerbations	Acute
Pathogenesis	Complement-mediateddestruction of CD59/55(–) RBCs	Immune lysisof normal red cells
Diagnosis	Flow cytometry to display a CD55/CD59(–) red cell population	Test forDonath-Landsteinerantibody
Triggers	Infections, alcohol, exercise, stress, or medications	Triggered by viral infection

Sold by @itachibot
If you purchased this from someone,
you may have been scammed.

Solution to Question 17:

Microangiopathic hemolytic anemia (MAHA) is not seen in prosthetic heart valves. It is associated with macroangiopathic hemolytic anemia.

MAHA refers to cell fragmentation that occurs in microcirculation. Microvascular lesions lead to luminal narrowing due to the deposition of fibrin and platelets. These produce shear stress that mechanically injures passing red cells. Hence, it is a type of non-immune hemolysis (Coombs-negative). On peripheral blood smear, schistocytes are seen.

Common causes of MAHA include:

- Thrombotic thrombocytopenic purpura
- Hemolytic uremic syndrome
- Disseminated intravascular coagulation
- Vasculitis
- Malignant hypertension
- APLA syndrome
- Drug-induced hemolysis
- Eclampsia

Solution to Question 18:

The best prognosis in atypical HUS is seen in membrane cofactor protein (MCP) gene mutation.

Familial HUS or atypical HUS (aHUS) is the only inherited cause of extracorpuscular hemolytic anemia. It can occur due to mutations in any one of several genes encoding complement regulatory proteins. Mutations in the following genes are seen:

Sold by @itachibot

• Membrane cofactor protein - best prognosis

• Complement factor B - worst prognosis

• Complement factor H - most common

• Complement factor I - least common

In these patients, infection or other triggers cause complement activation through the alternative pathway. This causes hemolysis and damage to endothelial cells in the kidney. It is usually self-limited.

The standard treatment is plasma exchange. Eculizumab has recently been shown to be beneficial in atypical HUS.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 19:

Deferoxamine has poor oral bioavailability and is administered parenterally (SC/IV).

Oral formulations of both deferiprone and deferasirox are available.

The adverse effects of iron chelators include:

• Deferiprone - neutropenia, and agranulocytosis.

• Deferasirox - transaminitis and mild, asymptomatic rise in serum creatinine

• Deferoxamine - occasional cataracts, deafness, and local skin reactions, but is usually well-tolerated

Solution to Question 20:

Naked eye single tube red-cell osmotic fragility test (NESTROFT) is the most sensitive test to detect thalassemia in the population. Hence, it is an effective screening test.

The principle is based on the decreased osmotic fragility of red cells in thalassemia. It is performed as follows:

• Mix EDTA blood and 0.36% buffered saline solution.

• Shake the tube and leave it at room temperature for 30 minutes.

• Shake the tube again and immediately hold the tube in front of a piece of paper with text.

The image below shows a positive NESTROF test (test tube marked as T1) as the line is not visible through the test tube. The test tube marked as C shows the control (Please note that a negative NESTROF test would show visible lines similar to the appearance of the control).

NESTROFT	Inference	Reason
Lines visible (negative)	Normal	Hemolysis has occurred
Lines not seen (positive)	Thalassemia	Stable RBC, unable to lyse

Sold by @itachibot

The following tests are specific and used to confirm the diagnosis of thalassemia:

• Hemoglobin electrophoresis

• High-performance liquid chromatography (HPLC)

• Genetic testing

Note: Eosin 5' maleimide (EMA) flow cytometry is the most specific test for the diagnosis of hereditary spherocytosis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 21:

This clinical scenario is consistent with β-thalassemia minor. The diagnosis can be established by hemoglobin electrophoresis.

This condition presents with the following features:

• Microcytes (N: 80-96 fL)

• Hypochromic (N: 27-33 pg)

• Mild anemia

• Target cells

• Normal iron profile

Hemoglobin electrophoresis findings are as follows:

• ≥4% of HbA2

• ≤5% of HbF

Sold by @itachibot

Solution to Question 22:

Acute pain crisis is not associated with sickle cell trait. Acute pain crisis is classicaly seen in sickle cell disease.

Manifestations of sickle cell trait include:

• Often asymptomatic.

• Hyposthenuria

• Painless hematuria - due to papillary necrosis

• Recurrent UTIs

• Risk of renal medullary carcinoma - rare

• Increased risk of CKD

• Sloughing of papillae with urethral obstruction is very rare

• Anemia and painful crises are rare.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 23:

Acute pain crisis is the most common clinical presentation of sickle cell anemia.

These are intermittent episodes of vaso-occlusion in connective and musculoskeletal structures. This results in ischemia resulting in acute pain and tenderness, fever, tachycardia, and anxiety. The management includes vigorous hydration, evaluation for underlying causes, and aggressive analgesia.

Other manifestations of sickle cell anemia include:

• Hand-foot syndrome - painful infarcts of the digits and dactylitis

• Priapism - infarction of the penile venous outflow tracts

• Chronic lower leg ulcers - ischemia and superinfection in the distal circulation

• Splenic sequestration crisis - life-threatening complication due to acute venous obstruction that requires emergency transfusion and/or splenectomy

• Aplastic crisis - commonly secondary to parvovirus B19 infection

• Acute chest syndrome - sickling within the lung, producing pain and temporary pulmonary dysfunction. Treatment may include lifelong transfusion support, using partial exchange transfusion

• Stroke - common in children

Sold by @itachibot

Myeloproliferative Disorders and Aplastic Anemia

Question 1:

According to WHO 2022 classification, which of the following is not a chronic myeloproliferative neoplasm?

a) Chronic myeloid leukemia

b) Chronic eosinophilic leukemia

c) Essential thrombocythemia

d) Chronic lymphocytic leukemia

Question 2:

What is the most common myeloproliferative neoplasm?

a) Chronic myeloid leukemia

b) Primary myelofibrosis

c) Essential thrombocytosis

d) Polycythemia vera

Question 3:

During a routine health checkup, the following lab reports were obtained for a young male patient. What will be your next step in the evaluation?

a) Perform bone marrow aspiration

b) Measure RBC mass

c) Measure arterial oxygen saturation

d) Look for JAK2 mutation

Question 4:

A 45-year-old man is found to have a hemoglobin level of 18 g/dl, a hematocrit of 54%, and an elevated erythropoietin level. Which of the following is the least likely diagnosis?

a) Carbon monoxide poisoning

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) Cerebellar hemangioblastoma

c) Renal cell carcinoma

d) Polycythemia vera

Question 5:

What is the most common cytogenetic abnormality in patients with polycythemia vera?

a) JAK6 mutation

b) JAK2 mutation

c) JAK8 mutation

d) JAK4 mutation

Question 6:

Which of the following statements about polycythemia vera is true?

a) Risk of thrombosis strongly correlates with the degree of thrombocytosis

b) Generalized pruritus is a consequence of mast cell activation by JAK 2 mutation

c) It is protective against H.pylori infection

d) Acute myeloid leukemia is the most common cause of mortality

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following would not be seen in patients with polycythemia vera?

a) Elevated platelet count

b) Elevated serum lactate dehydrogenase

c) Low to normal erythropoietin levels

d) Low WBC count

Question 8:

A 40-year-old man presents with a 1-month history of severe itching every time he takes a hot shower. He also reports a burning sensation in his fingers and frequent headaches. His hemoglobin is 18 g/dL and his EPO levels are low. Which of the following can provide symptomatic relief in this patient?

Sold by @itachibot

a) Dabrafenib

b) Nilotinib

c) Eculizumab

d) Ruxolitinib

Question 9:

You are a first-year medicine resident. While examining a patient, you note that his spleen is enlarged 8.5 cm below his costal margin. You would include all of the following in your differential diagnosis except:

a) Essential thrombocythemia

b) Chronic myeloid leukemia

c) Primary myelofibrosis

d) Polycythemia vera

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 10:

A 24-year-old woman, who was diagnosed with essential thrombocythemia, is being planned for cytoreductive therapy. What first-line drug should she be started on?

a) Anagrelide

b) Busulfan

c) Interferon alpha

d) Hydroxyurea

Question 11:

A 60-year-old man was referred with severe anemia and hepatosplenomegaly. Peripheral smear showed teardrop cells, nucleated RBCs, and giant abnormal platelets. Bone marrow aspirate could not be obtained despite multiple attempts, hence a bone marrow biopsy was done. The findings are given below. What is the most likely diagnosis?

Sold by @itachibot



a) Polycythemia vera

b) Essential thrombocythemia

c) Primary myelofibrosis

d) Chronic myeloid leukemia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Bone marrow aspirate of a patient with pancytopenia revealed a hypercellular marrow. Which of the following conditions is unlikely?

a) Paroxysmal nocturnal hemoglobinuria

b) Fanconi anemia

c) Megaloblastic anemia

d) Myelodysplastic syndrome

Question 13:

A 20-year-old male patient with the following exam findings is found to have pancytopenia. What is the most likely diagnosis?

Sold by @itachibot

a) Fanconi anemia

b) Shwachman-Diamond syndrome

c) Diamond-Blackfan syndrome

d) Dyskeratosis congenita

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 14:

All of the following can cause acquired aplastic anemia except:

a) Cyclophosphamide therapy

b) Radiation exposure

c) Paroxysmal nocturnal hemoglobinuria

d) Herpes simplex infection

Question 15:

A 25-year-old lady is being evaluated for recurrent epistaxis. On examination, the spleen is not palpable. Investigations reveal Hb of 6 g/dL, platelet count of 18,000/µL, and WBC count of 3000/µL. The results of a bone marrow biopsy are given below. What is the best method of management in this patient?

Sold by @itachibot



a) Administer hematopoietic growth factors

b) Umbilical cord blood transplantation

c) Hematopoietic stem cell transplantation

d) Antithymocyte globulin with cyclosporine

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 16:

What is the most common cytogenetic abnormality seen in patients with myelodysplastic neoplasm?

a) 7q deletion

b) 5q deletion

c) Monosomy 5

d) Monosomy 7

Question 17:

A patient with myelodysplastic syndrome was found to have 5q deletion. What is the drug of choice for this patient?

a) Cyclosporine

b) Cyclophosphamide

c) Thalidomide

d) Lenalidomide

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	d
3	b
4	d
5	b
6	b
7	d
8	d
9	a
10	d
11	c
12	b
13	d
14	d
15	c
16	b
17	d

If yo... you... else, ...d.

Detailed Explanations

Solution to Question 1:

Chronic lymphocytic leukemia (CLL) is a lymphoid neoplasm. It is not a myeloproliferative neoplasm.

Myeloproliferative neoplasms (MPN) are a group of disorders that cause an increased number of red blood cells, white blood cells, or platelets. According to 5th edition WHO classification (2022), myeloproliferative neoplasms consist of the following:

- Chronic myeloid leukemia (CML)
- Polycythemia vera (PV)
- Primary myelofibrosis (PMF)
- Essential thrombocythemia

Sold by @itachibot

• Juvenile myelomonocytic leukemia (JMML)

• Chronic neutrophilic leukemia (CNL)

• Chronic eosinophilic leukemia

• Myeloproliferative neoplasm, not otherwise specified

Solution to Question 2:

The most common myeloproliferative neoplasm is polycythemia vera.

It is a hematopoietic stem cell disorder that is characterized by uncontrolled proliferation of the red blood cells in the body. It is also frequently associated with an increase in white blood cells and platelets.

Solution to Question 3:

This patient has increased Hb and normal WBCs and platelets. The next step is to measure RBC mass to evaluate for polycythemia.

A person has to be evaluated for polycythemia if there is:

• Increased hemoglobin: >16.5 g/dL in men or >16.0 g/dL in women, (or)

• Increased hematocrit: >49 percent in men or >48 percent in women

This condition is due to either a true increase in the number of RBCs (absolute polycythemia) or an apparent increase due to a decrease in the plasma volume (relative polycythemia). Both of these conditions can be differentiated by measuring RBC mass (amount of total blood volume that is occupied by RBCs).

In patients with absolute polycythemia, erythropoietin (EPO) levels must be measured.

• Primary polycythemia - low or normal EPO levels

• Secondary polycythemia - elevated EPO levels

Solution to Question 4:

Polycythemia vera is associated with low or normal erythropoietin (EPO) levels.

Polycythemia refers to an increase in the number of red blood cells (RBC) in the body. It is classified as:

Autonomous production of erythropoietin occurs in renal cell carcinoma (RCC), hepatocellular carcinoma, cerebellar hemangioblastoma.

Relative (spurious) polycythemia refers to an apparent increase in RBC or hemoglobin count due to a decrease in the plasma volume. It is commonly seen in dehydration.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

	Primarypolycythemia (polycythemia vera)	Secondarypolycythemia
Pathophysiology	Caused by mutations in the hematopoietic stem cells	Hypoxia (COPD,carbon monoxidepoisoning)Autonomous EPO production
Erythropoietin (EPO)	Normal or low	High
WBCs, and platelet count	High (due to increased production of all cell lines)	Normal

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 5:

The most common cytogenetic abnormality in polycythemia vera is the JAK2 mutation.

JAK2 mutations can be found in the following neoplasms:

• Polycythemia vera

• Essential thrombocythemia

• Primary myelofibrosis

Solution to Question 6:

Generalized pruritus seen in polycythemia vera (PV) is a consequence of the activation of mast cells by JAK 2 mutation.

Other options:

Option A: The risk of thrombosis strongly correlates with erythrocytosis, not the degree of thrombocytosis.

Option C: In polycythemia vera, there is an increased risk of H.pylori infection.

Option D: Thrombosis is the major cause of mortality in polycythemia vera. The incidence of acute leukemia in patients not exposed to chemotherapy or radiation therapy is low.

PV is a clonal hematopoietic stem cell disorder in which red cells, granulocytes & platelets accumulate in the absence of a physiological stimulus. The most common mutation seen in PV is the JAK2 mutation.

Clinical features of polycythemia vera are:

• Thrombosis is the most common manifestation and is the major cause of mortality. Uncontrolled erythrocytosis causes increased viscosity of the blood causing thrombosis of the vessels. Patients may present with stroke, transient ischemic attacks, myocardial infarction, and Budd-Chiari syndrome.

• Aquagenic pruritis is generalized itching following a warm bath or a shower due to increased mast cells and histamine.

Sold by @itachibot

• Erythromelalgia is erythema and pain in the fingers, hands, and feet due to increased platelet stickiness.

• Splenomegaly due to increased destruction of RBCs in the spleen

• Gout is due to the excessive proliferation of the marrow cells leading to elevated uric acid levels.

Solution to Question 7:

Low WBC count is not a feature of polycythemia vera. It is associated with an elevated WBC count.

Common laboratory findings of polycythemia vera are:

• Elevated hemoglobin concentration, hematocrit levels, and RBC count

• Elevated WBCs and platelet counts

• Low to normal erythropoietin levels

• Elevated serum lactate dehydrogenase

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

This clinical scenario is suggestive of polycythemia vera. In these patients, ruxolitinib, a Janus-kinase inhibitor is used to provide symptomatic relief.

The treatment of polycythemia vera is as follows:

• Therapeutic phlebotomy - definitive treatment

• Anticoagulants - in patients with thrombosis

• Symptomatic relief:

• Ruxolitinib

• Hydroxyurea

• Interferon-alpha

• PUVA therapy

Solution to Question 9:

Enlargement of the spleen > 8 cm below the costal margin or weight > 1000 g is called massive splenomegaly. It is not a feature of essential thrombocythemia (ET), as it causes only mild splenomegaly.

The causes of massive splenomegaly include:

• Chronic myeloid leukemia

Sold by @itachibot

• Primary myelofibrosis

• Polycythemia vera

• Chronic lymphocytic leukemia

• Lymphoma

• Hairy cell leukemia

• Gaucher's disease

• Sarcoidosis

• Autoimmune hemolytic anemia

ET is a neoplasm in which the body produces too many platelets, resulting in a platelet count > 4.5 lakhs/µL. It occurs due to the mutations in the JAK2, CALR, or MPL genes.

The majority of the patients are asymptomatic and are diagnosed incidentally. Some patients may present with thrombotic events or rarely, hemorrhages. The following findings are seen:

• Peripheral smear - abnormally large platelets

• Bone marrow biopsy - increased number of megakaryocytes, with hyperlobulated nuclei

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

The first-line drug for cytoreductive therapy in patients with essential thrombocythemia is hydroxyurea.

Treatment of essential thrombocythemia is as follows:

• No symptoms or risk of thrombosis:

• Observation alone

• Low dose aspirin may be considered

• High risk of thrombosis (cardiovascular risk factors, prior history of thrombosis)

• Low dose aspirin

• Cytoreduction with hydroxyurea, busulfan, or interferon-alpha

Solution to Question 11:

The image shows the deposition of numerous collagen fibers in the bone marrow, which indicates marrow fibrosis. These clinical findings and history of a dry tap on bone marrow aspiration are diagnostic of primary myelofibrosis.

In this condition, the neoplastic proliferation of hematopoietic stem cells results in increased and abnormal proliferation of the megakaryocytes. These megakaryocytes release excessive amounts of platelet-derived growth factor (PDGF) and TGF-β. Increased growth factors lead to the deposition of collagen in the bone marrow which eventually leads to fibrosis.

Sold by @itachibot

The clinical features include:

• Early phase:

• Thrombocytosis (large, abnormal platelets) and leukocytosis

• Hypercellularity of bone marrow

• Late phase:

• Pancytopenia

• Hypocellularity and fibrosis of bone marrow (dry tap)

• Teardrop-shaped RBCs and leukoerythroblastosis (premature release of nucleated erythroid and early granulocytes into blood)

• Hepatosplenomegaly - due to extramedullary hematopoiesis.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 12:

Fanconi anemia results in pancytopenia with hypocellular bone marrow. Hence, it is not likely in this patient.

Causes of pancytopenia with hypocellular bone marrow include:

• Inherited aplastic anemia

• Fanconi anemia

• Dyskeratosis congenita

• Shwachman-Diamond syndrome

• Acquired aplastic anemia

Causes of pancytopenia with hypercellular bone marrow include:

• Primary bone marrow disorders

• Paroxysmal nocturnal hemoglobinuria (PNH)

• Myelodysplastic syndrome

• Myelofibrosis

• Systemic diseases

• Megaloblastic anemia

• Alcoholism

• Sarcoidosis

• Tuberculosis

• Brucellosis

Solution to Question 13:

Sold by @itachibot

The most likely diagnosis is dyskeratosis congenita. This condition is characterized by pancytopenia along with the clinical triad of:

• Reticular skin pigmentation

• Nail dystrophy

• Oral leukoplakia

Option A: Fanconi anemia is an autosomal recessive disorder characterized by pancytopenia, congenital anomalies, and an increased risk of malignancies.

Option B: Shwachman-Diamond syndrome presents with bone marrow failure and exocrine pancreatic insufficiency.

Option C: Diamond-Blackfan syndrome presents with low RBC counts due to congenital erythroid aplasia.

Note: Fanconi anemia is different from Fanconi syndrome. Fanconi syndrome is a disorder of renal tubular function (type 2 RTA) that results in increased excretion of glucose, amino acids, bicarbonates, and phosphates.

Solution to Question 14:

Herpes simplex infection is not a cause of acquired aplastic anemia.

Secondary causes of acquired aplastic anemia include:

• Radiation

• Toxins - Benzene

• Drugs

• Chemotherapy drugs (including cyclophosphamide)

• Sulfonamides

Sold by @itachibot

• Chloramphenicol

• Infections:

• 1-2 months after an attack of seronegative viral hepatitis

• HIV, EBV, CMV

• Parvovirus B19

• Eosinophilic fasciitis

• Paroxysmal nocturnal hemoglobinuria

• Graft vs host disease

Solution to Question 15:

The clinical and laboratory findings, as well as bone marrow hypocellularity are diagnostic of aplastic anemia. In this young patient, the best method of management is hematopoietic stem cell transplantation.

Aplastic anemia presents with pancytopenia and resulting bleeding manifestations, anemia, and frequent infections. Bone marrow biopsy reveals hypocellularity with fatty infiltration of the bone marrow. The management is as follows:

• Hematopoietic stem cell transplantation - best method of management in young patients

• Antithymocyte globulin with cyclosporine - elderly patients and patients without HLA-matched donors

• Eltrombopag - a thrombopoietin analog used as add-on therapy

Solution to Question 16:

The most common cytogenetic abnormality seen in myelodysplastic neoplasm is the 5q deletion.

Myelodysplastic syndromes/neoplasm (MDS) are a group of disorders in which abnormal stem cells produce increased numbers of immature blood cells. These immature cells are defective and die prematurely resulting in lower numbers of mature blood cells. It is characterized by pancytopenia and a high risk of acute myeloid leukemia (AML).

The cytogenetic abnormalities identified in this condition include:

• 5q deletion

• 3q deletion

• 7q deletion

• Monosomy 7

• Monosomy 5

5th edition WHO Classification of MDS (2022):

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

1. MDS with defining genetic abnormalities:

• MDS with a low blast and isolated 5q deletion (MDS-5q): <5% blasts in the bone marrow or <2% blasts in the peripheral blood.

• MDS with a low blast and SF3B1 mutation (MDS-SF3B1): Detection of ≥15% ring sideroblasts may substitute for SF3B1 mutation.

• MDS with biallelic TP53 inactivation (MDS-biTP53): <20% blasts in the bone marrow or peripheral blood.

2. MDS, morphologically defined:

• MDS with low blasts: <5% blasts in the bone marrow or <2% blasts in the peripheral blood.

• MDS-hypoplastic: ≤25% bone marrow cellularity

• MDS with increased blasts (IB):

• MDS-IB1: 5–9% of blasts in bone marrow or 2–4% in peripheral blood

• MDS-IB2:10-19% of blasts in bone marrow or 5-19% in peripheral blood or Auer rods

• MDS-fibrosis: 5-19% of blasts in bone marrow or 2–19% in peripheral blood

3. Childhood myelodysplastic neoplasm:

• Childhood MDS with low blasts:<5% blasts in the bone marrow or <2% blasts in the peripheral blood

• Childhood MDS with increased blasts: 5-19% blasts in the bone marrow or 2-19% blasts in the peripheral blood.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 17:

Lenalidomide is the drug of choice in patients with myelodysplastic syndrome (MDS) associated with 5q deletion.

It is a thalidomide derivative with the following benefits in these patients:

• Patients achieve transfusion independence

• Normal or near-normal hemoglobin levels

• Normalization of cytogenetics

Adverse effects include:

• Myelosuppression (worsening thrombocytopenia and neutropenia, necessitating blood count monitoring)

• Deep vein thrombosis and pulmonary embolism

However, hematopoietic stem cell transplant remains the definitive treatment.

Sold by @itachibot

Platelet Disorders

Question 1:

What is the lifespan of a platelet?

a) 24 hours

b) 5 days

c) 7 days

d) 2 weeks

Question 2:

Which of the following would you not expect to see in patients with a platelet disorder?

a) Petechiae

b) Ecchymosis

c) Hemarthrosis

d) Gastrointestinal bleeds

Question 3:

Which of the following is a disorder of platelet adhesion?

a) Glanzmann's thrombasthenia

b) Bernard Soulier syndrome

c) Wiskott Aldrich syndrome

d) Hermansky Pudlak syndrome

Question 4:

Patients with which of the following conditions would have a low platelet count?

a) Platelet storage pool disorder

b) Wiskott-Aldrich syndrome

c) Henoch-Schonlein purpura

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) Hemophilia

Question 5:

Immune-mediated destruction of platelets is seen in patients with which of the following conditions?

a) Bernard-Soulier syndrome

b) Hermansky-Pudlak syndrome

c) Thrombotic thrombocytopenic purpura

d) Drug-induced thrombocytopenia

Question 6:

A young man comes to you with complaints of bleeding gums and the following skin rash. He reports that he had diarrhea last week after eating at a roadside eatery. History is significant for deep vein thrombosis 6 months ago that was treated with heparin. His CBC report revealed a platelet count of 15,000/µL, normal RBC and WBC count. His peripheral smear was normal. What is the likely diagnosis?

a) Hemolytic uremic syndrome

b) Thrombotic thrombocytopenic purpura

c) Immune thrombocytopenic purpura

d) Heparin-induced thrombocytopenia

Sold by @itachibot
If you bought this from someone else, it has been scammed.

Sold by @itachibot

Question 7:

Two weeks after an upper respiratory tract infection, a 5-year-old presents with ecchymosis and petechiae all over the body. Abdominal examination revealed no hepatosplenomegaly. The platelet count is 15,000/µL and the coagulation profile is normal. Which of the following is false regarding this condition?

a) Peripheral blood smear is normal

b) Bone marrow biopsy shows increased megakaryocytes

c) Bleeding into the joints is a common manifestation

d) The condition resolves spontaneously in 80% patients

Question 8:

A 50-year-old woman comes to the ER after developing a spontaneous nosebleed. Over the past 1 week, she has noticed blood on her toothbrush while brushing her teeth. She was diagnosed with hepatitis C three months ago. Investigations reveal a platelet count of 32,000/µL and a bleeding time of 9 minutes . Which of the following is false regarding the treatment of this patient?

a) She requires regular platelet transfusions until platelet count is normalized

b) Prednisone or anti-D immunoglobulins can be used as the first-line treatment

c) Splenectomy may be done if she relapses after medical management

d) Romiplostim and eltrombopag can be used if there is no response to splenectomy

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 9:

Which of the following is the best treatment for the relapse of immune thrombocytopenic purpura?

a) IV immunoglobulins

b) Steroids

c) Splenectomy

d) Blood transfusion

Question 10:

Sold by @itachibot

It is your first day as an intern in the medicine ward. The senior resident tells you to check the platelet count of a patient who is being monitored for thrombocytopenia. He was started on a new medication the previous night. What drug was most likely prescribed?

a) Phenytoin

b) Ampicillin

c) Abciximab

d) Amiodarone

Question 11:

Patients with all of the following usually present with bleeding except:

a) Heparin-induced thrombocytopenia

b) Drug-induced thrombocytopenia

c) Inherited thrombocytopenia

d) Infection-induced thrombocytopenia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Which of the following cannot be used in the treatment of patients with heparin-induced thrombocytopenia?

a) Danaparoid

b) Lepirudin

c) Argatroban

d) Enoxaparin

Question 13:

The CBC of a 33-year-old woman revealed a low platelet count. However, further evaluation revealed this to be an in vitro sampling error. Which of the following is the likely cause of this finding?

a) EDTA

b) Sodium citrate

c) Heparin

d) Finger stick sample

Sold by @itachibot

Question 14:

Which of the following is false regarding patients with Bernard-Soulier syndrome?

a) Platelet aggregation tests with epinephrine and ADP are normal

b) Ristocetin aggregation test is normal

c) Peripheral smear shows large platelets

d) Mild thrombocytopenia is seen

Question 15:

Which of the following is the most common inherited bleeding disorder?

a) Hemophilia A

b) Hemophilia B

c) Immune thrombocytopenia purpura

d) Von Willebrand disease

Question 16:

Which of the following is called autosomal hemophilia?

a) Type 2M von Willebrand disease

b) Type 2N von Willebrand disease

c) Type 2A von Willebrand disease

d) Type 2B von Willebrand disease

Question 17:

Which of the following patients has Heyde's syndrome?

a) A man with aortic stenosis with gastrointestinal bleeding

b) A woman with aortic stenosis with intracranial bleed

c) A man with aortic regurgitation with gastrointestinal bleeding

d) A woman with aortic regurgitation with intracranial bleed

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 18:

Which of the following laboratory findings is seen in a patient with von Willebrand's disease?

a) Normal ristocetin aggregation test

b) Increased bleeding time

c) Increased prothrombin time

d) Decreased activated partial thromboplastin time

Question 19:

A 28-year-old woman is being evaluated for prolonged bleeding following a tooth extraction. She also reports heavy bleeding during menses and recurrent gum bleeds. Labs show increased aPTT, normal PT, and increased bleeding time. Which of the following is the first-line treatment for this patient?

a) Tranaxemic acid

b) Factor VIII concentrate

c) Von Willebrand factor concentrate

d) Desmopressin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 20:

Which of the following features is not seen in patients with Glanzmann's thrombasthenia?

a) Abnormal platelet aggregation on adding ADP

b) Decreased platelet count

c) Normal von Willebrand factor levels

d) Normal platelet aggregation to ristocetin

Question 21:

A physician decides to start a patient with stroke on antiplatelet medication. All of the following drugs would be effective except:

a) Clopidogrel

b) Ticlopidine

c) Ticagrelor

Sold by @itachibot

d) Pagoclone

Question 22:

A 24-year-old woman underwent evaluation for bleeding manifestations and severe anemia. Investigations revealed combined autoimmune hemolytic anemia and idiopathic thrombocytopenic purpura. What is the likely diagnosis?

a) Evans syndrome

b) May-Hegglin anomaly

c) Fechtner syndrome

d) Rosenthal syndrome

Question 23:

What is the most common side effect experienced by patients taking enteric-coated aspirin?

a) Gastrointestinal symptoms

b) Aspirin sensitivity

c) Aspirin tolerance

d) Medication overuse headache

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 24:

Which of the following is incorrect about the mechanism of action of dipyridamole?

a) Increases levels of cyclic AMP in platelets

b) Blocks the reuptake of adenosine

c) Inhibits phosphodiesterase-mediated cyclic AMP degradation

d) Increases intracellular adenosine levels

Question 25:

A 62-year-old man diagnosed with myocardial infarction is planned for percutaneous intervention. He was diagnosed with diabetic nephropathy 2 months ago. The cardiologist decides to start him on aspirin and an additional antiplatelet drug. Which of the following can be used for this patient?

Sold by @itachibot

a) Abciximab only

b) Abciximab or eptifibatide

c) Tirofiban or eptifibatide

d) Tirofiban only

Question 26:

Plasmapheresis is the mainstay of treatment in patients with which of the following?

a) Autoimmune hemolytic anemia

b) Disseminated intravascular coagulation

c) Thrombotic thrombocytopenic purpura

d) Hemolytic uremic syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 27:

Which of the following is not seen in patients with thrombotic thrombocytopenic purpura?

a) Hemolytic anemia

b) Focal neurological deficits

c) Venous thromboembolism

d) Elevated serum creatinine

Question 28:

Which of the following is incorrect regarding patients with thrombotic thrombocytopenic purpura?

a) Clinical features overlap with hemolytic uremic syndrome

b) PT and aPTT are prolonged

c) It is common in patients with HIV infection

d) It can be associated with use of quinine

Question 29:

What is the most common genetic abnormality seen in patients with recurrent history of venous thrombosis?

Sold by @itachibot

a) Factor V Leiden mutation

b) Prothrombin mutation

c) Protein C deficiency

d) Antithrombin III deficiency

Question 30:

Which of the following qualitative platelet disorders are inherited as autosomal dominant?

a) Glanzman thrombasthenia

b) Bernard Soulier syndrome

c) Chediak-Higashi syndrome

d) Platelet storage pool syndrome

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	b
5	d
6	c
7	c
8	a
9	c
10	c
11	a
12	d
13	a
14	b
15	d
16	b
17	a
18	b

Sold by @itachibot

19	d
20	b
21	d
22	a
23	a
24	d
25	a
26	c
27	c
28	b
29	a
30	d

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

The average lifespan of platelets in circulation is 7–10 days.

Thrombopoietin (TPO) is the regulator of platelet production. A reduction in platelet and megakaryocyte mass increases the level of TPO, which then stimulates platelet production.

The shelf life of donor platelets is 5 days. It is stored on a special agitator at 20-24 °C. The half-life of transfused platelets is 3-5 days.

Solution to Question 2:

Hemarthrosis is highly suggestive of a coagulation disorder, not a platelet disorder.

Bleeding diathesis can occur due to:

• Platelet disorders - mucocutaneous bleeding like petechiae, purpura, ecchymoses, epistaxis, menorrhagia, gastrointestinal (GI) bleeds

• Coagulation disorders - hemarthrosis, muscle hematomas, and bleeding into soft tissues

• Disorders of the vessel wall (Henoch-Schönlein purpura, scurvy) - purpura and ecchymosis

Sold by @itachibot



Solution to Question 3:

Bernard Soulier syndrome is a disorder of platelet adhesion. It occurs due to an inherited defect in the GPIb-IX complex.



Solution to Question 4:

Thrombocytopenia is a feature of Wiskott-Aldrich syndrome.

It is an X-linked recessive condition caused by mutations in the WASP gene. It affects T-cells and platelets and presents with the following features:

• Eczema

Sold by @itachibot

• Thrombocytopenia - along with decreased platelet size, impaired platelet aggregation response

• Immunodeficiency - low IgM, high IgE, defective regulatory T cells causing recurrent bacterial infections.

Option A: Platelet storage pool disorder is a disorder of defective platelet granule formation.

Option C: Henoch-Schonlein purpura is a type of small vessel vasculitis in children and young adults.

Option D: Hemophilia is a coagulation disorder, hence the platelet count is not affected.

Solution to Question 5:

Immune-mediated destruction of platelets is seen in drug-induced thrombocytopenia. Common drugs that cause this are quinidine, heparin, sulfa compounds.

Drug-dependent antibody binding can only be demonstrated in the presence of the drug in the laboratory assays. The thrombocytopenia normally occurs within 21 days of initial exposure and usually gets resolved after 7-10 days once the drug is stopped.

Immune-mediated destruction of platelets is seen in:

• Drug-induced thrombocytopenia

• Infection-induced thrombocytopenia

• Idiopathic/ immune thrombocytopenic purpura (ITP)

Option A: Bernard-Soulier syndrome is a disorder of platelet adhesion. There are no autoantibodies against platelets.

Option B: Hermansky-Pudlak syndrome is an inherited disorder with defective granule formation in platelets. It affects platelet activation and aggregation.

Option C: Thrombotic microangiopathies like thrombotic thrombocytopenic purpura (TTP) and hemolytic uremic syndrome (HUS) lead to thrombocytopenia due to increased platelet consumption.

Solution to Question 6:

The given clinical scenario and image showing petechiae are suggestive of immune thrombocytopenic purpura.

Immune thrombocytopenic purpura (ITP) is an acquired disorder with autoantibodies against platelet membrane glycoproteins IIb-IIIa or Ib-IX. It can either be primary (idiopathic) or secondary to autoimmune conditions like SLE or infections like HIV and hepatitis C.

It presents with features of mucocutaneous bleeding. CBC shows isolated thrombocytopenia and peripheral smear does not reveal any hemolysis.

Options A and B: HUS and TTP are microangiopathies. They are associated with hemolysis and schistocytes on peripheral smear.

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

Option D: Heparin-induced thrombocytopenia typically presents within 5-10 days of treatment. Hence, it is not the likely diagnosis in this patient.

Acute ITP	Chronic ITP
Common in children	Common in adults (females > males)
Associated with an antecedent viral infection	Not associated with antecedent infection
Spontaneous remission in 80 % cases	Spontaneous remission is not usually seen



Solution to Question 7:

This scenario is suggestive of acute immune thrombocytopenic purpura. Bleeding into the joints is seen in coagulation disorders and is not a manifestation of this condition.

Immune thrombocytopenic purpura or idiopathic thrombocytopenic purpura (ITP) is an acquired disorder in which there is an immune-mediated destruction of platelets. It can be acute or chronic.

Acute ITP is common in children and is usually preceded by viral infection. It presents with mucocutaneous bleeding (epistaxis, petechiae, ecchymosis). Hepatomegaly, splenomegaly, lymphadenopathy, fever and jaundice are absent. The following laboratory features are seen:

• Thrombocytopenia (commonly <20,000 / µL)

• Bleeding time - prolonged

• PT/aPTT/INR - normal

• Bone marrow - normal/increased megakaryocytes

• Peripheral smear - normal

Sold by @itachibot

This condition resolves spontaneously in 80% of the cases. Treatment is as follows:

• Platelet count < 30,000 with risk factors for bleeding (antiplatelet drugs/anticoagulants or concomitant bleeding disorders): Oral glucocorticoids or IVIG

• Severe bleeding symptoms: IV methylprednisolone ± IVIG

• Life-threatening bleeding (like intracranial hemorrhage): platelet transfusion + IV methylprednisolone + IVIG

Solution to Question 8:

This clinical scenario in an adult patient is suggestive of chronic ITP (idiopathic thrombocytopenic purpura). She does not require platelet transfusion. It is recommended only for patients with severe bleeding manifestations (intracranial/ gastrointestinal) or platelet count < 5000 µL.

Chronic ITP can either be primary (idiopathic) or secondary to autoimmune conditions like SLE or infections like HIV and hepatitis C. Depending on the clinical features, the management is as follows:

	Clinical features	Management
1.	Minor bleeding manifestations (and/or)Platelet count > 5 000/ µL	Prednisone(OR)Anti-Dimmunoglobulintherapy - effective only in Rh-positive patients (OR)Intravenous immunoglobulin (IVIG)
2.	Severe ITP (any of the following):Significant GI / intracranial bleedingPlatelet count < 5 000/µLSigns of impending bleeding (retinal or large oral mucosal bleeds)	Platelet transfusion (AND)High-dose glucocorticoids (AND)IVIG / Anti-D immunoglobulins + Rituximab
3.	Relapse after first-line therapy	Splenectomy
4.	Unresponsive to all other therapy or relapse after splenectomy	Thrombopoietin receptor agonists -romiplostim(subcutaneous),eltrombopag(oral)

Solution to Question 9:

Splenectomy is the treatment of choice in a case of immune thrombocytopenic purpura (ITP) relapse after first line therapy with glucocorticoids. The relapse usually occurs once they are tapered. Vaccination against encapsulated organisms (pneumococcus, meningococcus, H. influenzae) is advised before splenectomy is performed.

It is important to note that in some patients ITP resolves by itself, and observation along with intermittent IV immunoglobulins and anti - Rh(D) administration or administration of a

Sold by @itachibot

thrombopoietin receptor agonist (eltrombopag, romiplostim) can be a viable option before splenectomy is considered.

The treatment of ITP is as follows:

ITP is an acquired disorder characterized by the immune-mediated destruction of platelets and possibly inhibition of platelet release from the megakaryocyte. Secondary causes of ITP include systemic lupus erythematosus (SLE), HIV-AIDS, and Hepatitis C.

ITP usually presents with mucocutaneous bleeding, very low platelet count, and normal peripheral blood cells. Patients typically exhibit ecchymoses and petechiae or they may incidentally discover thrombocytopenia during a routine complete blood count. Rarely, life-threatening bleeding may occur. The presence of wet purpura (blood blisters in the mouth) and retinal hemorrhages can serve as warning signs of potentially life-threatening bleeding.

	Clinical features	Management
1.	Minor bleeding manifestations (and/or)Platelet count > 50 000/ µL	Prednisone(OR)Anti-D immunoglobulin therapy- effective only in Rh-positive patients (OR)Intravenous immunoglobulin (IVIG)
2.	Severe ITP (any of the following):Significant GI / intracranial bleedingPlatelet count < 50 000/µLSigns of impending bleeding (retinal or large oral mucosal bleeds)	Platelet transfusion (AND)High-dose glucocorticoids (AND)IVIG / Anti-D immunoglobulins + Rituximab
3.	Relapse after first-line therapy	Splenectomy
4.	Unresponsive to all other therapy or relapse after splenectomy	Thrombopoietin receptor agonists -romiplostim(subcutaneous),eltrombopag(oral)

Solution to Question 10:

Abciximab can cause a drop in platelet count within 24 hours of initial exposure. Hence, these patients must be monitored for thrombocytopenia. This is due to naturally-occurring antibodies that cross-react with the drug bound to the platelets.

Thrombocytopenia due to other drugs typically occurs several days after exposure (average 21 days) and resolves in 7-10 days after drug withdrawal. Drug-induced thrombocytopenia can occur due to the following mechanisms:

• Bone marrow suppression - chemotherapeutic drugs

• Drug dependent antibodies that react with platelet surface antigens - quinine and sulfonamides

• Naturally occurring antibodies that cross-react with drugs bound to the platelets - platelet Gp IIb/IIIa inhibitors

Sold by @itachibot

Some of the drugs that cause thrombocytopenia are:

• Antibiotics - ampicillin, cephalosporins, piperacillin, vancomycine

• Analgesics - ibuprofen, acetaminophen

• Anti-epileptics - phenytoin, carbamazepine

• Heparin

• Amiodarone

Solution to Question 11:

Heparin-induced thrombocytopenia (HIT) is not associated with bleeding. On the contrary, it is associated with an increased risk of thrombosis.

This condition occurs due to the production of antibodies against the heparin-PF4 complex (platelet factor 4, a platelet-specific protein). These antibodies can activate platelets and result in thrombosis. It is seen 5-14 days after exposure to heparin.

These antibodies can be detected using ELISA or serotonin release assay. This condition is managed by discontinuing heparin and using alternative anticoagulants.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 12:

Low molecular weight heparin (LMWH) such as enoxaparin cannot be used in the treatment of heparin-induced thrombocytopenia (HIT). This is because HIT antibodies (against anti-PF4-heparin complex) may cross-react with LMWH.

HIT can occur following exposure to unfractionated heparin and less commonly with LMWH.

The treatment of HIT is as follows:

• Discontinue heparin

• Anticoagulation with non-heparin anticoagulants even in the absence of thrombosis, since there is a high risk of thrombosis.

• Direct thrombin inhibitors: Argatroban, lepirudin, bivalirudin

• Danaparoid (heparinoid)

Solution to Question 13:

This scenario is suggestive of pseudothrombocytopenia. It is seen in blood samples collected with EDTA as the anticoagulant.

EDTA causes a reduction in the calcium content of the collected blood sample. This triggers in vitro platelet clumping and agglutination by antibodies (usually IgG), as shown below:

Sold by @itachibot



When a low platelet count is obtained with an EDTA sample, the peripheral blood smear should be evaluated. The platelet count must also be determined in blood collected in sodium citrate or heparin collection tubes or fresh fingerstick blood sample.

Sold by @itachi
If you purchased this from someone else,
you may have been scammed.

Solution to Question 14:

The ristocetin aggregation test is abnormal (no aggregation) in patients with Bernard-Soulier (BS) syndrome.

It is an inherited disorder of platelet function or qualitative platelet disorder due to deficiency of the Gp Ib-IX receptor. This is the platelet membrane receptor for the von Willebrand factor (vWF). The binding of the vWF to the GpIb-IX-V complex on the platelet membrane is essential for platelet adhesion to the vessel wall.

This condition presents with mild thrombocytopenia and peripheral smear shows large platelets.

Platelet aggregation tests are performed to measure platelet activation and aggregation in vitro. Various platelet agonists like ADP, epinephrine, collagen, and ristocetin can be used to perform these tests. In Bernard-Soulier syndrome, the following responses are seen:

• With ristocetin - No response/no aggregation, as ristocetin specifically binds to the platelet membrane via the Gp Ib-IXa receptor

• With ADP or epinephrine - Normal response, as Gp IIb-IIIa receptor is not involved in this process

Solution to Question 15:

Von Willebrand disease (vWD) is the most common inherited bleeding disorder.

Sold by @itachibot

vWD occurs due to deficient or abnormal von Willebrand factor (vWF). vWF is synthesized by endothelial cells and megakaryocytes (gene located on chromosome 12p). It is then stored in endothelial storage vesicles, Weibel-Palade bodies. Metalloprotease ADAMTS13 cleaves ultra-large multimers of VWF which are responsible for platelet adhesion.

The functions of vWF and the corresponding symptoms include:

• Adhesion of the platelets to the exposed subendothelium - symptoms of both platelet and bleeding disorders

• Prolongs the half-life of factor VIII by binding to it in circulation - hemophilia A like symptoms (rare)

Sold by @itachibot
This from someone else,
scammed.

Solution to Question 16:

Autosomal hemophilia is also called type 2N von Willebrand disease (vWD), as the clinical presentation is similar to hemophilia.

The types of vWD are:

In pseudo vWD, the vWF is normal. However, platelets have an increased affinity for vWF. It presents similar to vWD 2B with thrombocytopenia.

Type	Inheritance	Mechanism	Important characteristics
1	AD	Partial quantitative deficiency of VWF	Most commontype
3	AR	Complete quantitative deficiency	Most severe VWD
2A	AD	Loss of VWF multimers (increased cleavage by ADAMS13)	Platelet adhesion defect
2B	AD	Increased binding of VWF to platelets causing VWF-platelet complex destruction	Thrombocytopenia
2M	AD	Decreased affinity of VWF to platelets	Platelet adhesion defect, very rare
2N	AR	Decreased affinity of VWF to Factor VIII	Autosomal hemophilia

Solution to Question 17:

Heyde's syndrome is aortic stenosis with gastrointestinal bleeding, due to angiodysplasia of the gastrointestinal tract.

The shear stress on blood passing through the stenotic aortic valve produces a change in the von Willebrand factor making it susceptible to serum proteases. Hence, large multimer forms are lost,

Sold by @itachibot

leading to an acquired type 2 von Willebrand disease (vWD). The condition is reversed with the replacement of the aortic valve.

Other causes of acquired vWD include:

• Multiple myeloma

• Non-Hodgkin's lymphoma

• Polycythemia vera

• Systemic lupus erythematosus

Solution to Question 18:

Bleeding time is increased in von Willebrand disease (vWD). It is a combination of a platelet adhesion defect with factor VIII deficiency.

The laboratory findings in vWD include:

• Platelet count is normal (except type 2B)

• Bleeding time is prolonged

• Ristocetin aggregation is abnormal

• Other platelet aggregation tests are normal

• Prothrombin time (PT) is normal

• aPTT is prolonged

• VWF activity is decreased

• Factor VIII activity is decreased

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 19:

This clinical scenario is suggestive of von Willebrand's disease. The preferred first-line treatment is desmopressin (DDAVP).

This drug acts on endothelial V2 receptors causing the release of VWF and factor VIII from endothelial stores. DDAVP can be given intravenously or by an intranasal spray. The side effects include tachyphylaxis and hyponatremia.

VWF concentrate and factor VIII concentrate are reserved for patients with severe disease or prior to major procedures. Antifibrinolytic therapy (tranexamic acid or ε-aminocaproic acid) may be used for the prevention and treatment of mucosal bleeding.

Solution to Question 20:

Thrombocytopenia is not seen in Glanzmann's thrombasthenia.

Sold by @itachibot

Glanzmann's thrombasthenia is a primary defect of platelet function (aggregation). It occurs due to congenital absence of the GpIIb/IIIa receptor. It presents as superficial mucocutaneous bleeding with normal platelet count and single isolated platelets without clumping on peripheral smear. The levels of vWF are normal.

Normally, release of ADP causes platelets to change shape and bind to fibrinogen through the GpIIb/IIIa receptor. In Glanzmann's thrombasthenia, absence of the receptor causes abnormal aggregation (absent or reduced) on adding ADP. However, aggregation in response to ristocetin is normal, as it is dependent on GpIb and vWF.

Secondary platelet dysfunction can be seen in the following conditions:

• Uremia

• Paraproteinemia

• Myeloproliferative disorders

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 21:

Pagoclone is an anxiolytic medication. It is not an antiplatelet drug.

The thienopyridine drugs are ADP receptor or P2Y12 antagonists that inhibit ADP mediated platelet aggregation. They include the following drugs:

• Clopidogrel - commonly used

• Ticagrelor - reversible drug with a shorter t1/2, does not require activation, may cause mild, self-limited dyspnea

• Prasugrel - 10 times more potent than clopidogrel, rapid onset of action, better absorption, more predictable

• Ticlopidine

Clopidogrel and prasugrel are prodrugs that require activation by cytochrome P450 enzymes. CYP2C19 polymorphisms can cause reduced platelet inhibition with clopidogrel and increased risk of cardiovascular events.

Solution to Question 22:

Combined autoimmune hemolytic anemia (AIHA) and idiopathic thrombocytopenic purpura (ITP) is called Evans syndrome.

Options B and C: May-Hegglin anomaly and Fechtner syndrome are autosomal dominant hereditary conditions presenting with bleeding, sensorineural hearing loss, glomerulonephritis, and/or eye abnormalities. The severity of the bleeding symptoms is out of proportion to the moderately decreased platelet count.

Option D: Rosenthal syndrome or hemophilia C refers to factor XI deficiency.

Sold by @itachibot

Solution to Question 23:

The most common side effect of aspirin is gastrointestinal symptoms. The use of enteric-coated aspirin does not reduce these side effects.

Aspirin irreversibly acetylates and inhibits platelet cyclooxygenase (COX)-1, which causes reduced synthesis of thromboxane A2. At high doses, it also inhibits COX-2, which results in the anti-inflammatory effects of aspirin.

The major GI side effects include dyspepsia, erosive gastritis, peptic ulcers with bleeding and perforation. The overall risk of major bleeding with aspirin is 1–3% per year, which increases 2-3 fold if there is concurrent use of other antiplatelet drugs or anticoagulants. The risk of bleeding can be reduced by eradication of Helicobacter pylori infection and giving proton pump inhibitors (PPIs).

Aspirin sensitivity refers to bronchospasm or urticaria/angioedema seen immediately after administration of aspirin. It is also called Samter's triad (asthma, recurrent nasal polyps, aspirin sensitivity). In patients with aspirin resistance, there is failure of aspirin to produce its normal platelet inhibitory functions.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 24:

Dipyridamole increases extracellular adenosine levels.

It is an antiplatelet drug that increases levels of cyclic AMP (cAMP) in platelets by:

• Blocking the reuptake of adenosine and increasing extracellular adenosine levels

• Inhibiting phosphodiesterase-mediated cAMP degradation

This increase in cyclic AMP reduces intracellular levels of calcium. This inhibits platelet activation and aggregation.



Sold by @itachibot

Dipyridamole is a weak antiplatelet used in combination with aspirin for the prevention of stroke in patients with TIA.

Side effects include gastrointestinal and vasodilatory complaints (headache, flushing, dizziness, hypotension). Due to it's vasodilatory effects, it is contraindicated in patients with symptomatic coronary artery disease.

Solution to Question 25:

Abciximab is safe in patients with renal impairment. It belongs to the group of GpIIb-IIIa receptor antagonists. These are parenteral drugs that are used in the management of patients with acute coronary syndrome.

There are three important drugs in this group:

Immune-mediated thrombocytopenia and bleeding are the most common side effects. Thrombocytopenia can occur within 24 hours of initiation.

Feature	Abciximab	Eptifibatide	Tirofiban
Gp IIb/IIIa antagonist	Non-specific	Specific	Specific
Plasma half-life	Short (min)	Long (2.5 h)	Long (2.0 h)
Platelet-bound half-life	Long (days)	Short (s)	Short (s)
Renal clearance	No	Yes	Yes

Solution to Question 26:

Plasmapheresis is the mainstay of the treatment of patients with thrombotic thrombocytopenic purpura (TTP).

The best response to this therapy is seen in patients with ADAMTS13 autoantibodies. Plasmapheresis is continued until the resolution of platelet count and signs of hemolysis for at least 2 days.

Option A: The first line of treatment of autoimmune hemolytic anemia is glucocorticoids. The second line includes rituximab and splenectomy.

Option B: Treatment of disseminated intravascular coagulation is primarily supportive with measures to treat the underlying cause.

Option D: Treatment of HUS is primarily supportive. Dialysis is usually required. In some varients of HUS plasma infusion/exchange can be effective by replasing the complement-regulatory proteins.

Sold by @itachibot

Solution to Question 27:

Large vessel arterial or venous thromboemboli are not seen in thrombotic thrombocytopenic purpura (TTP). This condition is associated with microvascular thrombosis.

TTP occurs due to deficiency or autoantibodies against the metalloprotease ADAMTS13, which normally cleaves multimers of vWF. In this condition, the vWF molecules persist causing pathogenic platelet adhesion and aggregation. This causes localized platelet thrombi and fibrin deposits at various sites.

It presents with the following pentad of clinical features:

• Microangiopathic hemolytic anemia - mechanical shearing of RBCs as they pass through platelet rich-microthrombi

• Thrombocytopenia - deposition of platelets in microthrombi

• Renal failure - microthrombi in the renal vasculature

• Neurologic findings - headache, confusion, transient focal neurologic deficits

• Fever

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 28:

In patients with thrombotic thrombocytopenic purpura (TTP), the PT and aPTT are normal, as consumptive coagulopathy is not seen.

TTP and HUS are thrombotic microangiopathies with overlapping clinical features. TTP can be primary (ADAMTS13 deficiency) or secondary (associated with infections like HIV and drugs like quinine).

Note: In disseminated intravascular coagulation (DIC), thrombocytopenia and microangiopathy are seen. It is associated with consumption of clotting factors and fibrinogen resulting in prolonged PT and aPTT.

Solution to Question 29:

Factor V Leiden mutation is the most common inherited cause of venous thrombosis. Venous thrombosis typically manifests as deep vein thrombosis and pulmonary thromboembolism.

Hypercoagulable states or thrombophilias can be inherited or acquired:

• Inherited causes in decreasing order of prevalence are as follows:

• Factor V Leiden mutation

• Prothrombin mutation

• Protein C deficiency

Sold by @itachibot

- Protein S deficiency
- Antithrombin III deficiency
- Acquired causes include:
- Surgery
- Malignancy
- Antiphospholipid syndrome
- Trauma
- Pregnancy
- Long-haul travel
- Obesity
- Oral contraceptives/hormone replacement
- Myeloproliferative disorders
- Polycythemia vera

Sold by @itachibot
If you have purchased this from someone else, you have been scammed.

Solution to Question 30:

Platelet storage pool disease (SPD) is an example of an autosomal dominant qualitative platelet disorder. It refers to a group of disorders characterized by a deficiency of platelet granules causing secretion defects.



Bernard Soulier syndrome, Glanzman thrombasthenia, and Chediak-Higashi syndrome are inherited as autosomal recessive traits.

Sold by @itachibot

Acute Leukemia

Question 1:

What is the most common type of leukemia in adults?

a) Acute myeloid leukemia

b) Adult acute lymphoblastic leukemia

c) Chronic myeloid leukemia

d) Chronic lymphoid leukemia

Question 2:

Which of the following is not a risk factor for acute myeloid leukemia?

a) Down's syndrome

b) Bloom syndrome

c) Fanconi anemia

d) Evan's syndrome

Question 3:

A 2-year-old girl was referred with recurrent bacterial infections. Parents report that their first child also had similar features and died at the age of 18 months from myeloid leukemia. Investigations revealed markedly reduced absolute neutrophil count. What is the likely diagnosis?

a) Bloom syndrome

b) Kostmann disease

c) Diamond-Blackfan syndrome

d) Chediak-Higashi syndrome

Question 4:

A 60-year-old man is being evaluated for suspected acute myeloid leukemia. He reports a fast-growing swelling in his mouth for the past 1 week. On examination, there is a greenish

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

gingival mass with a necrotic surface. This presentation is likely associated with which of the following?

a) AML M2

b) AML M3

c) AML M7

d) AML M5a

Question 5:

Patients suffering from which type of AML are more prone to develop disseminated intravascular coagulation?

a) Monocytic

b) Myelomonocytic

c) Monoblastic

d) Acute promyelocytic

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Question 6:

Oral cavity examination of a 56-year-old man with suspected leukemia revealed the following. What subtype of AML is he likely to be suffering from?

a) M7

b) M5

Sold by @itachibot

c) M3

d) M2

Question 7:

Which of the following would be seen on flow cytometry of a patient with acute megakaryoblastic leukemia?

a) CD113

b) CD 117

c) CD 61

d) CD 13

Question 8:

Which of the following is true about the WHO classification of acute myeloid leukemia?

a) The clinical features are not considered to define the subtype

b) The cutoff value of blast cells for diagnosis is 20%

c) It is only based on morphological features of myeloid precursors

d) Chromosomal translocations along with > 20% blasts are needed for diagnosis

Question 9:

A 50-year-old man is referred for evaluation of severe anemia, persistent fever, and recurrent epistaxis. Lab investigations are given below. What is the diagnosis?

a) Acute lymphoblastic leukemia

b) Acute myeloblastic leukemia

c) Preleukemia

d) Aleukemic leukemia

Question 10:

Patients with all of the following are classified as AML with recurrent cytogenetic abnormalities except:

a) t(16,16)

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) NPM1 mutation

c) CEBPA mutation

d) t(9,22)

Question 11:

Which of the following cytogenetic abnormalities is seen in patients with acute promyelocytic leukemia?

a) t(15;17)

b) t(16;16)

c) RUNX1-RUNX1T1

d) CBF-β mutation

Sold by @Itachibot
If yc ased, this from someone else,
e been scammed.

Question 12:

A 65-year-old woman presented with fatigue and fever for the past 2 months. Investigations revealed pancytopenia and the following peripheral smear findings. What is the most likely diagnosis?



a) AML

b) ALL

c) CML

d) CLL

Sold by @itachibot

Question 13:

Which of the following patients with AML has a better prognosis than the others?

a) A 48-year-old man newly diagnosed with AML with a family history of lymphoma

b) A 60-year-old woman who developed AML following chemotherapy for ovarian cancer

c) A 71-year-old man with a longstanding history of myelodysplastic syndrome

d) A 67-year-old woman evaluated with a 4-month history of pancytopenia prior to diagnosis

Question 14:

A pathologist is examining the peripheral smear of a patient with suspected leukemia. All of the following features are suggestive of a myeloblast except:

a) Auer rods

b) Prominent nucleoli

c) Phi bodies

d) Pelger-Huet anomaly

Question 15:

A patient presents to the ER with a sudden-onset fever and the following painful rash over her face and upper limbs. CBC reveals neutrophilia. This presentation is associated with all of the following conditions except:

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Acute myeloid leukemia

b) Crohn's disease

c) Systemic lupus erythematosus

d) Aplastic anemia

Question 16:

What is the most commonly used complete remission induction regimen in patients with AML?

a) All-trans-retinoic acid and arsenic trioxide

b) L- asparginase

c) Cytarabine with anthracyclines

d) Cytarabine with alkylating agents

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Question 17:

A 60-year-old man came with generalized bone pain, and easy bruisability for the past 6 months. His WBC count was 60,000 cu. mm and bone marrow aspiration revealed the following. Immunohistochemistry was positive for MPO and cytogenetic studies showed presence of t(15:17). What is the preferred treatment regimen?



a) Cytarabine

b) Arsenic trioxide

Sold by @itachibot

c) All transretinoic acid

d) Arsenic trioxide and all transretinoic acid

Question 18:

An 18-year-old girl with the diagnosis of acute promyelocytic leukemia was treated medically. She developed fever and tachypnea and a chest X-ray showed pulmonary infiltrates. What drug should she be given next?

a) Cytarabine

b) Dexamethasone

c) Doxorubicin

d) Methotrexate

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

A 47-year-old man with a diagnosis of acute myeloid leukemia with a blood type O negative blood group presents to the transplant clinic to discuss proceeding with an allogeneic stem cell transplant. Which of the following would be an optimal donor?

a) His identical twin brother

b) Umbilical cord transplant

c) His HLA identical 50-year-old brother who is otherwise healthy and is blood type O+

d) An HLA identical matched unrelated donor who is blood type O–

Question 20:

Individuals in which age group have the highest incidence of T-cell acute lymphoblastic lymphoma?

a) 20-25 years

b) 1-10 years

c) 10-19 years

d) 26-30 years

Question 21:

Sold by @itachibot

Children with Down's syndrome are at an increased risk of developing all of the following except:

a) Acute lymphoid leukemia

b) Acute myeloid leukemia

c) Transient myeloproliferative disorder

d) Myelodysplastic disorder

Question 22:

An 8-year-old boy was brought with fever, weight loss, and easy bruisability. Initial examination revealed severe pallor and significant lymphadenopathy. Peripheral smear revealed the following cells, which were also seen on bone marrow studies. Further investigations revealed CD10+ cells. What is the likely diagnosis?



a) T-cell ALL

b) Pro B-cell ALL

c) Pre B-cell ALL

d) Mature B-cell ALL

Question 23:

Which of the following indicate a good prognosis in children diagnosed with acute lymphoblastic leukemia?

Sold by @itachibot

a) 3, 4, 5

b) 1, 2, 3

c) 2, 4, 5

d) 1, 2, 4

Question 24:

A 3-year-old boy with Down's syndrome was brought with recurrent infections and bruising over his legs. Further workup was diagnostic of acute lymphoblastic leukemia and he was started on chemotherapy. Which of the following must be added on to prevent metastases to the central nervous system?

a) L-Asparginase

b) Prednisolone

c) Cytarabine

d) Methotrextate

Sold by @itachibot
If you purchased this from someone else you have been scammed.

Answer Key

Question No.	Correct Option
1	d
2	d
3	b
4	a
5	d
6	b
7	c
8	b
9	b
10	d
11	a
12	a
13	a
14	d
15	d

Sold by @itachibot

16	c
17	d
18	b
19	c
20	c
21	d
22	c
23	a
24	d

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Chronic lymphoid leukemia (CLL) is the most common type of leukemia in adults.

The common types of leukemia in various age groups is as follows:

• Acute myeloid leukemia (AML) - most common acute leukemia in adults

• Acute lymphoid leukemia (ALL) - most common cancer in childhood. The most common subtype of ALL is B-cell ALL

• Non-Hodgkin's lymphoma (NHL) - most common lymphoid malignancy in adults

Solution to Question 2:

Evan's syndrome does not predispose to acute myeloid leukemia (AML).

The following conditions predispose to AML:

Genetic conditions:

• Down's syndrome:

• Young children <4 years

• Typically of the acute megakaryocytic subtype and associated with GATA1 mutations

• Excellent clinical outcomes

• Defective DNA repair in conditions like ataxia telangiectasia, Fanconi anemia, and Bloom syndrome

Environmental factors:

• Drugs:

• Alkylating agents - cyclophosphamide, melphalan, dacarbazine, platinum compounds, after 4-6 years of exposure

Sold by @itachibot

• Topoisomerase II inhibitors - etoposide, after 1-3 years of exposure.

• Radiation - high dose radiation, therapeutic radiation in patients exposed to alkylating agents

• Chemicals - benzene, petroleum products, pesticide

Hematological disorders with increased risk of AML:

• Myeloproliferative disorders

• Paroxysmal nocturnal hemoglobinuria

• Aplastic anemia

• Myelodysplastic syndrome

• Multiple myeloma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

This clinical scenario of congenital neutropenia is suggestive of Kostmann syndrome. It is associated with an increased risk of acute myeloid leukemia (AML).

Kostmann syndrome occurs due to mutations in the G-CSF receptor and neutrophil elastase. This causes bone marrow failure specifically in the myeloid lineage. It presents in young children as recurrent bacterial infections (such as otitis media, cellulitis) and oral ulcers.

Option A: Bloom syndrome occurs due to defects in DNA repair. It is not associated with neutropenia but has a higher risk of AML.

Option C: Diamond-Blackfan syndrome refers to congenital erythroid aplasia. It is also associated with a higher risk of AML.

Option D: Chediak-Higashi syndrome is due to defective phagolysosome formation. It is not associated with neutropenia.

Solution to Question 4:

This clinical scenario is suggestive of a chloroma. It is commonly associated with the M2 subtype of acute myeloid leukemia, also called AML with maturation.

Chloroma is also called myeloid sarcoma, extramedullary myeloid tumor, or granulocytic sarcoma. It is an extramedullary tumor mass consisting of myeloid blasts, with or without maturation. It is called chloroma due to the peculiar green color, caused by myeloperoxidase. The sites affected include orbit (most common), skull, oral cavity, paranasal sinuses, sternum, ribs, vertebrae, and pelvis.

Solution to Question 5:

Sold by @itachibot

Disseminated intravascular coagulation (DIC) is commonly seen in patients with acute promyelocytic leukemia (APL).

Solution to Question 6:

The given image shows swollen and spongy gums, indicating gum infiltration. This is a characteristic feature of acute monocytic leukemia (AML-M5).

Infiltration of the gingivae, skin, soft tissues or meninges with leukemic blasts is characteristic of the monocytic subtypes and those with 11q23 chromosomal abnormalities.

Less commonly, infiltration is also seen in acute myelomonocytic leukemia (AML-M4).

Solution to Question 7:

On flow cytometry, the immunophenotype of acute megakaryoblastic leukemia is CD41 and/or CD61 positive (platelet-specific antigens).

Flow cytometry is useful in establishing the different subtypes of AML:

Phenotype	CD markers
Myeloblastic	CD11b,13,33,117
Promyelocytic(M3)	CD13,33
Myelomonocytic(M4)	CD11b,13,14,33
Monocytic(M5)	CD11b,11c,13,14,33
Erythroid(M6)	Glycophorin,spectrin
Megakaryoblastic(M7)	CD41,61
Basophilic	CD11,13,33,123
Mast cell	CD13,33,117

Solution to Question 8:

The cutoff value of blast cells for the diagnosis of AML is 20% according to the WHO classification. This classification is based on the clinical features, cytogenetic and molecular abnormalities in addition to morphology. This is in contrast to the previously used French-American-British (FAB) classification which was based on only morphology of myeloid precursors.

The major changes in the WHO classification in contrast to FAB classification are:

If you purchased this from someone else, you've been scammed.
Sold by @itachibot

• Blast cutoff for diagnosis of AML is 20% (30% in FAB classification)

• Specific chromosomal rearrangements such as t (8;21), inv(16), t(16;16) and t (15;17) can define AML even with < 20% blasts

• Immunophenotyping (flow cytometry) is used to differentiate AML from ALL and to identify some subtypes of AML

• Myeloid dysplasia with blasts between 20-30% can benefit from therapy for myelodysplastic syndrome (decitabine, 5-azacytidine, cedazuridine)

Solution to Question 9:

The given clinical scenario is suggestive of acute myeloblastic leukemia (AML).

According to the WHO criteria, the diagnosis of AML requires any one of the following:

• Blast cell count ≥20% either in the peripheral blood or bone marrow

• Presence of any one of the following cytogenetic abnormalities:

• t(15;17)

• t(8;21)

• Inversion 16 or t(16;16)

• t(9;22)

• NPM1 mutation (Nucleophosmin mutation)

• CEBPA mutation

• FLT3-ITD

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

The chromosomal translocation t(9,22) is called the Philadelphia chromosome. It is seen in patients with CML.

AML with recurrent genetic abnormalities includes the following chromosomal aberrations:

• t(15;17)

• t(8;21)

• Inversion 16 or t(16;16)

Two gene mutations have been added to the criteria of AML with recurrent genetic abnormalities in addition to chromosomal aberrations. Both gene mutations are associated with a good prognosis.

• NPM1 mutation: Nucleophosmin mutation

• CEBPA mutation

Sold by @itachibot

Solution to Question 11:

t(15,17) is associated with acute promyelocytic leukemia (APL).

Solution to Question 12:

The given clinical scenario and peripheral smear showing blast cells with peroxidase granules are suggestive of acute myeloid leukemia (AML). Peroxidase granules are dark blue-stained granules seen in the cytoplasm on peroxidase staining.

Myeloperoxidase cytochemical activity is negative in chronic myeloid leukemia (CML) and blast crisis.

Solution to Question 13:

Young age of onset is associated with a good prognosis in patients with AML. Family history does not affect the prognosis.

Poor prognosis is seen in the following cases:

• Advancing age (usually defined as > 60 years)

• Previous hematologic disorders - myeloproliferative neoplasms and myelodysplastic syndrome

• Longstanding cytopenias >3 months before diagnosis

• Therapy-related AML (chemotherapy for other cancers)

• Chromosomal aberrations with poor outcomes

Solution to Question 14:

Pelger-Huet anomaly is not a feature of myeloblasts. Auer rods and Phi bodies are considered pathognomonic of AML.

Myeloblasts have the following characteristics as shown in the following image:

• Large, uniform population of cells

• Prominent nucleoli (yellow arrow in the image)

• Primary cytoplasmic granules

• Auer rods - needle-shaped azurophilic intracytoplasmic inclusion bodies formed by fusion of lysosomes (black arrow)

• Phi-bodies - fusiform or spindle-shaped rods similar to Auer rods that require special stains for hydroperoxidases

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Pseudo-Chediak-Higashi syndrome - Giant lysosomes forming intracytoplasmic inclusion bodies



Pelger-Huet anomaly refers to neutrophils with bilobed nuclei that occur as a benign, inherited condition. Pseudo-Pelger-Huet anomaly refers to an acquired defect seen in CML, myelodysplastic syndrome.

Note: Chediak-Higashi syndrome is a rare autosomal recessive disorder with abnormal lysosomal trafficking (LYST gene). Presents with oculocutaneous albinism and immunodeficiency.

If you purchased this from so... else, you may have been scammed. Sold by @ita...

Solution to Question 15:

This clinical presentation with painful edematous plaques is suggestive of febrile neutrophilic dermatosis or Sweet syndrome. It is not associated with aplastic anemia. It is associated with the following conditions:

• Acute myeloid leukemia (AML) - most common

• Inflammatory bowel disease (IBD)

• Solid malignancies of the genitourinary tract

• Systemic lupus erythematosus (SLE)

• Drugs - all-trans-retinoic acid, granulocyte colony-stimulating factor (G-CSF)

• Idiopathic Sweet syndrome - common in women following respiratory tract infection

This condition presents with the following triad:

• Fever

• Neutrophilia

• Dense dermal infiltrate of neutrophils:

• Pinkish-red to reddish-brown edematous painful plaques

• Occur over the head, neck, extremities

Sold by @itachibot

Solution to Question 16:

The most commonly used complete remission induction regimen in patients with AML is cytarabine and an anthracycline (e.g., daunorubicin, idarubicin).

The treatment of AML is as follows:

• Induction chemotherapy with a 7+3 regimen:

• Cytarabine - continuous intravenous infusion for 7 days

• IV daunorubicin on days 1,2,3

• Side effects include neutropenia, sepsis

• Post-remission therapy:

• High dose cytarabine

• Allogenic hematopoeitic stem cell transplant

Novel agents include gemtuzumab ozogamicin, a CD33 monoclonal antibody.

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Solution to Question 17:

The presence of blast cells with numerous Auer rods, MPO (myeloperoxidase) positive cells, and t(15:17) is diagnostic of acute promyelocytic leukemia (APL). The preferred treatment regimen is arsenic trioxide and all-trans-retinoic acid (ATRA).



APL is characterized by t(15; 17) translocation resulting in the PML-RAR-alpha fusion gene (promyelocyte-retinoic acid receptor alpha). It is a highly curable subtype of leukemia. The treatment is as follows:

• Induction therapy:

Sold by @itachibot

• ATRA + anthracycline (daunorubicin or idarubicin) or cytarabine (OR)

• ATRA + arsenic trioxide

• Consolidation therapy:

• Post remission anthracycline-based chemotherapy

Solution to Question 18:

The clinical scenario describes a patient of APL (acute promyelocytic leukemia) developing fever, tachypnea, and pulmonary infiltrates while on medical treatment. This is suggestive of APL differentiation syndrome, an adverse effect of all-trans-retinoic acid (ATRA). The management involves glucocorticoids like dexamethasone.

Along with glucocorticoids, the other management of APL differentiation syndrome includes chemotherapy for cytoreduction and supportive measures. In severe cases, temporary discontinuation of ATRA is necessary.

ATRA is an oral drug that induces the differentiation of leukemic cells bearing the t(15;17) chromosomal translocation. While ATRA reduces the incidence of DIC associated with cytarabine and daunorubicin used earlier in APL, it can cause APL differentiation syndrome due to the adhesion of differentiated neoplastic cells to the pulmonary vasculature endothelium.

In low-risk APL patients with a low leukocyte count at presentation, a combination of ATRA and arsenic trioxide (ATO) has been found to be superior and is the current standard of care.

Solution to Question 19:

In the case of acute myeloid leukemia (AML), allogeneic stem cell transplants are the treatment of choice. His HLA identical 50-year-old brother who is blood type O positive would be the best choice.

Other options:

Option A: An identical twin brother would not be the donor of choice. The therapeutic benefit of stem cell transplant in hematological malignancies is due to the graft-versus-leukemia (GvL). Donor T lymphocytes attack malignant hematopoietic stem cells of the recipient that remain after high-dose chemotherapy. Despite having a low rate of graft versus host disease (GvHD), stem cells from an identical twin do not have a desirable degree of GvL effect, leading to higher rates of recurrence post-transplant.

Option B: Autologous (umbilical cord transplant) is not recommended as higher rates of relapse have been reported.

Option D: An HLA identical sibling is preferred over an unrelated HLA identical donor because there is a greater chance of matching for minor histocompatibility complex (mHC). This reduces the risk of GvHD. ABO compatibility is also important to consider, but a minor ABO mismatch (Rh mismatch) doesn't have any impact on hemolysis, graft-versus-host disease, or survival.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Solution to Question 20:

The highest incidence of T-ALL (T-cell acute lymphoblastic lymphoma) is at 10-19 years (adolescence) when the thymus reaches the maximum size.

On the other hand, B-ALL which is more common has a peak incidence at 3 years (1-10 years). At this age, the number of normal bone marrow pre-B cells (the cell of origin) is at its greatest.

Solution to Question 21:

Down's syndrome is not associated with myelodysplastic disorder. These patients are at increased risk of developing the following conditions:

- Acute lymphoid leukemia (ALL) - most commonly B-cell ALL
- Acute myeloid leukemia (AML) - most commonly M7/megakaryoblastic subtype
- Transient myeloproliferative disorder
- Associated with GATA1 mutations.
- Seen in 10% of neonates, resolves in first 3 months of life
- Require close follow-up as 20-30% may develop AML by 3 years of life

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 22:

This clinical scenario and image showing blast cells in a young child are suggestive of acute lymphoblastic leukemia (ALL). The presence of CD10 is diagnostic of pre-B-cell ALL.

CD10 is also called the common ALL antigen (CALLA). It is the specific immunohistochemical marker for pre-B cell ALL. This is the most common type of ALL and is associated with a good prognosis. However, CD10 is negative in infantile leukemia, which is associated with translocation (4;11).

The immunohistochemistry of ALL is as follows:

Other abnormalities in ALL include:

Type	CD marker
pro-B cell	CD34,CD19, CD79a
pre- B cell	CD19, CD79a,CD10, Tdt, Cyt oplasmic Ig
Mature B c ell	CD19, CD79a,CD20, CD21,C D22, Surface IgM and D
T cell	CD 2,3,4,5,6,7

Sold by @itachibot

	Good prognosis	Bad prognosis
Cytogenic	t(12,21)	t(9,22), t(4,11)Burkitt lymphoma-associated ALL / ALL-L3 - t(2,8) , t(8,14), t(8,22)
Molecular	HyperdiploidyHOXNOTCH 1 mutations	Hypodiploidy

Solution to Question 23:

In children with ALL, a good prognosis is indicated by low WBC count, female sex, and hyperdiploidy.

Other abnormalities in ALL include:

	Good prognosis	Bad prognosis
Cytogenic	t(12,21)	t(9,22), t(4,11)Burkitt lymphoma-associated ALL / ALL-L3 - t(2,8) , t(8,14), t(8,22)
Molecular	HyperdiploidyNOTCH 1HOX	Hypodiploidy

Solution to Question 24:

Intrathecal methotrexate is used for CNS prophylaxis in acute lymphoblastic leukemia (ALL). It is essential in this condition as nearly 1/3 of patients develop CNS involvement at relapse without prophylactic therapy.

The treatment of ALL is as follows:

Induction (4 weeks):

• L-Asparaginase:

• Single-dose

• Side effects - pancreatitis, thrombosis

• Vincristine:

• Weekly

• Side effects - neuropathy, SIADH

• Prednisolone

Consolidation: Intrathecal methotrexate

Maintenance (24-36 months):

If you purchased this from @itachibot someone else, you may have been scammed.

Sold by @itachibot

- 6-Mercaptopurine
- Methotrexate

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Plasma Cell Disorders

Question 1:

What is the most common symptom seen in patients with multiple myeloma?

a) Easy fatigability

b) Bone pain

c) Oliguria

d) Fever

Question 2:

Which of the following is true about multiple myeloma?

a) Increased monoclonal IgG1 causes hyperviscosity syndrome

b) Monoclonal gammopathy of undetermined significance mostly progresses to multiple myeloma

c) There is decreased anion gap

d) Radioisotope bone scanning can be used for diagnosis

Question 3:

A 72-year-old man presented with complaints of fatiguability and chronic lower back pain. Laboratory findings are as follows. What is the most likely diagnosis?

a) Multiple myeloma

b) Monoclonal gammopathy of unknown significance

c) Smoldering multiple myeloma

d) Non-secretory myeloma

Question 4:

What is the most common cause of renal failure in patients with multiple myeloma?

a) Light chain deposition

b) Amyloidosis

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

c) Recurrent infections

d) Hypercalcemia

Question 5:

A patient was brought to the hospital with dehydration and generalised weakness. He was diagnosed with multiple myeloma 2 years ago. Investigations revealed metabolic acidosis and hypokalemia. Urine analysis showed a pH of 5.1, glycosuria, and proteinuria. What is the most likely diagnosis?

a) Renal tubular acidosis type 1

b) Adult Fanconi's syndrome

c) Renal tubular acidosis type 4

d) Alport syndrome

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

What is the immunophenotype of tumor cells in multiple myeloma?

a) CD19

b) CD38

c) CD 20

d) CD 5

Question 7:

A 45-year-old patient presented with multiple fractures and bone pain. Serum electrophoresis revealed elevated serum M protein. What does it most likely to constitute?

a) Ig M

b) Ig G

c) Albumin

d) Lamda light chains

Question 8:

Sold by @itachibot

A 60-year-old man presented with fatigue and back pain since 3 months. Investigations were done and are as follows. What is the most likely diagnosis?

a) Multiple myeloma

b) Lymphoplasmacytic lymphoma

c) Monoclonal gammopathy of undetermined significance

d) Non-secretory myeloma

Question 9:

A 69-year-old business tycoon is diagnosed with multiple myeloma. He is worried about his business empire and asks you how long he will survive. Which of the following is the best predictor for this?

a) Durie-Salmon staging system

b) Percentage of circulating plasma cells

c) Beta-2 microglobulin levels

d) Plasma cell labeling index

Question 10:

In patients with multiple myeloma, the presence of which of the following is associated with a poor prognosis?

a) t (11,14)

Sold by @itachibot
If you purchased this from
you may have been scamm

Sold by @itachibot

b) t (6,14)

c) 17p deletion

d) 13q deletion

Question 11:

An X-ray skull of a patient with marked hypercalcemia is shown below. Which of the following laboratory parameters is consistent with his diagnosis?



a) High alkaline phosphatase and normal ESR

b) High alkaline phosphatase and elevated ESR

c) Normal alkaline phosphatase and normal ESR

d) Normal alkaline phosphatase and elevated ESR

Question 12:

A 68-year-old male patient presented with multiple fractures. He has been suffering from recurrent infections for the past 6 months. Bone marrow biopsy showed the following. What is the treatment of choice in this condition?

Sold by @itachibot



a) Lenalidomide + bortezomib + dexamethasone

b) Thalidomide + cyclophosphamide

c) Rituximab + cyclophosphamide

d) Bortezomib + cyclophosphamide + dexamethasone

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 13:

Which of the following is a proteasome inhibitor?

a) Lenalidomide

b) Rituximab

c) Cyclophosphamide

d) Bortezomib

Question 14:

A 64-year-old male patient is admitted with complaints of weakness of the extremities, blurry vision, and nose-bleeds. On examination, there is lymphadenopathy and dilatation of the retinal veins. Peripheral smear is given below. Which of the following is false regarding this condition?

Sold by @itachibot



a) Bone marrow >10% lymphoplasmacytoid cells

b) Lytic bone lesions

c) M component in serum >30g/L

d) Increased viscosity of blood

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 15:

A 19-year-old boy presented with progressive shortness of breath, and easy fatiguability for 3 years. His older brother died of cardiac arrest during a football match a few years ago. Further evaluation revealed hypertrophic cardiomyopathy. Biopsy and staining of subcutaneous fat showed the following. What mutation is likely in this patient?



Sold by @itachibot

a) Leukocyte chemotactic factor

b) Transthyretin

c) Light chain

d) Beta 2 microglobulin

Question 16:

A 60-year-old man presented with progressive weight loss, generalized edema. On evaluation, he had nephrotic range proteinuria, macroglossia, and raccoon eyes. What is the most probable diagnosis?

a) Solitary plasmacytoma

b) Waldenstrom's macroglobulinemia

c) Amyloidosis

d) IgA nephropathy

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 17:

Which of the following is the most commonly involved organ in amyloidosis?

a) Heart

b) Kidneys

c) Liver

d) Spleen

Question 18:

What is the leading cause of death in patients with primary amyloidosis?

a) Renal failure

b) Cardiac disease

c) Liver failure

d) Recurrent infections

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	b
2	c
3	a
4	d
5	b
6	b
7	b
8	d
9	c
10	c
11	d
12	a
13	d
14	b
15	b
16	c
17	b
18	b

if you purcha... you may have... else, ...d.

Detailed Explanations

Solution to Question 1:

Bone pain is the most common symptom in multiple myeloma.

Multiple myeloma is a plasma cell malignancy in which there is an increased proliferation of monoclonal plasma cells. There is an increased production of monoclonal paraprotein (M protein).

Clinical features of multiple myeloma:

• Bone pain (most common):

• Due to proliferation of tumor cells, activation of osteoclasts that destroy bone, and suppression of osteoblasts that form new bone.

• Persistent localized pain in a patient with myeloma usually signifies a pathological fracture.

• Fever due to recurrent bacterial infections (second most common)

• Acute kidney injury due to:

• Hypercalcemia (most common)

Sold by @itachibot

• Amyloid deposits

• Hyperuricemia

• Recurrent infections

• Infiltration of the kidney by myeloma cells

• Normocytic-normochromic anemia due to:

• Replacement of normal marrow by expanding tumor cells

• Inhibition of hematopoiesis by factors made by the tumor

• Reduced production of erythropoietin by the kidney

Myeloma defining events (MDE) include the presence of one or more features such as hypercalcemia, renal failure, anemia and lytic bone lesions.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

There is decreased anion gap in multiple myeloma due to the M protein.

Anion gap in the serum is calculated as Na■ – (Cl– +HCO3–).

The M protein is cationic, resulting in the retention of chloride causing decreased anion gap.

Only 1% of patients with monoclonal gammopathy of undetermined significance (MGUS) go on to develop multiple myeloma. Hyperviscosity symptoms in multiple myeloma occurs due to excessive production of M proteins, usually of IgG3 or IgA subtype. Waldenstrom macroglobulinemia (IgM myeloma) is the most common cause for hyperviscosity syndrome due to excess IgM paraprotein in these cases.

Radioisotopic bone scanning is less useful in diagnosis because the bone lesions are lytic in nature and are rarely associated with osteoblastic new bone formation.

Solution to Question 3:

In this scenario, the patient has increased M protein (>30g/l), bone marrow clonal plasma cells >10%, and evidence of end-organ damage (fatiguability and back pain). The most likely diagnosis is multiple myeloma.

Diagnostic criteria for plasma cell disorders:

Note: Solitary plasmacytoma of bone has only a single area of bone destruction due to clonal plasma cells, no M protein, bone marrow appears normal, and no end-organ damage.

Plasma cell leukemia is an aggressive plasma cell proliferative disorder with >20% abnormal plasma cells in the peripheral blood.

Sold by @itachibot

Criteria	Monoclonal gammopathy of undetermined significance (MGUS)	Smoldering multiple myeloma (asymptomatic myeloma)	Multiple myeloma	Non-secretory myeloma
M protein (>30g/l)	No	Yes	Yes	No
Bone marrow plasma cells (>10%)	No	Yes	Yes	Yes
End organ damage(CRAB) S.Calcium >11 mg/dlRenal, S. Creatinine >2 mg/dlAnemia, Hemoglobin <10g/dlLyticbone lesions	No	No	Yes	Yes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

Hypercalcemia is the most common cause of renal failure in multiple myeloma.

The second most common cause of renal failure in multiple myeloma is light chain cast nephropathy. The most common type of myeloma causing renal failure is light chain myeloma.

Mechanisms involved in acute kidney injury are:

- Hypercalcemia
- Amyloid deposits
- Hyperuricemia
- Recurrent infections
- Infiltration of the kidney by myeloma cells
- Tubular damage by excretion of light chains (Fanconi's syndrome)

Solution to Question 5:

In this patient with hypokalemia and urine pH of 5.1, type 2 renal tubular acidosis (adult Fanconi's syndrome) is the likely diagnosis. It is commonly associated with multiple myeloma.

Normally, light chains are filtered, reabsorbed in the proximal tubules, and catabolized. In multiple myeloma, there is an increased amount of light chains and the tubular cells are overloaded.

This leads to proximal tubular damage either directly from light chain toxic effects or indirectly from the release of intracellular lysosomal enzymes.

Clinical features of renal tubular acidosis type 2:

- Metabolic acidosis
- Urine pH <5.3
- Proteinuria (only light chains)

Sold by @itachibot

• Glycosuria

• Aminoaciduria

Alport's syndrome is characterised by hematuria, hearing loss and eye abnormalities.

Solution to Question 6:

The immunophenotype of tumor cells in multiple myeloma is CD38 and CD138.

Multiple myeloma is derived from plasma cells in which B cell markers (CD19, CD20) are lost.

Immunophenotype of tumor cells in lymphoplasmacytic lymphoma / Waldenstrom's macroglobulinemia is CD19, CD20, surface IgM+, CD38. A lymphoplasmacytic cell retains B cell markers and carries plasma cell markers.

Solution to Question 7:

In the above clinical scenario, the patient most likely has multiple myeloma as there are pathological fractures and elevated M protein. Serum M protein is likely to constitute IgG.

The serum M component is IgG in 53% of patients, IgA in 25%, and IgD in 1%.

Serum IgM is elevated in Waldenstrom's macroglobulinemia (lymphoplasmacytic lymphoma).

Solution to Question 8:

In the given clinical scenario, the patient has normal SPEP and clonal plasmacytosis. The most likely diagnosis is non-secretory myeloma.

Non-secretory myeloma is characterised by the absence of M protein in the serum or urine. Bone marrow clonal plasmacytosis > 10% and myeloma defining symptoms.

Option A: In multiple myeloma, SPEP demonstrates a sharp peak (church spire or M band) in the gamma globulin region, as shown below.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Option B: Lymphoplasmacytic lymphoma (Waldenstrom's macroglobulinemia) also demonstrates a localized peak (M band) in the gamma globulin region. Immunofixation is required to identify the type of immunoglobulin.

Option C: Monoclonal gammopathy of undetermined significance (MGUS) does not present with myeloma-defining symptoms. Clonal bone marrow cells will be less than 10%

Solution to Question 9:

Serum β2-microglobulin levels are the best modality to predict the median survival period.

Albumin levels alone cannot be used to predict the median survival period. It is used in combination with β2-microglobulin for risk stratification.

Patients with β2-microglobulin levels <3.5 mg/L and albumin levels > 3.5 g/dl have a median survival period of 5 years. Those with levels >3.5 mg/L and albumin < 3.5 g/dl have a survival period of fewer than 2 years.

Durie-Salmon staging is unable to predict outcomes with newer agents. Hence it is no longer used.

Factors associated with poor prognosis:

- Plasma cell labeling index
- Chromosomal aberrations
- Circulating plasma cells
- Performance status
- Highly elevated LDH
- Extramedullary disease
- De novo plasma cell leukemia

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 10:

17p deletion is associated with a poor prognosis in multiple myeloma.

Chromosomal abnormalities in multiple myeloma are classified as follows:

Standard risk (6-7 years survival):

• t(11;14) (most common)

• t(6;14)

• 13q deletion

High risk (2-3 years survival):

• 17p deletion

• t(4;14)

• t(14;16)

• t(14;20)

Solution to Question 11:

In the above clinical scenario, hypercalcemia along with the skull demonstrating typical 'punched out' lesions is characteristic of multiple myeloma. Normal alkaline phosphatase and elevated ESR are consistent with multiple myeloma.

Multiple myeloma is a malignant plasma cell dyscrasia with the proliferation of plasma cells derived from a single clone. It is associated with an increased osteoclast activity with suppression of osteoblast activity, causing punched-out lytic lesions in the bone and hypercalcemia.

Signs and symptoms:

• Bone pain is the most common symptom

• Pathologic fractures

• Recurrent bacterial infections

• Renal failure

• Normocytic normochromic anemia, elevated ESR, hypercalcemia, normal ALP (ALP is a marker of bone growth and hence is not increased in multiple myeloma)

• Radiography: Punched-out lytic lesions on X-ray. The lesions are purely osteolytic with little or no osteoblastic activity. Hence alkaline phosphatase (ALP) is normal. The most common site of lytic lesion in multiple myeloma is the vertebral column.

• Electrophoresis: M spikes

• Urinalysis: Bence Jones proteinuria

• Bone marrow biopsy: Plasma cells

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Solution to Question 12:

In the above clinical scenario, the symptoms and bone marrow biopsy are characteristic of multiple myeloma. Lenalidomide + bortezomib + dexamethasone is the treatment of choice in this condition.

This combination achieves close to a 100% response. Hence it is one of the preferred induction regimens in transplant-eligible multiple myeloma patients.

Other pharmacological agents are used as second-line and have a 90% response rate. Chemotherapy followed by autologous stem cell transplantation prolongs overall survival.

The image below shows a bone marrow biopsy with an increased number of plasma cells.



Solution to Question 13:

Bortezomib is a proteasome inhibitor.

It has a rapid response and is effective in poor-prognosis myeloma. Side effects include neuropathy (both peripheral and autonomic). Subcutaneous and weekly administration can reduce neuropathic side effects.

Other proteasome inhibitors include carfilzomib (second-generation) and ixazomib, an oral proteasome inhibitor, used in relapsed disease.

Option A: Lenalidomide is an immunomodulatory analogue of thalidomide.

Option B: Rituximab is an anti-CD-20 monoclonal antibody.

Option C: Cyclophosphamide is an alkylating chemotherapeutic agent.

Solution to Question 14:

Sold by @itachibot

In the above clinical scenario, the hyperviscosity symptoms and rouleaux formation on peripherals smear point to a diagnosis of Waldenstrom's macroglobulinemia. It does not cause lytic bone lesions.

Waldenstrom's macroglöbulinemia is a B-cell malignancy affecting lymphoplasmacytoid and plasma cells. It is characterized by high levels of circulating IgM antibodies.

Clinical features include:

• Hyperviscosity symptoms

• Recurrent infections

• Epistaxis

• Visual disturbances

• Peripheral neuropathy - due to myelin-associated glycoprotein causing demyelination.

• Lymphadenopathy

• Hepatosplenomegaly

• Ophthalmoscopic examination may reveal vascular segmentation and dilation of the retinal veins.

Investigations:

• Bone marrow shows >10% infiltration with lymphoplasmacytic cells

• Serum M component is >30 g/L (3 g/dL). The size of the IgM paraprotein results in little renal excretion, and only 20% of patients excrete light chains.

The image below shows rouleaux formation.



Sold by @itachibot
If you have purchased this from someone else, you have been scammed.

Solution to Question 15:

Sold by @itachibot

The biopsy showing amyloid deposits stained by congo red, along with family history points to a diagnosis of familial amyloidosis (AF). It is most commonly due to the mutation of the transthyretin gene (TTR).

Amyloidosis is a group of protein misfolding disorders characterized by the accumulation of abnormal proteins called amyloid fibrils in various organs.

Solution to Question 16:

The patient has non-specific symptoms, which along with macroglossia and raccoon eyes are suggestive of amyloidosis.

Clinical features include:

• Fatigue

• Weight loss

• Nephrotic range proteinuria due to renal amyloidosis and is associated with hypoalbuminemia

• Cutaneous ecchymoses around the eyes are called the 'raccoon-eye' sign. Due to easy bruising as a result of amyloid deposits or deficiency of clotting factor X.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 17:

The kidneys are the most frequently involved organ in amyloidosis and are affected in 70–80% of patients.

The heart is the second most commonly affected organ (50–60% of patients).

The spleen is frequently involved, and there may be functional hyposplenism in the absence of significant splenomegaly.

Liver involvement causes cholestasis.

Macroglossia, a pathognomonic sign of AL amyloidosis is seen in only 10% of patients.

Solution to Question 18:

Cardiac disease is the leading cause of death from primary amyloidosis.

Sold by @itachibot

Chronic Myeloid Leukemia and Chronic Lymphoid Leukemia

Question 1:

Which of the following patients are at risk of developing chronic myeloid leukemia?

a) A middle-aged man whose mother died of chronic myeloid leukemia

b) An oil-refinery worker with occupational benzene exposure for 20 years

c) A woman who underwent radiotherapy for carcinoma cervix

d) An elderly man who received chemotherapy for Hodgkin's lymphoma

Question 2:

Which of the following is true about the Philadelphia chromosome?

a) Unbalanced translocation between short arm of chromosomes 9 and 22

b) Unbalanced translocation between long arm of chromosomes 9 and 22

c) Balanced translocation between short arm of chromosomes 9 and 22

d) Balanced translocation between long arm of chromosomes 9 and 22

Question 3:

What is the most common sign in patients with chronic myeloid leukemia?

a) Lymphadenopathy

b) Hepatomegaly

c) Splenomegaly

d) Weight loss

Question 4:

Leukocyte alkaline phosphatase score is decreased in patients with all of the following except:

a) Chronic myeloid leukemia

b) Paroxysmal nocturnal hemoglobinuria

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Sickle cell anemia

d) Polycythemia vera

Question 5:

In patients with chronic myeloid leukemia, which of the following are predominantly elevated in the peripheral smear?

a) Segmented neutrophils

b) Metamyelocytes

c) Myelocytes

d) Band forms

Question 6:

A 55-year-old man is found to have the following findings on a routine CBC. A subsequent peripheral smear is performed as shown below. A bone marrow aspiration study is planned for this patient. All of the following are likely to be seen on bone marrow except:



a) Hypercellular bone marrow

b) Myeloid: Erythroid ratio of 15:1

c) Marrow blasts >20 %

d) Sea-blue histiocytes

If you bought this from someone else, it has been scammed. Sold by @itachibot

Sold by @itachibot

Question 7:

A 48-year-old man presents to the clinic with easy fatiguability and weight loss. After a detailed workup, he is diagnosed with chronic myeloid leukemia. What is the initial treatment of choice?

a) Allogeneic stem cell transplantation

b) Monoclonal antibody against C5

c) Tyrosine kinase inhibitor

d) CD20 inhibitor

Question 8:

A 61-year-old with CML failed to respond to imatinib therapy. Further evaluation revealed a T315I mutation. What drug should he be started on?

a) Bosutinib

b) Nilotinib

c) Omacetaxine

d) Ponatinib

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 9:

Which of the following can be used to assess the prognosis in patients with chronic myeloid leukemia?

a) Rai and Binet staging system

b) Ann Arbor staging system

c) Hasford scoring system

d) Matutes scoring system

Question 10:

All of the following patients with chronic myeloid leukemia are in the accelerated phase except:

a) Patient 1

b) Patient 2

Sold by @itachibot

c) Patient 3

d) Patient 4

Question 11:

What is the classical immunophenotypic picture seen in patients with chronic lymphocytic leukemia?

a) CD19+, CD20+, CD21+, CD22+, CD 23+, CD5 +

b) CD19+, CD20+, CD21+, CD22+, CD23-, CD5+

c) CD79b+, CD19+, CD10+, CD20+, CD22+

d) CD79b+, CD19+, CD20+, CD22+, CD5-, CD10-

Sold by @itachibot

Question 12:

A 68-year-old man with fever is found to have a WBC count of 1,50,000/cu.mm. His peripheral smear study is given below. Which of the following is true about this condition?



a) It arises from the myeloid lineage

b) Radiation exposure is a risk factor

c) It arises from the T lymphocytes

d) Patient may have autoimmune hemolytic anemia

Question 13:

Sold by @itachibot

What is the treatment of choice for a 71-year-old asymptomatic woman diagnosed with chronic lymphocytic leukemia?

a) Fludarabine only

b) Fludarabine + Chlorambucil + Rituximab

c) Chlorambucil + Bendamustine

d) Expectant management

Question 14:

Which of the following is true about Richter syndrome?

a) Lymphadenopathy is absent

b) Transformation to an aggressive lymphoma

c) Decreased serum lactate dehydrogenase

d) Has a good prognosis

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	d
3	c
4	d
5	c
6	c
7	c
8	d
9	c
10	a
11	a
12	d
13	d
14	b

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Exposure to a high dose of ionizing radiation is the only known risk factor for developing chronic myeloid leukemia (CML). This may be in the form of radiation therapy for malignancy or other conditions and nuclear accidents. It is dose-dependent and low-dose radiation exposure is not a risk factor.

There are no familial associations, infections, or toxins associated with the development of CML.

Solution to Question 2:

Philadelphia chromosome is a balanced translocation between the long arm of chromosomes 9 and 22. It is present in the majority of patients with chronic myeloid leukemia (CML).

In a balanced translocation, a part of one chromosome is shifted to another chromosome and vice versa. In this case, the ABL gene present in chromosome 9 is translocated next to the BCR gene (breakpoint cluster region) on chromosome 22. It leads to the formation of shortened chromosome 22, which has been named the Philadelphia chromosome after the place of discovery.



Solution to Question 3:

Splenomegaly is the most common physical examination finding in patients with chronic myeloid leukemia (CML).

The majority of the patients with CML are asymptomatic. They are often incidentally diagnosed during routine blood tests when they have elevated white blood cells count. In later stages, the following clinical features develop:

Sold by @itachibot

• Anemia - easy fatigability, exercise intolerance

• Increased number of WBCs and platelets - hyperviscosity, leading to MI and stroke

• Increased cell metabolism - night sweats, weight loss, and hyperuricemia (gout)

• Splenomegaly - abdominal pain, early satiety

• Hepatomegaly, lymphadenopathy, skin or subcutaneous lesions

Solution to Question 4:

Leukocyte alkaline phosphatase (LAP) score is increased in polycythemia vera and leukemoid reactions.

The biochemical abnormalities in CML include:

• Low LAP score

• High levels of:

• Vitamin B12

• Uric acid

• Lactic dehydrogenase

• Lysozyme

Note: Leukemoid reaction refers to extreme elevations in WBC > 50,000/mm3 due to reactive causes outside the marrow. This includes severe infection, trauma, Down's syndrome.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

Myelocytes will be predominantly elevated in the peripheral smear of the patients with chronic myeloid leukemia (CML).

The clinical hallmark of CML is the uncontrolled production of mature and maturing granulocytes. All of the stages of development of granulocytes can be noted on the peripheral smear in varying numbers:

• Myeloblast - large cells with a round or oval nucleus, cytoplasm is dark and has no granules

• Myelocyte - nucleus is oval and is eccentrically placed.

• Metamyelocyte - nucleus is like a kidney bean-shaped

• Band forms - nucleus is in the shape of C or U.

• Segmented neutrophils- nucleus has 2-5 lobes.

All the cells except the myeloblast have a granular cytoplasm. Due to the varied appearance of the cells, it is called garden party appearance or college girl's appearance.

Sold by @itachibot



Sold by @itachi... ...else, ...this from some... ...been scammed.

Solution to Question 6:

The image shows a left-shift with a predominance of segmented neutrophils, myelocytes, metamyelocytes that is suggestive of chronic myeloid leukemia (CML). Marrow blasts are usually < 5% in the majority of CML patients.



In CML, the following bone marrow findings are noted:

- Markedly hypercellularity - due to increased granulocytic precursors
- Myeloid: erythroid ratio (M/E ratio) is significantly increased
- Megakaryocytes - normal or increased
- Erythroid precursors - usually decreased

Sold by @itachibot

• Sea blue histiocytes - macrophages with abundant wrinkled, green-blue cytoplasm.

• Bone marrow fibrosis may occur due to the deposition of reticulin.

• Pseudo-Gaucher cells - macrophages with wrinkled tissue paper appearance, as shown below. Due to the increased turnover of the cells, there is an increased accumulation of glucocerebroside within the macrophages.



If you purchased this ... else, you may have been ...

Solution to Question 7:

The initial treatment of choice in patients with chronic myeloid leukemia (CML) is tyrosine kinase inhibitors. These include imatinib, nilotinib, dasatinib, bosutinib, and ponatinib.

Allogeneic stem cell transplant is considered in patients who are negative for Philadelphia chromosome, or who are non-responsive to the tyrosine kinase inhibitors.

Solution to Question 8:

Ponatinib is effective in patients with CML with T315I mutations. It is a third-generation tyrosine kinase inhibitor.

The T315I mutation in BCR-ABL clones is specifically resistant to therapy with imatinib, dasatinib, nilotinib, and bosutinib but is sensitive to the third-generation agent ponatinib. It is also used in the treatment of accelerated and blast phases.

Solution to Question 9:

Hasford scoring system can be used to assess prognosis in patients with chronic myeloid leukemia (CML).

Sold by @itachibot

The prognosis of CML can be assessed by:

• Sokal score which is based on:

• Age

• Spleen size

• Platelet count

• Percentage of blast cells in the blood

• Hasford score - Similar to Sokal score, but additionally uses the percentage of eosinophils and basophils in the blood

Option A: Rai and Binet staging system is used for staging chronic lymphocytic leukemia (CLL).

Option B: Ann Arbor staging system is used in Hodgkin and non-Hodgkin lymphomas.

Option D: Matutes scoring system is used for the diagnosis of CLL by flow cytometry.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

A blast count ≥ 20% in peripheral blood is indicative of a blast crisis, not the accelerated phase of chronic myeloid leukemia (CML).

CML can be divided into three phases:

• Chronic stable phase - Blasts < 10% in peripheral blood and bone marrow

• Accelerated phase - any of the following:

• Blasts comprising 10-19% of white blood cells (WBCs) in peripheral blood and/or bone marrow

• Peripheral blood basophils ≥20%

• Thrombocytopenia or thrombocytosis not responsive to therapy

• Additional cytogenetic abnormalities in Philadelphia chromosome cells

• Blast crisis:

• Blasts ≥ 20% of peripheral blood WBCs or bone marrow

• Extramedullary blast proliferation

Majority of the patients present in the chronic stable phase. Without effective treatment, CML in the chronic phase will eventually move into an accelerated phase at first and then into the blast phase.

Solution to Question 11:

The classical immunophenotypic feature seen in patients with chronic lymphocytic leukemia (CLL) is CD19+, CD20+, CD21+,CD22+, CD23+, CD5+. It is significant for coexpression of the B cell marker CD19 with the T cell marker CD5.

Mantle cell lymphoma has the same immunophenotype but with absent CD23.

Sold by @itachibot

B cell precursors	Derived lymphomas/leukemia	Immunophenotype
Pro-B cell	Pro-B cell ALL	CD34, CD19, CD79a
Pre-B cell	Pre-B cell ALL	CD19, CD79a, CD10, Tdt, Cytoplasmic Ig
Mature B cell	CLL/SLL	CD19, CD79a,CD 20, CD21, CD22, surface IgM, CD 23, CD 5
Mantle zone	Mantle zone lymphoma	CD19, CD79a,CD 20, CD21, CD22, surface IgM, CD5
Germinal center/Follicular center	Follicular lymphomaDiffuse large B cell lymphomaBurkitt lymphoma	CD 79b, CD 19, CD 20, CD22, CD10, CD 23, Bcl-2CD10, Cd 23, Bcl-6CD10, CD23,Bcl-6
Memory B cells	Extranodal marginal zone lymphoma	CD19, CD79a,CD 20, surface Ig

Sold by @itachibot
If you paid for this from someone else, you've been scammed.

Solution to Question 12:

The smear shows small round lymphocytes with scant cytoplasm and the presence of a smudge cell. The most probable diagnosis is chronic lymphocytic leukemia (CLL). Patients with CLL may have autoimmune hemolytic anemia (AIHA).



Chronic lymphocytic leukemia/ small lymphocytic lymphoma refers to the monoclonal proliferation of mature B lymphocytes. It is more common in men than women. Increasing age and family history are important risk factors. CLL is one of the few leukemias not linked to radiation exposure.

The majority of the patients are asymptomatic and are diagnosed during routine blood tests. Some patients may have painless lymphadenopathy and splenomegaly. Increased B cells may

Sold by @itachibot

produce autoantibodies leading to autoimmune complications like autoimmune hemolytic anemia and immune thrombocytopenia (ITP).

Solution to Question 13:

Expectant management is the treatment of choice for asymptomatic patients with chronic lymphocytic leukemia (CLL). This is because there is no clear survival benefit with chemotherapy in asymptomatic patients.

Chemotherapy and monoclonal antibodies are used in the presence of the following features:

- Progressive marrow failure - worsening anemia, thrombocytopenia
- Progressive splenomegaly
- Progressive lymphadenopathy
- Presence of constitutional symptoms like weight loss, fever, night sweats

Important drugs used in the management of CLL are fludarabine, cyclophosphamide, bendamustine, and rituximab.

Solution to Question 14:

Richter's syndrome is the transformation of chronic lymphocytic leukemia (CLL) to aggressive lymphoma, most commonly diffuse large B cell lymphoma (DLBCL). It has a poor prognosis.

Patients develop the following features:

- Fever, and weight loss
- Rapidly enlarging lymph nodes
- Elevated serum lactate dehydrogenase

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Lymphomas

Question 1:

What is the most common non-Hodgkin lymphoma in India?

a) Burkitt's lymphoma

b) Diffuse large B-cell lymphoma

c) Mantle cell lymphoma

d) Follicular cell lymphoma

Question 2:

Which of the following is not caused by the Epstein-Barr virus?

a) Hodgkin's lymphoma

b) Primary diffuse large B cell lymphoma of the CNS

c) Post–organ transplant lymphoma

d) Primary effusion lymphoma

Question 3:

Which of the following types of non-Hodgkin's lymphoma and its corresponding grade are correctly matched?

a) Small lymphocytic lymphoma - High grade

b) Diffuse large B cell lymphoma - Low grade

c) Follicular lymphoma - Intermediate grade

d) Burkitt's lymphoma - High grade

Question 4:

What is the most common extranodal site of involvement in patients with non-Hodgkin's lymphoma?

a) Central nervous system

b) Stomach

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Small intestine

d) Skin

Question 5:

Which of the following is used to assess the prognosis in patients with non-Hodgkin's lymphoma?

a) Ann Arbor staging

b) Rai and Binet score

c) International Prognostic Index

d) Hasford scoring

Question 6:

A 63-year-old man presented with a 2-month history of fatigue, fever, and night sweats. On examination, generalized painless lymphadenopathy and splenomegaly were noted. Investigations revealed the following. What is the likely diagnosis?

a) Small lymphocytic lymphoma

b) Follicular cell lymphoma

c) Burkitt's lymphoma

d) Mantle cell lymphoma

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 7:

Which of the following is incorrectly matched?

a) Option 1

b) Option 2

c) Option 3

d) Option 4

Question 8:

A 59-year-old came to the OPD after noticing a swelling on her neck. On examination, painless lymphadenopathy was noted in the cervical, axillary and inguinal regions with no

Sold by @itachibot

hepatosplenomegaly. CBC was normal. Lymph node biopsy showed effacement of nodal architecture with centrocytes and centroblasts. Which of the following immunophenotypic panels will confirm this diagnosis?

a) CD19, CD 23, CD5

b) CD19, CD22, CD5

c) CD19, CD79a, CD10, TdT

d) CD 19, CD79a, CD 20, CD10, CD 23, Bcl-2

Question 9:

Which of the following statements is false about diffuse large B-cell lymphoma?

a) It is a high-grade aggressive lymphoma

b) It is commonly associated with BCL-6 chromosomal rearrangements

c) It is associated with CNS lymphoma and intravascular lymphoma

d) It does not present with extranodal involvement

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 10:

A physician is in Africa, working with the Doctors Without Borders group. He encounters a 7-year-old boy with the following finding. His parents say that he had a jaw swelling for 2 months that failed to resolve with antibiotics. A biopsy of the swelling reveals a starry sky appearance. Which of the following statements is true about this patient's condition?

Sold by @itachibot

a) It is associated with n-myc dysregulation

b) Tumor cells show CD21 expression

c) Expectant management is the treatment of choice

d) It is associated with HIV infection

Question 11:

The lymph node biopsy of a patient with lymphoma is given below. It shows two different cell populations labeled as A and B. Which of the following statements is false?



a) Cells labelled A will test positive for CD14, CD40, CD11b & CD64

b) The fraction of cells with positive Ki67 staining is 100%

c) Cells labelled B will test positive for CD19, CD10, BCL-6 & CD43

d) Cells labelled B represent the stars in starry sky appearance

Question 12:

Patients with which of the following conditions are at risk of developing salivary gland MALToma?

a) Helicobacter pylori gastritis

b) Hashimoto's thyroiditis

c) Sjogren's syndrome

d) Celiac disease

Sold by @itachibot

Question 13:

Which of the following is not a T-cell lymphoma?

a) Hairy cell leukemia

b) Mycosis fungoides

c) Anaplastic large cell lymphoma

d) Null cell lymphoma

Question 14:

The peripheral smear of a patient who presented with generalized lymphadenopathy and hepatosplenomegaly is given below. What is the likely diagnosis?



a) Mantle cell lymphoma

b) Peripheral T cell lymphoma

c) Anaplastic large cell lymphoma

d) Adult T cell leukemia/lymphoma

Question 15:

A 55-year-old man presented with fatigue and dragging sensation in the left upper abdomen. Physical examination revealed massive splenomegaly and no lymphadenopathy. CBC revealed

Sold by @itachibot

pancytopenia. Bone marrow aspirate yielded a dry tap. Immunohistochemical staining of the tumor cells tested positive for CD19, CD22, CD11c, CD25, CD103 and CD123. What is the likely diagnosis?

a) Myelofibrosis

b) Splenic marginal zone lymphoma

c) Multiple myeloma

d) Hairy cell leukemia

Question 16:

Which of the following best distinguishes hairy cell leukemia from other B cell malignancies?

a) Hairy cells on a Wright-stained peripheral blood smear

b) Annexin A1 on bone marrow specimens

c) Positive tartarate-resistant acid phosphatase stain

d) Intracytoplasmic inclusion bodies called ribosome-lamellar complexes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

Patients with hairy-cell leukemia are at a higher risk of infection with which of the following?

a) Parvovirus B19

b) Mycoplasma pneumoniae

c) Mycobacterium avium intracellulare

d) Cryptococcus neoformans

Question 18:

A 45-year-old man presented with progressive fatigue and weight loss for 3 months. CBC showed pancytopenia. On peripheral smear, the following cells were seen, which also showed positive staining with TRAP. What is the drug of choice for this patient?

Sold by @itachibot



a) Vemurafenib

b) Pentostatin

c) Interferon α

d) Cladribine

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

What subtype of Hodgkin's lymphoma is most commonly seen in India?

a) Nodular sclerosing

b) Mixed cellularity classical

c) Lymphocyte-rich classical

d) Lymphocyte depleted classical

Question 20:

Which of the following is not a classical form of Hodgkin lymphoma?

a) Nodular sclerosing

b) Lymphocyte depleted

c) Nodular lymphocyte predominant

d) Lymphocyte-rich

Sold by @itachibot

Question 21:

What is the most commonly observed clinical presentation in patients with Hodgkin's lymphoma?

a) Supraclavicular lymphadenopathy

b) Axillary lymphadenopathy

c) Hepatosplenomegaly

d) Mediastinal mass

Question 22:

A 29-year-old man with a 3-week history of fever is found to have multiple rubbery, enlarged lymph nodes. A biopsy reveals abnormal cells with bilobed nuclei that are positive for CD15 and CD30. Which of the following is associated with this man's condition?

a) Syndrome of inappropriate ADH secretion

b) Lambert-Eaton syndrome

c) Subacute cerebellar degeneration

d) Superficial migratory thrombophlebitis

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Question 23:

The lymph node biopsy of a 49-year-old woman with painless lymphadenopathy revealed the following finding. What is the likely diagnosis?



Sold by @itachibot

a) Lymphocyte-depleted Hodgkin's lymphoma

b) Lymphocyte-rich Hodgkin's lymphoma

c) Nodular sclerosis Hodgkin's lymphoma

d) Mixed cellularity Hodgkin's lymphoma

Question 24:

Which of the following statements is true about Hodgkin's lymphoma?

a) Reed-Sternberg cell is specific for Hodgkin's disease

b) Classical Hodgkin's lymphoma is associated with CD20 positive cells

c) Central nervous system is the most common site of extranodal disease

d) PET-CT is the most reliable technique for staging Hodgkin's lymphoma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 25:

A 68-year-old man presented with fever, night sweats, and weight loss for 2 months. On examination, non-tender, rubbery lymph nodes were palpable in the supraclavicular and inguinal regions on both sides. Biopsy showed the presence of Reed-Sternberg cells. How would you stage this patient according to the Ann Arbor Staging?

a) Stage IIA

b) Stage IIB

c) Stage IIIA

d) Stage IIIB

Question 26:

A 49-year-old man is diagnosed with stage IIIB Hodgkin's lymphoma. Lab investigations reveal a hemoglobin of 10.8 g/dl, WBC count of 14,000/ µL, and an absolute lymphocyte count of 450/ µL. Which of the following are poor prognostic factors in this patient?

a) 2, 3, 5

b) 1, 2, 5

c) 3, 4, 5

d) 1, 2, 4

Sold by @itachibot

Question 27:

A 48-year-old man with fever, night sweats, and weight loss is diagnosed with stage IIB Hodgkin's lymphoma. What is the treatment of choice for this patient?

a) R- CHOP regimen

b) ABVD regimen

c) ABVD regimen + Field radiation

d) R-CHOP regimen + Field radiation

Question 28:

Which of the following lymphoid neoplasms is incorrectly matched with its corresponding etiology?

a) H. pylori infection - Extra nodal marginal cell lymphoma

b) Gluten-sensitive enteropathy - Intestinal T-cell lymphomas

c) Breast implants - Anaplastic large cell lymphoma

d) Human immunodeficiency virus - Lymphoplasmacytic lymphoma

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	b
2	d
3	d
4	b
5	c
6	d
7	d
8	d
9	d
10	b
11	d
12	c
13	a

Sold by @itachibot

14	d
15	d
16	b
17	c
18	d
19	b
20	c
21	a
22	c
23	c
24	d
25	d
26	b
27	c
28	d

Sold by @itachibot If you purchased this from elsewhere, you may have been scammed.

Detailed Explanations

Solution to Question 1:

Diffuse large B-cell lymphoma (DLBCL) is the most common form of non-Hodgkin lymphoma (NHL) in adults and children over the age of 15 years, worldwide, and in India.

Burkitt's lymphoma is the most common NHL in children under the age of 15 years.

Solution to Question 2:

Human herpesvirus 8 (HHV-8) is the etiological factor for primary effusion lymphoma, not Epstein-Barr virus.

Lymphoid malignancies associated with Epstein-Barr virus include:

• Burkitt's lymphoma

• Post-organ transplant lymphoma

• Primary diffuse large B cell lymphoma of the CNS

• Hodgkin's disease

• Extranodal NK/T cell lymphoma, nasal type

Primary effusion lymphoma is a variant of diffuse large B-cell lymphoma that presents with pleural effusions, usually without apparent tumor mass lesions. It is most common in the setting

Sold by @itachibot

of immune deficiency disease, especially AIDS.

Solution to Question 3:

Burkitt lymphoma is a fast-growing, high-grade type of non-Hodgkin lymphoma (NHL).

NHL was earlier classified on the basis of its morphology and clinical behaviour. However, this classification is not currently used. Based on the clinical behavior, NHL is classified as:

• Low grade - CLL/SLL (small lymphocytic lymphoma), follicular lymphoma

• Intermediate grade - mantle cell and marginal cell lymphoma

• High grade - diffuse large B-cell lymphoma (DLBCL), Burkitt's lymphoma

The most aggressive NHL is DLBCL, whereas the most rapidly progressive tumor is Burkitt's lymphoma. Most pediatric NHLs are high-grade and aggressive.

Solution to Question 4:

The gastrointestinal tract is the most common extranodal site in patients with non-Hodgkin's lymphoma (NHL). The stomach is most commonly involved, followed by the small intestine.

The skin is the second-most common site. Mycosis fungoides is a mature T-cell (CD4+) NHL with skin involvement with eventual visceral and nodal spread. When malignant T-cells in the skin enter circulation, it is called Sezary syndrome. It presents as erythroderma (diffuse erythema and scaling). Histologically, there are Sézary-Lutzner cells (with cerebriform nuclei) and Pautrier microabscesses in epidermis.

CNS is the most common extranodal site in patients with HIV infection. Primary CNS lymphoma (histologically, DLBCL) is one of the AIDS-defining malignancies.

Solution to Question 5:

The prognosis of patients with non-Hodgkin's lymphoma is best assessed using the international Prognostic Index (IPI). It consists of 5 clinical risk factors for poor prognosis:

• Age >60 years

• Elevated levels of serum lactate dehydrogenase (LDH)

• Performance status ≥ 2 (ECOG) or ≤70 (Karnofsky)

• Ann Arbor stage III or IV

• >1 site of extranodal involvement

Option A: Ann Arbor staging is used for anatomic staging to develop a treatment plan for Hodgkin's and non-Hodgkin's lymphoma.

Option B: Rai and Binet score is used for staging of CLL.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option D: Hasford scoring is used to assess the prognosis of CML.

Solution to Question 6:

The clinical scenario is suggestive of mantle cell lymphoma. It is diagnosed by its characteristic immunophenotype - CD19+, CD20+, CD21+, CD22+, CD23(-), CD 5+.

Comparison between mantle cell lymphoma and chronic lymphocytic leukemia/small lymphocytic lymphoma (CLL/SLL):

	Mantle cell lymphoma	CLL/SLL
CD5	Positive	Positive
CD23	Negative	Positive
Cyclin D1	Positive	Negative
FMC7	Positive	Negative
CD10, BCL6	Negative	Negative

Solution to Question 7:

Translocations seen in Burkitt's lymphoma are t (8;14), t (2;8) and t (8;22).

Solution to Question 8:

This clinical scenario and the presence of centrocytes and centroblasts on biopsy are suggestive of follicular lymphoma. Immunophenotyping will be positive for CD19, CD79a, CD20, CD10, CD23, Bcl-2.

Follicular lymphoma presents at a median age of 59 years with no sex predilection. The patients most commonly develop painless generalized lymphadenopathy. Systemic symptoms are absent. A lymph node biopsy is usually adequate for the diagnosis of follicular lymphoma. Characteristic features following lymph node biopsy include:

• The normal nodal architecture is effaced by nodular expansions of tumour cells

• Centrocytes - small lymphocytes with cleaved nuclei

• Centroblasts - larger cells with vesicular chromatin and prominent nucleoli

Sold by @itachibot

Solution to Question 9:

Diffuse large B-cell lymphoma (DLBCL) commonly (>50%) presents with extranodal involvement at diagnosis, with the most common sites being the GI tract and bone marrow.

DLBCL is the most common non-Hodgkin's lymphoma. It is a high-grade aggressive tumour. It is associated with BCL-6 chromosomal rearrangements and t(14;18). It is commonly associated with CNS lymphoma, pleural effusion lymphoma, and intravascular lymphoma. It is considered an AIDS-defining malignancy.

AIDS-defining malignancies include:

- Kaposi sarcoma
- Primary central nervous system lymphoma (DLBCL)
- Non-Hodgkin's lymphoma
- Invasive cervical cancer

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

This clinical scenario is suggestive of the endemic type of Burkitt's lymphoma. In this condition, the tumor cells are CD21+.

Burkitt's lymphoma is an aggressive condition and is the most common NHL in children. This condition is endemic in African countries and is associated with Epstein-Barr virus (EBV) infection. It is associated with c-myc dysregulation, especially in cases in bone marrow involvement.

Patients require aggressive management with high-dose chemotherapy.

Solution to Question 11:

The given image shows the characteristic starry sky appearance of Burkitt's lymphoma. The reactive macrophages (labelled as A) represent the stars in starry sky appearance. On the other hand, the homogenous tumour cells (labelled as B) represent the sky.

The neoplastic cells are homogeneous, medium-sized B cells with a high growth fraction. These cells are positive for CD19, CD10, BCL-6 & CD43.

Reactive macrophages are scattered throughout the tumour. Their pale cytoplasm in a background of blue-staining tumour cells gives the tumor a so-called starry sky appearance. Reactive leukocytes and macrophages are not stained by special stains. They are positive for CD14, CD40, CD11b & CD64.

The growth fraction of these lymphomas is very high, with frequent mitotic figures. The fraction of Ki-67+ (MIB-1+) stained cells is nearly 100%.

Sold by @itachibot

Solution to Question 12:

Sjogren's syndrome is implicated in salivary gland MALToma.

MALT lymphomas are commonly associated with an underlying autoimmune or inflammatory process such as:

• Sjogren's syndrome - salivary gland

• Hashimoto's thyroiditis - thyroid

• Helicobacter pylori gastritis - gastric (most common)

• Celiac disease - small intestine

The most common translocation associated with MALToma is (11,18). It has a good prognosis.

Solution to Question 13:

Hairy cell leukemia arises from a late, activated memory B cell exhibiting post-germinal center features.

Mycosis fungoides and anaplastic large cell lymphoma (also called null cell lymphoma) are both T-cell lymphomas.

Solution to Question 14:

The given peripheral smear shows tumor cells with multilobulated nuclei called cloverleaf or flower cells. This is characteristic of adult T-Cell leukemia/lymphoma.

This condition occurs in adults infected by human T-cell leukemia retrovirus type 1 (HTLV-1). It is a rapidly progressive disease that is fatal within months to one year following diagnosis. It is characterized by the following clinical features:

• Skin lesions

• Generalized lymphadenopathy

• Hepatosplenomegaly

• Peripheral blood lymphocytosis,

• Hypercalcemia

Solution to Question 15:

This clinical scenario is suggestive of hairy cell leukemia (HCL).

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

In this condition, most patients present with symptoms related to cytopenias or splenomegaly. On bone marrow aspiration, a dry tap is obtained due to HCL-induced marrow fibrosis and the interlocking cell borders of hairy cells (distinct from myelofibrosis).

Tumor cells share certain characteristics of memory B cells. CD11c, CD25, CD103, CD123 is the characteristic immunophenotype diagnostic of this condition. Pan-B markers such as CD19, CD20, and CD22 may also be positive.

Note: Splenic marginal zone lymphoma involves both white and red pulp of the spleen, whereas hairy cell leukemia is limited to the red pulp. All non-Hodgkin's lymphomas involve white pulp of spleen except:

• Hairy cell leukemia

• Hepatosplenic lymphoma

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

Annexin A1 on immunophenotyping of bone marrow specimens is the best method to differentiate hairy cell leukemia from other B-cell malignancies. It has 100% sensitivity and specificity.

Hairy cells on Wright-stained peripheral blood smear are an important diagnostic finding. Typical hairy cells are mononuclear cells with abundant pale blue-grey agranular cytoplasm with elongated (hairy) projections. Intracytoplasmic inclusion bodies called ribosome-lamellar complexes can also be seen in 50% of cases. However, these can be seen in other B-cell disorders also.

Tartrate-resistant acid phosphatase (TRAP) stain is an important tool and is positive in 95% of cases of hairy cell leukemia. However, it is not pathognomonic.

The following pattern of bone marrow involvement is near to pathognomonic for hairy cell leukemia (>99%):

• Diffuse infiltration of tumor cells with honeycomb appearance.

• Individual cells have a fried egg appearance such that the nucleus of each hairy cell is surrounded by a halo of cytoplasm, as shown below.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

Hairy cell leukemia is associated with infections of Mycobacterium avium-intracellulare. This is possibly related to frequent unexplained monocytopenia. Among them, infection with Mycobacterium kansasii is common.

Solution to Question 18:

The given image shows mononuclear cells with abundant pale blue-grey agranular cytoplasm with elongated (hairy) projections, which are TRAP-positive. This clinical scenario is suggestive of hairy cell leukemia. The drug of choice is cladribine.

Hairy cell leukemia is responsive to chemotherapy with interferon α, pentostatin, and cladribine. However, cladribine is the preferred drug of choice due to complete remission in the majority of patients, and higher rates of long-term disease-free survival. It is administered daily for 7 days by continuous intravenous infusion.

Chemotherapy-refractory patients have responded to vemurafenib, a BRAF inhibitor. Splenectomy is often associated with prolonged remission.

Solution to Question 19:

The most common subtype of Hodgkin's lymphoma in India is the mixed cellularity classical Hodgkin's lymphoma. In contrast, nodular sclerosis is most commonly seen in western countries.

The most common subtype of Hodgkin's lymphoma in:

• Developed/ western countries - nodular sclerosing

Sold by @itachibot

• Children - mixed cellularity

• Females - nodular sclerosis (but Hodgkin's lymphoma is more common in males overall)

• HIV-associated - mixed cellularity

• EBV-associated - lymphocyte-depleted

Solution to Question 20:

Nodular lymphocyte-predominant Hodgkin's lymphoma (HL) is a distinct clinicopathological entity from classical Hodgkin lymphoma as per the WHO classification.

The following image shows the Reed-Sternberg cell in classic HL.

	Classical HL	Nodular lymphocyte-predominant HL
Subtypes	Nodular sclerosisMixed cellularityLymphocyte-richLymphocyte depleted	-
Key difference	Do not express many of the genes that define normal B cells	Retain the immunophenotypic features of germinal center B cells
Immunophenotype	CD15+ CD30+CD20- CD45-	CD20+ CD45+CD15- CD30-
Neoplastic cell	Reed-Sternbergcell	Popcorncell or L&H (lymphocytic & histiocytic) cells
Common Presentation	Mediastinal lymphadenopathy	Peripheral lymphadenopathy



The following image shows the popcorn cell in nodular lymphocyte-predominant HL.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 21:

Supraclavicular or cervical lymphadenopathy is the most common clinical presentation of Hodgkin's lymphoma.

The clinical features of Hodgkin's lymphoma are as follows:

• Asymptomatic lymphadenopathy (non-tender with a rubbery consistency) is the most common presentation.

• Most common - cervical or supraclavicular

• Second-most common presentation - mediastinal mass

• Hepatosplenomegaly in some cases

• Pel-Ebstein fever:

• Refers to recurrent febrile and afebrile intervals, each phase lasting for days to weeks

• More common in older patients with mixed-cellularity Hodgkin’s lymphoma in an abdominal site

• Alcohol-induced pain is a rare, unusual presentation

Solution to Question 22:

This clinical scenario is suggestive of Hodgkin's lymphoma. The paraneoplastic syndrome associated with this condition is subacute cerebellar degeneration.

Other paraneoplastic syndromes associated with Hodgkin's lymphoma include:

• Minimal change disease (nephrotic syndrome)

• Vanishing bile duct syndrome

Sold by @itachibot

• Cholangitis

• Alcohol-induced pain - pain typically occurs within a few minutes after the ingestion of even a small amount of alcohol and is localized to areas involved by HL

Option A: Syndrome of inappropriate ADH secretion is associated with small cell lung carcinoma of the lungs.

Option B: Lambert-Eaton syndrome is associated with small cell lung carcinoma of the lungs.

Option D: Migratory superficial thrombophlebitis (Trousseau syndrome) is associated with adenocarcinoma of the pancreas and lungs.

Solution to Question 23:

The given image shows a lacunar Reed-Sternberg (RS) cell. It is identified by a folded or multilobed nucleus surrounded by an open space, which is an artifact created by disruption of the cytoplasm during tissue sectioning. It is seen in nodular sclerosis classical Hodgkin's disease (CHL).

The histological subtypes of classical Hodgkin's disease are as follows:

Subtype	Histopathology	Clinical correlation
Nodular sclerosis	Lacunar RS cellNecrosis with inflammatory infiltrate (eosinophils, macrophages, plasma cells)Fibrous bands dividing cellular areas into nodules	Mediastinal lymphadenopathy and superior vena cava syndromeExcellent prognosis
Mixed cellularity	RS cells with rich inflammatory infiltrate	Biphasic: Children and elderlyB symptomsAssociated with EBV (and HIV)Very good prognosis
Lymphocyte-rich	Background infiltrate richin T-cells	-
Lymphocyte- depleted	Reticular variant, paucity of background reactive cells	Associated with HIV (and EBV)Poor prognosis

Solution to Question 24:

PET-CT is the most reliable technique for staging Hodgkin's lymphoma.

Reed-Sternberg cells on histology, point to a clinical diagnosis of Hodgkin's lymphoma. However, they are not specific for Hodgkin's lymphoma. Reed-Sternberg cells can also be seen in the following conditions:

• Infectious mononucleosis

• Solid tissue cancers

Sold by @itachibot

• Large cell non-Hodgkin's lymphomas (most common in immunoblastic type)

Classical Hodgkin's lymphoma is associated with positive test results for CD15 and CD30. Nodular lymphocyte-predominant Hodgkin's lymphoma is associated with CD20 positive cells.

The central nervous system (CNS) is an uncommon site of extranodal disease in Hodgkin's lymphoma. Lung, liver, bone, and bone marrow are the most common extranodal sites for the disease.

Solution to Question 25:

In this scenario of Hodgkin's lymphoma, the patient presented with B symptoms (fever, night sweats, and weight loss) and shows lymph node involvement on both sides of the diaphragm are involved (supraclavicular and inguinal). According to Ann Arbor Staging, this patient is categorized as stage IIIB.

Ann Arbor Staging of Hodgkin's lymphoma	
Stage I	Involvement of asinglelymph node (I) or a single extra-lymphatic organ or site (IE)
Stage II	Involvement of 2 or more lymph node regions on thesame side of the diaphragm(II) or localized involvement of an extra-lymphatic organ or site and 1 or more lymph node regions on the same side of the diaphragm (IIE)
Stage III	Involvement of lymph node regions onboth sides of thediaphragm(III), which may be accompanied by:Involvement of the spleen (IIIS) [or]Localized involvement of an extra-lymphatic organ or site (IIIE) [or]Both (IIISE)
Stage IV	Diffuseor disseminated involvement of 1 or more extra-lymphatic organs or tissuesWith or without associated lymph node involvement

Sold by @itachibot

Ann Arbor Staging of Hodgkin's lymphoma	
Additionally classified as:A-AsymptomaticB -Any of the following symptoms:Absence or incidence of fever >38°C (100.4°F) for 3 consecutive daysDrenching night sweatsUnexplained loss of >10% of body weight in the 6 months	

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 26:

In this patient with advanced-stage Hodgkin's lymphoma, his age, gender, and absolute lymphocyte count all indicate poor prognosis.

Unfavourable or poor prognostic factors for advanced-stage HL (stages III and IV) include:

• Age > 45 years

• Male gender

• Haemoglobin (Hb) < 10.5 g/dL

• Serum albumin <4 g/dL

• White blood cell (WBC) count ≥15,000/ µL

• Absolute lymphocyte count (ALC) < 600/ µL

Unfavourable or poor prognostic factors for early Hodgkin's lymphoma (stages I and II) include:

• Large mediastinal lymphadenopathy

• Involvement of 4 or more sites

• Age >45

• ESR > 30 mm/hr

• B symptoms - unexplained fever, night sweats or unexplained weight loss

The selection of initial treatment for Hodgkin's lymphoma is usually based on presenting stage and prognostic factors.

Solution to Question 27:

The patient's age (> 45 years) and presence of B symptoms indicate an unfavourable prognosis. The treatment of choice in early-stage Hodgkin's lymphoma (stage I and II) with an unfavourable prognosis is ABVD regimen along with field radiation. ABVD is a combination of doxorubicin (adriamycin), bleomycin, vinblastine, and dacarbazine.

The treatment of Hodgkin's lymphoma is as follows:

Sold by @itachibot

• Early Hodgkin's lymphoma (favourable and unfavourable) - ABVD regimen + limited field radiation

• Advanced Hodgkin's lymphoma - extended course of chemotherapy with ABVD regimen.

The side effects of this treatment include:

• Doxorubicin - dilated cardiomyopathy

• Bleomycin- interstitial lung disease

• Vinblastine - SIADH, Peripheral neuropathy

• Radiation - hypothyroidism, Lhermitte's syndrome, infertility

Note: R-CHOP regimen is used in the treatment of Non-Hodgkin's lymphoma. It is composed of rituximab + cyclophosphamide, doxorubicin, vincristine (oncovin) and prednisone.

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Solution to Question 28:

Hepatitis C virus (HCV) is the etiological agent for lymphoplasmacytic lymphoma/ Waldenstrom's macroglobulinemia.

Human immunodeficiency virus (HIV) is associated with diffuse large B-cell lymphoma (DLBCL) and Burkitt's lymphoma.

Other acquired lymphoid neoplasms include:

• Gastric MALToma/extranodal marginal cell lymphoma - induced by Helicobacter pylori infection

• Intestinal T-cell lymphomas/enteropathy-associated T cell lymphoma - induced by gluten-sensitive enteropathy

• Anaplastic large cell lymphoma in scar tissue - can occur following breast implants

Sold by @itachibot

Coagulation Disorders

Question 1:

An isolated prolongation of prothrombin time was seen in a patient with chronic liver disease. What does this suggest?

a) Factor VII deficiency

b) Factor XII deficiency

c) Factor XI deficiency

d) Factor VIII deficiency

Question 2:

Prolonged aPTT and normal PT are seen in patients with all of the following except?

a) Hemophilia B

b) Lupus anticoagulant

c) Von Willebrand disease

d) Disseminated intravascular coagulation

Question 3:

Laboratory investigations done in a chronic alcoholic patient are given below. Which of the following clotting factors is most likely to be normal in this patient?

a) Factor II

b) Factor VII

c) Factor VIII

d) Factor XI

Question 4:

Which of the following is the most common disorder of coagulation?

a) Hemophilia A

b) Hemophilia B

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) von Willebrand disease

d) Hemorrhagic diathesis of liver disease

Question 5:

Which of the following conditions are matched correctly with their mode of inheritance?

a) Factor VII deficiency - X linked recessive

b) Factor VIII deficiency - autosomal recessive

c) Factor IX deficiency - X linked recessive

d) Factor V deficiency - autosomal dominant

Question 6:

The most common bleeding manifestation seen in severe hemophilia is:

a) Recurrent hematomas

b) Recurrent hemarthrosis

c) Hematuria

d) Intracranial hemorrhage

Question 7:

A 5-year-old boy was brought with the following finding. His mother says that he had banged his leg against a chair yesterday. He has had similar episodes in the past but they were never this severe. What lab values will you likely find in this boy?

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

a) Normal bleeding time, prolonged PT, prolonged aPTT

b) Normal bleeding time, prolonged PT, normal aPTT

c) Prolonged bleeding time, normal PT, prolonged aPTT

d) Normal bleeding time, normal PT, prolonged aPTT

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 8:

Which of the following is false about acquired hemophilia A?

a) Autoantibodies are formed against factor VIII

b) Bethesda assay confirms the diagnosis

c) It presents in childhood or early adulthood

d) It may be associated with lymphoproliferative malignancies

Question 9:

In a patient with hemophilia A, how much can infusion of one unit of factor VIII/kg increase the plasma levels of factor VIII?

a) 30%

b) 20%

c) 10%

d) 2%

Sold by @itachibot

Question 10:

What effect does desmopressin have when given to patients with coagulation disorders?

a) Increases factor VIII

b) Increases factor IX

c) Increases factor XI

d) Increases factor VII

Question 11:

A teenager with hemophilia A presented with multiple spontaneous bleeds. He has been receiving prophylactic factor VIII infusion on a regular basis. Investigations revealed low factor VIII levels and abnormal plasma mixing study. Which of the following can be used to confirm the diagnosis?

a) 1,3 beta D glucan assay

b) Platelet activation assay

c) Serotonin release assay

d) Bethesda assay

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 12:

Which among the following is wrong about DIC?

a) Increased schistocytes

b) Increased PT

c) Increased fibrinogen

d) Increased FDPs

Question 13:

Which of the following are risk factors that can cause disseminated intravascular coagulation in patients?

a) 1,4

b) 1,2,4,5

c) 2,5

Sold by @itachibot

d) 1,3,4,5

Question 14:

A 33-year-old woman with a twin pregnancy was admitted for normal delivery. After medical induction of labor, she started to bleed profusely from her rectum, vagina, and IV sites. She became hypotensive, tachycardic, and pulses were feeble. Which of the following laboratory findings will you see in her case?

	Platelets	D-dimer	Antithrombin III	Thrombin-antithrombin complex
1	Decreased	Increased	Decreased	Decreased
2	Increased	Decreased	Decreased	Decreased
3	Decreased	Decreased	Decreased	Decreased
4	Decreased	Increased	Decreased	Increased

©MARROW

a) Option 2

b) Option 3

c) Option 1

d) Option 4

Question 15:

The most sensitive test for disseminated intravascular coagulation is _____.

a) Fibrin degradation products level

b) PTT and thrombin time

c) D-dimer assay

d) Platelet count and fibrinogen levels

Question 16:

Sold by @itachibot

A 3-year-old child is brought to the emergency with petechiae, bruising, and abdominal distension. Examination revealed an abdominal mass. Investigations revealed schistocytes and thrombocytopenia. What is the most probable cause of the mass?

a) Neuroblastoma

b) Hepatoblastoma

c) Wilm's tumor

d) Hepatic hemangioma

Question 17:

Patients with protein C deficiency are at an increased risk of developing all of the following except:

a) Deep vein thrombosis

b) Purpura fulminans

c) Warfarin-induced skin necrosis

d) Spontaneous bleeding

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A patient is on prophylactic unfractionated heparin for deep vein thrombosis. Which of the following tests can be used to monitor its anticoagulant effect?

a) Prothrombin time

b) Anti-Factor Xa level

c) PT and Anti-Factor Xa level

d) No monitoring is required

Question 19:

Which of the following is false about low molecular weight heparin?

a) Higher risk of of bleeding than heparin

b) No laboratory monitoring is required

c) Lower risk of heparin induced thrombocytopenia

d) Protamine sulphate is only partially effective against low molecular weight heparin

Sold by @itachibot

Question 20:

Which of the following is a direct thrombin inhibitor?

a) Apixaban

b) Abciximab

c) Argatroban

d) Edoxaban

Question 21:

A patient presented with bleeding gums and skin discoloration. She was given heparin for DVT prophylaxis 5 days ago. Investigations showed thrombocytopenia. Which of the following is the most specific test to diagnose this condition?

a) 4T scoring

b) Platelet activation assay

c) Heparin-PF-4 complex antibody assay

d) Serotonin release assay

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 22:

A 69-year-old male patient with chronic kidney disease presented with bleeding gums and skin rash. He had undergone dialysis 1 week ago. Further investigations revealed heparin-induced thrombocytopenia to be the cause. Which of the following drugs can be given to him?

a) Lepirudin

b) Argatroban

c) Bivalirudin

d) Fondaparinux

Question 23:

A patient on warfarin develops severe gastrointestinal bleeding. What is the treatment of choice?

a) Vitamin K slow IV infusion

b) Fresh frozen plasma

Sold by @itachibot

c) Vit K and FFP

d) Discontinuation of warfarin only

Question 24:

Which of the following is not a direct oral anticoagulant ?

a) Dabigatran

b) Danaparoid

c) Rivoraxaban

d) Edoxaban

Question 25:

In patients with which of the following is dabigatran approved for use?

a) Elective hip arthroplasty requiring thromboprophylaxis

b) Total knee arthroplasty requiring thromboprophylaxis

c) Metallic heart valve requiring stroke prophylaxis

d) Non-valvular atrial fibrillation requiring stroke prophylaxis

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	d
3	c
4	d
5	c
6	b
7	d
8	c
9	d
10	a
11	d

Sold by @itachibot

12	c
13	d
14	d
15	a
16	d
17	d
18	b
19	a
20	c
21	d
22	b
23	c
24	b
25	d

Sold by @itachibot
purchased this from somewhere else,
may have been scammed.

Detailed Explanations

Solution to Question 1:

Isolated PT prolongation suggests factor VII deficiency. It can be an inherited or acquired condition (chronic liver disease).

Test	Used to assess	Activated onexposure to	Normalv alue
PT	Extrinsicpathway (factorVII) Commonpathway (factorsI, II, V, X)	Tissue factor	12-16s
aPTT	Intrinsicpathway (factorsVIII, IX, XI, XII)Commonpathwa y (factorsI, II, V, X)	Activators(silica,kaolin)	26-34s
TT	Fibrinogen levels	Thrombin	15-19s

Solution to Question 2:

Disseminated intravascular coagulation is associated with prolonged PT and prolonged aPTT.

Conditions with prolonged aPTT can be classified on the basis of the bleeding symptoms:

• No bleeding symptoms: deficiency of Factor XII, prekallikrein, HMW kininogen, lupus anticoagulant

Sold by @itachibot

• Mild bleeding symptoms: Factor XI deficiency

• Severe bleeding symptoms: Hemophilia, von Willebrand disease

Lupus anticoagulant is an antiphospholipid antibody that interferes with aPTT test and causes its prolongation. It is corrected by adding excess phospholipid.

The clinical manifestation of lupus anticoagulant is thrombosis. Confirmatory tests are dilute Russell's viper venom test and kaolin clotting time.

Solution to Question 3:

All clotting factors are likely to be abnormal except factor VIII.

The liver is the site of production of the following factors:

• Fibrinogen (factor I)

• Thrombin (factor II)

• Factor V

• Factor VII (shortest half-life, hence PT affected first)

• Factor IX

• Factor X

• Factor XI

Factor VIII is produced in the endothelial cells and factor XIII A-subunit is produced in the bone marrow.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 4:

Hemorrhagic diathesis of liver disease is the most common disorder of coagulation among the given options. It is the increased tendency of spontaneous bleeding associated with fulminant or end-stage liver disease.

Acquired deficiencies of plasma coagulation factors are more frequent than congenital disorders. The most common acquired disorders include:

• Hemorrhagic diathesis of liver disease

• Disseminated intravascular coagulation (DIC)

• Vitamin K deficiency.

Hemophilia A is the most common inherited disorder of coagulation. Hemophilia A is more common than Hemophilia B.

von Willebrand disease is the most common bleeding disorder.

Sold by @itachibot

Solution to Question 5:

Factor IX deficiency (hemophilia B) is inherited as an X-linked recessive trait.

Hemophilia is an X-linked recessive hemorrhagic disease due to mutations in:

• Factor VIII gene (hemophilia A or classic hemophilia) -80% more common than hemophilia B

• Factor IX gene (hemophilia B)

Clinically, hemophilia A and hemophilia B are indistinguishable.

All clotting factor deficiencies except hemophilia are inherited as autosomal recessive.

Inherited coagulation disorders treatment:

Blood components	Composition	Indications
Factor VIII concentrates	Factor VIII	Hemophilia A
Factor IX concentrates	Factor IX	Hemophilia B
Cryoprecipitate	FibrinogenFactor XIIIFactor VIIIVWF	Fibrinogen deficiency, factor XIII deficiency, von Willebrand disease
Fresh Frozen Plasma (FFP)	Coagulation factorsProtein C Protein SAntithrombin	Deficiency of prothrombin, factor V, factor VII, factor X, factor XI

Solution to Question 6:

Recurrent hemarthrosis is the most common bleeding manifestation of severe hemophilia. Hemophilia is said to be severe when the residual activity of factor VIII is <1%.

Hemophilia is an X-linked recessive hemorrhagic disorder and is caused by mutations in either the F8 gene (hemophilia A or classic hemophilia) or the F9 gene (hemophilia B). Clinically, it is impossible to distinguish between hemophilia A and hemophilia B as they present with similar symptoms. It can be classified based on residual activity of FVIII or FIX as severe (<1%), moderate (1–5%), or mild (6–30%)

Hemophilia presents early in life- bleeding after circumcision or when beginning to walk or crawl.

Clinical features:

• Acute hemarthrosis:

• Local pain, swelling

• Erythema

• Muscle contractures due to fixed positioning.

• Chronic hemarthrosis:

• Synovial thickening

• Synovitis (due to intraarticular blood)

Sold by @itachibot

The most common joint affected is the knee. One joint is affected more than the other. This is called the target joint. Muscle hematomas can lead to compartment syndromes.

Hematuria is common but self-limited. Life-threatening bleeds such as bleeding into oropharyngeal spaces, CNS, and retroperitoneum can occur.

Treatment: Without treatment, severe hemophilia may limit life expectancy. Factor replacement for hemophilia has been the mainstay of therapy.

• FVIII dose (IU) = Target FVIII levels – FVIII baseline levels × body weight (kg) × 0.5 unit/kg .

• FIX dose (IU) = Target FIX levels – FIX baseline levels × body weight (kg) × 1 unit/kg

The FVIII half-life of 8–12 h requires injections twice a day to maintain therapeutic levels, whereas the FIX half-life is longer, ~24 h so that once-a-day injection is sufficient.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

In the above clinical scenario, recurrent hemarthrosis in a male child is suggestive of hemophilia. Normal bleeding time, normal PT, prolonged aPTT is seen in hemophilia A and B.

Hemophilia A has decreased factor VIII activity and hemophilia B has decreased factor IX activity.

Note: Prolonged bleeding time, normal PT, prolonged aPTT is seen in von Willebrand disease.

Solution to Question 8:

Acquired hemophilia A occurs predominantly in older adults (median age- 60 years).

It occasionally occurs in pregnant or postpartum women.

It is an acquired immune-mediated disease characterized by autoantibodies against clotting factor VIII. It is associated with an underlying disease in 50% of patients. They include:

• Malignancies - lymphomas, prostate cancer

• Dermatologic diseases - pemphigoid

Solution to Question 9:

One unit of factor VIII/kg body weight increases the plasma levels of factor VIII levels by 2%. Therefore, for a rise in 1 % in the plasma, half unit/ kg will be required, that is 0.5 units/kg.

It is used in the treatment of hemophilia. One unit of FVIII is the amount of FVIII in 1 mL of plasma.

The dose required for replacement is calculated as :

• Factor VIII dose = (Target factor VIII levels - baseline factor VIII levels) x bodyweight x 0.5 units/ kg

Sold by @itachibot

• Factor IX (IU) = (Target FIX levels – FIX baseline levels) x bodyweight (kg) x 1 unit/kg

Cryoprecipitate is used when factor concentrates are not available. Each bag of cryoprecipitate contains 80 IU of Factor VIII.

Blood components	Composition	Indications
Factor VIII concentrates	Factor VIII	Hemophilia A
Factor IX concentrates	Factor IX	Hemophilia B
Cryoprecipitate	FibrinogenFactor XIIIFactor VIIIVWF	Fibrinogen deficiency, Factor XIII deficiency, von Willebrand disease
Fresh frozen plasma (FFP)	Coagulation factorsProtein C Protein SAntithrombin	Deficiency of prothrombin, factor V, factor VII, factor X, factor XI

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

Desmopressin causes a transient rise in factor VIII and von Willebrand factor (VWF) hence it is given in mild and moderate hemophilia A.

It does not improve factor VIII levels in severe hemophilia A patients because there are no stores to release. Repeated dosing leads to tachyphylaxis.

It does not increase factor IX levels.

Solution to Question 11:

Bethesda assay is the best test for the confirmation of factor VIII inhibitors.

The formation of inhibitor/ alloantibodies to FVIII or FIX is the major complication of hemophilia treatment. When this happens, patients do not respond to factor replacement at therapeutic doses.

aPTT measurement is done on a mixed sample. It is used to determine the presence of factor VIII inhibitors. aPTT is estimated using a 1:1 mix of patient's plasma and normal plasma. It is interpreted as:

• In hemophilia, aPTT is corrected with a 1:1 mix.

• In patients with inhibitors, aPTT with 1:1 mix is abnormally prolonged (inhibitor neutralizes the normal factors).

Bethesda assay is based on the same principle but is superior because it helps to quantify the type and specificity of the inhibitor.

Option A: 1,3 beta D glucan assay is designed to diagnose invasive fungal infections (IFIs).

Options B and C: It is used to diagnose heparin-induced thrombocytopenia.

Sold by @itachibot

Solution to Question 12:

In disseminated intravascular coagulopathy (DIC), fibrinogen levels are decreased due to increased bleeding.

Disseminated intravascular coagulation (DIC) is an acquired syndrome characterized by the widespread activation of blood clotting factors within blood vessels, resulting in a loss of clotting control. The primary mechanism of DIC involves the uncontrolled generation of thrombin through multiple pathways which leads to sustained coagulation activation, the formation of fibrin, and the consumption of clotting factors and platelets.

Infections are the most frequent cause of DIC.

Laboratory investigations in DIC:

• Decreased platelet count

• Prolonged PT/TT

• Prolonged activated partial thromboplastin time (aPTT)

• Decreased fibrinogen

• Elevated fibrin degradation products (D-dimers)

• Peripheral smear: microangiopathic hemolytic anemia in DIC with schistocytes (fragmented red blood cells)

The treatment involves platelet transfusion when platelet counts are below 10,000-20,000/µL. Replacement of fibrinogen and coagulation factors is with fresh frozen plasma (FFP) along with cryoprecipitate or fibrinogen concentrate.

In cases of low-grade DIC associated with solid tumors or acute promyelocytic leukemia low doses of continuous-infusion heparin has been effective.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 13:

Acute pancreatitis can cause disseminated intravascular coagulation (DIC) not chronic pancreatitis.

Common causes of DIC:

• Sepsis- gram-positive and gram-negative bacteria, viral, mycotic, parasitic, rickettsial

• Acute pancreatitis

• Trauma, burns

• Malignancies- acute promyelocytic leukemia, adenocarcinoma of pancreas and prostate

• Obstetric- abruptio placentae, amniotic fluid embolism, and septic abortion

• Snakebite

Sold by @itachibot

Solution to Question 14:

In the above clinical scenario, the patient seems to have disseminated intravascular coagulation (DIC) due to amniotic fluid embolism following medical induction of labor.

Disseminated intravascular coagulation (DIC) is a condition characterized by widespread intravascular fibrin formation. The central mechanism is the uncontrolled generation of thrombin by exposure of blood to pathologic levels of tissue factor.

Lab abnormalities in DIC:

- Low platelet count
- Low fibrinogen
- Prolonged PT and aPTT
- Increased D-dimer
- Increased fibrin-degradation products
- Decreased protein C
- Decreased antithrombin III
- Increased thrombin-antithrombin complex

Sold by @itachibot
this from someone else,
en scammed.



Solution to Question 15:

The most sensitive test for disseminated intravascular coagulation (DIC) is FDP (fibrinogen degradation products) level.

Sold by @itachibot

Diagnostic test	Diagnostic efficiency	Sensitivity	Specificity
FDP and D-dimer combination	95%	91%	94%
FDP	87%	100%	67%
PT/PTT and FDP combination	86%	91%	71%
D-dimer	80%	91%	68%
PT, PTT, thrombin time, platelet count, fibrinogen, and the presence of schistocytes	57%, 57%, 70%, 67%, 65%, and 51%, respectively	low	low

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Solution to Question 16:

This patient presents with petechiae, and abdominal distension with schistocytes and thrombocytopenia. This in child most likely has Kasabach-Merritt syndrome (secondary DIC). His abdominal mass is likely to be a hepatic hemangioma.

Kasabach-Merritt syndrome is a life-threatening complication that occurs in patients with giant hemangiomas. It is most commonly seen in infants.

Clinical features:

• Enlarged abdomen

• Hepatomegaly or jaundice

• Petechiae, bruising, and frank bleeding

• Painful lesions

• Anemia

It is characterized by profound thrombocytopenia and consumptive coagulopathy leading to a DIC-like picture.

Other options:

Option A: Neuroblastomas are tumors of the sympathetic ganglia and adrenal medulla that are derived from primordial neural crest cells. It is the most common extracranial solid tumor of childhood and the most frequently diagnosed tumor in infancy. Brain tumors are the most common solid tumors in childhood. They present as a flank mass that crosses the midline.

Option B: Hepatoblastoma is the most common childhood liver tumor. Histologically it is composed of small polygonal embryonal cells forming trabeculae, acini, or papillary structures in the developing liver.

Option C: Wilms tumor is the most common primary malignant renal tumor in children and is the second most common malignant abdominal tumor in children. The clinical presentation usually involves an asymptomatic unilateral abdominal mass, hypertension, abdominal pain, and gross painless hematuria. Constitutional symptoms of fever, anorexia, and weight loss can also be present at the time of diagnosis.

Sold by @itachibot

Solution to Question 17:

Spontaneous bleeding is not seen in protein C deficiency. It is characterised by an increased tendency for thrombus formation.

Protein C deficiency is a rare genetic disorder. It predisposes to the following thrombotic complications:

• Deep vein thrombosis (DVT):

• Adults with a mild form of the disease at higher risk of developing venous thromboembolism.

• Purpura fulminans:

• Severe form of DIC

• Thrombosis of extensive areas of the skin

• Affects predominantly young children following viral or bacterial infection.

• Warfarin-induced skin necrosis:

• Rare complication seen in patients with congenital or acquired deficiencies of protein C or S.

• Thrombosis of the skin leads to well-demarcated erythematous lesions with central necrosis on the thighs, buttocks, breasts, or toes(shown below).

Sold by @itachibot
this from someone else,
en scammed.
If yo

Solution to Question 18:

The anticoagulant effect of heparin can be monitored using anti-factor Xa level and activated partial thromboplastin time (aPTT). aPTT is prolonged as heparin inhibits intrinsic pathway factors.

Sold by @itachibot

Prothrombin time and INR are used to monitor the anticoagulant effect of warfarin (inhibits vitamin K dependent factor II, VII, IX, X. Factor VII has the shortest half-life).

No monitoring is required with the use of low molecular weight heparin (LMWH) and fondaparinux.

Solution to Question 19:

Low molecular weight heparin (LMWH) has a lower risk of bleeding than heparin.

In contrast to heparin which requires monitoring by aPTT and anti-factorXa, laboratory monitoring is not required for LMWH. It is also associated with lower risk of heparin induced thrombocytopenia.

Examples of LMWH:

• Dalteparin

• Enoxaparin

• Tinzaparin

• Nadroparin

Protamine sulphate is the only clinically approved antidote for unfractionated heparin. Protamine sulphate is only partially effective against low molecular weight heparin.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 20:

Argatroban is a parenteral direct thrombin inhibitor.

Apixaban and edoxaban are a novel oral anticoagulant (NOAC) that inhibits factor Xa. Abciximab is an antiplatelet drug that inhibits Gp IIb-IIIa receptor.

Solution to Question 21:

Serotonin release assay is the most specific test for the diagnosis of heparin-induced thrombocytopenia (HIT).

HIT is commonly diagnosed clinically by means of 4T scoring (Thrombocytopenia, Timing of platelet count fall, Thrombosis, Other causes for Thrombocytopenia).

Serotonin release assay is performed by quantifying serotonin release.

Platelets loaded with labeled serotonin are exposed to patient serum in the absence or presence of varying concentrations of heparin. If the patient serum contains the HIT antibody, heparin addition induces platelet activation and serotonin release.

Enzyme-linked assays such as platelet activation assay and heparin-PF-4 complex antibody assay are sensitive but not specific for HIT.

Sold by @itachibot

Solution to Question 22:

In the above clinical scenario, the patient requires direct thrombin inhibitors for the treatment of heparin-induced thrombocytopenia. Argatroban is safe in renal disease.

The parenteral direct thrombin inhibitors are:

• Argatroban - hepatic clearance, approved for the treatment of HIT

• Lepirudin - renal clearance, contraindicated in renal disease

• Bivalirudin- renal clearance, contraindicated in renal disease, shortest half-life

Fondaparinux is a factor Xa inhibitor. It is contraindicated in renal disease because it has renal clearance.

Solution to Question 23:

Patients on warfarin who present with severe bleeding are treated with both IV infusion of vitamin K and fresh frozen plasma regardless of the INR.

Warfarin is an oral anticoagulant that blocks the enzyme vitamin K epoxide reductase. This inhibits factors II, VII, IX, and X, proteins C and S. This can cause an initial procoagulant state.

Since the anticoagulant action of warfarin takes 2-3 days, patients require concomitant treatment (bridging therapy) with a rapidly acting parenteral anticoagulant (heparin, LMWH or fondaparinux) for at least 5 days.

Warfarin has a narrow therapeutic window, hence all patients on warfarin should have their INR determined every 3–4 weeks. The most common side effect of warfarin is bleeding.

Severe or life-threatening bleed includes:

• Hematuria

• Gastrointestinal bleeding

• Intracranial hemorrhage

• Intramuscular or intra-abdominal bleed

Management of supratherapeutic INR:

INR	Bleeding	Management
3.5-10	No	Hold warfarin until INR is therapeutic
>10	No	Oral vitamin K 2.5-5 mg

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

INR	Bleeding	Management
Any	Serious or life-threatening	10 mg vitamin K slow IV infusion + fresh frozen plasma/prothrombin complex concentrate

Solution to Question 24:

Danaparoid is not a direct oral anticoagulant (DOAC). It is a heparinoid administered via IV or SC routes.

DOACs are alternatives to warfarin, easier to administer with fixed doses, and no requirement for monitoring.

All DOACs are contraindicated in patients with metallic/prosthetic heart valves.

Direct oral anticoagulants (DOAC)	
Thrombin inhibitors (dabigatran)	FactorXainhibitors
No CYP2C9 polymorphismsA standard dose of 150 mg BD No PT/INR monitoringDialyzableMild GI bleeding	RivaroxabanApixabanEdoxabanBetrixaban

Solution to Question 25:

Stroke prophylaxis in non-valvular atrial fibrillation is the only indication for which all DOACs are approved. All DOACs are contraindicated in patients with metallic/prosthetic heart valves.

Novel oral anticoagulants:

Dabigatran is a direct oral anticoagulant (DOAC). DOACs are alternatives to warfarin, easier to administer with fixed doses, and no requirement for monitoring.

Only rivaroxaban and apixaban are licensed for use as thromboprophylaxis after elective hip or knee arthroplasty.

Thrombin inhibitors	Factor Xa inhibitors
Dabigatran (oral)Argatroban LepirudinDesirudinBivalirudin	RivaroxabanApixabanEdoxabanBetrixaban

Sold by @itachibot

Haemoglobinopathies

Question 1:

Which of the following is true about hemoglobin?

a) Hemoglobin tetramer is highly soluble.

b) Individual globin chains are soluble.

c) Each heme moiety can bind to two O_2 molecules.

d) Each heme moiety consists of Iron in the ferric state.

Question 2:

A patient with alpha thalassemia is susceptible to which of the following infections?

a) Plasmodium vivax

b) Plasmodium falciparum

c) Plasmodium ovale

d) Plasmodium malaria

Question 3:

The peripheral smear of a patient is given below. This abnormality is due to which of the following point mutations?



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) 5th glutamic acid is replaced by valine

b) 5th valine is replaced by glutamic acid

c) 6th glutamic acid is replaced by valine

d) 6th valine is replaced by glutamic acid

Question 4:

A 10-year-old boy complains of multiple episodes of acute severe pain in his fingers and toes which resolves on its own for the past 1 year. Which of the following conditions is he most likely suffering from?

a) Alpha-thalassemia

b) Beta-thalassemia

c) Sickle cell anemia

d) Von Willebrand disease type 1

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

What is the most common cause of osteomyelitis in patients with sickle cell disease?

a) Staphylococcus aureus

b) Streptococcus

c) Staphylococcus epidermidis

d) Salmonella

Question 6:

What will be seen on performing hemoglobin electrophoresis in a patient with suspected sickle cell anemia?

a) HbS and HbA

b) HbS and HbF

c) HbF and HbA

d) HbA2 and HbF

Sold by @itachibot

Question 7:

Which of the following does not increase HbF levels?

a) Hydroxyurea

b) 5-Azacytidine

c) Decitabine

d) Gemcitabine

Question 8:

Which of the following abnormal hemoglobins would be present in patients with congenital methemoglobinemia?

a) HbE

b) HbC

c) Hb Yakima

d) Hb M Iwata

Question 9:

Which of the following would you not expect in a patient with sickle cell anemia?

a) Reticulocytosis

b) Howell jolly bodies

c) Granulocytosis

d) Thrombocytopenia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	c
5	d

Sold by @itachibot

6	b
7	d
8	d
9	d

Detailed Explanations

Solution to Question 1:

The hemoglobin tetramer is highly soluble in the plasma. However, individual globin chains are insoluble. Unpaired globin precipitates can form inclusions that damage the cell and trigger apoptosis.

Each hemoglobin molecule consists of a tetramer of globin polypeptide chains. This includes a pair of α-like chains and a pair of β-like chains. Each globin chain enfolds a single heme moiety. Heme consists of a protoporphyrin IX ring complexed with a single iron atom in the ferrous state (Fe^{2+}). Each heme moiety can bind one O_2 molecule.

The common forms of hemoglobin include:

- HbA ($\alpha_2\beta_2$): Major adult hemoglobin
- HbF ($\alpha_2\gamma_2$): Fetal hemoglobin
- HbA2 ($\alpha_2\delta_2$): Minor adult hemoglobin

Solution to Question 2:

Patients with α-thalassemia are susceptible to infection with Plasmodium vivax.

Thalassemia provides natural protection against infection with the more lethal Plasmodium falciparum. There is no conclusive evidence on the relationship between alpha thalassemia and P. malariae or P. ovale infection.

Conditions protective against Plasmodium falciparum include:

- Sickle cell trait
- Hemoglobin C
- Hemoglobin SC
- Hemoglobin E
- Hereditary persistence of fetal hemoglobin (Hb F)
- Alpha and beta-thalassemia
- Band 3 mutation
- G6PD deficiency

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Solution to Question 3:

The given image shows sickled RBCs, suggestive of sickle cell anemia. It occurs due to the replacement of the 6th glutamic acid by valine in the beta-globin gene on chromosome 11.

This mutation is due to a single nucleotide substitution (from GAG to GTG). The resulting amino acid substitution causes hydrophobic sickling of RBCs.

Solution to Question 4:

The features of multiple episodes of acute severe pain in fingers and toes with spontaneous resolution are suggestive of dactylitis (hand-foot syndrome), which usually occurs in sickle cell anemia.

The sickled RBCs have a sticking tendency towards the endothelium of the small venules. These abnormalities provoke unpredictable episodes of vaso-occlusion. Dactylitis is characterized by severe pain affecting the bones of the hands, feet, or both. It is often the first symptom of sickle cell anemia in children.

Treatment - Administration of patient-controlled analgesia or a frequent fixed dose of opioids with rescue doses for breakthrough pain are preferred with frequent monitoring during the initial 48-72 hrs to prevent unexpected death due to arrhythmias or pulmonary embolism.

Other manifestations include:

• Splenic autoinfarction - repeated splenic microinfarcts

• Acute splenic sequestration - venous obstruction, needing emergency transfusion/splenectomy

• Painful ischemic crises - vaso-occlusion in connective and musculoskeletal tissue

• Renal papillary necrosis leading to isosthenuria

• Cerebrovascular strokes

• Chronic lower leg ulcers

• Priapism in males

α-thalassemia (Option A) and β-thalassemia (Option B) are characterized by pallor, jaundice, growth failure, hepatosplenomegaly, characteristic thalassemic facies, and hair-on-end appearance of the skull.

Option D: Von Willebrand Disease is characterized by predominantly mucosal bleeding symptoms like excessive bruising and epistaxis.

Solution to Question 5:

Salmonella is the most common organism causing osteomyelitis in sickle cell disease.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Staphylococcus aureus is the most common organism causing osteomyelitis in the general population.

Solution to Question 6:

HbS and HbF are detected on electrophoresis in patients with sickle cell anemia. HbA and HbA2 are typically absent.

Condition	Clinical abnormalities	Hemoglobin level (g/dl)	HemoglobinElectrophoresis
Sickle cell trait	Rare painless hematuria may be asymptomatic	Normal	Hb S/A:40/60
Sickle cell anemia	Vasoocclusive crises with infarction of spleen, brain, marrow, kidney, lung; aseptic necrosis of bone; gallstones; priapism; ankle ulcers	7–10	HbS/HbA: 100/0HbF: 2–25%

Sold by @itachibot
If you purchased this from
you may have been scam

Solution to Question 7:

Gemcitabine is an anti-metabolite and has no effect on HbF levels.

The following drugs increase HbF levels and hence can be used in the management of sickle cell anemia:

• Hydroxyurea - mainstay of therapy for patients with severe symptoms

• 5-azacytidine - associated with acute toxicity and carcinogenesis

• 5-deoxyazacytidine or decitabine - alternative with less toxicity

Solution to Question 8:

Hb M Iwata causes congenital methemoglobinemia.

Methemoglobin is generated by oxidation of the heme iron moieties to the ferric state, causing a characteristic bluish-brown muddy color resembling cyanosis. Methemoglobin has such high oxygen affinity that virtually no oxygen is delivered. Levels >50–60% are often fatal.

Congenital methemoglobinemia arises from globin mutations that stabilize iron in the ferric state (HbM Iwata) or mutations that impair the enzymes that reduce methemoglobin to hemoglobin (methemoglobin reductase, NADP diaphorase).

Acquired methemoglobinemia is caused by toxins that oxidize heme iron such as:

• Chloroquine

Sold by @itachibot

• Dapsone

• Sulfonamides

• Nitroglycerin

The following image shows chocolate-brown discoloration of blood in methemoglobinemia.

If you purchased this you may have been s else,

Solution to Question 9:

Thrombocytosis is seen in sickle cell anemia, not thrombocytopenia. Laboratory findings of sickle cell anemia include:

• Peripheral blood smear:

• Irreversibly sickled cells (5–50% of red cells)

• Reticulocytosis (10–25%)

• Nucleated red blood cells

• Howell–Jolly bodies and target cells (due to hyposplenism)

• Granulocytosis - WBC count of 12,000–15,000/mcL

• Reactive thrombocytosis

• High indirect bilirubin levels

Sold by @itachibot

Blood Bank & Transfusion Medicine

Question 1:

What is the 'A blood antigen' composed of?

a) Galactose attached to H substance

b) Glucose attached to H substance

c) N-acetyl galactosamine attached to H substance

d) N-acetyl glucosamine attached to H substance

Question 2:

Which chromosome contains the genes that determine the A and B blood groups?

a) 9

b) 7

c) 5

d) 1

Question 3:

Individuals with a lack of which of the following blood group antigens have immunity against Plasmodium vivax?

a) Kell

b) Lewis

c) Kidd

d) Duffy

Question 4:

Donated blood is screened for all of the following infections except:

a) Kala azar

b) Hepatitis C

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) Syphilis

d) Malaria

Question 5:

As the medical officer at a blood bank, you are supervising the separation of blood components of donated blood. The preservative used for the packed red cells is SAG-M. Which of the following criteria must be fulfilled in order to safely transfuse this to patients?

a) 1

b) 2

c) 3

d) 4

Question 6:

Packed red blood cell transfusion is started for a patient in the ward. It must be completed within:

a) 5 hours

b) 2 hours

c) 3 hours

d) 4 hours

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Cryoprecipitate is useful for the management of patients with all of the following conditions except:

a) Hemophilia A

b) Von Willebrand disease

c) Hypofibrinogenemia

d) Christmas disease

Question 8:

Sold by @itachibot

You are monitoring the falling platelet count of a 31-year-old man with dengue fever. You would recommend prophylactic platelet transfusion when the count drops below what level?

a) 5000 platelets /µL

b) 10000 platelets /µL

c) 15000 platelets /µL

d) 50000 platelets /µL

Question 9:

Which of the following blood components does not carry the risk of transmitting cytomegalovirus infection?

a) Packed red blood cells

b) Whole blood

c) Fresh frozen plasma

d) Random donor platelets

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

Which of the following reactions occurs most frequently in patients undergoing blood transfusions?

a) Acute hemolytic transfusion reaction

b) Febrile non-hemolytic transfusion reaction

c) Transfusion-related acute lung injury

d) Anaphylactic reaction

Question 11:

Thirty minutes after a blood transfusion is started, a patient develops chills and severe flank pain. Her temperature is 102.4 F, SpO2 is 96%, and her blood pressure is 90/60 mm Hg. She then begins bleeding around her IV catheter site. What is the most likely diagnosis?

a) Anaphylactic reaction

b) Acute hemolytic reaction

c) Febrile non-hemolytic reaction

d) Anamnestic antibody response

Sold by @itachibot

Question 12:

A 40-year-old man is being evaluated in the ward for fatigue and pallor. One week ago, he had undergone a blood transfusion following a road accident. Further evaluation reveals Hb of 8.1 g/dl and positive direct Coomb's test and history of previous blood transfusion. What is the likely mechanism of this condition?

a) ABO incompatibility reaction

b) Antibodies against donor leukocytes

c) Antibodies against donor IgA

d) Anamnestic antibody response

Question 13:

Which among the following options is the most common cause of death in patients undergoing blood transfusions?

a) Febrile non-hemolytic transfusion reaction

b) Acute hemolytic transfusion reaction

c) Delayed hemolytic transfusion reaction

d) Transfusion-related acute lung injury

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 14:

Following a normal vaginal delivery, a 30-year-old woman develops severe postpartum hemorrhage. Over the next 24 hours, she undergoes transfusion of 11 units of pRBCs. She is at risk for all of the following complications except:

a) Metabolic alkalosis

b) Hypercalcemia

c) Hyperkalemia

d) Hypothermia

Question 15:

A patient with thalassemia has a history of multiple blood transfusions, iron overload, and cardiac arrhythmia. She has now come for blood transfusion and during the process,

Sold by @itachibot

complains of backache and looks very anxious. What would you do next?

a) Stop the blood transfusion

b) Continue the transfusion but do an ECG

c) Stop the transfusion. Wait for patient to become normal and then start it again

d) Do clerical check and get ECG

Question 16:

A patient of hemophilia received multiple blood transfusions. Which of the following metabolic abnormalities can be seen in a patient?

a) Metabolic alkalosis

b) Respiratory alkalosis

c) Metabolic acidosis

d) Respiratory acidosis

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	a
3	d
4	a
5	b
6	d
7	d
8	b
9	c
10	b
11	b
12	d
13	d
14	b
15	a

Sold by @itachibot

16	a

Detailed Explanations

Solution to Question 1:

The blood group 'antigen A' is composed of N-acetyl galactosamine attached to the H substance.

H substance is formed by the addition of fucose to the glycolipid or glycoprotein backbone. It is present in all blood groups, even the O group.

A antigen = N-acetyl galactosamine + H substance

B antigen = galactose + H substance

However, in case of the Bombay phenotype, RBCs lack H antigen. Hence, these individuals have anti-A, anti-B, and anti-H antibodies. The only compatible blood donor is another person with the Bombay blood group.

Solution to Question 2:

The genes that determine the A and B blood groups are located on chromosome 9. The phenotypes are expressed in the Mendelian codominant manner.

Rh genes determining Rh blood grouping are present on chromosome 1.

Solution to Question 3:

Duffy antigens serve as receptors for Plasmodium vivax. Hence, the absence of the Duffy antigen is protective against Plasmodium vivax.

Hemoglobinopathies that are protective against Plasmodium falciparum include:

- Sickle cell trait
- Hemoglobin C
- Hemoglobin SC
- Hemoglobin E
- Hereditary persistence of fetal hemoglobin (Hb F)
- Alpha and beta-thalassemia

Solution to Question 4:

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

Donated blood is not routinely screened for kala-azar. It is screened for the following infections in the blood bank:

- Human immunodeficiency virus
- Hepatitis B
- Hepatitis C
- Malaria
- Syphilis

Solution to Question 5:

The shelf life of packed red cells (pRBCs) in the SAG-M solution is 42 days. The optimum temperature for storage of packed RBCs and whole blood collected in preservatives is +2° to +6° C. This temperature maintains the viability of the cells and prevents bacterial contamination.

Blood component	Storage temperature	Shelf-life
PRBCs	+2° to +6 °C	CPD-21 daysCPD-A-35 daysSAG M-42 days
Platelets	+20 to +24 °C	5 days
Fresh frozen plasma	-30 °C or lower	1 year
Cryoprecipitate	-30 °C or lower	1 years

Solution to Question 6:

Transfusion of pRBCs (packed red blood cells) or whole blood should be completed within 4 hours after the commencement. Transfusion of platelets, fresh frozen plasma, and cryoprecipitate should be completed within 30 minutes.

Solution to Question 7:

Cryoprecipitate does not contain factor IX. Hence, it is not useful for the management of patients with Christmas disease, which occurs due to deficiency of factor IX.

Components of cryoprecipitate are:

- Factor VIII
- Factor XIII
- Fibrinogen

Sold by @itachibot

• Fibronectin

• Von Willebrand factor

Therefore it can be used in patients with hemophilia A, factor XIII deficiency, hypofibrinogenemia, and von Willebrand disease.

Solution to Question 8:

The threshold for prophylactic platelet transfusion is 10,000/µL in order to prevent spontaneous bleeding. If the patient is planned for any invasive procedures, the target platelet level is 50,000/µL.

The shelf life of donor platelets is 5 days. It is stored on a special agitator at 20-24 °C. The half-life of transfused platelets is 3-5 days. One unit of pooled platelets should be transfused within 30 minutes because of the risk of bacterial proliferation.

Solution to Question 9:

Fresh frozen plasma (FFP) is an acellular component and generally does not transmit viral infections.

Cytomegalovirus (CMV) infects >50% of the general population and is transmitted by infected passenger WBCs in transfused PRBCs or platelet components. Cellular components that are leukocyte-reduced have a decreased risk of transmitting CMV.

Infectious complications of blood transfusion include:

• Viral infections: Hepatitis C virus, hepatitis B virus, HIV, cytomegalovirus

• Bacterial contamination: sudden and fulminant onset of fever, chills, septic shock, and disseminated intravascular coagulation (DIC)

• Platelet concentrates stored at room temperature: gram-positive and coagulase-negative staphylococci

• PRBCs: Gram-negative bacteria like Yersinia, Pseudomonas, Serratia, Acinetobacter, and Escherichia species

• Parasites: Malaria, Babesiosis, Chaga's disease

• Others: Creutzfeldt–Jakob disease (CJD)

Solution to Question 10:

Febrile non-hemolytic transfusion reaction (FNHTR) is the most frequently occurring transfusion reaction.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Solution to Question 11:

This clinical scenario is suggestive of acute hemolytic transfusion reaction (AHTR).

Solution to Question 12:

This clinical scenario is suggestive of delayed hemolytic transfusion reaction (DHTR). It is due to an anamnestic antibody response. This refers to the rapid production of antibodies in the blood after exposure to an antigen to which the recipient had been previously sensitized.

Solution to Question 13:

Transfusion-related acute lung injury (TRALI) is the most common cause of transfusion-related fatalities.

TRALI is due to the presence of anti-HLA class II antibodies in the donor plasma. These antibodies bind recipient leukocytes, which aggregate in the pulmonary vasculature and release mediators that increase capillary permeability. Multiparous women are frequently implicated donors. Recipients with a history of liver disease, smoking, alcohol use are at a higher risk.

It occurs within 6 hours of transfusion and results in hypoxia and signs of non-cardiogenic pulmonary edema with bilateral interstitial infiltrates on chest X-ray. It is managed with supportive care.

Note: According to the latest edition of Harrison's, transfusion associated circulatory overload (TACO) is the leading cause of death followed by TRALI.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 14:

Massive transfusion is defined as transfusion of at least 10 units of red cells in 24 hours or 4 units in 1 hour. Hypocalcemia, not hypercalcemia, is a complication of massive blood transfusion.

Blood is anticoagulated with sodium citrate and citric acid. It binds to calcium and magnesium, leading to low levels of both. Citrate is also metabolized to bicarbonate, causing metabolic alkalosis.

Solution to Question 15:

The patient is being transfused blood and has complaints of backache, this is suggestive of an acute hemolytic transfusion reaction.

The first step is to stop transfusion immediately and then collect blood samples for testing.

Acute hemolytic transfusion reactions typically occur within the first 24 hours following a blood transfusion, primarily due to ABO incompatibility.

Sold by @itachibot

It presents with symptoms such as hypotension, tachypnea, tachycardia, fever (1-2°C increase in temperature), chills, chest and back pain, hemoglobinuria, and hemoglobinemia. In severe cases, disseminated intravascular coagulation (DIC), acute renal failure, shock, and death can occur.

Management of immune-mediated acute hemolytic transfusion reactions is to provide supportive care, administer vigorous hydration with isotonic saline and use diuretics to maintain urine output.

Solution to Question 16:

Among the given options, the metabolic abnormality seen in a patient who has received multiple blood transfusions is metabolic alkalosis.

This is because the excess sodium citrate (anticoagulant) in the transfused blood gets converted to bicarbonate leading to metabolic alkalosis. Each mmol of citrate generates 3 mEq of bicarbonate (a total of 23 mEq of bicarbonate due to each unit of blood).

Metabolic acidosis can also be seen in multiple blood transfusions as citrate is acidic, but metabolic alkalosis is more common.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

HIV / AIDS - Epidemiology and Clinical Features

Question 1:

What is the approximate time interval between HIV infection & manifestation of AIDS?

a) 7.5 years

b) 10 years

c) 12 years

d) 5 years

Question 2:

Which of the following case scenarios depict a mode of transmission that carries the highest risk of acquiring HIV?

a) A 23-year-old drug addict sharing syringes during intravenous drug abuse

b) A 15-year-old thalassemic adolescent undergoing blood transfusion

c) A 30-year-old man engaging in unprotected sexual intercourse with multiple partners

d) A 27-year-old nurse accidentally injuring herself while drawing blood

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which statement regarding HIV infection is true?

a) HIV-2 is more pathogenic than HIV -1

b) HIV-1 subtype C is the most common form worldwide

c) HIV-2 is prevalent worldwide

d) Donated blood is not screened for HIV- 2

Question 4:

A 47-year-old phlebotomist suffered a needle-stick injury while drawing blood from a patient with HIV. Which of the following mutations in him can cause a resistance to acquiring HIV infection?

a) CXCR4 2q21 mutation

Sold by @itachibot

b) HLA B*35

c) CCR5 delta 32 mutation

d) DC-SIGN

Question 5:

A 19-year-old woman presents to the casualty with a 24-hour history of fever, sore throat, and diffuse erythematous macular rash. Past history is positive for unprotected intercourse with a new partner. What is true about the likely diagnosis?

a) Incubation period is around 1 week

b) Weight loss is not seen

c) Virus is transmissible

d) Opportunistic infections do not occur

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

What is the correct WHO staging for an HIV-infected child with unexplained persistent parotid enlargement?

a) Stage 1

b) Stage 2

c) Stage 3

d) Stage 4

Question 7:

An HIV-positive individual presents with the following finding. What is the diagnosis & WHO staging?

Sold by @itachibot

a) Oral candidiasis & WHO stage 2

b) Oral hairy leukoplakia & WHO stage 3

c) Oral candidiasis & WHO stage 3

d) Oral hairy leukoplakia & WHO stage 2

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 8:

What is the most common opportunistic infection in HIV-infected individuals?

a) Tuberculosis

b) Disseminated candidiasis

c) Herpes zoster

d) Pneumocystis jiroveci pneumonia

Question 9:

A previously untreated HIV-positive patient is diagnosed with tuberculosis and treatment is initiated for both the diseases. Two weeks into treatment, he suddenly complains of high fever, severe malaise and worsening cough. What is the likely explanation?

a) Resistance to anti-tubercular therapy

b) Concurrent treatment with anti-retroviral therapy

c) Bacterial endotoxin release due to death of TB bacilli

d) Hypersensitivity reaction to isoniazid

Sold by @itachibot

Question 10:

An HIV-positive man presents with a high-grade fever. Examination reveals the following finding. CSF shows reduced glucose, increased protein, and increased leukocytes. What is the most common cause of this condition in such patients?

a) Streptococcus pneumoniae

b) Herpes simplex virus

c) Toxoplasma gondii

d) Cryptococcus neoformans

Question 11:

AIDS-defining malignancies include all except :

a) Kaposi's sarcoma

b) Melanoma

c) Invasive cervical carcinoma

d) Non hodgkins lymphoma

Question 12:

Sold by @itachibot

A 45-year-old HIV-positive woman presented with fever and malaise. She gives a history of bleeding from the following lesions. Warthin-Starry staining of a biopsy sample taken from the lesion showed which organism?

a) Bartonella henselae

b) Human Herpes virus - 8

c) Bartonella bacilliformis

d) Ebstein Barr Virus

Sold by
If you purchased this fro
you may have been scam

Question 13:

An HIV-positive woman was rushed to the ICU following an episode of generalized tonic-clonic seizures. There was no history of fever, night sweats or weight loss. MRI scan is shown below. What is the most likely diagnosis?

Sold by @itachibot



a) Cryptococcal meningitis

b) Primary central nervous system lymphoma

c) Cerebral toxoplasmosis

d) Progressive multifocal leukoencephalopathy

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 14:

All of the following are AIDS defining illnesses except:

a) Encephalopathy attributed to HIV

b) Invasive cervical cancer

c) Mycobacterium tuberculosis of any site

d) Oral candidiasis

Question 15:

An HIV-positive woman was referred to the ophthalmology OPD for decreased visual acuity. She described her vision as blurry with apparent flies floating in her field of vision. Fundoscopy revealed the following finding. What is the CD4+T cell count at which majority of such cases occur?

Sold by @itachibot

a) < 200/µL

b) < 50/µL

c) < 150/µL

d) < 250/µL

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

A unemployed HIV-positive patient, resident of a missionary home, presents with 2 weeks of fever, cough and night sweats. He was devastated after getting fired from his job because of his diagnosis and has since been non-compliant to cART. His CD4 count is <10/ µL. CT scan shows the following finding. What is the likely diagnosis?



Sold by @itachibot

a) Cytomegalovirus infection

b) Candidiasis

c) Mycobacterium avium complex infection

d) Pneumocystis jiroveci pneumonia

Question 17:

A 50-year-old man with HIV presents with lab reports showing: Low albumin, massive proteinuria. Renal ultrasound shows large, echogenic kidneys. Blood pressure is normal. Which of the following is the most likely cause?

a) Minimal change disease

b) Focal segmental glomerulosclerosis

c) Membranoproliferative glomerulonephritis

d) IgA nephropathy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

Which of the following is the most common form of heart disease associated with HIV?

a) Dilated cardiomyopathy

b) Thrombotic endocarditis

c) Coronary heart disease

d) Pericardial effusion

Answer Key

Question No.	Correct Option
1	b
2	b
3	b
4	c
5	c
6	b
7	c

Sold by @itachibot

8	a
9	b
10	d
11	b
12	a
13	c
14	d
15	b
16	c
17	b
18	c

Sold by @itachibot this from someone else, unscammed.

Detailed Explanations

Solution to Question 1:

The time interval between primary HIV infection and the manifestation of AIDS is 10 years.



After exposure to the HIV virus, there occurs primary infection following an incubation period of 2-4 weeks, which is usually symptomatic in more than 50% of cases, following 3-6 weeks there is acute HIV syndrome due to wide dissemination of the virus, characterized by fever, sore throat, diarrhea, lymphadenopathy, and rash.

This peak viremia drops as the immune response develops, to reach a plateau about 3 months later. Following this there is a prolonged period of clinical latency, during which infected individuals are asymptomatic, lasting for a median of 10 years.

Sold by @itachibot

Following this period, chronic persistent infection develops which is the hallmark of HIV disease called Acquired immunodeficiency syndrome (AIDS).

Long-term non-progressors: a small part of untreated HIV-infected people with CD4 counts within the reference range for >10 years.

Elite controllers: Some long-term non-progressors have undetectable viral loads.

Solution to Question 2:

Blood transfusion carries the highest risk of acquiring HIV.

Salient features of transmission of HIV:

• The highest risk of acquiring HIV from an infected source is via blood transfusion > needle sharing during drug abuse > receptive anal intercourse > needle-stick injury > vaginal & oral sexual intercourse.

• The most common mode of transmission worldwide is sexual transmission (heterosexual in India) > mother to child > least common modes are blood transfusion & needle-stick injury.

• Male to female transmission > female to male transmission.

• Viral load is highest in blood > semen > vaginal fluid > breastmilk & saliva and minimal to absent in tears, sweat & urine.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 3:

HIV-1 subtype C is the most common form worldwide.

Human Immunodeficiency Virus (HIV) has two serotypes HIV-1 and HIV- 2 based on differences in the env (envelope) gene.

HIV-1 is further divided into 4 distinct groups, group M (major)/N/O/P. The M group comprises nine subtypes A, B, C, D, F, G, H, J, and K.

• Subtype C is the most common form worldwide & is more associated with heterosexual transmission.

• Subtype B is more associated with spread through blood and homosexual contact.

Options A, C: HIV- 2 is less pathogenic than HIV-1 and is predominantly confined to Africa.

Option D: Donated blood is screened for both HIV -1 & HIV-2.

Solution to Question 4:

CCR 5 delta 32 mutation causes resistance to HIV infection. It is a mutated allele that causes a lack of surface expression of the CCR5 co-receptor resulting in blockade of HIV entry and

Sold by @itachibot

resistance to HIV infection.

Homozygous alleles cause complete resistance to HIV infection & heterozygous alleles cause partial resistance.

CXCR4 and CCR5 are chemokine receptors (co-receptors) present on the surface of CD4+ T-helper cells & macrophages that aid in the fusion and entry of the HIV virus.

DC-SIGN is a type of cell surface lectin receptor to which the HIV virus has the affinity to bind. The expression of this type of receptor on dendritic cells (antigen-presenting cells) enhances the infectivity of HIV.

Solution to Question 5:

The given clinical scenario is suggestive of acute HIV syndrome. The virus is transmissible during this early infection period.

It is characterized by acute burst of viremia and wide virus dissemination and seeding of lymphoid organs (e.g., gut-associated lymphoid tissue), culminating in the destruction of CD4+ cells.

The p24 antigen capture assay is used as a screening test for HIV infection in patients suspected of having the acute HIV syndrome.

Option A: The incubation period for acute HIV syndrome is around 3 to 6 weeks.

Option B: The acute HIV syndrome is typical of an acute viral syndrome and is very similar to acute infectious mononucleosis. The clinical findings include fever, sore throat, and lymphadenopathy. Rash, headache, arthralgia, and anorexia/weight loss can also be seen.

Option D: Opportunistic infections can occur during this stage of infection, reflecting the immunodeficiency that results from reduced numbers of CD4+T cells.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 6:

Unexplained persistent parotid enlargement in an HIV-infected child is categorized as stage 2 of HIV disease according to WHO staging.

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you have been scammed.

Solution to Question 7:

Oral candidiasis (thrush) is categorized as WHO stage 3.

Oral candidiasis and oral hairy leukoplakia are oral lesions categorized as WHO clinical stage 3. They generally occur in patients with CD4+ T cell counts of <300/µL.

The image given below shows a patient with oral hairy leukoplakia.

	Oral candidiasis	Oral hairy leukoplakia
Cause	Candida albicans	Epstein-Barr virus
Appearance	White, plaque-like lesions with cheesy exudate, on an erythematous mucosa	White, frond-like lesions with hair-like growths from folds in the patches
Location	Posterior oropharynx	Lateral borders of tongue
Clinical features	Typically painful, can be easily brushed off	Typically painless, cannot be brushed
Investigation	KOH preparation demonstrates budding yeasts, hyphae, or pseudohyphae	Despite its name, it is not considered premalignant.
Treatment	Antifungals	Antivirals

Sold by @itachibot

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 8:

Tuberculosisis the most common opportunistic infection in HIV-infected individuals globally and in India.

In patients with relatively high CD4+ T cell counts, the typical pattern of pulmonary reactivation occurs (60–80%). In patients with lower CD4+ T cell counts, disseminated disease is seen.

Primary prophylaxis: Isoniazid and pyridoxine for 9 months for HIV-infected individuals meeting the following conditions:

• PPD skin-test reaction of >5 mm or,

• Positive IFN-γ release assay or,

• Close household contacts of persons with active tuberculosis

Solution to Question 9:

Concurrent treatment with anti-retroviral therapy is the most likely explanation in this case resulting in Immune Reconstitution Inflammatory Syndrome (IRIS). Its characteristics are:

• Temporary paradoxical exacerbation of symptoms (lymphadenopathy, fever or worsening of pulmonary infiltrations, pleural effusion) and radiographic manifestations of TB after starting ART. This occurs as a result of immune reconstitution due to simultaneous administration of ART and ATT.

• It is seen weeks to months following the initiation of combination antiretroviral therapy (cART).

• It is most common in previously untreated patients starting therapy with a CD4+ T cell count <50/µL who experience a sudden drop in viral load.

Sold by @itachibot

• The earlier ART is started and the lower the baseline CD4+ T-cell count, the greater the risk of IRIS.

• In severe cases, the use of immunosuppressive drugs such as glucocorticoids may be required to blunt the inflammatory component of IRIS while specific antimicrobial therapy takes effect.

Note: Hypersensitivity to isoniazid is extremely rare and the presentation includes maculopapular rash.

Solution to Question 10:

The given clinical scenario, along with a positive Kernig's sign, is suggestive of meningitis. Cryptococcus is the leading infectious cause of meningitis in patients with AIDS. It generally occurs in untreated patients, with CD4+ T cell counts <100/µL.

Most patients present with a picture of subacute meningoencephalitis. The CSF profile may show modest elevations in WBC or protein levels and decrease in glucose. The CSF opening pressure is usually elevated (yeast colonies obstruct the free flow of CSF causing increased CSF pressure).

The diagnosis of cryptococcal meningitis is made by identification of Cryptococcus neoformans yeast cells in CSF with india ink stain or by the detection of cryptococcal antigen using latex agglutination (CrAg).

Prophylaxis: Oral fluconazole when CD4+ T cell count is <100/µL.

Treatment:

• Induction - IV or liposomal amphotericin B with flucytosine for at least 2 weeks, continuing with amphotericin alone ideally until the CSF culture turns negative.

• Maintenance - Oral fluconazole until the CD4+ T cell count has increased to >200 cells/µL for 6 months in response to cART.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 11:

Melanoma is not an AIDS-defining malignancy.

AIDS-defining malignancies are those cancers that a human immunodeficiency virus (HIV) infected person is at a high risk of developing. When a HIV-infected person develops one of the AIDS-defining malignancies it suggests that he has AIDS.

AIDS-defining malignancies include the following:

• Kaposi sarcoma - It is a multicentric neoplasm of vascular origin and has a strong association with HHV - 8.

• Non-Hodgkin's lymphoma

• Invasive cervical cancer

Given below is the image of kaposi sarcoma:

Sold by @itachibot

Note : The FDA has approved pomalidomide, a thalidomide analogue to be included in treating adult patients with AIDS-related Kaposi's sarcoma after failure of highly active antiretroviral therapy and Kaposi's sarcoma in adult patients who are HIV-negative.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 12:

The given clinical scenario is suggestive of bacillary angiomatosis in an immunocompromised patient. Warthin-Starry stain would show Bartonella henselae as the causative organism in these skin lesions.

Bacillary angiomatosis is an infectious vascular proliferative lesion presenting as reddish-purple nodules similar to KS lesions (caused by HHV-8). The diagnosis is confirmed with a biopsy. The drug of choice for treatment is erythromycin.

The image given below shows the Warthin-Starry stained lesion showing tangled pleomorphic bacilli.

Sold by @itachibot



Option B: HHV 8 is strongly associated with Kaposi sarcoma.

Option C: Bartonella bacilliformis causes Oroya fever.

Option D: EBV causes infectious mononucleosis.

Note: Bartonella henselae causes cat-scratch disease, a necrotizing granulomatous disorder of lymph nodes, in immunocompetent hosts.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 13:

The given clinical scenario is suggestive of cerebral toxoplasmosis. MRI scan shows multiple ring-enhancing lesions.

Cerebral toxoplasmosis is one of the most common AIDS-defining opportunistic CNS infections and usually occurs in patients with CD4+ T cell counts <200/µL. It is usually a reactivation of latent tissue cysts.

It usually has multiple lesions in multiple locations which are associated with inflammation and central necrosis, and hence show ring enhancement on contrast MRI. Sometimes an eccentric nodular area of enhancement is seen within the ring; this so-called eccentric target sign is typical of cerebral toxoplasmosis. Surrounding edema is also present.

Treatment includes sulfadiazine/pyrimethamine and leucovorin for a minimum of 4–6 weeks.

Primary prophylaxis: It is given to patients with IgG antibody to toxoplasma and CD4+ T cell count <100/µL. A single double strength tablet of trimethoprim/sulfamethoxazole (TMP/SMX) is given.

Secondary prophyaxis/maintanace therapy: It is given when CD4+ T cell counts <200 cells/µL. Sulfadiazine/pyrimethamine and leucovorin is given. It is discontinued if the CD4+ T cell counts are maintained >200/µL for 6 months.

Option A: Cryptococcal meningitis presents with fever, nausea, vomiting, altered mental status, headache, and meningeal signs. Seizures and focal neurologic deficits are rarely seen. MRI

Sold by @itachibot

findings include dilated perivascular spaces and basal ganglia pseudocysts.

Option B: Primary CNS lymphoma also presents with seizures and focal neurological deficit. Symptoms of fever, night sweats and weight loss are present in 80% of the patients. On imaging, it is identified as a CT hyperdense enhancing supratentorial mass, with MRI T1 hypointense, T2 iso- to hypointense, vivid homogeneous enhancement as well as restricted diffusion.

Option C: Progressive multifocal leukoencephalopathy, caused by JC virus, is characterized by demyelination in subcortical white matter reflected on MRI as multiple, non-enhancing white matter lesions.

Solution to Question 14:

Oral candidiasis, although seen in AIDS patients, is not an AIDS-defining illness. Candidiasis of the esophagus, trachea, bronchi, and lungs are AIDS-defining illnesses.

AIDS-defining illnesses are specific opportunistic illnesses in patients with AIDS, which signify an advanced stage of illness (Stage 3).

A case of HIV can be classified into five stages as per current CDC guidelines (0, 1, 2, 3, and unknown).

Stage	Criteria	CD4+ count (>5 years)
0	A negative HIV test within 6 months of first diagnosis	NA
1	Based on CD4+ count	>=500
2	Based on CD4+ count	200-499
3	AIDS-defining opportunistic illness	<200
unknown	Unavailable CD4+ counts	unknown

Solution to Question 15:

The given clinical scenario, along with the pizza pie appearance on fundoscopy, is suggestive of cytomegalovirus (CMV) retinitis. The majority of such cases occur in HIV-infected patients with a CD4+T cell count <50/µL.

CMV retinitis is a necrotic retinal infection that results in a painless irreversible vision loss. Patients may complain of blurred vision, floaters (described by patient as flies or spiders in the field of vision), and scintillations. It is usually bilateral and is characterized by perivascular hemorrhage and exudates in the retina (pizza pie appearance or scrambled eggs and ketchup appearance).

If you bought this from someone else, you got scammed. Sold by @itachibot

Sold by @itachibot

Treatment:

• Induction - IV ganciclovir (3 weeks)

• Maintenance - Oral valganciclovir (till CD4 count > 100/µl for 6months)

Solution to Question 16:

The given clinical scenario is suggestive of mycobacterium avium complex (MAC) infection. It is an AIDS-defining opportunistic infection, occurring predominantly at CD4+ T cell counts of <50/µL.

The most common presentation is disseminated disease with fever, weight loss, and night sweats. The diagnosis is made by culture of blood or involved tissue.

CT scan shows peribronchial inflammation of the infected lung giving rise to a tree in a bud appearance (small round structures attached to linear structures) indicated by the arrowhead. In severe cases, there might be bronchiectasis and cavity formation.

Treatment consists of clarithromycin / azithromycin with ethambutol/ rifabutin. Primary prophylaxis with azithromycin is indicated in patients with CD4+ T cell counts <50/µL.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

This patient with HIV presents with massive proteinuria and hypoalbuminemia. The most likely cause is HIV-1-associated nephropathy (HIVAN) - a form of focal segmental glomerulosclerosis (FSGS).

Proteinuria is the hallmark of this disorder. Edema and hypertension are rare. A definitive diagnosis is obtained through renal biopsy.

Solution to Question 18:

The most common form of heart disease associated with HIV infection is coronary heart disease.

In patients with HIV infection, cardiovascular disease may be a direct consequence of HIV infection, a complication of anti-retroviral therapy or due to classical risk factors like smoking.

Exposure to HIV protease inhibitors and reverse transcriptase inhibitors have been associated with an increase in total cholesterol and/or risk of MI (Myocardial infarction).

Dilated cardiomyopathy associated with congestive heart failure (CHF) is referred to as HIV-associated cardiomyopathy. This generally occurs as a late complication of HIV infection and, histologically, displays elements of myocarditis.

Sold by @itachibot

HIV / AIDS - Management and Prophylaxis

Question 1:

Anti-retroviral therapy in HIV-infected individuals should be initiated at _____

a) CD4 count <200 cells/ul and/or WHO clinical stage 2/3

b) CD4 count <350 cells/ul and/or WHO clinical stage 3/4

c) CD4 count <250 cells/ul and/or WHO clinical stage 4

d) Any CD4 count regardless of WHO clinical stage

Question 2:

Which of the following is true about fourth-generation serological testing for HIV?

a) It is used to confirm the diagnosis of HIV

b) It detects HIV antibodies and antigen

c) It has a lower sensitivity compared to first-generation HIV testing

d) It cannot be used to detect HIV-2 antibodies

Question 3:

A 46-year-old man, who is a chronic alcoholic, presented with acute dull and boring abdominal pain. His history is significant for ongoing treatment with ART. Serum lipase levels were noted to be grossly elevated. What is the most likely causative drug?

a) Nevirapine

b) Didanosine

c) Emtricitabine

d) Stavudine

Question 4:

An HIV-positive patient is upset because he has a high viral load despite good compliance with ART. The physician decides to start him on a parenteral drug. Which of the following drugs will be helpful for this patient?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Emtricitabine

b) Elvitegravir

c) Enfuvirtide

d) Efavirenz

Question 5:

An HIV-positive patient has the following serology results. What is the management of this patient?

a) Tenofovir + emtricitabine + abacavir

b) Tenofovir + lamivudine + dolutegravir

c) Lamivudine + emtricitabine + nevirapine

d) Tenofovir + lamivudine + efavirenz

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which of the following antiretroviral drugs is not used in HIV & hepatitis B co-infection?

a) Tenofovir

b) Abacavir

c) Lamivudine

d) Emtricitabine

Question 7:

Screening of patients for HLA- B5701 is recommended before initiation of which of the following antiretroviral drugs?

a) Tenofovir

b) Abacavir

c) Zidovudine

d) Emtricitabine

Question 8:

Sold by @itachibot

Evaluation of a patient with uncontrolled type 2 diabetes mellitus and hypertension has revealed moderately increased albuminuria. He has recently been diagnosed with HIV infection. Combination ART preparations including which of the following drugs should be avoided?

a) Tenofovir

b) Lamivudine

c) Emtricitabine

d) Dolutegravir

Question 9:

A 36-year-old HIV-positive man on cART is presented with the following finding. What is the most likely cause?

a) Abacavir

b) Entecavir

c) Nelfinavir

d) Adefovir

Question 10:

A 54-year-old HIV-positive woman has a Hb of 8gm/dl. Combination ART preparations including which of the following drugs should be avoided?

Sold by @itachibot

a) Zidovudine

b) Tenofovir

c) Raltegravir

d) Emtricitabine

Question 11:

During his routine counselling, a war veteran on cART for the past 2 weeks tells his therapist that he wishes to discontinue his HIV treatment. He explains that since starting them, he has been having extreme dizziness and intense nightmares leading to repeated wakening in the night. What is the likely cause?

a) Tenofovir

b) Dolutegravir

c) Efavirenz

d) Lamivudine

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 12:

A 38-year-old man is tested for HIV and the following band pattern is seen on a standard western blot. Protein gp36 is present on the immunoblotting assay. To which class of antiretroviral drug would he be intrinsically resistant?

a) Nucleoside reverse transcriptase inhibitors

b) Nucleotide reverse transcriptase inhibitors

c) Non-nucleoside reverse transcriptase inhibitors

d) Integrase strand transfer inhibitors

Question 13:

An individual positive for HIV-1 infection was started on a combination ART regimen of tenofovir–emtricitabine–efavirenz. After 6 months, failure of treatment was established and attributed to efavirenz resistance. Which drug of the same class can be used instead?

a) Delavirdine

b) Nevirapine

c) Nelfinavir

d) Etravirine

Sold by @itachibot

Question 14:

A woman is diagnosed to be positive for HIV in her first trimester. What is the appropriate management?

a) Start ART immediately and continue till 6 weeks after delivery

b) Start ART after 1st trimester and continue life-long

c) Start ART immediately and continue life-long

d) Start ART after 1st trimester and continue till 6 weeks after delivery

Question 15:

A newly diagnosed HIV patient presented with fever, cough, and hemoptysis. His sputum sample showed acid-fast bacilli. What is the appropriate management for this patient?

a) Start ART if CD4<200

b) Start ART and then ATT after 2 weeks

c) Start ART and then ART after 2 weeks

d) Start ATT and ART simultaneously

Question 16:

What is the diagnostic test of choice for HIV in a baby born to an HIV-infected mother?

a) HIV DNA PCR

b) Cord blood ELISA

c) Western blot

d) Third generation ELISA

Question 17:

A 29-year-old HIV-positive woman presents to the OPD with fever, cough, and hemoptysis. Chest X-ray is given below. What is the diagnostic test for this condition in this patient?

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot



a) Cartridge based nucleic acid amiplification test

b) Sputum microscopy & AFB staining

c) Interferon gamma release assay

d) Bronchoalveolar lavage fluid cytology

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 18:

A 52-year-old man was admitted for multiple episodes of seizures following 7 months of neurocognitive decline. He also had weight loss, paranoia, and delusions. Clinical workup revealed HIV infection along with the following findings on MRI. What is the ideal timing of steroid therapy in this condition?



Sold by @itachibot

a) 1 day before performing a biopsy

b) Just before obtaining the sample for a biopsy

c) Only after obtaining a biopsy

d) There is no specific ideal timing

Question 19:

What is the WHO recommendation regarding the initiation of cotrimoxazole prophylaxis?

a) Any CD4 count or WHO clinical stage

b) CD4 count <500 and/or WHO stage 3/4

c) CD4 count <350 and/or WHO stage 3/4

d) CD4 count <150 and/or WHO stage 3/4

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Question 20:

A cachexic man presented with a non-productive cough and shortness of breath. He was diagnosed with HIV 6 months ago but is non-adherent to ART. Serum LDH is 350 U/L. Giemsa-stained smears are seen below. At what CD4 count should the prophylaxis be started according to the CDC recommendation to prevent this disease?



a) <100/mm3

b) <150/mm3

c) <400/mm3

Sold by @itachibot

d) <200/mm3

Question 21:

You suffer a bleeding puncture wound on your finger while assisting with surgery on a patient infected with HIV. What regimen should you take in this situation?

a) Zidovudine + Lamivudine for 4 weeks

b) Lamivudine + Tenofovir + Efavirenz for 4 weeks

c) Lamivudine + Tenofovir + Dolutegravir for 4 weeks

d) Tenofovir + Lamivudine + Nevirapine for 4 weeks

If you bought this from someone else, it was scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	c
5	b
6	b
7	b
8	a
9	c
10	a
11	c
12	c
13	d
14	c
15	c
16	a
17	a
18	c
19	c
20	d

Sold by @itachibot

21	c

Detailed Explanations

Solution to Question 1:

Antiretroviral therapy (ART) is recommended for all HIV patients, regardless of CD4 T lymphocyte cell count or clinical stage.

This is done to reduce the morbidity and mortality associated with HIV infection. A secondary goal of ART is to reduce the risk of HIV transmission.

Solution to Question 2:

The 4th generation HIV test combines the detection of antibodies against HIV-1 & HIV-2 with detection of the core p24 antigen of HIV-1. Inclusion of p24 antigen capture allows detection of HIV-1 infection even before seroconversion i.e., presence of antibodies in the blood.

The approximate time to positivity following infection is 15 to 20 days. Consequently, the window period between infection and detection of the infection is shortened.

Fourth-generation HIV testing is highly sensitive (99.9%) and is the CDC-recommended initial screening test for HIV. No further testing is required for specimens that are non-reactive on ELISA screening.

Generation	Serological test
First generation	IgG antibody
Second generation	IgG antibody
Third generation	IgM and IgG antibody
Fourth generation	IgM and IgG antibody and p 24 antigen

Solution to Question 3:

Didanosine is the most likely causative drug. The clinical scenario, suggestive of pancreatitis, is a dose-limiting side effect of didanosine and occurs more frequently in alcoholic patients and those with hypertriglyceridemia.

Option A: Nevirapine side-effects include severe liver problems, skin rash, and skin reactions.

Option C: Emtricitabine is contraindicated in pregnancy and young children.

Sold by @itachibot
If you purchased this from someone else, they have been scammed.

Sold by @itachibot

Option D: Stavudine can cause peripheral neuropathy.

Solution to Question 4:

Enfuvirtide is the antiretroviral drug administered parenterally, used only in the treatment of those who have evidence of HIV replication i.e., high viral load, despite ongoing antiretroviral therapy.

Enfuvirtide prevents viral fusion and entry by binding to the gp41 subunit of the HIV-1 envelope. It is not active against HIV-2. It is administered as a twice-a-day subcutaneous injection.

Note: Zidovudine is also available as a parenteral formulation.

Solution to Question 5:

The given clinical scenario is suggestive of an acute infection with hepatitis B virus. The preferred regimen in the management of HIV-1 and hepatitis B co-infection is tenofovir + lamivudine + dolutegravir.

This regimen is also the first-line management in adults, adolescents, pregnant women, and HIV-TB co-infection.

Antivirals that are effective against hepatitis C infection are sofosbuvir, grazoprevir and glecaprevir.

*Note: Pearl attached below has been updated according to the NACO ART guidelines 2021. Efavirenz has been replaced by Dolutegravir as per the new guidelines.

Solution to Question 6:

Abacavir is not used in HIV and hepatitis-B co-infection. It is used in the first-line ART regimen for patients with abnormal serum creatinine values.

Nucleoside reverse transcriptase inhibitor (NRTI) drugs like lamivudine, its closely related drug emtricitabine, and the nucleotide reverse transcriptase inhibitor (NRTI) tenofovir have activity against hepatitis B virus. These drugs are preferred in combination with other antiretrovirals for the treatment of HIV & HBV co-infections.

Solution to Question 7:

Screening of patients for HLA-B5701 is recommended before initiation of abacavir since it is associated with hypersensitivity.

The hypersensitivity syndrome includes fever, maculopapular rash, abdominal pain, gastrointestinal problems, malaise and fatigue. Hence, abacavir should only be used as a last

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

resort and with close monitoring in patients who are HLA-B5701-positive.

Solution to Question 8:

The given clinical scenerio is suggestive of early stages of diabetic and hypertensive kidney disease. So combinaton antiretroviral therapy (cART) preparations including tenfovir should not be prescribed to this patient.

Tenofovir is primarily eliminated by the kidneys and is contraindicated in patients with renal impairment.

Solution to Question 9:

The given clinical scenario is suggestive of fat redistribution resulting in the classic buffalo hump appearance. Nelfinavir is the most likely cause.

It is a protease inhibitor and its chronic administration can result in fat redistribution and metabolic syndrome.

Note: Atazanavir has the least propensity to cause lipodystrophy and metabolic syndrome.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Solution to Question 10:

Zidovudine should be avoided in this patient because the primary toxicity of zidovudine is bone marrow suppression. It can lead to anemia and neutropenia, which may require transfusions.

Solution to Question 11:

The patient is most likely prescribed a cART regimen including efavirenz.

Many patients treated with efavirenz note a feeling of light-headedness and dizziness following the initiation of therapy. They also complain of insomnia, vivid dreams and nightmares.

These symptoms are self-limiting and tend to disappear gradually after several weeks of therapy. If the sleep problems are persistent, stopping the drug is the only solution.

It is recommended to take efavirenz before bedtime to minimize side effects.

The CNS/ psychiatric adverse effects seen with the use of efavirenz are:

- Frank psychosis (depression, hallucinations, and/or mania)
- Impaired concentration
- Dysphoria
- Dizziness

Sold by @itachibot

• Vivid dreams

• Insomnia

*Note: Pearl attached below has been updated according to the NACO ART guidelines 2021. Efavirenz has been replaced by Dolutegravir as per the new guidelines.

Solution to Question 12:

The given clinical scenario is suggestive of HIV-2 infection. It is intrinsically resistant to non-nucleoside reverse transcriptase inhibitors (NNRTI) and also to fusion inhibitor enfuvirtide.

Western blot has a high specificity. It detects specific antibodies against multiple proteins of all 3 major HIV genes - gag, pol, and env.

Persons with HIV-2 will often have an indeterminate western blot pattern showing the presence of Gag bands and Pol bands, but an absence of Env bands. This pattern occurs because HIV-1 and HIV-2 share little similarity in the env gene. Moreover, the presence of gp36 points towards a diagnosis of HIV-2.

In India, NACO provides western blot confirmatory testing only to resolve indeterminate results on previous serological testing and not for routine confirmation after a positive result on screening tests.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 13:

Etravirine can be used instead of efavirenz.

Etravirine is a non-nucleoside reverse transcriptase inhibitor (NNRTI) currently approved for the treatment of HIV infection in combination with other agents in treatment-experienced patients.

Options A, B: Etravirine is active against strains of HIV that are resistant to other NNRTIs such as delavirdine, nevirapine, and efavirenz which all exhibit cross-resistance. Rilpivirine, an NNRTI, is also effective across a broad range of NNRTI-resistant viruses but shares cross-resistance with etravirine.

Option C: Nelfinavir is a protease inhibitor and hence, does not belong to the same class.

Solution to Question 14:

ART should be initiated immediately and continued lifelong in all pregnant and breastfeeding women with HIV. This is regardless of their WHO clinical stage and CD4 cell count.

*Note: Pearl attached below has been updated according to the NACO ART guidelines 2021. Efavirenz has been replaced by Dolutegravir as per the new guidelines.

Sold by @itachibot

Solution to Question 15:

In a patient with HIV and TB start ATT first and then start ART as soon as ATT is tolerated (between 2 weeks and 2 months) by the patient. This is done in order to avoid immune reconstitution inflammatory syndrome (IRIS).

Immune Reconstitution Inflammatory Syndrome (IRIS):

Patients with HIV-related TB may experience a temporary paradoxical exacerbation of symptoms and radiographic manifestations of TB after starting ATT. This occurs as a result of immune reconstitution due to simultaneous administration of ART and ATT.

It is characterized by high fever, lymphadenopathy, expanding intra-thoracic lesions, and worsening of chest X-ray findings. Diagnosis should be made only after ruling out treatment failure and other infections. For severe cases, prednisone 1-2mg/kg is used for 1-2 weeks and then gradually tapered.

However, in patients with CD4 <50/µL, the benefits of administering ART immediately after initiating ATT outweigh the risks of IRIS, and therefore ART should be started as early as possible.

Note: As per RNTCP guidelines, CBNAAT is the preferred diagnostic technique for testing TB in PLHIV (people living with HIV). Smear microscopy has poor sensitivity in detecting TB in PLHIV due to fewer organisms in sputum, and hence is not preferred.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

In a baby born to an HIV-infected mother, HIV infection is diagnosed using HIV DNA PCR. NACO recommends the use of a qualitative HIV-1 DNA PCR (NAT) in children aged <18 months for Early Infant Diagnosis (EID).

Antibody assays (e.g., ELISA) are not reliable in infants and young children because of the persistence of transplacentally acquired maternal antibodies. For this reason, HIV virologic testing must be performed using tests that detect HIV DNA or RNA to diagnose HIV infection in children.

Scheduled repeated testing is necessary because test sensitivity increases with time from HIV virus exposure and very recent infection will not be detectable. The cord blood should not be used for testing because of the possibility of contamination of the sample with maternal blood.

Solution to Question 17:

The given clinical scenario is suggestive of tuberculosis. Cartridge-based nucleic acid amiplification test (CBNAAT) is the diagnostic test for TB in HIV positive patients.

CBNAAT is a rapid molecular test that detects the DNA of TB bacteria in HIV-positive individuals. It uses sputum or any other biological specimen (except blood and blood-contaminated specimens). It can give a result in less than 2 hours. It can also detect the genetic mutations associated with resistance to the drug rifampicin.

Sold by @itachibot

Sputum microscopy has poor sensitivity for detecting TB in HIV-infected individuals due to fewer organisms in sputum.

Solution to Question 18:

The given clinical scenario is suggestive of primary CNS lymphoma (PCNSL). In this condition, glucocorticoids should be administered only after a biopsy has been obtained. This is because steroids have a cytolytic effect on lymphoma cells and hence may lead to nondiagnostic tissue.

On CT, PCNSL is identified as a hyperdense enhancing supratentorial mass. On MRI, vivid homogeneous enhancement, as well as restricted diffusion, is seen.

PCNSL is a type of non-Hodgkin's lymphoma. The workup usually includes:

- HIV testing
- CSF analysis
- MRI spine
- PET/CT to assess the extent of the disease
- Bone marrow biopsy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 19:

WHO recommends cotrimoxazole prophylaxis at CD4 count <350 and/or WHO stage 3/4.

Only in settings where malaria or severe bacterial infections are highly prevalent, co-trimoxazole prophylaxis should be initiated regardless of WHO clinical stage or CD4 counts.

Prophylaxis in HIV patients:

Organism	CD4 count below which prophylaxis is started	Drug given
Pneumocystis jirovecii	<200/mm3(Acc to CDC)<350 /mm3(Acc to WHO and NACO)	TMP-SMX (Cotrimoxazole)
Mycobacterium-avium complex	<50/mm3	Azithromycin
Toxoplasma gondii	<100/mm3	TMP-SMX (Cotrimoxazole)

Solution to Question 20:

The given clinical scenario is suggestive of pneumocystis pneumonia (PCP). The prophylaxis for PCP is started at <200/mm3 CD4 count according to the CDC recommendation. In PCP LDH is

Sold by @itachibot

elevated and smear stained with Giemsa stain shows trophic forms with dot-like nuclei and pale blue cytoplasm as well as intracystic bodies of Pneumocystis jiroveci.

Primary prophylaxis of PCP:

Indications	CD4+ T cell count <200/uL (CD4 cell %<15)Recent oroph aryngeal candidiasis and prio r PCPAny patient with unexp lained fever for > 2 weeks.
Drug of choi ce	Trimethoprim/sulfamethoxaz ole(TMP/SMX), one double-st rength tablet daily.Alternativ es- Dapsone/Pyrimethamine + Leucovorin or Atovaqone o r Aerosolised Pentamidine

Note : TMP-SMX also provides protection against toxoplasmosis and some bacterial respiratory pathogens. The prophylaxis is discontinued in those patients treated with ART who maintain good suppression of HIV (<50 copies/mL) and CD4+ T-cell counts >200/ µL for at least 3 months.

Sold by @itachibot If you purchased this from someone else, you may have been scammed!

Solution to Question 21:

Recommended post-exposure prophylaxis (PEP) is a three-drug regimen that includes lamivudine, tenofovir, and dolutegravir.

A regimen for HIV post-exposure prophylaxis with two ARV drugs is effective, but three drugs are preferred. The PEP must be initiated ideally within 2 hours and certainly within 72 hours.



Sold by @itachibot

COVID 19 - Epidemiology and Clinical Features

Question 1:

Coronaviruses are _____

a) Naked helically symmetrical viruses

b) Enveloped icosahedrally symmetrical viruses

c) Enveloped helically symmetrical viruses

d) Variably enveloped viruses with a complex symmetry

Question 2:

An elderly diabetic man presents with difficulty breathing, cough, and fever. He came back after visiting his son in New York, 10 days ago. His SpO2 is 89%. HRCT shows bilateral, multilobar ground-glass opacification. What is the receptor used by the likely causative organism to gain entry into the cells?

a) Angiotensin-converting enzyme 1

b) Angiotensin-converting enzyme 2

c) Dipeptidylpeptidase 2

d) Dipeptidylpeptidase 4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which viral structure determines the infectious nature and host specificity of SARS-CoV-2?

Sold by @itachibot



©MARROW

a) 2

b) 1

c) 4

d) 3

Question 4:

Which of the following scenarios is unlikely for transmission of SARS-CoV-2?

a) A gym trainer transmitted the virus to his client following a sudden, involuntary cough.

b) A librarian got infected from a reader through the share and exchange of books.

c) A resident transmitted the virus to his neighbour who entered the same apartment lift after a few hours.

d) A thalassemic girl got infected following routine blood transfusion.

Question 5:

A relative of a COVID-19 positive man, who came in his contact yesterday, reported to the flu clinic. Within how many days is he most likely to develop the disease?

a) 7-14 days

b) 2-3 weeks

c) 1-3 weeks

d) 1-14 days

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 6:

Covid 19 outbreak is a type of which of the following kinds of the epidemic?

a) Propagated epidemic

b) Point source, single exposure

c) Point source, multiple exposures

d) Slow epidemic

Question 7:

What is not included in the definition of a suspect case of SARS-CoV-2?

a) Myalgia

b) Coryza

c) Altered mental status

d) Convulsions

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

Which of the following is not considered a probable case of SARS-CoV-2?

a) An asymptomatic contact of a probable case with a positive COVID-19 rapid antigen test.

b) A person with acute onset fever and cough who is a contact of a confirmed case.

c) A suspect case with positive chest radiological findings suggestive of COVID-19 disease.

d) A person with recent onset of anosmia or ageusia in the absence of any other identified cause.

Question 9:

Which of the following people would you not classify as a high risk contact of SARS-CoV2?

a) A family member who lives in the same household as the patient

b) A doctor who examined the patient with a surgical mask and gloves

c) A passenger in the same bus as a symptomatic patient for 6 hours and the patient later tested positive

d) A sanitary worker who cleaned the linens of the patient

Sold by @itachibot

Question 10:

You would classify all of the following patients as having major risk factors for severe COVID-19 except:

a) A 19 year old female undergoing dialysis for CKD

b) A 49 year old male who has recently undergone liver transplantation

c) A 42 year old man who had malaria 2 months ago

d) A 65 year old male with no comorbidities

Question 11:

A woman presents with breathlessness and hemoptysis since yesterday. Examination findings are given below. CT scan shows localised ground-glass opacity in the left posterior lobe of the lung. What is the likely diagnosis?

a) Tuberculosis

b) Necrotising pneumonia

c) Viral pneumonitis

d) Invasive aspergillosis

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 12:

What is the most common lung pathology seen during the autopsy of COVID-19 patients?

a) Pulmonary infarction

b) Endothelial injury

c) Diffuse alveolar damage

d) Fibrin clots

Question 13:

You are posted as an intern in casualty. An elderly 60-year-old man is brought to you with complaints of chest pain. He has tested positive for COVID. Which of the following features will prompt you to get this patient admitted to dedicated COVID Health Centre?

a) A & B only

Sold by @itachibot

b) A, B, C, & E only

c) A, B & C only

d) A, B, C, D & E

Answer Key

Question No.	Correct Option
1	c
2	b
3	b
4	d
5	d
6	a
7	d
8	a
9	c
10	c
11	c
12	c
13	b

Detailed Explanations

Solution to Question 1:

Coronaviruses are enveloped helically symmetrical viruses.

Being enveloped contributes to their lipid solvent susceptibility. This is the reason alcohol-based hand sanitizers and soaps are used. The lipid membrane dissolution and protein denaturation by ethanol lead to membrane disruption and inhibition of metabolism.

The coronaviridiae have a single unsegmented ss (+) RNA, which is the largest genome among RNA viruses. They are about 120- 160 nm in size and contain a helical nucleocapsid. A total of seven human coronaviruses have been identified:

The image given below shows the coronavirus.

Sold by @itachibot

Name	Illness
SARS-CoV-2	COVID-19
SARS-CoV	Severe acute respiratory syndrome (SARS)
MERS-CoV	Middle East respiratory syndrome (MERS)
HCoV-NL63	Usually mild respiratory illness
HCoV-229E	
HCoV-OC43	
HKU1	



Solution to Question 2:

The given clinical vignette showing breathlessness, fever, and cough with a low oxygen saturation and ground-glass opacities on HRCT is indicative of COVID-19 caused by the SARS-CoV-2 virus. Angiotensin-converting enzyme 2 (ACE-2) is the specific functional receptor for SARS-CoV-2 in human cells.

ACE2 is expressed in nearly all human organs in varying degrees. In the respiratory system, ACE2 is mainly expressed by the type 2 alveolar epithelial cells in the highest quantity. Hence, the primary target of SARS-CoV-2 is type 2 alveolar cells.

The functional receptor for SARS-CoV (SARS 2002) is also ACE 2.

Note: Dipeptidylpeptidase 4 is the functional receptor for MERS-CoV (MERS 2012).

Sold by @itachibot

Solution to Question 3:

Spike protein (marked as 1 in the image) determines the infectious nature and host specificity of SARS-CoV-2.

The spike protein also referred to as the CoV S protein is one of the key components determining virus virulence, tissue tropism, and host range. The coronavirus spike protein mediates coronavirus entry into host cells. Each spike protein has two distinct subunits S1 and S2.

• S1 subunit is responsible for binding to the specific receptor (ACE 2) on the host cell surface facilitating viral attachment to the surface of target cells.

• S2 subunit drives the fusion of the viral and host cell membranes.

The host protease enzymes, mainly transmembrane protease serine-type 2 (TMPRSS2) cleave the spike protein. This activates the spike protein to induce virus-cell membrane fusion at the cell surface and allows the entry of coronaviruses into the host cell.



Solution to Question 4:

Blood transfusion is not a mode of transmission of SARS-CoV-2.

Option A: Droplet transmission

• A type of direct transmission

• Occurs when someone coughs, sneezes, speaks, or spits

• Virus-containing droplets from the infected person travel shortly in the air directly contact another person's conjunctiva, oral or nasal mucosa

• Limited to about 6 feet distance between the source and the host

Sold by @itachibot

• Particle size is 5-10 µm in diameter

Option B: Fomite transmission

• A type of indirect transmission

• Objects other than water or food contaminated by infectious agents - clothes, towels, linen, cups, spoons, pencils, books, toys, door handles, taps, etc.

• Transmission occurs when one touches the contaminated objects and touches his mouth, nose, or eyes

Option C: Airborne - droplet nuclei transmission

• A type of indirect transmission

• Droplet nuclei - dried residue of droplets of size <5µm that are generated by the evaporation of droplets coughed or sneezed into the air and by a variety of atomizing devices

• Remain infectious when suspended in the air for some time (typically hours)

• Transmission occurs when one inhales droplet nuclei/aerosols containing the virus in sufficient quantity

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 5:

The incubation period of SARS-CoV-2 infection ranges from 1-14 days so he is most likely to develop the disease after this period. The median incubation period is around 4-6 days.

The incubation period is defined as the time interval between invasion by an infectious agent and appearance of the first sign or symptom of the disease.

Solution to Question 6:

The outbreak of COVID-19 is an example of a propagated epidemic.

The propagated epidemic results from person-to-person transmission of the infectious agent over a prolonged duration of time. Other examples of this type include hepatitis A and polio.

Solution to Question 7:

All of the given symptoms are included in the definition of a suspect case of SARS-CoV-2/COVID-19 except convulsions.

As per WHO and MOHFW, a suspect case is defined as:

• A person who meets the clinical and epidemiological criteria-

• Clinical criteria:

• Acute onset of fever AND cough; OR

Sold by @itachibot

• Acute onset of any 3 or more of the following: fever, cough, general weakness/ fatigue, headache, myalgia, sore throat, coryza, dyspnoea, anorexia/nausea/vomiting, diarrhea, altered mental status.

• Epidemiological criteria: A person present in any of the following settings for 14 days prior to symptom onset-

• Residing or working in an area with a high risk of transmission of virus OR

• Residing or travel to an area with community transmission OR

• Working in any healthcare setting

• A patient with severe acute respiratory illness with-

• history of fever or measured fever of ≥38 C° along with cough

• onset within the last 10 days

• requiring hospitalization

• An asymptomatic person not meeting epidemiologic criteria with a positive SARS-CoV-2 antigen-RDT.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

An asymptomatic contact of a probable case with a positive COVID-19 rapid antigen test is considered as a confirmed case, and not a probable case.

Probable case is defined as:

• A patient who meets clinical criteria and is a contact of a probable or confirmed case, or linked to a COVID-19 cluster.

• A suspect case with chest imaging showing findings suggestive of COVID-19 disease.

• A person with recent onset of anosmia or ageusia in the absence of any other identified cause.

• Death, not otherwise explained, in an adult with respiratory distress preceding death and was a contact of a probable or confirmed case or linked to a COVID-19 cluster.

Confirmed case is defined as:

• A person with a positive Nucleic Acid Amplification Test (NAAT) including RT-PCR or any other similar test approved by ICMR.

• A person with a positive COVID-19 rapid antigen test and meeting either the probable case definition or suspect criteria.

• An asymptomatic person with a positive COVID-19 rapid antigen test who is a contact of a probable or confirmed case.

Solution to Question 9:

A person traveling in the same environment and not having significant exposure to the confirmed case is not classified as a high-risk contact.

Sold by @itachibot

A contact of COVID-19 is a person living in the same household or having direct physical contact with the patient/infectious secretions without appropriate PPE or a person who was in a closed environment or at a distance of within 1 meter of a COVID-19 case.

High-risk contacts:

• Touched body fluids of the patient

• Had direct contact with the body of the patient during the physical examination without proper PPE

• Touched or cleaned the linens, clothes, or dishes of the patient

• Lives in the same household as the patient

• Comes in close contact (within 1 meter) with the patient without necessary precautions

• A passenger in close proximity with the symptomatic patient for 6 hours and the patient later tested positive

Low-risk contacts:

• Shared the same space but does not have a high-risk exposure to the confirmed case.

• Travelled in the same environment but does not have a high-risk exposure to the confirmed case.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

A middle-aged man who had malaria 2 months ago is not at major risk of developing severe COVID-19 disease.

The major risk factors for severe COVID-19 are:

• Increasing age (>60 years)

• Underlying non-communicable diseases

• Cardiac disease

• Chronic lung disease

• Cerebrovascular disease

• Hypertension

• Diabetes

• Obesity

• Chronic kidney disease

• Immune suppression

• Cancer

Solution to Question 11:

Sold by @itachibot

The given clinical scenario is suggestive of COVID-19. Hemoptysis is one of the atypical manifestations of COVID-19 and it may be present with more common features such as breathlessness, fever, and cough. CT-finding of localized ground-glass opacity is also supportive of the diagnosis.

Other atypical manifestations of COVID-19 include:

• Respiratory-

• Pulmonary embolism

• Hemoptysis

• Cardiac-

• Myocardial infarction

• Myocarditis

• Pericarditis

• Pericardial effusion

• Neurological-

• Stroke

• Guillain■Barre Syndrome

• Viral encephalitis

• Toxic encephalopathy

• Acute hemorrhagic necrotizing encephalopathy

• Olfactory and taste disorders

• Gastrointestinal-

• Diarrhea

• Nausea

• Vomiting

• Abdominal pain

• Ocular-

• Conjunctivitis

• Epiphora

• Cutaneous-

• Erythematous rash

• Widespread urticaria

• Chickenpox■like vesicles

• Localized pruritic lesions

• Petechial skin rash

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Option A: Tuberculosis would have a more chronic history and radiologically appears as dense, homogeneous parenchymal consolidation in subpleural sites of lower and middle lobes.

Option B: In necrotising pneumonia, CT imaging is likely to show distinct areas of low attenuation with decreased parenchymal enhancement representing liquefaction.

Option D: Invasive aspergillosis is likely to occur in the setting of immunosuppression. The associated ground glass opacity, occurring due to hemorrhage, surrounds a pulmonary nodule and is known as the halo sign.

Solution to Question 12:

Diffuse alveolar damage is the most common lung pathology noted during the autopsy of COVID-19 patients.

COVID-19 related histopathological changes in the lungs include:

- Diffuse alveolar damage
- Hyaline membrane formation
- Desquamation of pneumocytes
- Fibrin deposits in lungs
- Inflammation with mononuclear cells

Note: Diffuse alveolar damage is not specific to COVID-19. It may be seen in all cases of acute respiratory distress syndrome (ARDS) including pneumonia, burns, near drowning, etc

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 13:

A, B, C & E are clinical criteria of admission under COVID-19 ward.



Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

COVID 19 - Investigations

Question 1:

During the swab collection for COVID-19 testing, the healthcare professional asks you to tilt your head approximately to what degree?

a) 30 degrees

b) 45 degrees

c) 70 degrees

d) 15 degrees

Question 2:

Which of the following is a false statement regarding sample collection for COVID-19 testing?

a) The nasal swab is placed in a different viral transport media tube as the throat swab.

b) Oropharyngeal sample is not collected using a calcium alginate swab.

c) Swab is rubbed over the tonsillar pillars and posterior oropharynx for oropharyngeal swab.

d) Swab is inserted parallel to the palate for nasopharyngeal swab.

Question 3:

The specimens collected for COVID-19 testing should be transported to the lab at what temperatures?

a) -30 degree C

b) -4 to -8 degree C

c) 2 to 8 degree C

d) 12 to 16 degree C

Question 4:

Which level of BSL (Biosafety level) facility is required during specimen handling for molecular testing of SARS CoV 2?

a) BSL 1

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) BSL 2

c) BSL 3

d) BSL 4

Question 5:

An elderly patient was rushed to the casualty following bouts of severe breathlessness. He reported a history of fever and cough for the past 3 days. On examination, his SpO2 was 85% and RR was 32/min. CT scan is shown below. What is the gold standard investigation for his diagnosis?



a) CLIA

b) ELISA

c) RAT

d) RT-PCR

Question 6:

Choose the correct statement regarding testing of COVID-19:

a) All newborns must be tested.

b) All pregnant women must be tested for COVID-19 in the first trimester.

c) No individual can voluntarily subject themselves to COVID-19 test.

d) A single positive RT-PCR test is considered confirmatory without repeat testing.

Sold by @itachibot

Question 7:

A woman, whose husband has tested positive for COVID-19, comes to the clinic with a negative report on the rapid antigen test. She is currently asymptomatic. What will be the next step?

a) Retest with rapid antigen test after 5 days

b) Retest with RT-PCR after 5 days

c) Retest with RT-PCR immediately

d) Retest with RT-PCR if symptoms develop

Question 8:

A man presents to the flu clinic with complaints of fever and sore throat for the past 4 days. COVID-19 is suspected and a paper-based test kit as shown below is used for testing. Identify the test.



a) TRUNAT

b) FELUDA

c) COVID Kavach ELISA

d) Q COVID-19 Ag Test

Question 9:

Sold by @itachibot

What is the name of the COVID-19 IgG antibody detection ELISA test kit developed by ICMR and NIV?

a) COVID Nischay ELISA

b) COVID Aarogya ELISA

c) COVID Kavach ELISA

d) COVID Sankalp ELISA

Question 10:

What is least likely to be seen in a patient suspected of COVID-19?

a) Elevated transaminases

b) Elevated fibrinogen

c) Lymphocytosis

d) Prolonged prothrombin time

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 11:

What is the most commonly seen pattern in chest CT of patients affected with COVID-19?

a) Interlobular septal thickenings

b) Air bronchograms

c) Ground glass opacity

d) Crazy paving pattern

Question 12:

Increased concentrations of all of the following are poor prognostic markers in a COVID-19 patient except:

a) Glycosylated ferritin

b) C-reactive protein

c) N-terminal pro-BNP

d) Procalcitonin

Question 13:

Sold by @itachibot

Which inflammatory marker is of prognostic significance in COVID-19 patients?

a) IL-6

b) ESR

c) NLR

d) PCT

Answer Key

Question No.	Correct Option
1	c
2	a
3	c
4	b
5	d
6	d
7	d
8	b
9	c
10	c
11	c
12	a
13	a

Detailed Explanations

Solution to Question 1:

For collecting nasal/nasopharyngeal/oropharyngeal specimens for COVID-19 testing, a person's head should be tilted back 70 degrees.

Guidelines for specimen collection:

• Use appropriate PPE and maintain proper infection control measures

• Restricted entry to visitors or attendants during sample collection

• Complete the requisition form for each specimen submitted

• Proper disposal of all waste generated

Sold by @itachibot

Recommended personal protective equipment (PPE) for the healthcare professional who is collecting the specimen or coming within 6 feet of the patient:

• N95 or higher-level respirator (or facemask if a respirator is not available)

• eye protection

• gloves

• gown

Solution to Question 2:

The nasal swab is placed in the same viral transport media tube as the throat swab.

Samples that can be used for COVID-19 testing are:

• Throat and nasal swab

• Oropharyngeal swab

• Nasopharyngeal swab

• Broncho-alveolar lavage

• Endotracheal aspirate

Combined throat and nasal swabs:

• Tilt patient's head back 70 degrees.

• Gently rotate the swab and insert less than one inch into the nostril. Rotate the swab several times against the nasal wall and repeat the same in the other nostril with the same swab.

• Place the tip of the swab into the transport media tube and cut off the applicator.

• For throat swab, take a second swab, insert into the mouth and swab the posterior pharynx and tonsillar areas.

• Place the tip of the swab into the same tube and cut off the applicator tip.

Oropharyngeal swab:

• Tilt patient's head back 70 degrees.

• Rub swab over both tonsillar pillars and posterior oropharynx and avoid touching the tongue, teeth, and gums.

• Use only synthetic fiber swabs with plastic shafts. Avoid calcium alginate swabs and swabs with wooden shafts.

• Place the swab immediately into sterile tubes with viral transport medium.

Nasopharyngeal swab:

• Tilt the patient's head back 70 degrees

• Insert flexible swab through the nares parallel to the palate (not upwards) until resistance is encountered

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Rub and roll the swab and leave it in place for several seconds to absorb secretions before removing

• Place the swab immediately into sterile tubes with viral transport medium.

Viral transport mediums:

• Hanks balanced salt solution with Ca and Mg and heat inactivated fetal bovine serum, gentamicin and amphotericin B (most commonly used)

• Sterile saline

• Phosphate buffered saline-glycerol medium

• Amies medium

Sold by @itachibot
If you purchased this from someone else, you've been scammed

Solution to Question 3:

Specimens collected for COVID-19 testing should be transported to the lab at a temperature of 2 to 8 degrees C as per the WHO guidelines. The ICMR/NIV guidelines recommend 4 degrees C for transport to the laboratory.

Both guidelines recommend storing specimens at -70°C or below if a delay in testing or shipping is expected.

Specimen	Sensitivity
Bronchoalveolar lavage fluid	93%
Sputum	72%
Nasopharyngeal swab	63%
Oropharyngeal swab	32%
Faeces	29%
Blood	1%

Solution to Question 4:

Specimen handling for molecular testing of SARS CoV 2 would require BSL-2 or equivalent facilities.

Microbiology and pathology laboratories performing diagnostic tests on a stool or respiratory specimen should handle potential SARS-CoV specimens using Biosafety Level 2 (BSL-2). Virus culture and initial characterization of viral agents recovered in cultures requires minimum BSL-3 facilities.

Note: Biosafety levels include a specific combination of work practices, safety equipment, and facilities that are designed to minimize the exposure of workers and the environment to infectious agents. There are four biosafety levels.

Sold by @itachibot

Solution to Question 5:

The given clinical scenario, along with CT showing bilateral multiple patchy consolidation and grid-like thickness of interlobular septa, is suggestive of COVID-19. The gold standard for its diagnosis is the identification of viral genome targets in the respiratory tract materials by reverse transcriptase polymerase chain reaction (RT-PCR) during the first week of symptoms.

RT-PCR involves the reverse transcription of the genetic material of the virus (RNA) to complementary DNA (cDNA), followed by amplification of some regions of the cDNA. Four regions of the SARS CoV2 genome are used as targets:

- RdRp gene (RNA-dependent RNA polymerase)
- Genes from structural proteins
- E (virus envelope)
- N (virus nucleocapsid),
- ORF1ab gene (open reading frame 1a and 1b)

CBNAAT (cartridge-based nucleic acid amplification test) uses small, chip-based, portable, battery-operated devices that detect target genetic material of SARS CoV2. Examples include TrueNat and Cepheid Xpert Xpress SARS CoV-2

Options A and B: Antibody testing- Antibodies of IgA, IgM, and IgG isotypes specific to different virus proteins are detected by enzyme-linked immunosorbent assay (ELISA) or chemiluminescence immunoassays (CLIA). CLIA is more sensitive compared to ELISA.

Option C: Rapid antigen test (RAT)- Antigen-based assays are used as point-of-care tests for early detection of SARS-CoV-2. They are used at the field level for early detection of infection and quick containment. Most of such tests have relatively moderate sensitivity but high specificity. They are used for diagnosis along with RT-PCR.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

A single RT-PCR/TrueNat/CBNAAT/RAT positive is considered confirmatory without repeat testing. If symptoms develop following a negative RAT, a repeat RAT or RT-PCR should be done.

Routine surveillance in containment zones and screening at points of entry for all symptomatic cases including:

- Healthcare and frontline workers
- Direct and high-risk contacts and individuals

Routine surveillance in non-containment areas for all symptomatic cases including:

- History of international travel in the last 14 days
- Contacts of a laboratory-confirmed case
- Healthcare and frontline workers involved in containment and mitigation activities

Sold by @itachibot

• Returnees and migrants within 7 days of illness

• All asymptomatic high-risk contacts

In hospital settings for all symptomatic cases including:

• All patients of Severe Acute Respiratory Infection (SARI)

• Patients presenting in a healthcare setting

• Neonates presenting with acute respiratory/sepsis-like illness

• Patients with atypical manifestations (stroke, encephalitis, hemoptysis, Guillain Barre Syndrome etc.) based on the doctor's discretion

• All pregnant women in/near labor hospitalized for delivery

• Asymptomatic high-risk patients who are hospitalized or seeking immediate hospitalization (eg. immune compromise, malignancy, transplant, chronic comorbidities, elderly > 65 years)

Testing on-demand for all individuals:

• Traveling to countries/states mandating negative COVID-19 test at the point of entry

• Who wish to get themselves tested

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Solution to Question 7:

Asymptomatic individuals with a negative rapid antigen test (RAT) need to be retested only if they develop symptoms of COVID-19 such as cough, fever, and difficulty in breathing.

Rapid antigen test, when positive, is reported as positive in both symptomatic and asymptomatic patients. When the test is negative, the patient is retested with another RAT or RT-PCR only if the patient is symptomatic.

Solution to Question 8:

The paper-based Covid-19 test kit developed by CSIR and IGIB and approved by the Drugs Controller General of India (DCGI) is named FELUDA.

FELUDA is an acronym for FNCAS9 Editor Linked Uniform Detection Assay. The name is inspired by a fictitious detective character in novels written by Satyajit Ray.

FELUDA uses CRISPR-Cas9 technology (CRISPR = Clustered Regularly Interspaced Short Palindromic Repeats). The CRISPR technology uses a highly specific CAS9 protein to find and bind to the target Covid signature. This is then coupled with paper-strip chemistry to elicit a visual readout on a paper strip. It is a very affordable, sensitive (96%), and specific (98%) strip test which can detect viral genetic material within 45 minutes.

Solution to Question 9:

Sold by @itachibot

COVID Kavach ELISA is an indigenous ELISA test kit for COVID-19 antibody detection developed by ICMR Delhi and NIV Pune.

It is reported to have a 98.7% sensitivity and 100% specificity and testing is possible even at the district level. As the ELISA kit has an inactivated virus, only minimal biosafety and biosecurity requirements as compared to the real-time RT-PCR test.

IgG antibodies usually start appearing after two weeks of the onset of infection. Therefore, the IgG test is not useful for detecting acute infection. IgG based antibody tests are used in sero-surveillance of the population.

Solution to Question 10:

Lymphocytosis is least likely to be seen in a patient with COVID-19. On the contrary, lymphopenia is likely to be seen.

Common laboratory abnormalities seen in COVID-19 are:

• Lymphopenia

• Mild thrombocytopenia

• Elevated aminotransaminase levels

• Elevated lactate dehydrogenase levels

• Elevated inflammatory markers - eg, ferritin, C-reactive protein, and erythrocyte sedimentation rate

• Markers of a hypercoagulable state - high fibrinogen, high D-dimer levels, increased fibrin degradation products, prolonged prothrombin time (PT) and activated partial thromboplastin time (aPTT)

The degree of these abnormalities tends to correlate with disease severity.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 11:

Ground glass opacity (GGO) is the most commonly seen pattern in chest CT of patients with COVID-19.

Some of the typical features seen in the CT scans of COVID-19 patients include:

• Ground glass opacities

• Consolidation

• Air bronchograms

• Interlobular septal thickenings

• Visible intralobular lines (crazy-paving pattern)

• Reversed halo sign

Sold by @itachibot

The image below shows an axial CT scan with multiple patchy areas of pure ground-glass opacities. GGO are hazy areas with slightly increased density in the lungs without obscuration of underlying bronchial and vascular margins.

The image given below shows a CT scan with bilateral GGO in the lower lobe and an air bronchogram (arrow) in the left subpleural area. Air bronchograms appear like air-filled bronchi within a highly attenuated parenchymal background. These are dilated bronchi filled with mucus plugs and do not truly contain air.

The image given below shows a CT scan with multiple consolidation areas and interlobular septal thickening (arrows) in both lungs. The pulmonary interstitial structures such as interlobular septa and intralobular lines can thicken and manifest as small linear opacities.

Sold by @itachibot



The image given below shows a CT scan with crazy-paving pattern (circled area). It is defined as the presence of GGOs with superimposed interlobular septal thickening giving the appearance of irregular paving stones.



The image given below shows a CT scan with multiple lesions with reversed halo sign (arrows). The reversed halo sign or atoll sign is the presence of a ring-like area of consolidation with a superimposed rounded GGO.

Sold by @itachibot



Sold by @itachi... if you purchased this from some... you may have been scammed.

Solution to Question 12:

Glycosylated ferritin is decreased in severe COVID-19 illness and hence, it is not a poor prognostic marker.

Interestingly, in contrast to ferritin which is actually elevated in individuals with COVID 19, the glycosylated ferritin is decreased. It is a potentially sensitive biomarker that can better discriminate severe COVID than ferritin alone.

Increased levels of the following laboratory markers is associated with poorer outcomes:

- D-dimer
- C-reactive protein
- Lactate dehydrogenase
- Cardiac troponin
- Ferritin
- Creatinine phosphokinase

Solution to Question 13:

Elevated pro-inflammatory cytokines like interleukin-6 (IL-6) are associated with critical and fatal COVID-19. It is specifically useful in predicting prognosis of such patients.

Increased levels of the following laboratory markers are associated with poorer outcomes:

- D-dimer
- C-reactive protein
- Lactate dehydrogenase

Sold by @itachibot

• Cardiac troponin

• Ferritin

• Creatinine phosphokinase

Options B, C and D: Inflammatory markers associated with increased severity of COVID-19 are:

• PCT (procalcitonin test)

• ESR

• WBC

• LMR (lymphocyte/mean platelet volume ratio)

• HsCRP (high-sensitivity C-reactive protein)

• NLR (neutrophil to lymphocyte ratio)

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

COVID 19 - Treatment

Question 1:

The airport authority sent a COVID-19 suspect patient to the designated district hospital for a consultation. He came positive on RT-PCR testing. As the duty doctor, you observe that he has mild symptoms that can be treated on an outpatient basis. What will be your recommendation?

a) Home isolation for 7 days from testing positive

b) Home quarantine for 10 days from testing positive

c) Home isolation for 10 days from testing positive

d) Home quarantine for 7 days from testing positive

Question 2:

A COVID-19 positive man is rushed to the casualty due to severe breathlessness. Examination findings are suggestive of pneumonia and his vitals are- PR: 90/min, BP: 134/88 mmHg, RR: 22/min and SpO2: 86%. What will you advise this patient?

a) Home isolation

b) Admit to COVID care center

c) Admit to ward

d) Admit to ICU

Question 3:

A man presents with fever and cough for 2 days. His wife recently tested positive for COVID-19 and has been under home isolation. Examination findings are given below. What is not recommended in his management?

a) Awake proning

b) Administration of paracetamol

c) Oxygen support

d) Prophylactic LMW heparin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Question 4:

A 42-year-old COVID-19 positive man under home isolation develops breathing difficulty. SpO2 is 86%. He is administered oxygen via a face mask with a reservoir bag with a flow rate of 10 L/min. After a trial of one hour, his SpO2 is 88%. What is the next step?

a) Increase flow rate to 15 L/min

b) High-flow nasal cannula oxygen therapy

c) Endotracheal intubation

d) No specific intervention necessary

Question 5:

An 84-year-old man is being treated in the ICU for sepsis induced respiratory failure secondary to COVID-19. What is the maximum tidal volume setting permissible for his mechanical ventilation? (PBW-predicted body weight)

a) 4 ml/kg PBW

b) 6 ml/kg PBW

c) 8 ml/kg PBW

d) 10 ml/kg PBW

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Question 6:

An elderly diabetic man with COVID-19 is desaturating on non-invasive ventilation. You intubate him and place him on a mechanical ventilator. He improves briefly but the SpO2 suddenly drops. You suspect a tubal block and perform closed suctioning. Despite this, he goes into cardiac arrest and you start CPR. You would not expect aerosol generation in:

a) Non-invasive ventilation

b) Manual ventilation

c) Closed suctioning

d) CPR

Question 7:

A 67-year-old patient with end-stage renal disease is tested positive for COVID-19. Which anticoagulation agent would you recommended for him?

Sold by @itachibot

a) Unfractionated heparin

b) LMWH

c) Warfarin

d) Dabigatran

Question 8:

Which case scenarios along with the drugs prescribed are inappropriate?

a) 1, 2, 3, 4, 5

b) 1, 2, 3, and 4 only

c) 3, 4 and 5

d) 1 and 2 only

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

A G2P1 woman presents to the clinic at 36 weeks of gestation. Her husband was COVID positive and now she has also been tested positive. What is true regarding her management?

a) C-section is the preferred mode of delivery

b) Awake proning is not recommended

c) CT scan is absolutely contraindicated

d) Aim to keep oxygen saturation more than 94% during labour

Question 10:

Which of the following is a criterion for discharge in a patient with mild COVID-19 infection?

a) Discharge after 10 days from testing positive & no fever for 3 days

b) Discharge after 7 days from testing positive & no fever for 3 days

c) Discharge if SpO2>98% on room air for 3 consecutive days

d) Discharge if repeat RT-PCR testing is negative

Question 11:

A 33-year-old delivery boy tested COVID-19 positive. He started experiencing worsening breathlessness and cough and rushed to the hospital when his pulse oximeter reading was

Sold by @itachibot

87%. Following treatment, he showed signs of improvement. What is correct before discharging him?

a) Discharge after 7 days from symptom onset & no fever for 3 days

b) Discharge if repeat RT-PCR testing is negative after recovery

c) Discharge if SpO2>98% on room air for 3 consecutive days

d) Discharge at the discretion of the treating medical officer

Question 12:

A 10-year-old boy with COVID-19 presents with sore throat, rhinorrhea and cough. He does not have fast breathing. What is recommended for this child?

a) 2 and 5

b) 3 only

c) 2 and 4

d) 1 only

Question 13:

FDA has recently sanctioned emergency use authorization to the first oral antiviral drug combination for the treatment of COVID-19. This drug combination contains:

a) Nirmatrelvir and Ritonavir

b) Molnupiravir and Ritonavir

c) Nirmatrelvir and Indinavir

d) Molnupiravir and Indinavir

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	d
3	a
4	b

Sold by @itachibot

5	c
6	c
7	a
8	b
9	d
10	b
11	d
12	d
13	a

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Detailed Explanations

Solution to Question 1:

The given scenario is a mild case of COVID-19 who can be recommended home isolation for a duration of 7 days from testing positive.

Isolation is the separation and restriction of movements or activities of persons who are sick, to prevent the transmission of disease.

Eligibility for home isolation:

• Clinically assigned as a mild or asymptomatic case

• Facility at their residence for self-isolation and for quarantining the family contacts

• Availability of a caregiver

• Must monitor his health regularly and inform his health status to the district surveillance officer

The patient can end home isolation after 7 days from testing positive and no fever for 3 successive days. There is no need for testing after the home isolation period is over.

The patient should seek medical help if he/she develops:

• Unresolved High-grade fever (more than 100° F for more than 3 days)

• Difficulty in breathing,

• Dip in oxygen saturation (SpO2 ≤ 93% on room air at least 3 readings within 1 hour) or respiratory rate >24/ min

• Persistent pain/pressure in the chest

• Mental confusion or inability to arouse

• Severe fatigue and myalgia

Quarantine separates and restricts the movement of people who were exposed to a contagious disease to see if they become sick.

Sold by @itachibot

Solution to Question 2:

The patient is suffering from severe COVID-19 and hence has to be admitted to ICU. The disease is said to be severe when there is RR>30/min, breathlessness OR SpO2<90% on room air.



Solution to Question 3:

The given clinical scenario is suggestive of moderate COVID-19 (SpO2 <94%, RR≥24). Awake proning should be avoided in patients with hemodynamic instability.

Awake proning in moderate COVID disease is recommended for any patient with severe respiratory distress. It improves oxygenation by keeping the alveolar units open. Timely proning of patients with SpO2 < 94% can help reduce mortality.

Steps for self-proning:

• 1 pillow below the neck

• 1 or 2 pillows below the chest through upper thighs

• 2 pillows below the shins

• Keep alternating between proning, laying on your right, sitting up, and laying on your left, in a lying position, ideally for not more than 30 minutes in each position with an upper limit of 2 hours.

• One may prone up to 16 hours a day, in multiple cycles, if tolerable.

Avoid proning:

• Hemodynamic instability

• For 1 hour after meals

• DVT, cardiac conditions

• Fractures of spine, femur, or pelvis

If you purchased this from elsewhere, you may have been scammed.
Sold by @itachibot

Solution to Question 4:

This patient has persistent hypoxia after an adequate trial of standard oxygen therapy and should be started on high-flow nasal cannula oxygen therapy.

When respiratory distress and/or hypoxemia of the patient cannot be alleviated after receiving standard oxygen therapy, high–flow nasal cannula oxygen therapy (HFNO), or non–invasive ventilation can be considered. Compared to standard oxygen therapy, HFNO reduces the need for intubation.

Patients on HFNO should be closely monitored, and endotracheal intubation should be done if the patient acutely deteriorates or does not improve after a short trial (about 1 hr).

Contraindications to HFNO includes patients with:

• Hypercapnia (exacerbation of COPD)

• Hemodynamic instability

• Multi-organ failure

• Abnormal mental status

Note: If an high flow nasal cannula (HFNC) or simple nasal cannula is used to administer O2, an N95 mask should be applied over it.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

A maximum tidal volume of up to 8 ml/kg PBW is allowed in a patient with ARDS who is being mechanically ventilated.

In patients with ARDS, mechanical ventilation should be implemented using lower tidal volumes (4–8 ml/kg PBW) and lower inspiratory pressures (plateau pressure <30 cmH2O).

In patients with moderate or severe ARDS, a higher PEEP is suggested. PEEP titration requires consideration of benefits (improving alveolar recruitment) vs. risks (end-inspiratory over distension leading to lung injury).

Lung protective ventilation strategy by ARDS net protocol recommendations:

Parameters	Values
Tidal volume	6ml/kg
RR	15-35/min
PEEP	5-15cm H2O
Target plateau pressure	< 30cm H2O
Target SpO2	88-95%

Sold by @itachibot

Parameters	Values
Target PaO2	55-80m mHg

Solution to Question 6:

Open suctioning of the respiratory tract is associated with significant amounts of aerosol generation and not closed suctioning.

Commonly performed aerosol-generating procedures are:

• Open suctioning of the respiratory tract

• Bronchoscopy

• Cardiopulmonary resuscitation (CPR)

• Non-invasive ventilation (e.g., BiPAP, CPAP)

• Endotracheal intubation and extubation

• Manual ventilation

It is recommended that healthcare providers should be protected with PPE while performing these procedures. It must be performed in an adequately ventilated single room with a minimum of 12 air changes/hour.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 7:

The given clinical scenario is suggestive of COVID-19 in a patient with end-stage renal disease. Unfractionated heparin is recommended for use in such patients.

Prophylactic anticoagulation is used in the management of moderate and severe COVID cases. A prophylactic dose of low molecular weight heparin or unfractionated heparin is used to prevent thromboembolism. In a COVID patient with end-stage renal disease, unfractionated heparin is the preferred agent.

Solution to Question 8:

The correct answer is 1, 2, 3, and 4.

• Tocilizumab is contraindicated in HIV-positive patients and those with active bacterial or fungal infection.

• Bamlanivimab is not authorized for patients with mild COVID-19 pneumonia who are not at a high risk for hospitalization.

• Remedisivir is contraindicated in postnatal lactating women.

Sold by @itachibot

• Convalescent plasma therapy is no longer recommended in the management of COVID patients.

Tocilizumab acts by IL-6 inhibition. It may be considered in moderate-severe COVID patients with progressively increasing oxygen requirements and mechanically ventilated patients not improving despite the use of steroids. Patients should be carefully monitored for secondary infections and neutropenia after treatment with tocilizumab. It is contraindicated in:

• HIV

• Active infections (systemic bacterial/fungal)

• Tuberculosis

• Active hepatitis

• Absolute neutrophil count < 2000/mm3

• Platelet count < 1,00,000/mm3

Bamlanivimab is a monoclonal antibody directed against the spike protein of SARS-CoV-2. The FDA has issued an emergency use authorization for bamlanivimab for the treatment of adult and pediatric mild to moderate COVID patients (12 years of age and older weighing at least 40 kilograms) who are at high risk for progressing to severe COVID-19 and/or hospitalization. Possible side effects include anaphylaxis and infusion-related reactions, nausea, diarrhea, dizziness, headache, itching, and vomiting. There is no specific contraindication for this drug.

Remdesivir is an antiviral drug that works by inhibiting RNA-dependent RNA polymerase. It may be considered in moderate and severe hospitalized COVID patients requiring supplemental oxygen/oxygen delivery through a high flow device/ noninvasive ventilation/ invasive mechanical ventilation/ ECMO. It is contraindicated in:

• AST/ALT > 5 times upper limit of normal

• Severe renal impairment (i.e., eGFR < 30ml/min/m2 or need for hemodialysis)

• Pregnancy or lactating females

• Children <12 years of age

Convalescent plasma therapy is no longer recommended in the management of COVID patients, as per the revised guidelines.

Dexamethasone is generally given to all patients of moderate and severe COVID-19. Patients with severe COVID-19 can develop a systemic inflammatory response that can lead to lung injury and multisystem organ dysfunction.

Solution to Question 9:

The given clinical scenario is suggestive of COVID-19 disease in a pregnant patient. The management should include an aim to keep oxygen saturation above 94% during labour in this patient. In such cases, maternal observations and assessment during labour should be continued as per standard practice, with the addition of hourly oxygen saturation.

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Sold by @itachibot

Option A: There is currently no evidence to favour one mode of birth over another. Mode of birth should not be influenced by the presence of COVID-19, unless the woman's respiratory condition demands urgent intervention for birth.

Option B: Awake proning is recommended. Although evidence is limited, there are reports that this is feasible with appropriate padding up to at least 28 weeks of gestation.

Option C: CT scan is relatively contraindicated. Pregnant women with abdominal shielding should not be excluded from chest CT if clinically recommended.

Solution to Question 10:

COVID-19 cases that are mild/very mild/presymptomatic can be discharged from home isolation after 7 days of testing positive and no fever for 3 successive days. There is no need for testing after the end of home isolation.

Criteria to diagnose asymptomatic COVID-19: Laboratory confirmed case with no symptoms with a saturation of more than 93% in room air.

Criteria to diagnose mild COVID-19: Laboratory confirmed case with upper respiratory tract symptoms with/without fever, without shortness of breath, and saturation of more than 93% in room air.

Patients are advised to isolate from other family members and monitor themselves at home, under proper medical guidance. Patients >60 years with co-morbidities, and immunocompromised patients are allowed to isolate only after receiving clearance from the designated medical officer.

Immediate medical attention is required if the patient experiences the following:

- A 100 °F fever persists for >3 days and does not resolve on medication.
- Difficulty in breathing
- Oxygen saturation ≤93% (at least 3 readings within 1 hour)
- Respiratory rate >24/min
- Persistent chest pain or pressure
- Mental confusion
- Severe fatigue or myalgia

Solution to Question 11:

The given clinical scenario is suggestive of severe COVID-19 disease. Severe cases of COVID-19 (and immunocompromised patients) are discharged based on their clinical recovery at the discretion of the treating medical officer.

Note: Mild and moderate cases do not need an RT-PCR test before discharge. But, at the time of discharge, they are advised to self-monitor and continue wearing masks for a duration of 7 days.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Solution to Question 12:

The presence of sore throat, rhinorrhea, cough with no fast breathing are suggestive of mild COVID-19 disease. Hence, home isolation along with supportive care is recommended in this child.



Solution to Question 13:

FDA recently sanctioned emergency use authorization to a combination of nirmatrelvir and ritonavir tablets (Paxlovid) for the treatment of mild-to-moderate COVID-19. It is the first oral antiviral for the treatment of COVID-19.

Nirmatrelvir is a peptidomimetic inhibitor of the SARS-CoV-2 main protease (Mpro). Inhibition of this enzyme makes the virus incapable of processing polyprotein precursors and thus prevents viral replication.

Ritonavir is an HIV-1 protease inhibitor. It inhibits the CYP3A-mediated metabolism of nirmatrelvir which results in increased plasma concentrations of nirmatrelvir. It is not active against SARS-CoV-2 Mpro.

Patients who tested positive with direct SARS-CoV-2 test can be administered this drug combination, provided they are:

• 12 years of age and older

• Weighing at least 40 kgs

• At high risk for progression to severe COVID-19.

Contraindications include hypersensitivity reactions to the active ingredients or any other components, co-administration with drugs highly dependent on CYP3A for clearance, co-administration with potent CYP3A inducers. Adverse reactions include dysgeusia, diarrhea,

If you purchased this, you may have been scammed.

Sold by @itachibot

hypertension, and myalgia.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

COVID 19 - Prophylaxis and Complications

Question 1:

Your worried friend calls you for advice after attending a family function. He informs you that his aunt, who also travelled to the function in the same car, tested positive for COVID-19. What is your advice for your friend?

a) Home isolation for 14 days

b) Home quarantine for 14 days

c) Home quarantine for 10 days

d) Home isolation for 10 days

Question 2:

A 50-year-old man recovering from COVID-19 presents as follows. His medical and personal histories are insignificant with no exposure to cold. What would you suspect?

a) Chilblains

b) COVID toes

c) Varicella

d) Eczema

If you bought this from someone else, you've been scammed. Sold by @itachibot

Sold by @itachibot

Question 3:

While donning a set of PPE, what is the order that you will follow?

a) 2→1→3→4

b) 1→2→3→4

c) 1→2→4→3

d) 2→1→4→3

Question 4:

While doffing, what is the first component of the PPE that you would remove?

a) Coverall

b) Shoe covers

c) Face shield

d) Gloves

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

You are a medical officer at the designated COVID hospital in your city and are required to don PPE every day at work. Which area of the PPE should you not touch while doffing?

a) Earpiece / ties of the mask

b) Inside the mask

c) Front of face shield

d) Back of isolation gown

Question 6:

After your COVID ICU shift, where would you dispose your face shield?

a) Red bin

b) Black bag

c) Yellow bin

d) White bag

Sold by @itachibot

Question 7:

What is recommended for cleaning and disinfecting a healthcare facility against COVID-19?

a) 1% formaldehyde

b) 1% glutaraldehyde

c) 35% isopropyl alcohol

d) 1% sodium hypochlorite

Question 8:

Identify the wrong match regarding COVID-19 vaccines:

a) Pfizer - mRNA based vaccine

b) Covaxin - Inactivated vaccine

c) Moderna - Recombinant DNA vaccine

d) AstraZeneca - Non-replicating viral vector vaccine

Question 9:

Four days after being extubated, an elderly diabetic man with COVID-19 developed rhinosinusitis. The tissue obtained from the sinuses showed the following. The physician wanted to start the patient on amphotericin B. However, it was not available. Which other drug can be given to this patient?



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

a) Voriconazole

b) Isavuconazole

c) Itraconazole

d) Flucytosine

Question 10:

A young girl, who recently recovered from COVID-19, presents with fever, abdominal pain and rashes for 4 days. On examination, strawberry tongue is observed and cervical nodes are found to be enlarged bilaterally. What is the likely diagnosis?

a) Kawasaki disease

b) Multisystem inflammatory syndrome in children

c) Bacterial sepsis

d) Toxic shock syndrome

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 11:

Which gene increases the risk of collapsing glomerulopathy in COVID-19 patients?

a) NPHS1

b) ACTN4

c) APOL1

d) TRPC6

Question 12:

As a healthcare worker working in a COVID hospital, you receive the following vaccine. Which of the following is not an adverse event related to this vaccine?

Sold by @itachibot



a) Cough

b) Tremors

c) Demyelinating disorder

d) Dizziness-giddiness

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

In which of the following patients, would you administer Covidshield vaccination?

a) 1, 2, 3

b) 1, 2, 4, 5

c) 2, 3, 4, 5

d) 3, 4, 5

Question 14:

A 24-year-old boy who had SARS CoV-2 infection 3 months back came with complaints of fatigue and shortness of breath for 3 days. The minimum duration of these symptoms to define it as post-Covid-19 condition according to WHO is ____

a) 2 months

b) 2 weeks

c) 4 months

d) 4 weeks

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	b
2	b
3	a
4	b
5	c
6	a
7	d
8	c
9	b
10	b
11	c
12	c
13	c
14	a

If you purch... you may ha... else, ...d.

Detailed Explanations

Solution to Question 1:

Home quarantine for 14 days is advised when a person comes in contact with a COVID-19 positive patient.

The home quarantined person:

• Should stay in a well ventilated single room and stay away from high-risk individuals

• Should not attend any religious or social gatherings

• Must wash hands often

• Must change mask every 6-8 hrs

• Their surroundings must be disinfected with 1% sodium hypochlorite solution

In case the person being quarantined becomes symptomatic, all his contacts should be home quarantined for 14 days and followed up for another 14 days or till the report turns out negative on lab testing.

Sold by @itachibot

Isolation is the separation and restriction of movements or activities of persons who are sick, to prevent the transmission of disease. Quarantine is the separation and restriction of movements or activities of persons who are not ill but are believed to have been exposed to infection, for the purpose of preventing transmission of diseases.

Solution to Question 2:

The given clinical scenario is suggestive of COVID toes. The image shows pernio-like lesions caused due to COVID-19.

Pernio / chilblains are skin lesions on the acral surfaces of the body characterised by the development of erythematous-violaceous or purpuric macules. They may be accompanied by edema and pruritus.

It is usually due to cold exposure or underlying hematological disorders but has also been reported in patients with COVID-19 disease. The underlying mechanism is thought to be an inflammatory process. The lesions appear late (postviral phase) with the patients exhibiting positive IgM/ IgG antibodies for SARS-CoV-2 and negative RT-PCR. However, pernio like lesions can also be seen in early stages of the disease when the patients are still infective.

There are no treatment guidelines but high-potency topical corticosteroids may be helpful if the lesions are causing discomfort. Other cutaneous manifestations include exanthematous rash, livedo reticularis, urticaria, MIS-C etc.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 3:

The correct sequence of donning a PPE is first pair of gloves → coverall → Shoe cover → N-95 mask.

Preparation for donning the PPE:

• To be performed only in the designated area

• The health care worker must be prepared for the shift duty by

• Taking a meal

• Using the washroom

• Drinking water prior to entering the donning area

• Remove all accessories, jewelry

• Change into clean hospital scrubs and closed washable footwear

• Perform hand hygiene with soap and water

Steps of donning/putting on the PPE:

• Hand hygiene

• Wear the first pair of gloves

• Sterile or unsterile

Sold by @itachibot

• Make sure the cuff of the gloves goes as far as possible

• Examine and wear coverall

• Examine the coverall for any damage

• Start by wearing the leg sleeves and then the arm sleeves

• Zip up and cover the zip with the flap attached

• Wear shoe cover

• Sit on the clean chair and wear shoe cover

• Pull them up till your calves

• Wear N-95 mask and wear hood

• Hold the mask like a cup in the hand with both the straps hanging out in the front

• Place the mask on the face, pull and wear the lower strap first placing it below the ears

• Then pull and wear the upper strap placing above the ears

• The air leakage should be minimum around the mask when forcibly exhaling

• Wear the hood covering the forehead and the cheeks

• Wear face shield/eye goggles over the hood and ensure the fit

• Wear the second pair of gloves - pull the gloves to cover the sleeve cuff and forearm as much as possible

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

The component of the PPE kit to be removed first is the shoe covers.

Preparation for doffing the PPE:

• To be performed only in the designated area

• Check for any leak or soiling

• Doffing room should have two chairs, dirty and clean

• All the PPE must be discarded in the red bin

• Hand hygiene must be performed after every step

Steps of doffing the PPE:

• Engage the observer and ask him to check the PPE - observer should check for any leak, tear, or contamination of PPE. If contaminated, then it should be disinfected with alcohol-based wipes.

• Disinfect the gloves

• Remove and discard shoe covers

• Sit on the dirty chair with legs apart

• Remove the shoe covers by slowly pulling the outer surface starting from the top and then pull from the toes end. Discard them

Sold by @itachibot

• Disinfect and remove outer pair of gloves

• While removing the outer gloves, inner gloves should not be torn or contaminated

• Pinch the first glove on the outer surface at the wrist and pull it inside out

• Hold this glove in the other hand like a ball

• Slide the thumb inside the other glove and remove inside out balling around the previous glove, making it into a single bag. Then discard it.

• Disinfect inner gloves

• Remove face shield/eye goggle - loosen the screw and remove the face shield by holding it from the back handle. Discard it.

• Disinfect inner gloves again

• Remove hood and coverall

• Slide back the hood by holding it from the top

• Separate the zip flap and unzip the coverall suit

• Holding the suit at the arms, slide it off the shoulder slightly, then pull the suit from the side of the waist

• Roll the arms out one by one

• By touching the inner surface of the suit carefully roll out each leg. Discard it

• Disinfect and remove inner gloves as previously described

• Perform hand hygiene and wear a new pair of gloves

• Remove and discard N-95 mask

• Do not touch the exposed surface of the mask

• Stoop forward and first remove the lower strap

• Then pull out the upper strap and remove the mask

• Disinfect gloves

• Sit on the clean chair and disinfect shoes using alcohol wipes

• Disinfect and remove gloves

• Perform hand hygiene

• Wipe the shoes on sodium hypochlorite soaked doormat at exit

• Exit the doffing area and take a shower

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

The front of the face shield is considered to be a contaminated area of the personal protective equipment (PPE). Hence, it should not be touched while doffing.

The areas that are considered clean are the parts that will be touched while removing a PPE:

Sold by @itachibot

• Inside the gloves

• Inside and back of the gown including the ties

• Ties or earpieces of the mask, goggles, and face shields

The areas that are considered contaminated are:

• Outside front and sleeves of the isolation gown

• Front of goggles, mask, respirator, and face shield

• Outside of the gloves

Solution to Question 6:

The face shield used by a healthcare worker during COVID-19 pandemic is disposed in the red bin.

Solution to Question 7:

1% sodium hypochlorite is recommended for cleaning and disinfection against COVID-19.

The sodium hypochlorite solution should be prepared fresh. Leaving the solution for a contact time of at least 10 minutes is recommended.

Note: Alcohol (e.g. isopropyl 70% or ethyl alcohol 70%) can be used to wipe down surfaces where the use of bleach is not suitable, e.g. metals.

Solution to Question 8:

The Moderna COVID-19 vaccine is also an mRNA-based vaccine.

Vaccines	Mechanism
Pfizer/BioNtech (Comirnaty)	mRNA based vaccine
Moderna	mRNA based vaccine
Oxford University/ Astra Zeneca/ Serum Institute of India (Covishield)	Non-replicating viral vector
Gamaleya (Sputnik V)/ Dr.Reddys	Non-replicating viral vector
Johnson and Johnson (Janssen Ad26.CoV2.S)	Non-replicating viral vector
Bharat biotech/NIV (Covaxin)	Inactivated vaccine
Zydus Cadila (ZyCoV-D)	Plasmid DNA vaccine
Novavax	Protein subunit

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

The Drugs Controller General of India (DCGI) has granted permission for restricted emergency use of the following vaccines against COVID-19:

• Covaxin (Bharat biotech/NIV)

• Covishield (Oxford University/ AstraZeneca/ Serum Institute of India)

• Sputnik V (Gamaleya Research Institute/ Dr. Reddy's Laboratories)

• ZyCoV-D (Zydus Cadila)

The FDA has issued an emergency use authorization for the following vaccines against COVID-19:

• Pfizer/BioNtech

• Moderna

• Johnson and Johnson

According to the new vaccine guidelines by NEGVAC (National Expert Group on Vaccine Administration for COVID-19):

• Vaccination is to be deferred by 3 months in the following persons:

• Individuals (proven positive) recovered from the disease.

• Recipients of monoclonal antibodies or convalescent plasma.

• 2nd dose is to be deferred by 3 months if infected with COVID after 1st dose.

• 4-8 weeks gap, after recovery of patients with other serious illnesses requiring hospitalization/ICU care.

• Blood donation: Wait for 14 days after receiving the vaccine/testing RT-PCR negative.

• No screening is required by rapid antigen test (RAT) prior to vaccination.

• Recommended for all lactating women.

COVID-19 vaccination during pregnancy:

• MoHFW has recently approved the vaccination of pregnant women against COVID-19.

• COVID-19 vaccine can be taken anytime during pregnancy and should be taken at the earliest.

• If any woman has been infected with COVID-19 during pregnancy, she should be vaccinated soon after delivery.

• COVID-19 vaccine is safe in pregnancy.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

The given clinical scenario is suggestive of mucormycosis. Intravenous administration of amphotericin B is the first line treatment. Other drugs which are approved for its management include isavuconazole and posaconazole. Surgical debridement is the last line management.

Immunocompromised states are important risk factors including diabetes mellitus and glucocorticoid treatment as well as patients on ventilation.

Sold by @itachibot

Diagnosis can be made by culturing the organisms from sterile sites like needle aspirate, pleural fluid, or a tissue biopsy specimen. The biopsy is most sensitive. It reveals characteristic wide, thick-walled, ribbon-like, aseptate hyphal elements that branch at right angles.

It can present as:

• Rhino-orbital-cerebral mucormycosis - (most common) the initial symptoms are nonspecific and include eye or facial pain and facial numbness followed by the onset of conjunctival suffusion and blurry vision. If untreated it can spread from the sinuses to the orbit, and then to the frontal lobe.

• Pulmonary mucormycosis - (second most common)

• Cutaneous mucormycosis - usually results after trauma

• Gastrointestinal mucormycosis - primarily occurs in premature neonates who have disseminated disease or necrotizing enterocolitis

• Disseminated mucormycosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

This is a case of multisystem inflammatory syndrome in children (MIS-C), associated with COVID-19 infection.

Diagnostic criteria for MIS-C:

Children and adolescents 0–18 years of age with fever ≥3 days

AND two of these:

• Rash or bilateral non-purulent conjunctivitis or mucocutaneous inflammation signs (palmoplantar erythema, lip hyperemia, periorbital erythema and edema, strawberry tongue, and malar erythema)

• Hypotension or shock.

• Features of myocardial dysfunction, pericarditis, valvulitis, or coronary abnormalities (including ECHO findings or elevated Troponin/NT-proBNP)

• Evidence of coagulopathy (PT, PTT, elevated D-Dimers)

• Acute gastrointestinal problems (diarrhoea, vomiting, or abdominal pain)

AND

Elevated ESR(>40 mm), C-reactive protein (>5 mg/L), or procalcitonin.

AND

No other obvious microbial cause of inflammation including bacterial sepsis, staphylococcal or streptococcal shock syndromes.

AND

Evidence of COVID-19 (RT-PCR, antigen test, or serology positive), or likely contact with patients with COVID-19.

Types of MIS-C:

Sold by @itachibot

• MIS-C with shock/multi-organ dysfunction

• Kawasaki phenotype

• MIS-C without shock

MIS-C is treated with IV methylprednisolone. The dose can be increased in patients with shock. If the condition does not improve for 48-72 hours, specialists can be consulted for the addition of biologicals to the treatment regimen.

Solution to Question 11:

APOL1 gene increases the risk of collapsing nephropathy in patients infected with COVID-19.

Glomerular lesions are reported in a minority of COVID-19 patients. They are associated with collapsing focal segmental glomerulosclerosis (FSGS) and are called COVID-associated nephropathy (COVAN). Such patients present with nephrotic-range proteinuria and acute kidney injury (AKI). It specifically affects individuals of African ancestry who are carriers of this gene variant.

Thrombotic microangiopathy is another uncommon finding among these patients. Histopathology is similar to those seen in other forms of collapsing glomerulopathy. But, tubulointerstitial lesions including microcystic tubular dilatation and tubular injury are typically associated with APOL1-related disease.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 12:

Demyelinating disorders are rare adverse events following Covishield, not Covaxin.

Composition of Covaxin:

• Inactivated coronavirus

• Aluminum hydroxide gel

• TLR 7/8 agonist

• 2-Phenoxyethanol and

• Phosphate Buffered Saline/ NKA1

Adverse effects:

Rarely, complications like demyelinating disorders, Guillain-Barre syndrome, and vaccine-induced immune thrombotic thrombocytopenia (VITT) are also seen following Covishield vaccine.

Covaxin	Covishield
Fever, cough, sweating	Fever and chills
Injection site pain and swelli ng	Injection site pain and swelli ng

Sold by @itachibot

Covaxin	Covishield
Headache, bodyache, fatigue	Headache, malaise, fatigue
Abdominal pain, nausea, vo miting	Nausea
Dizziness, giddiness, tremor	Myalgia, arthralgia

Solution to Question 13:

The 51-year-old man should receive COVID-19 vaccination at least 3 months after hematopoietic cell transplantation (and not 5 weeks) to maximize vaccine efficacy.

Graft-versus-host disease and immunosuppressive regimens like systemic corticosteroids can blunt the immune responses to vaccination. Delay of vaccination until immunosuppressive therapy is reduced can be considered.

Consider additional risk factors linked to adverse COVID-19 complications including, but not limited to:

• Patients with advanced age (eg, ≥65 years of age)

• Patients with comorbidities (eg, chronic pulmonary, cardiovascular, or renal disease)

• Social and demographic factors

COVID-19 vaccines should be prioritized over other vaccines. A gap of 14 days is recommended between COVID-19 vaccines and any other vaccine.

If COVID vaccine has been given first, then TT vaccine can be given on the same day. If TT vaccine have been given first, then wait for 14 days to give COVID vaccine.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 14:

The minimum duration of symptoms such as fatigue, shortness of breath to define it as post Covid-19 condition according to WHO is 2 months.

WHO clinical case definition of post Covid-19 - Post Covid-19 condition occurs in individuals with a history of probable or confirmed SARS CoV-2 infection, usually 3 months from the onset of Covid-19, with symptoms that last for at least 2 months and cannot be explained by an alternative diagnosis.

Common symptoms include

• Fatigue

• Shortness of breath

• Cognitive dysfunction.

Sold by @itachibot