MARROW ED8

Microbiology

Comprehensive Question Bank

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Contents

Chapter	Title	Page
1	General Microbiology	4
2	Components of Immune system	30
3	Structure and Functions of the Immune System & ...	55
4	Hypersensitivity	70
5	Immune Disorders	83
6	Streptococci and Enterococci	96
7	Staphylococci	115
8	Corynebacterium, Listeria and Actinomyces	128
9	Clostridium and Bacillus	147
10	Mycobacteria Tuberculosis	164
11	Other Mycobacteria	178
12	Escherichia, Proteus and Klebsiella	190
13	Shigella and Salmonella	207
14	Vibrio and Campylobacterales	218
15	Pseudomonas and Burkholderiales	238
16	Haemophilus	249
17	Miscellaneous Bacteria - Yersinia, Brucella, Ba...	262
18	Gram Negative Cocci	284
19	Rickettsia, Chlamydia and Mycoplasma	299
20	Spirochetes	316

Sold by @itachibot

Chapter	Title	Page
21	General Properties of Viruses	336
22	DNA Viruses	351
23	Hepatitis	373
24	HIV	390
25	Myxoviruses and Rhabdoviruses	399
26	Arboviruses and Picorna Viruses	416
27	Miscellaneous Viruses - Rubella, Coronaviruses,...	432
28	Superficial and Systemic Mycoses	447
29	Opportunistic Mycoses	473
30	General Parasitology	489
31	Protozoology - Amoebae, Ciliates & Flagellates	505
32	Protozoology - Sporozoa	535
33	Helminthology - Cestodes & Trematodes	553
34	Helminthology - Nematodes	575
35	Applied Microbiology	602

Sold by @itachibot

General Microbiology

Question 1:

Identify the condenser in the following image.

a) Above the stage

b) Below the stage

c) Near the eyepiece

d) Above the objective lenses

Question 2:

A 28-year-old sex worker presented with a painless ulcer on her vulva. On examination, inguinal lymphadenopathy is present. Which of the following methods has been used to demonstrate the causative agent as per the given image?

Sold by @itachibot



a) Dark-field microscopy

b) Phase contrast microscopy

c) Fluorescent microscopy

d) Electron microscopy

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which among the following statements is false?

a) Muramic acid is present in prokaryotes

b) Mitotic division is present in eukaryotes

c) 80S ribosome is present in prokaryotes

d) Multiple linear chromosomes present in eukaryotes

Question 4:

Which of the following methods will not help you visualize the bacterial cell wall?

a) Fluorescence microscopy

b) Electron microscopy

c) Differential staining

d) Simple staining

Sold by @itachibot

Question 5:

In which of the following phases of the bacterial growth curve does sporulation occur?



©MARROW

a) D

b) A

c) C

d) B

Question 6:

Which organism has the following cell wall structure?



©MARROW

Sold by @itachibot

a) Gram-positive organism

b) Gram-negative organism

c) Fungi

d) Mycobacteria

Question 7:

Which of the following helps in bacterial adhesion?

a) Cytoplasmic membrane

b) Mesosomes

c) Fimbriae

d) Lipopolysaccharides

Question 8:

Pasteurization is done at _____.

a) 73 degree Celsius for 20 minutes

b) 63 degree Celsius for 30 minutes

c) 72 degree Celsius for 30 seconds

d) 63 degree Celsius for 30 seconds

Sold by @Itachibot
If you purchased this from someone else, you may have been scammed.

Question 9:

Which of the following methods of sterilization involves exposure to 100 °C for 20 minutes on 3 successive days?

a) Autoclaving

b) Tyndallization

c) Pasteurization

d) Inspissation

Question 10:

Which of the following works by disrupting the cell membrane?

Sold by @itachibot

a) Lysol

b) Povidone-iodine

c) Thiomersal

d) Ethylene oxide

Question 11:

Which of the following methods is used in the test shown below?

a) Streak culture

b) Lawn culture

c) Stroke culture

d) Stab culture

Question 12:

In which of the following organisms will the following phenomenon not be observed in?

Sold by @itachibot



a) Bacillus anthracis

b) Klebsiella pneumoniae

c) Neisseria gonorrheae

d) Hemophilus influenzae

Sold by @itachi... purchased this from some... else, ...u may have been scammed.

Question 13:

A 32-year-old laborer working at a construction site presented with fever and hemoptysis. The sputum sample collected for examination showed the following. The smear will be stained by which of the following sequences?



a) Gentian violet - iodine - alcohol - saffranin

Sold by @itachibot

b) Methylene blue - malachite green- acetic acid-water

c) Methanol-methylene blue-acid-water

d) Carbol fuschin - acid-alcohol - methylene blue

Question 14:

Given are a few case scenarios. The etiological agents of which of the following are microaerophilic?

a) 2 and 5

b) 3 and 4

c) 1 and 4

d) 1 and 2

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Question 15:

The given image shows a culture medium with lactose fermenting and non-lactose fermenting bacteria. Which of the following is not a component of the culture medium?



a) Taurocholate

b) Peptone

c) Neutral red

d) Cysteine

Sold by @itachibot

Question 16:

Which of the following is an enrichment medium?

a) B

b) C

c) A

d) D

Question 17:

A microbiologist observes the following growth on a blood agar plate. Which of the following will not be the etiological agent?

MARROW
Sold by
purchased this fro
you may have been scam
else,
d.

Sold by @itachibot

a) Clostridium tetani

b) Corynebacterium diphtheriae

c) Proteus mirabilis

d) Bacillus cereus

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 18:

Bacterial transduction occurs by

a) Plasmids

b) Sex pili

c) Bacteriophage

d) Uptake of genetic material by other bacteria.

Question 19:

Which of the following principles is primarily used in polymerase chain reaction?

a) Transcription of DNA

b) It can amplify small amounts of DNA

c) It can amplify small amounts of RNA

d) Reverse transcription of RNA

Sold by @itachibot

Question 20:

Which of the following describes the reverse transcriptase enzyme used in RT-PCR correctly?

a) DNA-dependent RNA polymerase

b) RNA-dependent DNA polymerase

c) DNA-dependent DNA polymerase

d) RNA-dependent RNA polymerase

Question 21:

What are jumping genes known as?

a) Episomes

b) Plasmids

c) Transposons

d) Cosmids

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 22:

In which of the following situations will you use a sanitizer?

a) A, B & C only

b) A & B only

c) A, B C & D

d) C & D only

Question 23:

An orthopedician is performing the following procedure to visualize the anterior cruciate ligament in a patient. Which of the following will be used to sterilise the instrument used?

Sold by @itachibot



a) Ethylene oxide

b) Gamma rays

c) Hydrogen peroxide

d) Ultraviolet rays

Sold by @itachi... purchased this from some... else, If yo... ...ay have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	a
3	c
4	d
5	c
6	a
7	c
8	b
9	b
10	a
11	b
12	c
13	d
14	d

Sold by @itachibot

15	d
16	c
17	b
18	c
19	b
20	b
21	c
22	a
23	c

@itachibot someone else,

Detailed Explanations

Solution to Question 1:

This is a modern compound microscope, and the condenser is located below the stage.

The purpose of the condenser lens is to focus the light from the illuminator onto the specimen. Microscopes with a stage condenser lens render a sharper image than those with no condenser lens.

The simple microscope was invented first by Antonie van Leeuwenhoek, the father of microscopy and microbiology. It uses 1 lens for magnification. The compound microscope, introduced by Hans and Zacharias Janssen uses 2 lenses, an objective lens and an eyepiece, to provide better magnification power than the simple microscope.

Solution to Question 2:

Sold by @itachibot

A painless genital ulcer with inguinal lymphadenopathy with the presence of spiral organisms on microscopy is suggestive of syphilis. It is caused by Treponema pallidum, demonstrated by dark-field microscopy.

Dark-field microscopy images have a dark background with a refractile organism. The dark ground microscope has an optical system that enhances the contrast of unstained bodies. It uses reflected light and has a higher resolving power than the light microscope. Dark-field microscopy is the the most specific and sensitive technique to visualize Treponema pallidum when an active chancre or condyloma lata is present.

Other options:

Option B: Phase contrast microscopy improves the contrast to delineate structures within cells. A dark image is formed on a light background.

Option C: Fluorescent microscopy uses fluorescent dyes like auramine-rhodamine to appreciate Mycobacterial species and malarial parasites.

Option D: The electron microscope, invented by Ernst Ruska, helps in appreciating viral structures. It has a very high resolving power.

Sold by @itachibot
If you bought this from someone else, you have been scammed.

Solution to Question 3:

80S ribosome is present in eukaryotes, while 70S ribosome is present in prokaryotes.

80S ribosome has subunits 60S and 40S, while 70S has subunits 50S and 30S.



Solution to Question 4:

The cell wall of bacteria cannot be seen with simple stains or direct light microscopy.

Sold by @itachibot

The bacterial cell wall is present outside the plasma membrane and confers strength and shape to the cell. It is composed of peptidoglycans. The cell wall can be demonstrated by the following methods:

• Plasmolysis

• Microdissection

• Reaction with specific antibodies

• Mechanical rupture of the cell

• Differential staining procedures (E.g. Chance's method)

• Electron microscopy

• Fluorescence microscopy

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 5:

Sporulation occurs in the stationary phase of the bacterial growth curve, which is marked as C in the image.

The bacterial growth curve is obtained when a batch of bacteria is seeded in a liquid medium and incubated. It consists of 4 distinct phases:

• Lag phase (A) - The bacteria adapt to a new environment. They synthesize and accumulate enzymes and metabolic intermediates required for multiplication. Thus, there is no increase in cell number but cell size increases. Maximum cell size is obtained towards the end of this phase.

• Log or exponential phase (B) - The bacteria start dividing and their numbers increase exponentially with time. Cells are smaller and stain uniformly. The growth rate is constant.

• Stationary phase (C) - There is depletion of nutrients and accumulation of toxic products. Thus, cell division reduces and a state of equilibrium is attained, where the number of new cells formed equals the number that die. Growth rate is zero. Sporulation occurs during this phase. Many bacteria produce secondary metabolic products such as exotoxins and antibiotics.

• Death phase or the phase of decline (D) - Growth rate is negative. The population decreases due to cell death from nutritional exhaustion, toxin accumulation and the action of autolytic enzymes. Involution forms are common.

Sold by @itachibot



Sold by @itachi... chased this from some... else, ...ave been scammed. If yo...

Solution to Question 6:

The presence of teichoic acid and lipoteichoic acid with a thick peptidoglycan layer is a feature of gram-positive bacteria.

The cell wall of bacteria provides shape and ridigity to the bacterial cell. It is composed of peptidoglycans.

	Gram positive	Gram negative
Teichoic acid, lipoteichoic acid	Present	Absent
Thickness (Peptidoglycan)	Thicker	Thinner
Variety of amino acids	Few	Several
Aromatic and sulfur-containing amino acids	Absent	Present
Lipopolysaccharides and outer membrane	Absent	Present

Sold by @itachibot



Other options:

Option C: The cell wall of fungi consists of chitin, mannan, and polysaccharides (glucan).

Option D: The cell wall of mycobacteria have mycolic acid, which is responsible for acid fastness.

Solution to Question 7:

Fimbriae help in bacterial adhesion.

Fimbriae, also called pili, are hair-like surface appendage that projects from the cell surface as straight filaments. They are primarily seen in Gram-negative bacteria. They are polymers of pilin. Fimbriae are classified based on the function as ordinary pili - for adhesion - and sex pili - involved in bacterial conjugation. The functions of pili are given below:

• Bacterial conjugation through sex pili, which transfer of genetic material occurs.

• Bacterial adhesion of symbiotic and pathogenic bacteria to host cells using adhesins.

• Resistance to phagocytosis.

• Antigenic pilin proteins are responsible for the immune response.

• Twitching motility to bacteria.

Solution to Question 8:

Pasteurization is done at 63 degree Celsius for 30 minutes.

Pasteurization is the process of applying heat to milk to kill or stunt the growth of pathogenic bacteria. It was developed by Louis Pasteur. There are two methods for pasteurization:

• Holder method: Milk is heated at 63°C for 30 minutes

Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Flash process: Milk is heated to 72°C for 15–20 seconds

Once heated appropriately, the milk is cooled quickly to 13°C or lower. All the non-sporing pathogens (mycobacteria, brucellae, salmonellae) are killed in pasteurization. In the Holder method, heat resistant Coxiella burnetti may survive.

Note: Louis Pasteur pioneered the techniques of pasteurization and sterilization. He also first advanced the germ theory of disease which states that most diseases were caused by microorganisms. Furthermore, he disproved the theory of spontaneous generation and developed vaccines for rabies and fowl cholera. He is thus known as the father of modern microbiology.

Solution to Question 9:

The method of sterilization that involves exposure at 100°C for 20 minutes for 3 successive days is called tyndallization or intermittent sterilization.

Tyndallization, which utilizes steam, was used for sterilization of media containing gelatin, egg, serum, or sugar. This method may not kill spores of certain anaerobes and thermophiles. Autoclaving has replaced this method now.

Other options:

Option A: Autoclaving involves heating at 121°C for 15 minutes at a pressure of 15psi. This is the used for intruments, linen, laboratory ware.

Option C: In pasteurization, milk is heated at either 63°C for 30 minutes (the holder method) or at 72°C for 15–20 seconds (the flash process), followed by cooling quickly to 13°C or lower.

Option D: Heating an article at 80–85 °C for 30 minutes on 3 consecutive days is called inspissation.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Lysol works by disrupting the cell membrane.

Lysol is a phenol compound, which is obtained by distillation of coal tar. Phenols cause cell membrane damage and inactivate membrane-bound oxidases and dehydrogenases. This causes lysis and release of cell contents. Low concentrations of phenol can precipitate proteins. They are not inactivated by organic matter and are thus useful disinfectants.

The mechnisms of action of other disinfecting agents are:

• DNA damage

• Ionizing radiation

• Ultraviolet light

• Hydrogen peroxide

• DNA reactive chemicals like acriflavin.

• Protein modification

Sold by @itachibot

• Alcohol

• Chlorine

• Iodine

• Heavy metals (e.g. thiomersal)

• Hydrogen peroxide

• Formaldehyde

• Glutaraldehyde

• Ethylene oxide

• Acids and alkalis

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 11:

The given image shows antibiotic sensitivity testing which uses the lawn culture method.

Lawn or carpet culture is the method in which a uniform surface growth of bacterium is present on the medium. It is used for antibiotic sensitivity testing (disc method) and bacteriophage typing. Stab culture (option D) is used for demonstration of gelatin liquefaction and oxygen requirements of bacterium, for maintenance of stock cultures. Streak culture (option A) is routinely used for the isolation of bacteria in pure culture from clinical specimens. Stroke culture (option C) is used for providing pure growth of bacterium for slide agglutination and other diagnostic tests. Other methods of culture include:

• Liquid culture - These are preferred for inocula containing antibiotics and other inhibitory substances.

• Pour plate culture - It gives an estimate of viable bacterial count in a suspension.



Sold by @itachibot

Solution to Question 12:

The given image shows Quellung reaction (halo around the bacteria) and is seen in capsulated bacteria. Neisseria gonorrhoeae is not capsulated, therefore Quellung reaction will not be observed in it.

In Quellung reaction, a suspension of a capsulated bacterium is mixed with its specific anticapsular serum and examined under the microscope. The anticapsular antibodies bind to the capsule and thus the capsule becomes very prominent and appears swollen. Capsulated organisms can be remembered with the following mnemonic:

Some - Streptococcus pneumoniae

Killers - Klebsiella pneumoniae

Have - Haemophilus influenzae

Nice - Neisseria meningitidis

Capsules - Cryptococcus neoformans

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 13:

The given clinical vignette of fever and hemoptysis in a labourer with the sputum smear showing acid-fast bacilli is suggestive of tuberculosis. Mycobacterium tuberculosis, detected on sputum smear, is appreciated by Ziehl-Neelsen staining, which would includes carbol fuschin - acid - alcohol - methylene blue.

Acid-fast bacteria (like MTB) in a stained Ziehl-Neelsen (ZN) smear appear bright red against a blue background. Steps involved in ZN staining are:

• Place the sample on slide evenly spread it with rotational movement

• Air-dry the slide and fix it by passing it over a flame

• Apply carbol fuschin stain and heat to fix it.

• Wash with water to remove excess stain

• Decolourise with alcohol

• Apply Methylene blue/Malachite green stain

• Wash with water to remove excess stain

• View under the microscope

Kinyoun modified the classical ZN staining such that heat fixing is not required. It is called the cold method. It involves increasing the concentration of phenol acid and the duration of staining. The slide is then washed with water and decolourized with acid alcohol. Counterstaining with Loeffler's methylene blue or malachite green is done for a minute and viewed under a microscope. Acid-fast bacteria in a stained smear appear bright red against a blue background.

The image attached below summarizes the steps.

Sold by @itachibot



Option A: It depicts the order of Gram staining, which was discovered by Hans Christian Gram in 1884.

• Primary staining with para-rosaniline dye (crystal violet, methyl violet or gentian violet).

• Application of dilute solution of iodine.

• Decolorisation with an organic solvent (ethanol, acetone, aniline).

• Counterstaining with the dye of contrasting colour (carbol fuchsin, safranine, neutral red).

Option B: It includes the steps for performing Albert's stain which is used for detected Corynebacterium diphtheriae.

• A heavy smear of the bacterial isolate is prepared on a glass slide and allowed to dry without fixing.

• Albert's stain is applied for (3-4) minutes. Then it is washed with water and Albert's iodine is applied for one minute.

• Wash with water and let dry with blotting paper.

• Examine the slide under oil immersion.

Corynebacterium diphtheriae are slender bacilli with an arrangement resembling the letters V or L. They have metachromatic granules at the poles of bacilli which are also called polar bodies, volutin, or Babes-Ernst granules. Albert staining is useful in the demonstration of metachromatic granules. The presence of metachromatic granules helps to distinguish C.diphtheriae from diphtheroids which lack them. The images below show Corynebacterium diphtheriae.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 14:

The aetiological agents of 1 and 2 are microaerophilic.

The given case scenarios are;

1. A businessman with epigastric pain and positive urease test has peptic ulcer disease, caused by Helicobacter pylori. It is microaerophilic.

2. Diarrhea caused by Gram-negative, small, spirally curved rod-like, motile bacilli describes Campylobacter jejuni. It is microaerophilic.

3. Rice-water diarrhea and Gram-negative curved bacilli with 'fish-in-stream' appearance describes cholera caused by Vibrio cholerae. It is a facultative anaerobe.

4. Nasopharyngeal swab from a child with paroxysmal cough showing 'mercury drops' colonies describes Bordetella pertussis. It is an obligate aerobe.

5. Dry, rough, raised, irregular colonies on LJ medium describes Mycobacerium tuberculosis. It is an obligate aerobe.

Microaerophilic organisms grow best in low oxygen tension (2-10% oxygen).

Note: Mycobacteria, such as M. bovis, can grow at oxygen tensions ranging from atmospheric to microaerophilic conditions. However, M. tuberculosis is an obligate aerobe.

Solution to Question 15:

The given image shows MacConkey agar. Cysteine is not a constituent of MacConkey agar.

MacConkey agar is a differential or indicator medium. Lactose-fermenting organisms (Eg: E. coli) produce a pink color, and non-lactose fermenting organisms (Shigella sp., Salmonella sp.) do not change the color. Contents of MacConkey agar include peptone, lactose, agar, neutral red and

Sold by @itachibot

sodium taurocholate (Mnemonic- PLANT). MacConkey agar is also a selective medium i.e. the bile salts (sodium taurocholate) and dyes (neutral red) inhibit the growth of Gram-positive organisms and Gram-negative cocci.

Solution to Question 16:

The image 'A' represents alkaline peptone water, which is an enrichment medium used for Vibrio cholerae.

The other images represent:

Image B- Blood agar with P. aeruginosa colonies

Image C- Loeffler's serum slope, which is used to isolate C. diphtheriae

Image D- Chocolate agar with F. tularensis colonies

Enrichment media suppress the growth of commensal bacteria and allow for the growth of certain organisms. These media are used for samples with mixed flora e.g. stool. They are always liquid media, like alkaline peptone water. Other examples include tetrathionate broth and selenite-F broth for Salmonella sp. and Shigella sp.

Enriched medium are those media to which additional nutrients such as blood, serum, or egg are added. They're used to grow bacteria with specific nutritional requirements. Examples include blood agar to test the hemolytic property of bacteria, chocolate agar for isolation of H. influenzae and Loeffler's serum slope for isolation of C. diphtheriae.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

The above image shows a swarming type of growth on the agar plate. It will not be caused by Corynebacterium diphtheriae.

Swarming is the growth of motile bacteria in the form of a thin film on solid culture media. It is exhibited by the following:

• Gram-positive bacilli - Clostridium tetani (option A), Bacillus cereus (option D)

• Gram-negative bacilli - Proteus mirabilis (option C), Proteus vulgaris

• Gram-negative halophilic bacteria- Vibrio parahaemolyticus

Swarming can be inhibited by increasing the concentration of agar (6%), alcohol, sodium azide, boric acid, chloral hydrate, sulfonamide, and surface-active agents.

Solution to Question 18:

Bacterial transduction occurs by bacteriophages.

Transduction is a process of acquiring bacterial DNA from a bacteriophage (virus) that has incorporated bacterial DNA from previous host bacteria.

Sold by @itachibot

Other methods of exchange of genetic material are transformation and conjugation. Transformation involves the uptake of genetic material from the environment, which is released by other bacteria. Conjugation is the direct transfer of DNA from one bacterial cell to another bacterial cell through a sex pilus or bridge. The transferred DNA is a plasmid, a circle of DNA that is distinct from the main bacterial chromosome.



Solo
If you purchased this
you may have been
else,

Solution to Question 19:

The primary principle used in polymerase chain reaction (PCR) is the amplification of small amounts of DNA.

PCR is an automated process in which a single or a few copies of a DNA piece is amplified to generate of thousands to millions of copies of that particular DNA sequence. There are 3 basic steps are involved in PCR:

• DNA extraction from microorganism

• Amplification of extracted DNA - Denaturation (95°C), Primer annealing (55°C), Extension (72°C)

• Gel electrophoresis of the amplified product

PCR is used in rapid diagnostic tests for infections (e.g. HIV DNA), forensic investigations etc. The founder of PCR is Karry B Mullis.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 20:

The enzyme involved in RT-PCR is reverse transcriptase (RT) which has RNA-dependent DNA polymerase activity.

In reverse transcription-polymerase chain reation (RT-PCR), reverse transcriptase synthesizes DNA from the RNA template. RT-PCR can be used for the demonstration of poliovirus RNA, causing paralytic poliomyelitis, in CSF. It can also be used for the detection of viruses from nasopharyngeal swabs (SARS-CoV2, H1N1).

In real-time RT-PCR, fluorescent dyes or DNA probes are used to quantify the pathogen DNA load in the specimen. So, it can be used to quantify viral loads in HIV, hepatitis B, etc., and is also called quantitative PCR (qPCR).



Sold by @itachibot

Solution to Question 21:

Jumping genes are also called transposons.

Transposons are genetic elements that can move between chromosomal and extra-chromosomal DNA molecules. Insertion of a transposon leads to acquisition of new characteristics by the recipient DNA. Transposons contain several genes, including those necessary for their migration from one genetic locus to another. They may contain genes for specialised functions like antibiotic resistance. However, they do not contain genes for their replication.

Jumping genes were discovered by Barbara McClintock and she was awarded the Nobel Prize in Physiology or Medicine 1983 for the same.

Other options:

Options A and B: The cytoplasmic carriers of genetic information are termed plasmids or episomes.

Option D: A cosmid is a type of hybrid plasmid that contains a Lambda phage cos sequence. (Cos sites + plasmid = cosmids).

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 22:

You should use a sanitiser before and after examining the patient and after touching the patients' surroundings. The hands should be washed with soap and water thoroughly in case they are visibly soiled.

Sanitization is the process through which pathogenic organisms are reduced to safe levels on inanimate objects to reduce the likelihood of cross-infections. It is a component of hand-hygiene, which is demonstrated by the image below.



Sold by @itachibot

Note: Disinfectants like chlorhexidine and alexidine have a biguanide functional group. They are mostly used as oral disinfectants and hand products. They are bactericidal but not sporicidal. Mycobacteria are also resistant to these compounds.

Solution to Question 23:

The given image shows arthroscopy being performed. Arthroscopes are sterilized using plasma sterilization which utilises hydrogen peroxide.

In plasma sterilisation, radiofrequency energy is applied to create an electromagnetic field through which hydrogen peroxide vapours are passed. This generates plasma which contains free radicals of hydrogen and oxygen. Plasma denatures all organisms and is used sterilize arthroscopes and ureteroscopes. Sterilization control is monitored by Geobacillus stearothermophilus and Bacillus atrophaeus.

Option A: Ethylene oxide has both microbicidal and sporicidal activity. It is used for sterilisation heat-sensitive equipment like heart-lung machines, respirators, suture materials etc. Sterilization control is monitored by Bacillius atrophaeus and Bacillus subtilus.

Option B: Gamma rays cause breakage of DNA. It destroys spores and vegetative cells, but is not effective against viruses. It is referred to as cold sterilisation and is used in commercial plants to sterilize plastics, syringes, swabs. Sterilization control is monitored by Bacillus pumilius.

Option D: Ultraviolet rays are used to sterilise biosafety cabinets, entryways, operation theatres and laboratories.

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Sold by @itachibot

Components of Immune system

Question 1:

What type of immunity is being conferred to a neonate receiving the BCG vaccine?

a) Natural active immunity

b) Natural passive immunity

c) Artificial active immunity

d) Artificial passive immunity

Question 2:

A 33-year-old with a large dirty wound on his knee and abrasions on his palms presents to the casualty. He does not remember the last time he received a TT injection and was thus given tetanus immunoglobulin. Which of the following is false about this type of immunity?

a) Acquired immediately

b) Memory is absent

c) Long lasting

d) No negative phase

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A patient with chronic mucocutaneous candidiasis was administered transfer factor as treatment. What type of immunity does this result in?

a) Active immunity

b) Passive immunity

c) Combined immunity

d) Adoptive immunity

Question 4:

Which of the following is not a pattern recognition receptor?

a) NOD-like receptors

Sold by @itachibot

b) Toll-like receptors

c) RIG-like receptors

d) BMP-like receptors

Question 5:

A 23-year-old presents with diarrhea, fever, and vomiting. The given bacteria is implicated. Which toll-like receptor binds to the structure of the bacteria marked A?

©MARROW

a) TLR 2

b) TLR 4

c) TLR 5

d) TLR 9

Question 6:

Which of the following statements is true about T cell dependent antigens?

a) Their response doesn't show immunological memory

b) They activate only mature B cells

c) They activate B cells polyclonally

d) They activate only IgG antibodies

Sold by @itachibot

Question 7:

A 28-year-old woman presents with a fever of 103°F, hypotension, confusion, and a diffuse rash as shown below. 3 days back, she had uncontrollable epistaxis for which a nasal pack was placed. What is the likely mechanism of action of the microbial toxin that has caused her current illness?

a) Enhancement of phagocytosis

b) Increased antigen presentation by macrophages

c) Polyclonal activation of T cells

d) Uncontrolled activation of complement

If you purchased ... you may have be... else, ...d.

Question 8:

What are the changes in the variable regions of an immunoglobulin known as?

a) Idiotype

b) Isotype

c) Allotype

d) Epitope

Question 9:

Which of the following statements about immunoglobulins is false?

Sold by @itachibot

a) IgG is used to diagnose fetal infection

b) IgM has weak precipitation activity

c) IgG has four subclasses

d) IgM is mainly intravascular

Question 10:

Which of the following immunoglobulins acts as surface immunoglobulin on B cells?

a) Ig G

b) Ig A

c) Ig M

d) Ig E

Question 11:

A 3-year-old child presents with a history of repeated sinopulmonary infections caused by encapsulated organisms. Which of the following is most likely to be deficient in this case?

a) IgG1

b) IgG2

c) IgG3

d) IgG4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

Identify the class of immunoglobulin■ shown in the image below.

Sold by @itachibot



©MARROW

a) Ig A

b) Ig G

c) Ig M

d) Ig E

Question 13:

Which immunoglobulin is majorly responsible for the reaction demonstrated below?



a) Ig G

b) Ig A

Sold by @itachibot

c) Ig M

d) Ig E

Question 14:

You are counseling a new mother about the importance of breastfeeding. What forms the secretory component of the antibody that is maximally present in breast milk?

a) Plasma cell only

b) Epithelial cells of lining mucosa only

c) Plasma cell and oligosaccharides

d) Oligosaccharides and epithelial cells

Question 15:

When papain acts on a molecule of immunoglobulin, which of the following are the end products?

a) 2 Fc and 1 Fab fragments

b) 1 variable chain and 1 constant chain

c) 1 Fc and 2 Fab fragments

d) 1 Fab and 1 hypervariable region

Question 16:

Affinity maturation is due to:

a) Isotype switching

b) Somatic hypermutation

c) mRNA splicing

d) VDJ rearrangement

Question 17:

An immunologist is conducting an experiment to understand precipitation reactions. He notes that on adding excessive amount of antigen to a serum sample containing small amount of antibodies, the amount of precipitate formed is less, compared to adding smaller amount

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

of antigen to the same sample. What is this phenomenon called?

a) Pro–zone phenomenon

b) Zone of equivalence

c) Post–zone phenomenon

d) Flocculation phenomenon

Question 18:

Double diffusion in two dimensions is called as _____.

a) Oudin procedure

b) Oakley – Fulthrope procedure

c) Radial immunodiffusion

d) Ouchterlony procedure

Question 19:

A patient presented to the casualty with the following finding. A positive Nagler's reaction confirmed the presence of lecithinase-producing organisms. What is the principle behind this test?



a) Precipitation

b) Complement fixation test

c) Agglutination

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Sold by @itachibot

d) Neutralisation

Question 20:

A patient presents with a painless ulcer on his penis. A Wasserman test is done and is reactive. What is the principle of this test?

a) Precipitation

b) Complement fixation

c) Agglutination

d) Neutralisation

Question 21:

Which of the following activates the classical complement pathway?

a) Immune complexes

b) Mannose residues

c) Hydrolysis of C3

d) Endotoxins

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 22:

In the alternate complement pathway, which of the following functions as C3 convertase?

a) C4b2b

b) C3bB

c) C3bBb

d) C4b2a

Question 23:

A 23-year-old girl presents with the finding shown below. She has a history of multiple similar episodes in the past and reports that some of her family members have similar complaints. She has no itching. Which of the following is deficient in this patient?

Sold by @itachibot

a) C3b inactivator

b) C1, C2, C4

c) C1 inhibitor

d) C5-C8

Sold by @itachi... purchased this from some... else, ...may have been scammed.

Answer Key

Question No.	Correct Option
1	c
2	c
3	d
4	d
5	c
6	b
7	c
8	a
9	a
10	c
11	b
12	a
13	c
14	b

Sold by @itachibot

15	c
16	b
17	c
18	d
19	d
20	b
21	a
22	c
23	c

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The immunity conferred to a neonate receiving the BCG vaccine is artificial active immunity, as it involves administering a live attenuated vaccine form of Mycobacterium bovis to stimulate an adaptive immune response and develop antibodies, establishing long-term immunity.

Types of adaptive immunity:

• Natural active immunity: It results from either a clinical or an inapparent infection by a microbe.

• Artificial active immunity: It is the resistance induced by vaccines which are preparations of killed or attenuated microorganisms or their products.

• Natural passive immunity: It is the immunity passively transferred from mother to baby. In human infants, maternal antibodies are transmitted predominantly through the placenta, while in animals such as pigs, the transfer of antibodies occurs mainly through the colostrum.

• Artificial passive immunity: It is the resistance passively transferred to a recipient by the administration of antibodies. Hyperimmune sera of animal or human origin, convalescent sera, and pooled human immunoglobulin are used for this purpose.

• Combined immunization: A combination of the active and passive methods of immunization is used.

Examples of the above types of immunity:

Note: Adoptive immunity is immunity in which immunologically competent lymphocytes are injected. The transfer factor is composed of these immunologically competent lymphocytes. It can passively transfer cell-mediated immunity against those antigens to which the donor is sensitive.

Immunity	Description
Natural active	Clinical or subclinicalinfection
Artificial active	Vaccination

Sold by @itachibot

Immunity	Description
Natural passive	Transplacentaltransfer of antibody,colostrum
Artificial passive	Anti-tetanus serum, anti-diphtherial serum
Combined immunization	Antisera and vaccine are given at thesame time,e.g- post-exposure prophylaxis for rabies

Solution to Question 2:

Giving pre-formed immunoglobulin (e.g. Tetanus immunoglobulin) directly is an example of passive immunity, which is not long-lasting.

The differentiating features between active and passive immunity:

Negative phase: It is the time period during which the level of measurable immunity is lower than it was before the antigenic stimulus. This is because the antigen combines with any pre-existing antibody and lowers its level in circulation. It is seen at the beginning of the active immune response.

Secondary type response: If an individual who has been actively immunized against an antigen experiences the same antigen subsequently, the immune response occurs more quickly and abundantly than during the first encounter.

Active Immunity	Passive Immunity
Actively produced by the host's immune system	Passively received by the host
Induced by infection or immunogens	Readymade antibody is transferred
Immunological memory present	No memory
Immunity is effective after a lag period or latent period	Immediate immunity
Broad specificity	Narrow specificity
A negative phase may occur	No negative phase.
Booster effect on subsequent dose	The subsequent dose is less effective
Secondary type response seen	No secondary type of response
More effective	Less effective
Long-lasting (Years)	Short-lasting (Days to weeks)
Example: Vaccines	Example: Antisera

If you purchased this from someone else, it may have been scanned. Sold by @itachibot

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 3:

Adoptive immunity occurs when transfer factor is given.

Transfer factor is composed of an extract of immunologically competent lymphocytes such as T lymphocytes, NK cells, macrophages, or B cells. It has the ability to passively transfer cell-mediated immunity against antigens to which the donor is sensitive.

It has been attempted in the treatment of lepromatous leprosy and chronic mucocutaneous candidiasis.

Solution to Question 4:

Bone morphogenetic protein-like receptors are not pattern-recognition receptors.

Pattern recognition receptors (PRRs) are essential for the innate immune system. They identify pathogen-associated molecular patterns (PAMPs) common to classes of microbes and trigger immune responses. These receptors can be found on the cell surface, detecting extracellular pathogens, or within the endosomes and cellular matrix, detecting intracellular invading viruses.

PRRs activate pro-inflammatory pathways, promote phagocytosis in cells like macrophages, neutrophils and dendritic cells, and may act as secreted proteins binding directly to pathogens. Based on the localization and functional characteristics, PRRs are classified as:

• Membrane-bound PRRs:

• Toll-like receptors (TLR)- detect microbial carbohydrates and lipoproteins (Option B)

• C-type lectin receptors (CLR)- detect mannose residues

• Cytoplasmic (intracellular) PRRs:

Sold by @itachibot

• Nucleotide oligomerization (NOD)-like receptors (NLR) - NOD1 and NOD2 - Detects bacterial cell wall peptidoglycans (Option A)

• RIG-I-like receptors (RLRs) - detect viral RNA (Option C)

• Secreted PRRs

On the contrary, BMP receptors are multifunctional transmembrane serine-threonine kinase receptors that belong to the transforming growth factor β (TGF-β) superfamily, regulating cell growth, differentiation, and apoptosis with no role in pattern recognition. Also, there are no receptors termed "BMP-like receptors".

Note: A mutation in the Bone morphogenetic protein receptor 2 (BMPR2) gene enhances the proliferation and reduces the apoptosis of pulmonary vascular endothelial and smooth muscle cells, leading to hereditary pulmonary arterial hypertension (PAH).

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 5:

The given image and clinical history suggest a typhoid infection. The structure marked A is the flagella of bacteria. Toll-like receptor (TLR) 5 binds to the flagella of bacteria.

Toll-like receptors (TLRs) are transmembrane receptors that have an extracellular domain, that detect infecting pathogens.

• TLR 1 - Mycobacteria and Gram-negative bacteria

• TLR 2 - Mycobacteria and Gram-positive bacteria

• TLR 3 - Viral RNA

• TLR 4 - Gram-negative bacteria

• TLR 5 - Flagella

• TLR 9 - Bacterial DNA, some herpes viruses, malarial parasite

TLRs act through the NF-KB pathway and initiate inflammation.

Solution to Question 6:

T cell-dependent antigens activate only mature B cells.

Depending on their ability to induce the formation of antibodies, antigens are classified as T-cell dependent and T-cell independent antigens.

T cell-dependent antigens (TD)	T cell-independent antigens (TI)
Require helper T cell participation for the production of immune response	Directly stimulate antibody production by B cells
Structurally complex	Structurally simple

Sold by @itachibot

T cell-dependent antigens (T D)	T cell-independent antigens (TI)
Activate only mature B cells monoclonally	Activate both mature and im mature B cells polyclonally
B cells undergo affinity matu ration and class switch over	B cells do not undergo affinit y maturation and class switc h over
All classes of antibodies are a ctivated	Antibody response is restrict ed to IgM and IgG3
Rapidly metabolized	Slowly metabolized
Immunological memory pres ent	No immunological memory

Sold by @itachibot
If you purchased this from someone else, it may have been scanned.

Solution to Question 7:

This woman presented with fever, hypotension, and a diffuse erythematous rash, three days after a nasal packing was done. This is characteristic of toxic shock syndrome - commonly caused by staphylococcal toxic shock syndrome toxin-1 (TSST-1), which is a superantigen.

Superantigens act by causing polyclonal activation of T cells. They activate a large number of T cells irrespective of their antigen specificity.

In contrast to regular antigens which bind to the binding groove on T cells, superantigens bind outside the groove directly to the lateral aspect of the TCR β chain. This leads to a massive release of cytokines and polyclonal activation of T cells.

Clinically, this manifests as fever, hypotension, and skin exfoliation.



Examples of organisms that produce superantigens:

• Bacterial

Sold by @itachibot

- Staphylococcus aureus- Enterotoxins
- Group A streptococci - Pyrogenic exotoxin
- Mycoplasma arthritidis- mitogen 1
- Yersinia enterocolitica
- Yersinia pseudotuberculosis
- Viral
- Epstein–Barr virus (EBV)
- Cytomegalovirus (CMV)
- Rabies nucleocapsid
- HIV-encoded negative regulatory factor.
- Fungal- Malassezia furfur

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

Changes in the variable regions of heavy and light chains of an immunoglobulin are called an idiotype.

Other options:

Option B: Isotypes are the changes in the constant regions of the heavy chain of the immunoglobulin.

Option C: Allotypes refer to multiple alleles that exist for some genes. This leads to subtle amino acid differences that occur in some but not all members of the same species.

Option D: An epitope is the smallest unit of antigenicity and it is an antigenic determinant.

Solution to Question 9:

IgG cannot detect fetal infections because it is not produced by the fetus in significant amounts.

IgG is the only maternal immunoglobulin that is normally transported across the placenta and provides natural passive immunity to the newborn.

Solution to Question 10:

Monomeric IgM is the major antibody receptor on the surface of B lymphocytes for antigen recognition.

IgD is the other class of antibodies which acts as the B cell receptors.

Sold by @itachibot

Solution to Question 11:

Deficiency of IgG2 results in recurrent sinopulmonary infections with encapsulated organisms.

IgG is the only maternal immunoglobulin that is normally transported across the placenta and provides natural passive immunity to the newborn. IgG1 is the antibody maximally transported across the placenta, whereas IgG2 has the least transplacental movement.

Deficiencies of other IgG:

• IgG1- bacterial infections (diphtheria, tetanus)(Option A)

• IgG3- viral infections (Option C)

• IgG4- parasitic infections (Option D)

Solution to Question 12:

The above image shows a dimeric structure, characteristic of secretory IgA.

IgA immunoglobulin is the major antibody responsible for mucosal immunity. Its serum levels are usually low. It is predominantly present in the extravascular secretions. It is present in secretions such as milk, saliva, and tears, and in secretions of the respiratory, intestinal, and genital tracts.

When present in the serum, it exists as a monomer similar to IgG. However, when it is secreted, it exists as a dimer. The dimeric structure consists of two monomers, which are stabilized by two additional polypeptides:

• J chain, which stabilizes the molecule.

• Secretory component, which is incorporated into the IgA when transported through the epithelial cell.



If you purchased this from someone else, it may have been scammed.
Sold by @itachibot
MARROW

Sold by @itachibot

Solution to Question 13:

The image shows blood typing, which works on the principle of agglutination. IgM is mainly responsible for agglutination.

It is the most efficient immunoglobulin in agglutination, complement fixation, and other antigen–antibody reactions and is important also in defense against bacteria and viruses.

It is a pentamer, and hence is more effective than IgG in opsonization (C3b mediated), bactericidal action and bacterial agglutination but less effective in neutralization of toxins and viruses.

Reaction	IgG	IgM
Precipitation	Strong	Weak
Agglutination	Weak	Strong
Complement fixation	Moderate	Strong
Lysis	Weak	Strong

Sold to someone else, If you purchased this ... you may have been scammed.

Solution to Question 14:

The antibody maximally present in breast milk is IgA. The secretory component of IgA is formed by epithelial cells of lining mucosa.

IgA occurs in two forms- serum IgA (monomer) and secretory IgA.

Secretory IgA is a dimer formed by two monomer units joined together by a J-chain and a secretory component (secretory piece). Secretory IgA along with the J chain is synthesized by the plasma cells situated near the mucosal or glandular epithelium. The secretory component is synthesized by epithelial cells, not by plasma cells.

It is believed to protect IgA from denaturation by bacterial proteases.

Sold by @itachibot



Solution to Question 15:

When papain acts on a molecule of immunoglobulin, the end products are 1 Fc and 2 Fab fragments.

Enzymatic digestion of antibody:

- Papain- 2 Fab fragments and one Fc fragment
- Pepsin- One F(ab')2 fragment and many smaller fragments
- Mercaptoethanol- 2 H and 2 L chains



Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 16:

Affinity maturation is due to somatic hypermutation.

Affinity maturation is the process by which antibodies gain increased affinity, avidity, and anti-pathogen activity. It is a result of somatic hypermutation (SHM) of immunoglobulin genes in B cells, coupled to selection for antigen binding.

This process occurs in the germinal centers and proceeds for weeks after acute infection or vaccination, or for many cycles during chronic infection. The resulting antibodies can be highly mutated from their germline-encoded counterparts, with increases of several orders of magnitude in affinity for antigen compared to the corresponding naïve B cell receptors (BCRs).

Other options:

Option A: Isotype switching, also known as class switch recombination, is a process where the constant region of the antibody heavy chain is changed, leading to a switch in the antibody isotype (e.g., from IgM to IgG). This process is crucial for the functional properties of antibodies but is not directly involved in affinity maturation.

Option C: mRNA splicing is a cellular process that involves the removal of introns and the joining of exons to generate mature mRNA. While this process is essential for the production of functional antibodies, it is not specifically associated with affinity maturation. Affinity maturation primarily occurs through somatic hypermutation.

Option D: VDJ rearrangement is the process by which the variable regions of immunoglobulin genes are assembled from gene segments (V, D, and J segments). This process contributes to the diversity of antibodies but is distinct from affinity maturation. Affinity maturation occurs after VDJ rearrangement and is driven by somatic hypermutation in the variable regions.

Solution to Question 17:

Sold by @itachibot

This phenomenon is called post–zone phenomenon because it has excessive antigen in the test system, leading to weak or even absent precipitation.

Lattice hypothesis explains precipitation reactions. The amount of precipitate formed is greatly influenced by the relative proportions of antigens and antibodies.

• Prozone phenomenon is caused by excessive antibodies in the test system. (weak/absent precipitation- false-negative test)

• Zone of equivalence- antigen, and antibody are in optimum proportions and maximum precipitation occurs.

• Post–zone phenomenon is due to the presence of excessive antigen in the test system. (weaker/absent precipitation)



Flocculation: Instead of sedimenting, precipitate remains suspended as floculus.

Solution to Question 18:

Double diffusion in two dimensions is called the Ouchterlony procedure. Ex: Elek's test for testing toxigenicity in diphtheria bacilli.

The images below show the different types of immunodiffusion.

Sold by @itachibot

Single diffusion in one dimension-
Oudin procedure



©MARROW

Double diffusion in one dimension-
Oakley-Fulthorpe procedure



©MARROW

Single diffusion in two dimensions-
Mancini procedure, radioimmunodiffusion



©MARROW

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 19:

Nagler's reaction is an example of a neutralization test. It is done for the detection of lecithinase toxin produced by C. perfringens.

The alpha-toxin of C. perfringens has phospholipase activity. When grown on a medium containing egg yolk phospholipid, an area of opacity around the bacterial colonies is seen (Nagler's positive). The presence of antitoxin can inhibit this.

Other examples of neutralization tests are:

- Plaque inhibition test- neutralization of bacteriophages
- Schick test- neutralization of diphtheria toxin by antitoxin
- Anti-streptolysin O test- for beta-hemolytic Streptococci.



Sold by @itachibot

Solution to Question 20:

Wassermann's reaction is an example of a complement fixation test. It is a non-specific test for the diagnosis of syphilis.

An antigen combines with its (specific) antibody in the presence of complement, and the antigen-antibody complex adsorbs the complement. This reaction is not visible and requires the use of an indicator (sheep erythrocytes and the specific anti-erythrocyte antibody).



Solution to Question 21:

The classical pathway of complement is activated by immune complexes or antigen-antibody complexes. (antigen bound to IgM or IgG).

Lipopolysaccharide of gram-negative bacteria activates the alternative complement pathway.

The binding of mannose-binding lectin to mannose residues (on the surfaces of Salmonella, Neisseria, and Listeria) activates the lectin complement pathway.

Sold by @itachibot



Solution to Question 22:

C3 convertase in the alternate pathway is C3bBb.

In the classical pathway and lectin pathway, the C3 convertase is C4b2a.



Solution to Question 23:

This girl presents with features of hereditary angioneurotic edema, which is due to a deficiency of C1 inhibitor.

Hereditary angioneurotic edema is an autosomal dominant disorder that is more common than complement deficiency states. A deficiency of C1 inhibitor leads

Sold by @itachibot

to unregulated activation of kallikrein. This increases the production of vasoactive peptides such as bradykinin. It causes episodes of edema affecting the skin and mucosal surfaces such as the larynx and the gastrointestinal tract. This may result in life-threatening asphyxia or nausea, vomiting, and diarrhea after minor trauma or emotional stress.

Note -

• C1, C2, C4 deficiency - collagen vascular disorders.

• C5-C8 deficiency - recurrent bacteremia with Neisseria

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Structure and Functions of the Immune System & Immune Response

Question 1:

In which of the following areas do B lymphocytes reside?

a) Spleen follicles

b) Lymph node paracortex

c) Periarteriolar lymphoid sheath

d) Thymic medulla

Question 2:

A mother brings her 1-year-old child to the hospital with complaints that he turns blue while breastfeeding. It resolves when he assumes a squatting position. On examination, a cleft palate is seen. Labs show hypocalcemia. Which of the following abnormalities is expected to be seen in the lymphoid organs?

a) Decreased size of germinal centres of lymph nodes

b) Decreased size of paracortical areas of lymph nodes

c) Increased size of splenic follicles

d) Increased size of the peri arteriolar lymphoid sheath of spleen

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 3:

Match the following specialized macrophages with the tissues they are found in.

a) I-A; II-C; III-D; IV-E; V-B

b) I-C; II-B; III-D; IV-E; V-A

c) I-D; II-C; III-A; IV-B; V-E

d) I-B; II-C; III-D; IV-E; V-A

Question 4:

Which of the following is the most potent stimulator of naive T-cells?

Sold by @itachibot

a) Mature dendritic cells

b) Follicular dendritic cells

c) B lymphocytes

d) NK cells

Question 5:

Nude mice are not resistant to xenograft due to the absence of _____

a) B cell

b) T cell

c) Both B and T cell

d) NK cell

Question 6:

Which of the following statements about T-cell receptors is false?

a) In 95% of T cells, the TCR consists of ° and β polypeptide chains.

b) γδ TCR recognize peptide antigens presented by MHC molecules.

c) Each TCR is non-covalently linked to six polypeptide chains, which form CD3 complex.

d) The linked CD3 complex and ζ chains initiate activating signals.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

A 52-year-old requires a kidney transplant due to chronic kidney disease. Her 2 children match her HLA type 4/6 and 5/6 respectively. On what chromosome is this match protein sequence located?

a) Chromosome 5

b) Chromosome 6

c) Chromosome 7

d) Chromosome 8

Question 8:

Which of the following statements is false regarding MHC molecules?

Sold by @itachibot

a) MHC class II presents peptide antigen to CD4 T cells

b) The action of MHC II occurs via endocytic pathway

c) MHC Class II presents endogenous antigens

d) MHC II is present on antigen presenting cells

Question 9:

You are observing flow cytometry being performed at a laboratory for MHC class 1 antigen. Identify the cells that lack this antigen.

a) Platelets

b) T lymphocytes

c) B lymphocytes

d) Red blood cells

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A patient newly diagnosed with HIV is found to have a CD4+ count of 78cells/μL. Which of the following is not a function of these cells?

a) Immunogenic memory

b) Produce immunoglobulins

c) Activate macrophages

d) Activate cytotoxic cells

Question 11:

Which of the following statements is false regarding the surface markers of T and B cells?

a) T cells express CD 3

b) T cells express CD 10

c) B cells express CD 19

d) B cells express CD 21

Question 12:

Which of the following markers is used for the identification of memory T cells?

Sold by @itachibot

a) CD45RA

b) CD45RB

c) CD45RC

d) CD45RO

Question 13:

In a patient with influenza, there is decreased expression of MHC proteins on the surface of the virus-infected cells. Which component of the immune system acts on such cells?

a) Cytotoxic T cells

b) Macrophages

c) NK cells

d) Helper T cells

Sold by @Itachibot
If you purchased this from someone else you have been scammed.

Question 14:

A 9- year-old boy was brought with colicky pain, nausea, vomiting. Lab investigations revealed eosinophilia and stool examination shows the following egg. Which interleukin is secreted in this scenario?

a) IL-1

b) IL-3

c) IL-6

Sold by @itachibot

d) IL-5

Question 15:

Macrophages are the major source of _____.

a) IL-1

b) IL-5

c) IL-7

d) IFN-γ

Question 16:

Which of the following mediators of inflammation is an example of a C-X-C chemokine?

a) Lipoxin LXA– 4

b) Interleukin IL– 8

c) Lymphotactin-° XCL1

d) Fractalkine CX3CL1

Question 17:

Which of the following is not an antigen-presenting cell?

a) Helper T cells

b) Macrophages

c) Fibroblasts

d) Dendritic cells

Question 18:

A surgeon takes the following 2 images of a patient's arm 6 weeks apart. Which of the following mediators promote the process shown below?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) TGF-β

b) IFN-°

c) TNF-°

d) IFN-β

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

A patient's plasma has high levels of interferon beta. He most likely has an infection due to _____

a) Virus

b) Bacteria

c) Fungi

d) Mycoplasma

Question 20:

A newborn diagnosed with a hepatoblastoma is planned for liver transplantation. Her father is found to be an HLA match, and he donates part of his liver. What type of transplantation is described here?

a) Autograft

b) Allograft

c) Xenograft

Sold by @itachibot

d) Isograft

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	a
5	b
6	b
7	b
8	c
9	d
10	b
11	b
12	d
13	c
14	d
15	a
16	b
17	a
18	a
19	a
20	b

Detailed Explanations

Solution to Question 1:

B lymphocytes reside in the splenic follicles.

Lymphocytes are part of the adaptive immune system. They include B lymphocytes and T lymphocytes. B lymphocytes mature in the bone marrow and then migrate to secondary lymphoid organs - follicles of spleen and in follicles of lymph nodes. T lymphocytes are produced in the bone marrow but mature in the thymus. They are also present in the periarteriolar lymphoid sheaths (PALS) of spleen and paracortex of lymph node.

Sold by @itachibot

In a lymph node, follicles may contain a central region called germinal centre. This contains B cells that have been stimulated by an antigen.





Solution to Question 2:

Turning blue on breastfeeding with resolution on squatting describes a tet-spell or a cyanotic spell, indicating a congenital cardiac defect. Cleft palate, cardiac defects and hypocalcemia (parathyroid hypoplasia/aplasia) suggest a diagnosis of DiGeorge syndrome. In this disorder, a decreased size of paracortical areas of lymph nodes is noted.

DiGeorge syndrome is characterised by thymic aplasia, causing a deficiency of T cells and absent cell-mediated immunity. Thus, paracortical areas of lymph nodes and periarteriolar sheaths of the spleen, where T cells reside, are reduced in size. B cells are usually not affected.

Sold by @itachibot

Solution to Question 3:

The correct answer is I-B; II-C; III-D; IV-E; V-A.

Macrophages can be divided into:

1. Blood macrophages (monocytes) are the largest of the myeloid cells found in the peripheral blood. They survive for 3 days.

2. Tissue macrophages present in different organs. They survive for months.

The main functions of macrophages are phagocytosis and antigen presentation.

The specialised (tissue-specific) macrophages present in different tissues are listed below:

- Peripheral blood: Monocytes
- Placenta: Hofbauer cells
- Liver: Kupffer cells
- Brain: Microglial cells
- Kidney: Mesangial cells
- Lungs: Alveolar macrophages
- Bone: Osteoclasts
- Connective tissue: Histiocytes

If you purchased this from someone else, you have been scammed. Sold by @itachibot

Cell	Tissue
I. Hofbauer cells	B. Placenta
II. Histiocytes	C. Connective tissue
III. Kupffer cells	D. Liver
IV. Microglia	E. Brain
V. Mesangial cells	A. Kidney

Solution to Question 4:

Mature dendritic cells are the most potent stimulator of naive T cells.

Naive CD4+ T cells are activated when they come in contact with two ligands - MHC class II and a costimulary membrane protein (eg: CD28). Both of these are present in antigen-presenting cells, the most potent of which is the mature dendritic cell. Other professional antigen-presenting cells include macrophages and B lymphocytes (option C).

Other options:

Sold by @itachibot

Option B: Follicular dendritic cells are functionally different from dendritic cells. They are not antigen-presenting-cells and do not activate naive T cells.

Option D: NK cells do not stimulate naive T cells.



Solution to Question 5:

Nude mice are not resistant to xenograft due to absence of T cell.

A xenograft is a tissue graft or organ transplant from a donor of a different species.

Nude mice lack body hair, hence the name. They lack a thymus and thus cannot generate mature T lymphocytes. The absence of T-lymphocytes makes them unable to mount adaptive immune responses requiring helper T cells and cytotoxic T cells. The absence of functioning T cells also prevents the rejection of allografts & xenografts.

Solution to Question 6:

γδ TCR recognize different type of antigens without the requirement of major histocompatibility complex (MHC)

Each T cell recognizes a specific bound antigen by means of an antigen-specific T-cell receptor (TCR). The TCR is a heterodimer and is of two types:

• αβ TCR (95%) recognize peptide antigens that are presented by MHC molecules on the surfaces of antigen-presenting cells.

• γδ TCR (5%) recognize peptides, lipids, and small molecules, without a requirement for display by MHC proteins. They have particular localization patterns in the mucosal tissues, which protect against microbes that try to enter through epithelia.

Sold by @itachibot

TCRs are non-covalently bound to six polypeptide chains or three dimers (a δε pair, a γε pair, and a ζζ or ζη pair), forming the CD3 complex and the ζ dimer. The CD3 complex and ζ dimer convey signals received by the receptor to the interior of the cell.

Solution to Question 7:

HLA complex of genes is located on the short arm of chromosome 6 (chromosome 6p21).

The human leukocyte antigen (HLA) system or complex is a gene complex encoding the major histocompatibility complex (MHC) proteins. In humans, the MHC molecules are called human leukocyte antigens (HLA) because they were initially detected on leukocytes by the binding of antibodies. The combination of HLA alleles in each individual is called the HLA haplotype. The function of MHC molecules is to display peptide fragments of protein antigens for recognition by antigen-specific T cells.

The HLA system is highly polymorphic. Each individual's HLA alleles differ from the other. It has to be checked before organ transplantation. The HLA complex consists of 3 separate clusters of genes:

• Class I– A, B and C loci

• Class II– DR, DQ and DP loci

• Class III– Complement C2 and C4 of classical pathway and properdin factor B of the alternate pathway, heat shock proteins(HSP), TNF ° and β.

If you purchased this from someone else, it may have been scammed. Sold by @itachibot

Solution to Question 8:

MHC class II presents exogenous antigens while MHC class I presents endogenous antigens.



Sold by @itachibot

Solution to Question 9:

Red blood cells are anucleated cells that lack MHC class I antigens.

MHC class I antigens are present on platelets and all nucleated cells, except sperm cells. They are responsible for the presentation of peptide antigen to CD8 T cells.

The human major histocompatibility complex (MHC), or the HLA complex, is a cluster of genes whose products are responsible for antigen presentation to T cells, transplant rejection, etc. HLA complex of genes is present on the short arm of chromosome 6. The genes are clustered in three regions named MHC region-I, II and III.

MHC class II antigens are present only on antigen-presenting cells (macrophages, dendritic cells, B cells). They present the peptide antigen to CD4 T cells. APCs also contain MHC class I antigens.

MHC region III comprises of genes that code for complement factors like C2, C4, C3 convertase, factor B and properdin, heat shock protein, tumor necrosis factor and steroid 21-hydroxylases.

Solution to Question 10:

CD4+ is a marker of T helper and memory cells. They do not produce immunoglobulins.

CD4+ T cells are activated when an antigen-presenting cell expressing MHC-II binds to them. The activated CD4+ T cell differentiates into T helper (TH) cells and regulatory T cells. The TH cells are further differentiated into TH1, TH2 and TH17 cells. Helper T cells have the following functions:

• Activation and proliferation of cytotoxic T cells

• Regulation of monocyte-macrophage system

• Activation of B cell to produce plasma cells

• Production of interleukins

Memory cells (Tm) have both CD4+ and CD8+ markers. They provide anamestic immune response and are thus responsible for immunogenic memory.

Immunoglobulins are produced by the plasma cells.

Solution to Question 11:

T cells do not express CD10. CD10 is a marker of pre-B cell and germinal center B cells.

Given below is a list of cell surface markers and the cells they are expressed on:

• CD3: All T cells

• CD4: Helper T cells

• CD8: Cytotoxic T cells

• CD4, CD25: Regulatory T cells

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

• Ig, CD19, CD20, CD21: B Cells

• CD10: Pre-B Cells, germinal center B cells

• CD40, MHC-II: Macrophages

• CD16, CD56: NK Cells

• CD34: Hematopoietic stem cells

Note : Pan T antigens are antigens found on all T cells and include CD2, CD3, CD5 and CD7.

Solution to Question 12:

CD45RO is used as a marker for the identification of memory T cells.

CD45RA is a marker for virgin or nai■ve T helper cells.

Solution to Question 13:

Natural killer cells (NK cells) act on virus-infected cells with reduced MHC expression.

NK cells are a subtype of lymphocytes that kill damaged or infected cells. NK cells express CD16 and CD56. Their action is inhibited when MHC I is expressed on normal cells. This prevents NK Cells from killing normal cells. In the presence of a viral infection or malignancy, MHC expression of a cell is reduced. Thus, NK cells get activated and kill virus-infected/tumor cells. NK cells also secrete cytokines (IFN-γ) which activate macrophages (option B) for destruction of ingested microbes.

Other options:

Option A: Cytotoxic T cells act when cells expressing MHC I present endogenous antigens.

Option D: Helper T cells act when cells expressing MHC II (antigen-presenting cells) present exogenous antigens.

Solution to Question 14:

A child with colicky abdominal pain, nausea, vomiting with eosinophilia likely has a parasitic infection with Ascaris lumbricoides (egg shown in the image). Parasitic infections trigger the release of IL-5.

In a parasitic infection, Th2 cells are stimulated and they produce IL-4 and IL-5. IL-4 stimulates the differentiation of B-cells into IgE-secreting plasma cells, while IL-5 stimulates the maturation and release of eosinophils from the bone marrow. The secreted IgE coats the parasite. The activated eosinophils bind to the IgE-coated parasite and kill them through the release of major basic protein.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Solution to Question 15:

Macrophages are the major source of IL-1.

Macrophages are phagocytic cells that secrete the following molecules:

• Enzymes - Lysozyme, proteases, elastases, collagenases, plasminogen activator.

• Cytokines - Interleukins 1, 6, 8, 10, 12; TNF-° ; TGF-β; prostaglandins.

• Platelet-derived growth factor (PDGF), Platelet activated factor (PAF).

• Colony-stimulating factor (CSF), Angiotensin-converting enzyme (ACE)

• Complements - C2, C4.

• Reactive O2 species (ROS)- H2O2, NO, OH.

Interferon-gamma (IFN-γ) is secreted by T-cells. It is one of the most potent macrophage activators.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

Interleukin-8 is an example of a C-X-C chemokine.

Chemokines are chemotactic (chemoattractant) cytokines. They recruit leukocytes to sites of inflammation. The C-X-C or alpha chemokines have a total of four cysteines, of which the first two separated by one amino acid. represented by 'X'.

Solution to Question 17:

T cell is not an antigen-presenting cell.

Antigen presenting cells contain MHC class II molecules on their surface. They digest pathogens at the site of infection and display those exogenous antigens to T lymphopcytes, thus leading to T lymphocyte activation. Examples of antigen-presenting cells include dendritic cells, macrophages, B cells, glial cells (brain), fibroblasts (skin), thymic epithelial cells and vascular endothelial cells.

Solution to Question 18:

The image shows wound healing. TGF-β (transforming growth factor-β) is responsible for wound healing.

TGF-β is produced by B and T cells and it inhibits their proliferation. It acts as a growth factor for fibroblasts and promotes wound healing. It stimulates the production of collagen, fibronectin and proteoglycans. It also promotes class switching of B cells to IgA. IL-10 is also involved in wound healing and suppression of inflammation.

Sold by @itachibot

Option B: IFN-° inhibits viral replication, activates NK cells, and regulates the activity of macrophages and T cells.

Option C: TNF-° is involved in tumour cytotoxicity, phagocytic cell activation, and antiviral and antiparasitic effects. It is an acute phase protein.

Option D: IFN-β has antiviral activity.

Solution to Question 19:

Interferon-beta is secreted during infection due to viruses.

Interferons are classified as cytokines. Interferons are non-toxic, non-antigenic and species-specific molecules. They are inactivated by proteases. They are stable at pH over 2-10, except INF-γ (labile at pH 2) and are non-dialysable and non-sedimentable.

There are three types of interferons - ° , β and γ. IFN-° or leukocyte interferon is produced by leucocytes. It has antiviral activity, is immunoregulaotry and antiproliferative. IFN-β or fibroblast interferon is produced by fibroblasts and epithelial cells. It has antiviral activity and is thus increased in viral infections. It enhances MHC class 1 and NK cell activity. IFN-γ or immune interferon is produced by T-lymphocytes, NK cells . It activates macrophages, increases neutrophil function, enhances both MHC class 1 and class II expression and has anti-proliferative (anti-tumour) functions.

Solution to Question 20:

The transplant of an organ/tissue from a father to child is an example of an allograft.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Hypersensitivity

Question 1:

What route of administration of the shocking dose of an antigen is most effective in causing an anaphylactic reaction?

a) Intradermal

b) Intravenous

c) Subcutaneous

d) Oral

Question 2:

Which type of reaction is the Schultz-Dale phenomenon?

a) Type II hypersensitivity reaction

b) Type I hypersensitivity reaction

c) Type III hypersensitivity reaction

d) Type IV hypersensitivity reaction

Question 3:

The presence of which of the following antibodies is demonstrated by the Prausnitz-Kustner reaction?

a) IgM

b) IgG

c) IgE

d) IgA

Question 4:

A child, upon ingestion of shellfish, developed a tingling sensation in the mouth, swelling of the face, lips, and tongue, and also complained of abdominal pain and dizziness. What type of reaction is this?

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

a) IgE mediated reaction

b) T cell mediated cytotoxic reaction

c) IgG mediated reaction

d) IgA mediated hypersensitivity reaction

Question 5:

Two days after penicillin administration, a patient developed antibodies that resulted in hemolysis. Which type of hypersensitivity reaction is this?

a) Type I

b) Type II

c) Type III

d) Type IV

Question 6:

A skin transplant was done from a sister to her brother. After a few years, a brother-to-sister skin transplant was done, but rejection occurred. What is this phenomenon known as?

a) Schwartzman reaction

b) Theobald Smith phenomena

c) Eichwald Silmser effect

d) Schultz Dale phenomena

Question 7:

A 9-year-old girl is brought with complaints of sensitivity to light and profuse lacrimation. On inspection, the given finding is seen on her upper tarsal conjunctiva. What type of hypersensitivity reaction is this?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Type I

b) Type II

c) Type II and III

d) Type IV

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 8:

Which type of hypersensitivity reaction is involved in the hyperacute rejection of renal transplants?

a) Type IV

b) Type III

c) Type II

d) Type I

Question 9:

Which of the following statements is not true about anaphylaxis?

a) It is a type I hypersensitivity reaction

b) Large amount of histamine released

c) Primary mediators like IL 4, IL 5 and GM-CSF are released

d) Mediated through allergen specific IgE

Sold by @itachibot

Question 10:

While studying the patient data on hypersensitivity reactions over the last 2 months, you are tasked with classifying the cases by type. Who among the following patients falls under type II hypersensitivity?

a) A patient with berylliosis

b) Eczematous reaction on the hand of an HCW wearing powdered gloves

c) A patient with heat intolerance, sinus tachycardia, and proptosis

d) Glomerulonephritis in a patient with systemic lupus erythematosus

Question 11:

A 22-year-old hiker presents with an itchy rash on her wrist. She gives a history of getting brushed with poison ivy on a trek the previous day. What cells are involved in this reaction?

a) T cells

b) B cells

c) Macrophages

d) Langerhans cells

Question 12:

A 45-year-old hypertensive man complains of fever, migraine, loss of appetite, and myalgia. On examination, splinter hemorrhages and palpable purpura are noted. On CT angiography,

@itachibot

Sold by @itachibot

aneurysm and stenosis of medium-sized vessels are observed. What is the type of hypersensitivity reaction associated with this disease?

a) Type I hypersensitivity

b) Type II hypersensitivity

c) Type III hypersensitivity

d) Type IV hypersensitivity

Question 13:

Which type of hypersensitivity reaction is implicated in the autoimmune disease shown below?

a) Type I

b) Type II

c) Type III

d) Type IV

Question 14:

A 24-year-old patient comes with recurrent episodes of sneezing, rhinorrhea, and watering of the eyes. She says she has these symptoms throughout the year. Which of the following immunoglobulins will be elevated in her case?

a) IgA

Sold by @itachibot

b) IgM

c) IgE

d) IgG

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	a
5	b
6	c
7	a
8	c
9	c
10	c
11	a
12	c
13	b
14	c

Detailed Explanations

Solution to Question 1:

The intravenous route of administration is the most effective route of administration for shocking dose in causing anaphylaxis.

Anaphylaxis is a type I hypersensitivity syndrome. There are 2 clinical syndromes which are characterised as:

• Anaphylaxis - acute, potentially fatal systemic form

• Atopy - chronic, recurrent, non-fatal, localized form

Priming dose: For induction of hypersensitivity reactions, the host should have contact with the antigen, leading to priming of the appropriate B or T lymphocytes.

Shocking dose: Subsequent contact with allergen causing manifestations of hypersensitivity.

Sold by @itachibot

Effectiveness: intravenous > intraperitoneally~subcutaneously > intradermally (least)



Solution to Question 2:

The Schultz–Dale phenomenon is a type I hypersensitivity reaction.

It describes in-vitro contraction of smooth muscles that have been exposed to antigens, demonstrating immediate hypersensitivity reaction in some human and animal tissues.

Other type I hypersensitivity reactions are the Prausnitz-Kustner reaction and the Theobald Smith phenomenon.

Solution to Question 3:

The presence and activity of IgE antibodies are demonstrated by the Prausnitz–Küstner (P–K) reaction.

In this method, the serum is injected from an allergic person into the skin of a nonallergic individual. Then a suspected antigen is later injected into the same site. A positive P-K reaction is a wheal and flare reaction seen in the non-allergic individual. This indicates that the suspected antigen is causing hypersensitivity. Because of the danger of transmitting viral hepatitis and AIDS, this test is no longer used.

Solution to Question 4:

This child who developed tingling in the mouth, and swelling of the lips and face after eating shelf life is an example of an IgE mediated type of hypersensitivity reaction.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Anaphylaxis is a severe, potentially life-threatening allergic reaction that occurs rapidly and affects the whole body. It is characterized by a sudden release of histamines and other chemicals from the immune system in response to an allergen, such as certain foods, medications, insect stings, or latex. The onset of symptoms can occur within minutes to a few hours after exposure to the allergen.The primary mediators are histamine and serotonin. The secondary mediators are prostaglandins and leukotrienes.

Clinically, anaphylaxis can present with a combination of symptoms that typically include:

• Respiratory Symptoms: Patients often experience wheezing and respiratory distress due to bronchoconstriction. Other signs may include shortness of breath, coughing, and a feeling of tightness in the chest.

• Skin Symptoms: Urticaria (hives) and angioedema (swelling of the skin and mucous membranes) are common. These skin reactions usually develop rapidly, often within minutes to two hours of exposure to the allergen.

• Cardiovascular Symptoms: Hypotension (low blood pressure) is a key indicator of anaphylaxis. Patients may also experience tachycardia (increased heart rate) and, in severe cases, shock.

• Gastrointestinal Symptoms: Abdominal pain, vomiting, diarrhea, and cramps can occur, particularly if the anaphylaxis is triggered by food allergens.

• Neurological Symptoms: Dizziness, lightheadedness, or loss of consciousness can occur due to the sudden drop in blood pressure and reduced oxygen supply to the brain.



Solution to Question 5:

Development of antibodies resulting in hemolysis after penicillin administration is an example of a type II reaction. This is a drug-induced type II hypersensitivity reaction.

Penicillin in high doses can result in immune-mediated hemolysis via the hapten mechanism. In this, the drug attaches to surface of RBC membrane. Antibodies are produced that are targeted against this combination of penicillin and RBCs.

Sold by @itachibot

Complement is activated by the attached antibody leading to the removal of red blood cells by the spleen.



Solution to Question 6:

The above scenario describes an Eichwald-Silmser effect, which is a unilateral sex-linked histo-incompatibility.

While transplants between same-sex siblings are successful, grafts from female to male are also successful, but an exception is seen when the donor is a male and the recipient is a female. Such grafts are rejected as the grafted male tissue (XY) will have antigens determined by the Y chromosome which will be absent in the female (XX) recipient.

Solution to Question 7:

Spring catarrh or vernal keratoconjunctivitis is a type I hypersensitivity reaction.

It is a bilateral, seasonally recurrent, chronic conjunctivitis due to type I hypersensitivity allergic reaction to an external antigen. The antigen reacts with IgE, this stimulates mast cell degranulation leading to the release of histamine and other mediators of the allergic response.

Spring catarrh is one of the several forms of allergic conjunctivitis.

Solution to Question 8:

Hyperacute graft rejection is an example of type II hypersensitivity reaction.

It is mediated by preformed antibodies. It is a rare event that occurs in kidney transplantation.

Sold by @itachibot
If you purchased this from
you may have been scammed.
else,

Sold by @itachibot

Graft rejection	Time taken for rejection	Immune mechanisms involved	Type of hypersensitivity
Hyperacute	Minutes to hours	Preformed antibodies	II
Acute	Weeks to months	Cytotoxic T-cell mediated	IV
Chronic	Months to years	Chronic DTH mediated and Ab mediated	IV

Solution to Question 9:

IL4, IL5, GM-CSF are secondary mediators and not primary mediators.

Anaphylaxis is a type I reaction and is mediated through IgE.

Primary mediators are preformed contents of mast cells and basophil granules such as histamine, serotonin. They cause vasodilation and bronchoconstriction. Secondary mediators are newly formed upon stimulation by mast cells, basophils, and other leucocytes (such as leukotrienes IL3, IL4, IL5, IL6; GM-CSF, and prostaglandins) which lead to inflammatory cell infiltrates.



Solution to Question 10:

Heat intolerance, sinus tachycardia, and proptosis of the eye suggest a diagnosis of Grave's disease. It is a type II hypersensitivity reaction that results in the binding of anti-thyroid-stimulating hormone (anti-TSH) antibodies to TSH receptors and stimulates the thyroid cells to produce thyroid hormones.

Type II hypersensitivity reactions involve combining the of IgG antibodies with the antigenic determinants on the surface of cells leading to cytotoxic effects.

Sold by @itachibot



Other options:

Option A: Berylliosis is a type IV hypersensitivity reaction.

Option B: Eczematous reaction on the hand of an HCW wearing powdered gloves (contact dermatitis) is a type IV hypersensitivity reaction.

Option D: Glomerulonephritis in a patient with SLE (systemic lupus erythematosus) is a type III hypersensitivity reaction.

Solution to Question 11:

The history and image suggest a diagnosis of contact dermatitis, a type IV hypersensitivity reaction mediated by T cells. These are delayed hypersensitivity reactions and are induced by sensitized T cells, which on contact with the specific antigen releases cytokines that produce biological effects on leucocytes, macrophages, and tissue cells.

Two types of delayed hypersensitivity are:

• Tuberculin type

• Contact dermatitis type

Solution to Question 12:

The most likely diagnosis is polyarteritis nodosa. It is an example of a type III hypersensitivity reaction.

In type III hypersensitivity, immune complexes may deposit in tissues or blood vessels, and elicit an acute inflammatory reaction by activating the complement system. The inflammatory reaction causes tissue injury.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

The pathogenesis of systemic immune complex disease can be divided into three phases:

• Formation of immune complexes

• Deposition of immune complexes

• Inflammation and tissue injury.

Examples of type III hypersensitivity reactions:

• Post-streptococcal glomerulonephritis

• SLE

• Polyarteritis nodosa

• Reactive arthritis

• Serum sickness

• Arthus reaction

• Farmer's lung

• Hypersensitivity pneumonitis

• Type II lepra reaction

• Henoch Schonlein purpura

d by @itachibot
s from someone else,
scammed.



Solution to Question 13:

The disease shown in the image is myasthenia gravis. It is an example of type II hypersensitivity.

Type II hypersensitivity reactions are mediated by antibodies directed towards self-antigens- cell surface or extracellular matrix.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 14:

The clinical vignette describes a case of chronic allergic rhinits. IgE is elevated in these cases.

Perennial allergic rhinitis, a type of chronic rhinitis is a year-round problem and is often caused by indoor allergens such as dust and animal dander in addition to pollens that may exist at the time. Symptoms tend to occur regardless of the time of the year.

Sold by @itachibot

Immune Disorders

Question 1:

A 6-month-old child has a history of recurrent skin and respiratory tract infections. Investigations are carried out and a disorder of phagocytosis is suspected. Which of the following will not be a differential diagnosis in this case?

a) DiGeorge syndrome

b) Leukocyte adhesion deficiency

c) Chediak-Higashi syndrome

d) Chronic granulomatous disease

Question 2:

A 1-year-old boy presents with meningococcal meningitis. He has a history of recurrent bacterial chest and ear infections. Family history reveals that he has a male cousin (maternal side) who died in infancy from infection. The physician suspects X-linked agammaglobulinemia. Which of the following is false regarding his condition?

a) It is caused by a mutation of the BTK gene

b) Both humoral and cell mediated immunity are affected

c) All classes of immunoglobulin levels are reduced

d) Lymph node biopsy shows depletion of follicles

Question 3:

An immunologist orders a nitroblue tetrazolium test on a child suspected to have a primary immunodeficiency disease. What condition is she screening the child for?

a) Tuftsin deficiency

b) Wiskott-Aldrich syndrome

c) Chronic granulomatous disease

d) Chediak-Higashi syndrome

Question 4:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

A 1-year-old boy is brought with a history of recurrent pneumonia. He also has eczema and a history of prolonged bleeding after circumcision. Peripheral smear shows thrombocytopenia with small platelets. Which of the following is most likely to be deficient?

a) IgM

b) IgG

c) IgA

d) IgE

Question 5:

An 8-year-old boy was brought to the pediatric OPD with multiple cold abscess lesions as shown below on his face, chest, and back. The child has a history of recurrent■ respiratory infections. On examination, he has atopic excoriations over the skin folds. What is the most probable diagnosis?

a) Hyper IgM syndrome

b) Wiscott Aldrich syndrome

c) Hyper IgE syndrome

d) X-linked agammaglobulinemia

Question 6:

Which of the following is a combined immunodeficiency disorder associated with both B-cell and T-cell defects?

Sold by @itachibot

a) Louis–Bar syndrome

b) DiGeorge syndrome

c) Chediak-Higashi syndrome

d) Tuftsin deficiency

Question 7:

A 2-year-old boy presents with reports showing fungal pneumonia, the causative organism is shown below. Parents give a history of recurrent bacterial infections. Laboratory analysis show markedly elevated IgM and low levels of IgG. Which of the following is the most likely cause of his illness?



a) Defective cytokine signaling in T-cell precursors

b) Autosomal-dominant mutation in the STAT3 gene

c) Deficiency in CD40 ligand or CD40

d) Defect in the NADPH oxidative pathway

Question 8:

A 5-year-old boy is brought with complaints as shown below. The parents report that these complaints have been present for the last 2 years. Which of the following is found to be the most effective treatment for this disease?

Sold by @itachibot

a) 1 and 3

b) 1 and 2

c) Only 2

d) 2 and 3

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 9:

An infant with a history of delayed separation of the umbilical cord now presents with recurrent pneumonia. The total leukocyte count is 20,000/microlitre. Which of the following genetic defects is most likely present?

a) Mutation of Bruton tyrosine kinase gene

b) Low levels of NADPH oxidase and negative DHR test

c) Reduced phagocyte surface expression of sialyl-Lewis x

d) Excessive IgM with reduced IgG and IgA

Question 10:

Which of the following statements is true about severe combined immunodeficiency disorders?

a) Reticular dysgenesis of de Vaal is a mild form of combined immunodeficiency

b) They are inherited as autosomal dominant disorders

c) Adenosine deaminase deficiency is associated with chondrocyte abnormalities

Sold by @itachibot

d) The defect in Swiss type agammaglobulinemia is at the level of multipotent hemopoeitic stem cell

Question 11:

Which of the following is an immunologically privileged site?

a) Seminiferous tubule

b) Area postrema

c) Optic nerve

d) Loop of Henle

If you bought this from someone else, @itachibot

Answer Key

Question No.	Correct Option
1	a
2	b
3	c
4	a
5	c
6	a
7	c
8	d
9	c
10	c
11	a

Detailed Explanations

Solution to Question 1:

DiGeorge syndrome is not a disorder of phagocytosis.

It is due to a developmental defect involving the endodermal derivatives of the 3rd and 4th pharyngeal pouches. Cell-mediated immunity is defective due to decreased T-cells. Affected individuals also have depressed graft rejection, hypocalcemia, and cardiac defects.

Sold by @itachibot

Disorders of phagocytosis:

• Leukocyte adhesion deficiency

• Chediak–Higashi syndrome

• Chronic granulomatous disease

• Tuftsin deficiency

• Cyclic neutropenia

Other options:

Option B: Leukocyte adhesion deficiency leads to defective neutrophil migration to sites of infection, peripheral neutrophilia, and recurrent infections without pus.

Option C: Chediak–Higashi syndrome is an autosomal-recessive disorder due to defects in lysosomal transport protein (LYST), leading to defective microtubule polymerization and defective phagolysosome formation. There is neutropenia and recurrent infections.

Option D: Chronic granulomatous disease is due to a defect in NADPH oxidase resulting in the absence of respiratory burst. It presents with recurrent infections with catalase-positive microbes.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 2:

X-linked or Bruton's agammaglobulinemia is primarily a disease of humoral immunity. Cell-mediated immunity is not affected.

It is an inherited immunodeficiency disease caused by mutations in the gene coding for Bruton tyrosine kinase (BTK). B cells do not differentiate and mature. They are arrested in the pre-B cell stage.

There are extremely low IgG levels (below 100 mg/dL) and an absence of other immunoglobulin classes. The tonsils and lymph nodes are usually atrophic. Lymph node biopsy reveals depletion of cells of B cell-dependent areas (follicles). Babies born with this disorder have virtually no peripheral B cells (<1%) and suffer from recurrent bacterial infections.

Solution to Question 3:

The Nitroblue Tetrazolium Test (NBT) is used as a screening test for chronic granulomatous disease (CGD) by assessing the reactive oxygen species (ROS) production in phagocytes.

This principle relies on the color change of nitroblue tetrazolium, which transforms from colorless/yellow to deep blue when it encounters ROS. A higher nitroblue test score indicates that the cell is more efficient in producing ROS.

In chronic granulomatous disease, there is an NADPH oxidase defect. Hence, the phagocyte cannot produce reactive oxygen species or radicals required for bacterial killing, which results in recurrent infections. Here, the leukocytes fail to reduce nitroblue tetrazolium during phagocytosis. Hence, a negative test (NBT remains colourless to yellow) is seen in CGD.

A positive test result means neutrophils turn blue, as shown below:

Sold by @itachibot



A negative test means nitroblue tetrazolium remains colourless/yellow, as shown below:



Solution to Question 4:

The history here is highly suggestive of Wiskott-Aldrich syndrome (WAS), a condition characterized by decreased IgM levels.

Wiskott-Aldrich syndrome (WAS) is an X-linked recessive immunodeficiency disorder. It is caused by mutations in the Wiskott-Aldrich syndrome protein (WASp) that primarily affects the T lymphocytes. Normally, WASp regulates the actin cytoskeleton and thus helps in cell adhesion, migration, and the formation of synapses between antigen-presenting and target cells. In WAS, there is a progressive loss of T lymphocytes in the peripheral blood resulting in defects in cellular immunity.

Sold by @itachibot

The clinical features of WAS are a triad of recurrent bacterial infections, eczema, and bleeding due to thrombocytopenia (with small platelets). Diagnosis is based on intracellular immunofluorescence analysis of WAS protein (WASp) expression in blood cells. IgM levels in the serum are low, but levels of IgG are usually normal. IgA and IgE levels are normal or elevated. Reduced-sized platelet is a typical feature of WAS.

The image below shows an 18-month-old child with WAS, showing multiple petechiae and hematoma.

If you purchased this, your may have be... else, ...d.

Solution to Question 5:

A child with multiple abscesses (as shown in the image above) with a history of atopy and recurrent respiratory infections suggests immunodeficiency, and the most probable diagnosis is Hyper IgE syndrome.

The hyper-IgE syndromes are rare primary immunodeficiency syndromes characterized by recurrent severe staphylococcal abscesses of the skin, lungs, and other sites, eczema, eosinophilia, and marked elevation of serum IgE levels. They occur in 2 forms:

• Autosomal dominant - involves mutations in the gene encoding STAT-3 (also known as Job syndrome)

• Autosomal recessive - involves mutations in DOCK8.

In Job syndrome, mutation of the STAT3 gene leads to IgE overproduction without modulation. Loss of modulation is due to the absence of anti-inflammatory cytokines IL-6, IL-10, and IFN-gamma. Lack of IL-6 leads to deficiency of CD4 + Th17 cells, making them susceptible to various infections.

It usually presents as a triad of atopic dermatitis, recurrent skin staphylococcal infections, and recurrent pulmonary infections that occur early in childhood. The eczematoid rash affects the scalp and face, gets superinfected with Staphylococcus aureus and forms cold abscesses. Candida infections are common. Recurrent staphylococcal pneumonia with pneumatocoeles is also

Sold by @itachibot

commonly seen. Other non-immunological features are:

• Dysmorphic facial features that appear in early adolescence and are absent in the autosomal recessive form

• Craniosynostosis

• Arnold-Chiari type 1

• Hyperextensible joints

• Scoliosis

• Pathological fractures

• Retention of primary teeth

• Vascular complications

• Ocular complications

Diagnosis is based on measurements of serum IgE levels >2000U/mL where IgA, IgG and IgM concentrations are generally normal, in adjunct with the clinical features seen in Job syndrome. It is associated with peripheral blood eosinophilia.

The treatment primarily focuses on preventing infections through the use of antimicrobials, including antibiotics and antifungals, along with immunoglobulin replacement therapy.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

Louis–Bar syndrome or ataxia-telangiectasia is a combined immunodeficiency disorder.

Ataxia-telangiectasia is a syndrome complex that comprises neurologic, immunologic, endocrinologic, and cutaneous disorders. The gene involved in this condition is the mutation in the ATM gene mapped to the long arm of chromosome 11(11q22-23). The mode of inheritance is autosomal recessive. There is defective DNA repair and increased sensitivity to ionizing radiation.

Clinical features:

• Progressive cerebellar ataxia

• Oculocutaneous telangiectasias

• Chronic sinopulmonary disease

• High incidence of malignancy - adenocarcinomas and lymphoreticular tumors (lymphoma, leukemia, and Hodgkin's disease)

• Humoral and cellular immunodeficiency

• Ovarian dysgenesis and chromosomal abnormalities

• Hypoplastic thymus

• Ataxia and chorioathetoid movements start in infancy.

The onset of telangiectasia is around 3 to 6 years and involves the face, conjunctiva and other parts of the body. By 12 years of age, affected children are usually confined to wheelchairs due to progressive ataxia.

Sold by @itachibot

Death occurs due to sinopulmonary infection in early life or due to malignancy in the second or third decade. They lack IgA and IgE. Cell-mediated immunity is also defective.

The image below shows ocular telangiectasia in a patient with A-T.

Other options:

Option B: DiGeorge syndrome is a cellular immunodeficiency disorder associated with T-cell deficiency.

Options C and D: Chediak-Higashi syndrome and tuftsin deficiency are disorders of phagocytosis.

Sold by @itachibot
If you purchased this from anywhere else, you may have been scammed.

Solution to Question 7:

The given history of recurrent bacterial infections and fungal pneumonia caused by Pneumocystis jirovecii (as shown in the image above), along with elevated IgM and low IgG levels, are suggestive of hyper-IgM syndrome, an inherited deficiency in the CD40 ligand or CD40.

Hyper-IgM syndrome is a primary immunodeficiency disorder that is inherited as:

• X-linked form (most common): CD40 ligand (CD40L) defect on T-helper cells, preventing B-cell activation, class switching and memory cells formation.

• Autosomal recessive form: Affect CD40 on B-cells or enzymes like activation-induced cytidine deaminase (AID), a DNA-editing enzyme expressed in B cells.

The B-cell response to T-independent antigen is unaffected. CD40–CD40L interactions are also required for IL-12 secretion. So, defects in this pathway result in increased susceptibility to intracellular pathogens.

In normal immune response, B-cells expressing CD40 on their surface interact with CD40L on T-cells to undergo class switching from IgM to various other classes (IgG, IgA, IgE) depending on the type of antigen. Thus, failure of this class switching leads to an increase in IgM with extremely low levels of IgG and no IgA or IgE.

Sold by @itachibot

Clinical features include recurrent pyogenic infections caused by low levels of opsonizing IgG antibodies. Patients are highly susceptible to pneumonia caused by Pneumocystis jirovecii.

The illustration below depicts the defects in components of T-cell & B-cell interactions that give rise to primary immunodeficiency.



Other options:

Option A: Defective cytokine signalling in T-cell precursors is seen in X-linked Severe Combined Immunodeficiency (SCID). It is caused by a mutation in the common γ-chain (γc) subunit of cytokine receptors. This defect disrupts Interleukins (IL-2, IL-4, IL-7, IL-9, IL-15 and IL-21) signalling, leading to low T-cell and NK-cell numbers. Although B-cell numbers are normal, these are non-functional due to the absence of T-cell help, resulting in impaired antibody synthesis.

Option B: Mutation in the STAT3 gene is seen in an autosomal dominant form of Hyper IgE syndrome (also known as Job syndrome), characterized by recurrent staphylococcal skin abscesses, recurrent pulmonary infections often with pneumatoceles and atopic dermatitis. Here, serum IgE levels are markedly elevated (due to IgE overproduction without modulation), and IgA, IgG and IgM concentrations are generally normal.

Option D: A defect in NADPH oxidative pathway is seen in chronic granulomatous disease, leading to failure of the oxidative burst in phagocytes, predisposing individuals to catalase-positive microbial infections.

Solution to Question 8:

This 5-year-old boy presenting with a 2-year history of thick, white-coated tongue and dystrophic nail changes, as shown in the above image, suggests chronic mucocutaneous candidiasis, for which transfer factor therapy and amphotericin B have been reported to be the effective treatments.

Chronic mucocutaneous candidiasis (CMC) is characterized by deficient cell-mediated immunity to candida, usually due to defects in IL-17. Mutations affecting the IL-17 signalling pathway lead to

Sold by @itachibot

impaired IL-17 production and defective TH17 function, compromising mucosal and skin defence mechanisms and increasing susceptibility to fungal infections.

Patients develop persistent fungal infections of the skin, nails, oral cavity, and genitals, primarily caused by Candida species. Despite chronic mucocutaneous involvement, these patients typically do not develop invasive candidiasis. A key feature of CMC is the absence of delayed hypersensitivity to candida antigens. However, circulating antibodies to them are present in high titres. They do not show increased susceptibility to other infections but may develop endocrinopathies, autoimmune disorders, and vascular abnormalities.

Transfer factor therapy and Amphotericin B have been reported to be effective for the treatment of chronic mucocutaneous candidiasis.

If you purchased this from someone else, you may have been scammed
Sold by @itachibot

Solution to Question 9:

This history of delayed separation of the umbilical cord with recurrent pneumonia and peripheral leucocytosis is highly suggestive of leucocyte adhesion defect (LAD). In this defect, there is reduced phagocyte surface expression of sialyl-Lewis X.

LAD is an autosomal recessive disorder affecting leukocyte function. Neutrophils have reduced adhesion, mobility, and phagocytic ability. There are three main types of LAD:

• LAD 1- Defective β2 integrin common chain (CD18), leading to abnormal neutrophil chemotaxis and defective tight adhesion.

• LAD 2- Reduced phagocyte surface expression of sialyl-Lewis X (fucosylated protein), leading to abnormal neutrophil rolling and tethering.

• LAD 3- Defect in regulatory protein (kindlin 3, also known as Fermt 3) involved in activating the ligand affinity for β2 integrins, leading to abnormal neutrophil chemotaxis and adhesion (milder than type 1).

LAD is characterised by recurrent bacterial and fungal infections, reduced inflammatory responses despite elevated blood neutrophilia, and delayed wound healing and delayed umbilical cord loss. in patients with LAD type 3, there may be bleeding as β2 integrin on platelets is non-functional.

Typical signs of inflammation may be absent, pus does not form with few neutrophils on biopsy of infected tissues along with massive hyperleukocytosis (>30,000/μL) in the blood. Early allogeneic hematopoietic stem cell transplantation is used in the treatment of severe LAD.

Solution to Question 10:

Adenosine deaminase deficiency is associated with chondrocyte abnormalities.

Severe combined immunodeficiency disorders are associated with severe deficiency of both humoral and CMI. They are inherited in both X-linked and autosomal recessive modes. There are different forms of this disorder:

Sold by @itachibot

• Reticular dysgenesis is the most serious form of combined immunodeficiency. Here the defect is at the level of the multipotent hemopoietic stem cell.

• Adenosine deaminase (ADA) deficiency is associated with enzyme deficiency. The condition is associated with chondrocyte abnormalities which can be discerned radiologically.

• Swiss-type agammaglobulinemia is associated with lymphocytopenia and severe defect in CMI. The basic defect is at the level of the lymphoid stem cell.

Solution to Question 11:

The seminiferous tubules are an immunologically privileged site.

Immunologically privileged sites can tolerate the introduction of antigens without eliciting an immune response. This allows a higher likelihood of graft survival, even without perfect HLA matching. As a result, an allograft can be placed in these sites without endangering a rejection reaction.

The unique immune tolerance in these regions is primarily due to the lack of effective lymphatic drainage, which prevents antigen presentation to the immune system, and the presence of blood-tissue barriers that restrict immune cell infiltration. Low vascularity contributes to immune privilege by minimising antigen exposure and immune cell recruitment.

Examples of immune-privileged sites are the anterior chamber of the eye, cornea, testes, and brain.

Note: Although part of the brain, the blood-brain barrier is absent at the area postrema, therefore it is not an immune-privileged site.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Streptococci and Enterococci

Question 1:

What is the Lancefield grouping of β-hemolytic-streptococci based on?

a) T antigen

b) M protein

c) Group-specific cell wall polypeptide

d) Group-specific C carbohydrate antigen

Question 2:

An elderly diabetic lady presented with swelling and pain of her right leg following a skin biopsy. She was febrile at the time of examination and hemorrhagic bullae with necrosis of the surrounding skin were noted on her leg. Gram staining of the pus collected from the bullae is shown below. Which of the following is the most likely organism causing this condition?



a) Clostridium histolyticum

b) Streptococcus pyogenes

c) Staphylococcus aureus

d) Clostridium perfringens

Sold by @itachibot
If you purchased this from someone else, it has been scammed.

Sold by @itachibot

Question 3:

A 7-year-old child was brought in by her mother to the OPD with complaints of fever, sore throat, and a scarlatiniform rash. A throat swab was taken and sent to the lab for culture on blood agar. Which of the following factors is responsible for the zones of hemolysis seen in the culture below?

a) Lipoteichoic acid

b) Streptolysin S

c) Streptokinase

d) Hyaluronidase

If you purchased this, you may have been scammed.

Question 4:

Which of the following organisms produces superantigen?

a) Staphylococcus epidermidis

b) Streptococcus pyogenes

c) Clostridium perfringens

d) Vibrio cholera

Question 5:

Which of the following types of streptococcal pyrogenic exotoxins is bacteriophage mediated?

Sold by @itachibot

a) type A & type B

b) type C & type D

c) type B & type D

d) type A & type C

Question 6:

Which of the following is a streptococcal toxin responsible for connective tissue breakdown?

a) Hyaluronidase

b) Streptococcal Pyrogenic Exotoxin (SPE)

c) Streptolysin O

d) Serum opacity factor

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which streptococcal antigen cross reacts with myocardium?

a) Peptidoglycan

b) Cell wall M protein

c) Hyaluronic acid

d) Cell wall C carbohydrate

Question 8:

A 12-year-old child presented with a sore throat for 3 days. Which of the following medium is used to culture the throat swab?

a) Blood Agar

b) Lowenstein-Jensen medium

c) Stewart medium

d) Chocolate Agar

Question 9:

Which of the following organisms show beta-hemolysis on blood agar, resistance to bacitracin, and a positive CAMP test?

Sold by @itachibot

a) Streptococcus pyogenes

b) Streptococcus agalactiae

c) Pneumococcus

d) Enterococcus

Question 10:

A child presents to the OPD with an infective skin lesion over his leg. On microscopic examination of the pus from the lesion, gram-positive cocci in chains are visualized and pus culture shows β-haemolytic colonies on blood agar. Which of the following tests can be performed to identify the causative organism?

a) Bile solubility

b) Optochin sensitivity

c) Bacitracin sensitivity

d) Catalase positive

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 11:

Which of the following bacteria is associated with the causation of colorectal carcinoma, rectal polyp, endocarditis and septicaemia and is pyrrolidinyl arylamidase test negative?

a) Streptococcus equisimilis

b) Streptococcus gallolyticus

c) Enterococcus

d) Streptococcus pyogenes

Question 12:

Which of the following organisms can survive up to 60°C, is PYR positive, and can grow on media with 6.5% NaCl?

a) Streptococcus pyogenes

b) Streptococcus agalactiae

c) Enterococcus feacalis

d) Streptococcus pneumoniae

Sold by @itachibot

Question 13:

A patient admitted to an ICU spiked a fever and his blood culture was positive for gram-positive cocci in chains, which are catalase-negative. He was being given ceftazidime and amikacin via a central venous line for the past week. Following this, you started him on vancomycin but his blood culture remained positive for the same organism even after 2 weeks of therapy. Which of the following is the most likely organism causing the infection?

a) Streptococcus pyogenes

b) Viridans streptococci

c) Enterococcus faecium

d) Coagulase-negative Staphylococcus

Question 14:

Oral examination of a young patient shows the following finding. Which bacteria belonging to viridans group is most commonly implicated in this condition?

a) Streptococcus mutans

b) Streptococcus mitis

c) Streptococcus sanguis

d) Streptococcus salivarius

Question 15:

If yo... Sold by @itachibot ...s from someone else, ...scammed.

Sold by @itachibot

A 49-year-old man presented with high-grade fever, cough and shortness of breath. On examination, he was febrile, tachypneic and tachycardic. A chest X-ray was performed which is shown in the image. What is the single most likely causative organism?

a) Staphylococcus aureus

b) Streptococcus pyogenes

c) Streptococcus pneumoniae

d) Haemophilus influenzae

Sold by
If you purchased this fro
you may have been scam
else,
d.

Question 16:

A 70-year-old male presented with fever, chills and cough associated with chest pain. His sputum sample was sent for culture on blood agar which showed the presence of ° -hemolytic colonies the following day. Which of the following best describes the organism likely implicated here?

a) Gram-positive cocci in short chains, catalase negative and bile resistant

b) Gram-positive cocci in pairs, catalase negative and bile soluble

c) Gram-positive cocci in clusters, catalase positive and coagulase positive

d) Gram-positive cocci in clusters, catalase positive and coagulase negative

Question 17:

Quellung reaction is used to demonstrate_____.

Sold by @itachibot

a) Capsule

b) Cell wall

c) Cell membrane

d) Pilli

Question 18:

Which of the following factors is responsible for the draughtsman appearance of pneumococcal colonies?

a) Pneumolysin

b) Autolysin

c) Capsular polysaccharide

d) Pneumotoxin

Question 19:

In a splenectomized patient, there is an increased risk of infection by all capsulated organisms. Which of the following is least likely to cause an infection in such patients?

a) Streptococcus pneumoniae

b) Klebsiella pneumoniae

c) Haemophilus influenzae

d) Staphylococcus aureus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 20:

Which of the following statements is true about the pneumococcal vaccine?

a) Pneumococcal polyvalent vaccine PPSV23 is indicated in splenectomized patients.

b) Pneumococcal polyvalent vaccine PCV13 is contraindicated in children less than 2 years.

c) Pneumococcal polyvalent vaccine PPV 23 gives protection for 10 years.

d) Pneumococcal polyvalent vaccine is contraindicated if age > 65 years.

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	d
2	b
3	b
4	b
5	d
6	a
7	b
8	a
9	b
10	c
11	b
12	c
13	c
14	a
15	c
16	b
17	a
18	b
19	d
20	a

Detailed Explanations

Solution to Question 1:

Lancefield grouping of β-hemolytic-streptococci is based on group-specific carbohydrate C antigen present on the cell wall. He classified them into 20 serological types from Group-A to Group V except for I and J.

Group A β-hemolytic streptococci (Streptococcus pyogenes) are further classified by Griffith typing based on protein antigens (M, T, R) present on the cell surface.

Sold by @itachibot



Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 2:

The given clinical vignette is strongly suggestive of necrotizing fasciitis which is most commonly caused by Streptococcus pyogenes (Group A β-hemolytic streptococcus).

Necrotizing fasciitis is a rapidly progressive inflammatory infection of the fascia, with secondary necrosis of the subcutaneous tissues. It has also been referred to as Meleney ulcer, acute dermal gangrene, hospital gangrene, suppurative fasciitis, and synergistic necrotizing cellulitis. The source of the infection is either the skin, with organisms introduced into the tissue through trauma, or the bowel flora, with organisms released during abdominal surgery.

The onset of symptoms is usually quite acute and is marked by severe pain at the site of involvement, malaise, fever, chills, and a toxic appearance. The affected limb is edematous with the overlying skin turning dusky with mottled erythema and the development of hemorrhagic bullae.

This is a surgical emergency as the condition is rapidly fatal. Treatment of the condition includes:

• Surgical intervention is the major therapeutic modality in cases of necrotizing fasciitis.

• Antibiotics- penicillin (1-4 million U IV q4h) plus clindamycin (600-900 mg IV q8h).

The following image shows surgical debridement being done for a case of necrotising fasciitis.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

The given case scenario o a patient with fever, sore throat, and a scarlatiniform rash is suggestive of scarlet fever caused by Streptococcus pyogenes. Streptolysin S is the factor responsible for the zones of hemolysis seen around streptococcal colonies on blood agar.

Streptococci produces two hemolysins:

• Streptolysin O is called so because it is oxygen-labile. It is also heat-labile and it has specific cardiotoxic and leucotoxic activity.

• Streptolysin S is called so because it is soluble in serum. It is also oxygen-stable hemolysin. It is a protein but is not antigenic.

The following image shows blood agar with colonies of streptococci with hemolytic zones around them:



Sold by @itachibot

Solution to Question 4:

Streptococcus pyogenes produces streptococcal pyrogenic exotoxin (SPE), which is a superantigen.

Superantigens are protein toxins that stimulate large numbers of T cells, irrespective of antigen specificity. They bind to the class II MHC molecule outside the peptide-binding groove and Vβ region of the TCR. Some examples of superantigens are staphylococcal enterotoxins, toxic shock syndrome toxin, and group A streptococcal pyrogenic exotoxin.

The below image shows the binding of superantigen to T-cell receptor.



Solution to Question 5:

Streptococcal pyrogenic exotoxins type A and type C are bacteriophage-mediated toxins.

The streptococcal pyrogenic exotoxin(SPE) is so named because the primary effect of the toxin is the induction of fever. It is also known as erythrogenic toxin because its intra-dermal injection into susceptible individuals produced an erythematous reaction (Dick Test).

There are 3 types of streptococcal pyrogenic exotoxins namely:

• type A - phage mediated.

• type B - chromosome mediated.

• type C - phage mediated.

Solution to Question 6:

Sold by @itachibot

Hyaluronidase is the streptococcal toxin responsible for connective tissue breakdown.

Hyaluronidase or spreading factor is expressed by non-capsulated strains of Streptococcus pyogenes, such as M type 4 and 22. It acts by cleaving hyaluronic acid, which is present in the connective tissue, epithelial, and neural tissues. This results in the spread of infection along the intercellular space.

Solution to Question 7:

Antibodies to the cell wall M antigen present on the Streptococcus cross-reacts with a similar myocardial antigen. This cross-reaction with myocardium is the cause of acute rheumatic fever.

Solution to Question 8:

Blood agar is the preferred culture media as it is used for the culture and isolation of Streptococcus pyogenes which is the most common cause of sore throat.

The selective medium for Streptococcus pyogenes is Crystal violet blood agar and the transport media for this organism is Pikes medium.

Solution to Question 9:

Streptococcus agalactiae is a group B streptococci that shows β-hemolysis on blood agar, is resistant to bacitracin, and gives a positive CAMP test.

The CAMP test is used to identify group B β-hemolyitic-streptococci based on their ability to form CAMP factor. This factor causes synergistic hemolysis with β-hemolysin formed by Staphylococcus aureus, when Streptococcus agalactiae is inoculated perpendicular to a streak of Staphylococcus aureus on blood agar. This produces the characteristic arrow-head zone of hemolysis.

The below image shows a positive CAMP test.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



The following table provides important laboratory characteristics which differentiate Streptococcus pyogenes from Streptococcus agalactiae.

Species	Lancefield Group	Typical Hemolysis	Important Lab Characteristics
S.pyogenes	A	β	Bacitracin sensitive, PYR test positive
S.agalactiae	B	β	Bacitracin resistant, PYR test negative, Hippurate hydrolysis positive, CAMP test positive

Solution to Question 10:

β hemolytic colonies on blood agar with gram-positive cocci in chains seen on microscopy suggest the organism is a β-hemolytic streptococcus. Bacitracin sensitivity is used to differentiate between group A and group B β-hemolytic streptococci.

Group A: Bacitracin-sensitive (S. pyogenes)

Group B: Bacitracin-resistant (S. agalactiae)

Options A & B: Bile solubility and optochin sensitivity are used to differentiate Streptococcus pneumoniae and viridans streptococci. They are both ° hemolytic on blood agar but some species of viridans streptocci are non hemolytic.

Option D: Catalase test is used to differentiate streptococcus and staphylococcus. Staphylococcus appear as clusters of gram positive cocci on microscopy and not chains.

Sold by @itachibot

Solution to Question 11:

Streptococcus gallolyticus is associated with the causation of colorectal CA, rectal polyp, endocarditis, and septicemia and is pyrrolidinyl arylamidase test (PYR test) negative.

Streptococcus gallolyticus, formerly called Streptococcus bovis, is a non-enterococcal Group D streptococci. It is bile insoluble, esculin hydrolysis positive, PYR negative and shows no growth in 6.5% NaCl.

Solution to Question 12:

Among the given options, Enterococcus feacalis can survive up to 60°C, is PYR test positive, and can grow in the presence of 6.5% NaCl.

Enterococci have the following features:

• Formation of tiny deep pink colonies on MacConkey's medium

• Survives at 600C for 30 minutes.

• Growth in the presence of 40 % bile, 6.5% sodium chloride at pH 9.6.

• PYR (Pyrrolidonyl Arylamidase) positive

• Esculin hydrolysis is positive.

• Usually nonhemolytic, but may show ° or β hemolysis.

Option A: Streptococcus pyogenes is a delicate organism, easily destroyed by heat (54°C for 30 min), PYR positive.

Option B: Streptococcus agalactiae is PYR test negative.

Option D: Streptococcus pneumoniae are readily destroyed by heat (52°C for 15 min).

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 13:

The given scenario is suggestive of vancomycin-resistant enterococci (VRE) infection.

Gram-positive cocci in chains that are catalase-negative suggest that the organism belongs to Streptococcaceae family. Positive culture after 2 weeks of vancomycin therapy suggests resistance to vancomycin.

Option A: Streptococcus pyogenes are sensitive to vancomycin and are not commonly implicated in central venous line infections.

Option B: Viridans streptococci are usually mouth commensals and are sensitive to vancomycin.

Option D: Coagulase-negative staphylococci are gram-positive cocci in clusters which are catalase-positive. Hence, they can be ruled out.

Sold by @itachibot

Solution to Question 14:

The above image shows dental caries which are commonly caused by Streptococcus mutans. It belongs to the viridans group/° -hemolytic streptococci.

Streptococcus mutans breaks down the dietary sucrose producing acid and a tough adhesive dextran. The acid damages dentine while the dextrans bind together with food debris, epithelial cells, mucus, and bacteria to form dental plaques, leading to formation of dental caries.

Solution to Question 15:

The given clinical history and the X-ray showing right lower lobe consolidation is strongly suggestive of lobar pneumonia, most commonly caused by Streptococcus pneumoniae (>95 % of cases).

Lobar pneumonia is an acute exudative inflammation of an entire pulmonary lobe. The below image shows the histopathological picture of lobar pneumonia.



Solution to Question 16:

The given clinical scenario and the growth of alpha-hemolytic colonies from the sputum of the patient are strongly suggestive of pneumococcal pneumonia. Pneumococci are gram-positive cocci in pairs, catalase negative, and bile soluble.

The image below shows Gram positive cocci in pairs. bile solubility, and catalase reaction.

Sold by @itachibot



The image below shows positive bile solubility test shown by Streptococcus pneumoniae.



The below image shows catalse test.

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 17:

Quellung reaction is used to demonstrate the capsule of streptococci.

Quellung reaction (Quellung = swelling):

• A suspension of pneumococci is mixed on a slide with a drop of type-specific antiserum and a loop full of methylene blue solution.

• In the presence of homologous antiserum, the capsule becomes apparently swollen, sharply delineated and refractile.

• The Quellung test can be done directly with sputum from acute pneumonia cases.

The below image shows the swollen capsule of pneumococci following the addition of homologous antiserum.



Sold by @itachibot

Solution to Question 18:

The factor responsible for the draughtsman appearance of pneumococcal colonies is autolysin.

When pneumococci are incubated on blood agar for a prolonged duration, the resultant colonies are surrounded with concentric rings and have a central umbonation. This appearance of pneumococcal colonies is known as carrom coin appearance or draughtsman appearance.

Autolysin is an amidase enzyme produced by pneumococci that cleaves peptidoglycan, leading to the autolysis of cells. It is also responsible for the bile solubility of pneumococci.

The below image shows the characteristic draughtsman appearance of pneumococcal colonies.



Solution to Question 19:

In splenectomized patients, there is an increased risk of infection by capsulated organisms such as Streptococcus pneumoniae, Haemophilus influenzae, and Klebsiella pneumoniae. Staphylococcus aureus does not have a capsule and so poses no increased risk to these individuals.

The phagocytosis of capsulated organisms is prevented by their capsule. Thus, the body's main defence against these organisms is the spleen.

Certain components called opsonins attach to these capsulated organisms, and then they are delivered to the spleen, where their uptake is facilitated by receptors present on the surface of splenic phagocytes. After splenectomy, this defence mechanism is lost leaving patients susceptible to infection from these bacteria.

Solution to Question 20:

Sold by @itachibot

Pneumococcal polyvalent vaccine PPSV23 is indicated in splenectomised patients.

There are two kinds of polyvalent pneumococcal vaccines:

• Pneumococcal polysaccharide vaccine - PPSV23

• It is indicated in splenectomized patients, sickle cell disease, immunocompromised patients, and patients above 65 years of age.

• CDC currently recommends the administration of one additional dose of PPSV23 at least 5 years after the last dose.

• Children younger than 2 years old should not get this vaccine since they fail to mount an adequate response.

• Pneumococcal conjugate vaccine-

• It includes PCV10 and PCV13.

• It is indicated in children less than 2 years.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Staphylococci

Question 1:

A child injures his knee while playing. A few days later, the wound on his knee shows signs of inflammation. The wound is cultured and its Gram-stained smear is given below. Further testing shows that the cultured organism is coagulase-positive. Identify the organism.



a) Streptococcus pyogenes

b) Staphylococcus aureus

c) Staphylococcus epidermidis

d) Staphylococcus saprophyticus

Question 2:

The specimen obtained by performing incision and drainage of a breast abscess was sent for culture on nutrient agar. An image of the culture plate is given below. Microscopy of the isolated organism revealed gram-positive cocci arranged in clusters. Which of the following is a false statement regarding this organism?

Sold by @itachibot



a) Pigment production occurs optimally at 35°C.

b) Golden yellow pigmentation is due to β-carotene

c) Pigment production is enhanced by adding milk to culture media

d) Confluent growth presents a characteristic 'oil-paint appearance'

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

Which factor forms the basis of co-agglutination reaction shown by Staphylococcus aureus?

a) Protein A

b) Clumping factor

c) Beta hemolysin

d) Coagulase

Question 4:

Which of the following toxins exhibits hot-cold phenomenon?

a) ° -hemolysin

b) β-hemolysin

c) γ-hemolysin

d) δ-hemolysin

Sold by @itachibot

Question 5:

Which of the following pairs of toxins produced by Staphylococcus aureus are called synergohymenotropic toxins?

a) ° and β toxins

b) β and Panton-Valentine-toxins

c) γ and Panton-Valentine-toxins

d) β and δ toxins

Question 6:

Which of the following staphylococcal infections is not toxin mediated?

a) Food poisoning

b) Toxic shock syndrome

c) Septicemia

d) Staphylococcal scalded skin syndrome

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

A college student with a history of intravenous drug abuse presents with a fever for the last 3 weeks. Echocardiography is performed which reveals large friable vegetations on the cusps of the tricuspid valve. Which of the following organisms is the most common cause for this condition?

a) Staphylococcus aureus

b) Staphylococcus epidermidis

c) Streptococcus gallolyticus

d) Streptococcus viridans

Question 8:

By which of the following mechanisms is penicillin resistance commonly transferred to Staphylococcus aureus?

a) Change in PBP

b) Conjugation

Sold by @itachibot

c) Transduction

d) Production of β-lactamase

Question 9:

Contact based isolation is employed for which of the following diseases?

a) Diphtheria

b) MRSA

c) Mumps

d) Asthma

Question 10:

An outbreak of methicillin-resistant Staphylococcus aureus infection was reported in the surgical ward of a hospital. Which of the following precautionary measures would you recommend to best curb its spread?

a) Administration of vancomycin empirically to all the patients

b) Frequent fumigation of wards

c) Wearing masks before performing invasive procedures

d) Washing of hands before & after treating the patients

Question 11:

A 32-year-old man presented with multiple tender, erythematous fluctuant skin lesions as shown in the image. You suspect it to be a case of botryomycosis. Which of the following is the most likely organism implicated in this condition?

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

a) Madurella mycetomatis

b) Actinomyces israelii

c) Staphylococcus aureus

d) Staphylococcus lugdunensis

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 12:

A 14-year-old boy presents with a history of vomiting within 3 hours of consumption of food at a party. Which of the following organism is most likely to cause these symptoms?

a) Staphylococcus aureus

b) Clostridium botulinum

c) Clostridium perfringes

d) Salmonella typhi

Question 13:

An 18-year-old girl presented with burning micturition and fever with chills for three days. Urine culture grew bacteria which were coagulase-negative and novobiocin-resistant. Which of the following is the most likely organism implicated here?

a) Staphylococcus aureus

b) Staphylococcus hominis

c) Staphylococcus epidermidis

Sold by @itachibot

d) Staphylococcus saprophyticus

Question 14:

Choose the incorrect statement regarding Micrococci.

a) It is modified oxidase positive.

b) It is catalase positive.

c) It is rarely associated with disease.

d) It is arranged in chains.

@itachibot someone else, med. If yo

Answer Key

Question No.	Correct Option
1	b
2	a
3	a
4	b
5	c
6	c
7	a
8	c
9	b
10	d
11	c
12	a
13	d
14	d

Detailed Explanations

Solution to Question 1:

An infected wound culture showing Gram-positive cocci arranged in clusters which are coagulase-positive suggest that the cultured organism is Staphylococcus aureus.

Sold by @itachibot

Staphylococci are Gram-positive cocci that are catalase-positive. They grow in clusters because the cells divide successively in three perpendicular planes, with the daughter cells remaining attached to one another following each successive division. Among the 45 different species of Staphylococcus, Staphylococcus aureus is the only one which is coagulase positive.

Other options:

Option A: Streptococci are catalase negative. They are Gram-positive cocci arranged in chains.

Options C & D: S. epidermidis and S. saprophyticus are both coagulase negative.



Different bacteria with their typical arrangements are given below:

• Staphylococcus: Clusters

• Streptococcus: Chains

• Pneumococcus: Pairs/lanceolate shaped

• Micrococcus: Tetrads

Solution to Question 2:

Gram-positive cocci arranged in clusters that produce golden-yellow pigment on nutrient agar is suggestive of Staphylococcus aureus. The pigment production occurs optimally at 22°C.

Staphylococcus aureus is a Gram-positive coccus. On nutrient agar, its colonies are large (2-4 mm in diameter), convex, smooth, shiny, opaque and easily emulsifiable. The colonies appear golden yellow due to the presence of staphyloxanthin (carotenoid). Pigment production occurs optimally at 22°C and in aerobic conditions. It is enhanced when 1% glycerol or milk is incorporated in the medium.

On a nutrient agar slope, there is confluent growth of Staphylococcus, which appears as a characteristic oil-paint appearance.

Sold by @itachibot

Solution to Question 3:

Protein A forms the basis of the co-agglutination reaction shown by Staphylococcus aureus.

Protein A bearing staphylococci are coated with IgG antiserum. Protein A binds to the Fc terminal of IgG, leaving the Fab terminal free to combine with its specific antigen. The corresponding antigen is then added to this mixture which causes agglutination of the bacteria. This is known as the co-agglutination reaction. It is used for streptococcal, gonococcal or hemophilus typing.



Solution to Question 4:

β-hemolysin exhibits a hot-cold phenomenon.

β-hemolysin is a sphingomyelinase that is hemolytic only for sheep cells but not human cells. It initiates hemolysis at 37oC but the hemolysis is evident only after chilling. This is called the hot-cold phenomenon.

Solution to Question 5:

γ-toxin and Panton-Valentine-toxin (leucocidin) of Staphylococcus aureus are called synergohymenotropic toxins.

Synergohymenotropic (synergo - working together, hymenotropic - form pores) toxins are a family of bicomponent toxins that act through the synergistic activity of 2 non-associated secretory proteins - class F and class S proteins. Gamma hemolysin has two components necessary for its hemolytic action. It is encoded by three genes HlgA and HlgC (belonging to class S) and HlgB (belonging to class F), resulting in 2 pairs of class S and F components - HlgA and HlgB, or HlgC and HlgB.

Sold by @itachibot

Panton-Valentine Leucocidin is composed of two components- S and F - encoded by two contiguous and cotranscribed genes, lukS-PV and lukF-PV. It is associated with the majority of community-associated methicillin-resistant Staphylococcus aureus (CA-MRSA).

Solution to Question 6:

Staphylococcal septicemia is not toxin mediated.

Septicemia occurs due to the invasion of bacterial organisms into the bloodstream. This causes an inflammatory response to bacterial products, leading to various manifestations seen in sepsis - fever, hypotension, alkalosis.

Toxin mediated staphylococcal infections include food poisoning caused by enterotoxin, toxic shock syndrome caused by toxic shock syndrome toxin and staphylococcal scalded skin syndrome caused by exfoliative/epidermolytic toxin.

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 7:

Fever with vegetations on the tricuspid valve points to a diagnosis of right-sided infective endocarditis. Staphylococcus aureus is the most common cause of right-sided infective endocarditis among intravenous drug abusers.

Infective endocarditis is the microbial infection of heart valves or endocardium, leading to the formation of vegetations (thrombotic debris and microbial organisms).The following image shows a graphical illustration of vegetations seen in infective endocarditis.



Solution to Question 8:

Sold by @itachibot

Penicillin resistance in Staphylococcus aureus is determined by plasmids that are most commonly transferred by transduction.

Penicillin resistance in Staphylococci most commonly occurs due to bacterial production of beta-lactamase, which inactivates penicillins by splitting their beta-lactam ring. These enzymes are inducible and are plasmid-mediated. They confer resistance to penicillin G, ampicillin, piperacillin etc.

The genetic transfer of antibiotic resistance can occur in the following ways:

• Transformation: After death or cell lysis, some bacteria release their DNA into the environment. Other bacteria, generally of the same species, can come into contact with these free DNA fragments, take them up and incorporate them into their DNA by recombination.

• Transduction: This method involves the transfer of DNA, episome or plasmid from one bacterium to another with the use of a bacteriophage.

• Conjugation: This mechanism requires the presence of a special plasmid called the F plasmid that codes for the sex pilus. It is the most common method of transfer of antibiotic resistance in most other bacteria.

Note: Change in penicillin binding proteins (PBPs) and production of β-lactamases are modes of penicillin resistance seen in S. aureus but the question asks for methods of transfer of drug resistance.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 9:

Contact based isolation is employed for patients who are infected with MRSA (methicillin-resistant Staphylococcus aureus).

The three categories of transmission-based precautions include:

• Contact-based isolation is used for infections, diseases, or germs that are spread by touching the patient or items in the room. Examples include MRSA, vancomycin-resistant enterococci (VRE), diarrheal illnesses, open wounds, and respiratory syncytial virus (RSV).

• Droplet-based isolation is employed for diphtheria and mumps.

• Airborne precautions are required to protect against conditions like measles, severe acute respiratory syndrome (SARS), varicella (chickenpox), and Mycobacterium tuberculosis.

Solution to Question 10:

Among the given options, washing hands before and after treating the patients is the most effective way to curb the spread of methicillin-resistant Staphylococcus aureus (MRSA) infections.

In addition to this, personnel with active S. aureus lesions and carriers should be excluded from areas having patients at highest risk for severe staphylococcal infections like newborn nurseries, intensive care units, operating rooms, and cancer chemotherapy wards. In such individuals, the application of topical antiseptics such as mupirocin to nasal or perineal carriage sites may diminish the shedding of MRSA.

Sold by @itachibot

Solution to Question 11:

The most common cause of botryomycosis is Staphylococcus aureus.

Botryomycosis is a rare chronic bacterial granulomatous disease that usually involves the skin and rarely viscera. It is most commonly caused by Staphylococcus aureus but can also be caused by Pseudomonas, Escherichia coli, Proteus, and Streptococci. Patients present with chronic, suppurative, granulomatous skin lesions like nodules, abscesses and sinuses.

Actinomycetoma, eumycetoma and botryomycosis are difficult to distinguish clinically but are readily distinguished with microbiologic testing or biopsy. Granules of botryomycosis consist of loose masses of cocci. Organisms can also be seen in wound discharge or histologic specimens.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

Vomiting 2 hours after the consumption of food is likely food poisoning by Staphylococcus aureus.

Staphylococcus aureus can grow in carbohydrate and protein food sources, producing enterotoxins. There are 15 types, all of which are superantigens. They are preformed toxins that are relatively heat stable. They stimulate the vagus nerve, vomiting centre of the brain and intestinal peristaltic activity. Type A enterotoxin is responsible for most cases of food poisoning.

The common sources of infection are the foods that are not cooked after handling. These include sliced meats, puddings, pastries, and sandwiches. Ingestion of such contaminated food causes food poisoning within 30 minutes to 6 hours after consumption. Patients present with nausea, vomiting, stomach cramps, and diarrhoea.

Solution to Question 13:

Burning micturition and fever with chills in a young woman points to a diagnosis of urinary tract infection (UTI). The most likely organism implicated here is Staphylococcus saprophyticus as it is coagulase-negative and novobiocin-resistant.

Staphylococcus saprophyticus is a part of the normal flora of human skin and periurethral area and thus is a common cause of urinary tract infection (UTI) in young women.

Sold by @itachibot



Option A: Staphylococcus aureus is a coagulase-positive bacterium. It is an uncommon cause of UTI.

Options B and C: Staphylococcus hominis and Staphylococcus epidermidis are coagulase-negative and novobiocin-sensitive. They are usually considered contaminants if found on urine culture but they may also cause UTIs in children with underlying urinary tract abnormalities.

Solution to Question 14:

Micrococci are arranged in pairs, tetrads or clusters of eight.

Micrococci are Gram-positive cocci that are often mistaken for S. aureus. However, they are larger in size and mostly occur in pairs, tetrads or clusters of eight. They form smaller colonies in culture in comparison to S. aureus. Hugh-Leifson oxidation-fermentation test can be used to differentiate between micrococci (oxidative pattern) and staphylococci (fermentative pattern).

Micrococci are catalase-positive and modified oxidase-positive. They are ordinarily non-pathogenic but can rarely cause infections in an immunocompromised host. They are part of normal bacterial flora in the body.

The following image shows the tetrad arrangement of micrococci:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Different bacteria with their typical arrangements are given below:

1. Staphylococcus: Clusters
2. Streptococcus: Chains
3. Pneumococcus: Pairs/lanceolate shaped
4. Micrococcus: Tetrads

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Corynebacterium, Listeria and Actinomyces

Question 1:

A father brought his 9-year-old unimmunized child to the pediatric OPD with complaints of fever, sore throat and difficulty in swallowing. The findings of oropharyngeal examination are shown in the image below. Which of the following is the most likely causative organism of this condition?

a) Gram-negative diplococci

b) Spiral gram-negative bacteria

c) Club-shaped gram-positive bacilli

d) Gram-positive cocci in chains

Question 2:

A patient has complains of fever with chills. A throat examination reveals swollen glands and a thick, whitish membrane over his tonsils. Albert staining is to be done. What are the constituents of Albert's stain?

a) Carbol fuschin and methylene blue

b) Carbol fuschin and malachite green

c) Toluidine and malachite green

d) Toluidine and methylene blue

Sold by @itachibot

Question 3:

Which of the following is not a selective medium for Corynebacterium diptheriae?

a) Tinsdale medium

b) Loeffler's serum slope

c) Potassium tellurite agar

d) McLeod's medium

Question 4:

A throat swab taken from a patient who presented with fever, dysphagia and cervical lymphadenopathy was sent for culture on potassium tellurite agar. The resultant colonies had a poached egg appearance. Which among the following is the most likely organism implicated here?

a) Corynebacterium diphtheriae gravis

b) Corynebacterium diphtheriae mitis

c) Corynebacterium diphtheriae intermedius

d) Corynebacterium pseudotuberculosis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

The toxin produced by the organisms shown below acts via?



Sold by @itachibot

a) Increase in cAMP

b) Increase in cGMP

c) ADP ribosylation of EF2

d) RNA glycosidase

Question 6:

Which of the following bacteria gain toxigenicity by lysogenic conversion?

a) Corynebacterium diphtheriae

b) Enterotoxigenic E. coli

c) Shigella dysenteriae

d) Klebsiella pneumoniae

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following factors does not affect the toxin production of Corynebacterium diphtheria?

a) Concentration of iron in the medium

b) Tox Phage

c) Biotype

d) Concentration of zinc in the medium

Question 8:

What is Elek's gel precipitation test used to test?

a) Susceptibility to Corynebacterium diphtheriae

b) Biotype of Corynebacterium diphtheriae

c) Toxigenicity of Corynebacterium diphtheriae

d) Antibody titer against Corynebacterium diphtheriae

Question 9:

What diphtheria antibody titer is considered optimally protective?

Sold by @itachibot

a) >0.1IU/ml

b) >0.01IU/ml

c) >0.05 IU/ml

d) >0.5 IU/ml

Question 10:

A 5-year-old boy presented with fever, sore throat, dysphagia, and a tonsillar pseudomembrane. Which of the following complications will you not anticipate in him?

a) Asphyxia

b) Otitis media

c) Myocarditis

d) Pupillary paralysis

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 11:

Which of the following features is not typically seen in diphtheroids?

a) They possess few or no metachromatic granules

b) They are usually arranged in parallel rows

c) Most of them do not produce toxins

d) They do not ferment sucrose

Question 12:

A 45-year-old fisherman presents with pain and a lesion in his hand similar to what is shown in the image. What is the causative organism for this condition?

Sold by @itachibot

a) Streptococcus pyogenes

b) Corynebacterium xerosis

c) Erysipelothrix rhusiopathiae

d) Bacillus anthracis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

A 1-day-old newborn developed severe sepsis and died after 1 day despite aggressive treatment. On examination, multiple granulomas are present over the skin. The organism isolated from the blood sample shows the motility depicted below. What is the most likely causative organism?



Sold by @itachibot

a) Treponema pallidum

b) Vibrio cholera

c) Listeria monocytogenes

d) Clostridium

Question 14:

A 45-year-old man presents to the OPD with complaints of fever with chills, headache, abdominal pain, and diarrhoea. He gives a history of consuming raw refrigerated cucumbers the previous day. What is the most likely etiologic organism?

a) Staphylococcus aureus

b) Bacillus cereus

c) Listeria monocytogenes

d) Vibrio cholerae

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Question 15:

Multiple abscesses with discharging sinuses, demonstrating sulphur granules in pus, are caused by

a) Actinomyces

b) Nocardia

c) Salmonella

d) Tularemia

Question 16:

Which of the following is a false statement regarding Nocardia?

a) They are aerobes

b) They are acid-fast

c) They are commensals

d) They produce urease

Question 17:

Sold by @itachibot

Identify the organism from the given image.



a) Mycobacterium leprae

b) Mycobacterium tuberculosis

c) Actinomyces israelli

d) Nocardia asteroides

Sold by @itachibot
If you purchased this from somewhere else,
you may have been scammed.

Question 18:

A patient presents with a mobile, warm to touch nodule in the neck. On the anaerobic culture of the biopsy material from this swelling, it showed colonies as shown in the image which had gram-positive branching rods on staining. Which of the following is the drug of choice in this condition?

Sold by @itachibot



a) Doxycycline

b) Cotrimoxazole

c) Penicillin

d) Surgical drainage

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 19:

Which of the following toxins act by inhibiting elongation factor -2?

a) 1 & 4

b) 2 & 4

c) 3 & 4

d) 4 & 5

Question 20:

The most common clinical form of actinomycosis is

a) Abdominal

b) Cervicofacial

c) Pelvic

d) Thoracic

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	c
2	c
3	b
4	b
5	c
6	a
7	d
8	c
9	a
10	d
11	d
12	c
13	c
14	c
15	a
16	c
17	d
18	c
19	a
20	b

If yo... you... else, ...d.

Detailed Explanations

Solution to Question 1:

Fever, sore throat and difficulty in swallowing with the presence of a dirty white pseudomembrane on the posterior pharynx is strongly suggestive of diphtheria caused by Corynebacterium diphtheriae. They are described as club-shaped gram positive bacilli.

Corynebacterium diphtheriae are gram-positive, non-sporing, non-capsulated and non-motile bacilli. They possess polymetaphosphate granules that serve as storage granules and are called metachromatic granules. These give the bacilli a beaded or club-shaped appearance on microscopic examination.

Sold by @itachibot

Other options:

Option A: Gram-negative diplococci describes Neisseria menigitidis and N. gonorrhoeae.

Option C: Spiral gram-negative bacteria describes Campylobacter spp. and Helicobacter spp..

Option D: Gram-positive cocci in chains describes Streptococci. They typically cause acute exudative pharyngitis characterised by purulent exudate over the tonsils.

If you purchased this from some else, you may have been scammed.
Sold by @itachi

Solution to Question 2:

The given scenario is suggestive of diphtheria caused by Corynebacterium diphtheria. This organism is stained using Albert's stain which contains toluidine blue 'O' and malachite green. Both these dyes have an affinity for acidic cytoplasm as they are basic dyes.

Albert's, Neisser's, and Ponder's stain are used to demonstrate the metachromatic granules(Volutin / Babes Ernst granules) of Corynebacterium diphtheria.

Corynebacterium diphtheria appears as green-coloured bacilli with purple-blue-coloured metachromatic granules in cuneiform arrangement.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Loeffler's serum slope is an enriched medium for the growth of Corynebacterium diphtheriae. It is not a selective medium as other organisms can also grow on this medium.

The selective media for Corynebacterium diphtheria are potassium tellurite agar, Tinsdale medium and McLeod's medium (tellurite blood agar). Tellurite inhibits the growth of other bacteria. C. diphtheria reduces the tellurite to metallic tellurium, giving its colonies a grey black colour. On

On Loeffler's serum slope, colonies appear as small white opaque discs that may acquire a yellow ting. The advantages for growing C. diphtheriae on Loeffler's serum slope are rapid growth in 6-8 hrs and that its the best medium for metachromatic granules production. The disadvantage of using this medium is that on incubation beyond 6-8 hrs, it supports the growth of commensals as well.

Solution to Question 4:

Poached egg colonies on potassium tellurite agar indicate Corynebacterium diphtheriae mitis.

Macleod classified diphtheria bacilli based on colony morphology on tellurite blood agar into the following three types:

• Gravis: They form daisy head colonies and are variably hemolytic. They ferment glycogen and starch.

• Intermedius: They form frog's egg colonies and are non-hemolytic. They do not ferment glycogen and starch.

• Mitis: They form poached egg colonies and are hemolytic. They do not ferment glycogen and starch.

Sold by @itachibot

The following image shows the various colony morphologies of C. diphtheriae.



Solution to Question 5:

The given image shows gram-positive rods with club-shaped ends, beaded appearance, and a Chinese-letter pattern of arrangement indicative of Corynebacterium diphtheriae.

The toxin produced by the bacteria - diphtheria toxin - inhibits polypeptide chain elongation and protein synthesis by inactivating the elongation factor-2 (EF-2) by ADP ribosylation. It has two fragments - fragment B helps in binding of the toxin to host cells and fragment A inactivates EF-2.



Solution to Question 6:

Sold by @itachibot

Lysogenic or phage conversion is seen in Corynebacterium diphtheriae.

The toxigenicity of diphtheria bacillus depends on the presence of the corynephages (tox+). When the bacillus is cured of the phage, its toxigenic capacity is lost.

A bacteriophage is a virus that parasitises bacteria. It can carry host DNA and transfer it to another bacterium. In lysogenic conversion, the viral DNA is integrated with the bacterial DNA and thus can code for new characteristics, like toxin production. The conversion of non-toxigenic bacterial strains to toxigenic by infecting them with bacteriophages is known as lysogenic conversion.



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

Toxin production of Corynebacterium diphtheriae depends on the concentration of iron in the medium, presence of tox phage, and its biotype. Concentration of zinc in the medium has no effect on this process.

An increase in iron concentration (option A) upregulates the DT (diphtheria toxin) repressor gene, thereby decreasing the toxin production. The optimum level of iron in the medium for toxin production is 0.14 µg/millilitre, while a concentration of 0.5 µg/millilitre inhibits the formation of toxins.

The gene for diphtheria toxin is coded by bacteriophage (corynebacteriophage tox+) and thus the presence of phages (option B) is integral to toxin production. Toxin production also depends on the biotype (option C) - gravis, intermedius, mitis and belfanti - of the bacilli. All four biotypes are toxigenic.

Diphtheria toxin is also produced by Corynebacterium ulcerans and Corynebacterium pseudotuberculosis.

Solution to Question 8:

Sold by @itachibot

Elek's test is used to test the toxigenicity of Corynebacterium diphtheriae.

Elek's gel precipitation test is an in vitro immunoprecipitation (immunodiffusion) test used to determine whether or not a strain of Corynebacterium diphtheriae is toxigenic. In this test, a rectangular strip of filter paper impregnated with diphtheria antitoxin is placed on the surface of a 20% normal horse serum agar. Narrow streaks of the C. diphtheriae strains are made at right angles to the filter paper strip with positive and negative controls. If the strain produces toxins, the toxin will diffuse in the agar and produce a line of precipitation (arrowhead) where it meets the antitoxin at optimum concentration.

The test is shown in the image below.



Option A: For checking susceptibility to diphtheria, Schick test is used.

Option B: The biotype of diphtheria can be determined based on colony morphology and biochemical reactions.

Option D: The antibody titer against diphtheria can be measured by various serological methods like passive haemagglutination, latex agglutination test, toxoid enzyme-linked immunosorbent assay (Toxoid-ELISA), and toxin-binding inhibition ELISA (ToBI-ELISA).

Solution to Question 9:

Diphtheria antibody titer is considered optimally protective if the titre is >0.1 IU/mL.

Diphtheria is primarily the result of the action of diphtheria toxin, not bacterial invasion. Hence, antibodies to the diphtheria toxin is a good measure of protection against diphtheria. These antibodies can be induced by immunisation with diphtheria toxoid vaccine. The titers of IgG antibody against diphtheria toxoid vaccine reflect various levels of protection against the disease as follows:

• <0.01 IU/mL: No protection

• 0.01 -0.1 IU/mL: Minimal protection

Sold by @itachibot

• >0.1 IU/mL: Optimal protection

Note: The question asks optimally protective titer and not minimum effective titer which is >0.01 IU/mL.

Solution to Question 10:

A child with fever, sore throat, dysphagia, and a tonsillar pseudomembrane likely has diphthteria. Pupillary reactions remain normal and pupillary paralysis is not a complication seen with diphtheria.

The complications of diphtheria include asphyxia due to the mechanical airway obstruction by the pseudomembrane. Myocarditis and polyneuropathy are toxic complications of diphtheria. Patients may develop post diphtheritic paralysis involving palatine and ciliary nerves. Paralysis of the diaphragm and neurogenic bladder dysfunction may also occur. Acute circulatory failure, septic complications, such as pneumonia and otitis media and renal failure may also occur.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 11:

Diphtheroids ferment both sucrose and glucose, whereas Corynebacterium diphtheriae ferments only glucose.

Sucrose fermentation is done to differentiate between C. diphtheriae and diphtheroids.

Diphtheroids:

• They are corynebacteria resembling C. diphtheriae which occur as normal commensals in the throat, skin, conjunctiva and other areas.

• In general, diphtheroids stain more uniformly than diphtheria bacilli and possess few or no metachromatic granules.

• Most of them do not produce toxins.

• They tend to be arranged in parallel rows (palisades) rather than the cuneiform pattern.

• The common diphtheroids include Corynebacterium xerosis.

Solution to Question 12:

The above image shows a raised, well-circumscribed, violaceous lesion with the absence of pus. Thus, it is erysipeloid which is a skin infection caused by Erysipelothrix rhusiopathiae.

Erysipeloid usually occurs on the fingers by direct inoculation at the site of a cut or abrasion, in people who handle animals, fish or animal products. Pain, swelling and erythema develop and the condition has been called seal finger and whale finger. Pus is usually absent at the infection site, which helps differentiate it from staphylococcal and streptococcal skin infections. It resolves without treatment in 3-4 weeks.

Sold by @itachibot

Erysipelothrix rhusiopathiae is a slender, non-motile, non-sporing Gram positive bacillus. It is catalase, oxidase, and indole negative. It causes disease in multiple land animals like swine, turkeys and sheep.

Note: Erysipelas is caused by Streptococcus pyogenes. It presents as erythematous raised lesions with sharply demarcated edges.

Solution to Question 13:

Early onset sepsis with skin granulomas and organisms showing tumbling motility points to a diagnosis of granulomatosis infantiseptica caused by Listeria monocytogenes.

Listeria monocytogenes shows slow tumbling motility when grown at 25 °C. But at 37°C, it is non-motile because the peritrichous flagella are produced by the bacillus optimally between 20°C and 30°C. This is known as differential motility as motility pattern changes with a change in temperature. Listeria moves from cell-to-cell using host cell actin framework.

Listeria infection can manifest as febrile gastroenteritis, bacteremia, neurolisteriosis and granulomatosis infantiseptica.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 14:

Fever, headache, abdominal pain, diarrhoea following consumption of refridgerated food is suggestive of food poisoning caused by Listeria monocytogenes.

Listeria monocytogenes can survive at refrigerator temperatures (4°C), under conditions of low pH and high salt. So it is able to overcome food preservation and safety barriers, making it an important foodborne pathogen.

Healthy persons exposed to L. monocytogenes in food may develop a mild, self-limiting febrile gastroenteritis lasting 1–3 days. This develops after an incubation period of 6–48 hours. Symptoms include fever, chills, headache, myalgias, abdominal pain, and diarrhea. Immunocompromised individuals can develop meningoencephalitis, bacteremia, or focal infections.

The drug of choice is high dose intravenous ampicillin.

Solution to Question 15:

Multiple abscesses with discharging sinuses, demonstrating sulfur granules in pus, are characteristic of actinomycosis caused by Actinomyces.

Actinomycosis is a chronic granulomatous infection characterized by indurated swellings in the connective tissue with suppuration and discharge of sulfur granules. The most common causative agent is Actinomyces israelli.

Sold by @itachibot

Various clinical forms of actinomycosis include cervicofacial, thoracic, abdominal, and pelvic actinomycosis. Among them, oral-cervicofacial actinomycosis is most common, presenting as a soft tissue swelling, abscess, or mass lesion involving the angle of the jaw.

Solution to Question 16:

Nocardia are free-living organisms and not commensals.

Nocardia are aerobic Gram-positive filamentous bacilli. They are non-sporing. They are considered weakly acid-fast, determined by Kinyoun's acid-fast staining technique. They produce urease and are catalase positive. They are found in the soil. Nocardia asteroides is the Nocardia species which is most commonly involved in human disease (cutaneous and systemic). The paraffin bait technique is used for the isolation of Nocardia.

Solution to Question 17:

The organism shown in the image is Nocardia asteroides.

Nocardia species are weakly acid-fast filamentous organisms with hyphae-like branching on microscopy. They are aerobic, gram-positive, catalase-positive, and urease-positive. They are weakly acid-fast because they stain with carbolfuchsin and retain when decolourized with 1–4% sulfuric acid instead of the stronger acid-alcohol decolorant.

Infection follows traumatic inoculation of the organism into the subcutaneous tissue from soil or vegetable sources. They can present with a characteristic powdery substance that appears as gritty or necrotic material on the skin or within soft tissues. Accompanying this are multiple discharging sinuses, which are abnormal openings that form as the body attempts to expel infected material. These sinuses can drain purulent, often foul-smelling discharge typically associated with localised abscesses, and the discharge may be viscous, sometimes containing necrotic tissue.

Other options:

Options A and B: Mycobacteria species are acid-fast but not have branching filaments.

Option C: Actinomyces species are not acid-fast. They are Gram-positive arranged in a sun-ray appearance.

Solution to Question 18:

The given image shows molar tooth colonies that are characteristic of Actinomyces species. The given clinical scenario of a mobile, warm to touch nodule in the neck describes cervicofacial actinomycosis also known as 'lumpy jaw syndrome'. Penicillin is the drug of choice for actinomyces infection.

Actinomyces are Gram-positive anaerobic bacilli that produce molar tooth colonies on incubation for 72 hours on brain-heart infusion agar. They cause a chronic suppurative and granulomatous infection called actinomycosis. It can present as one of the following clinical types:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

• Cervicofacial type- Fluctuant mass in the jaw area producing draining fistulas.

• Thoracic type- Pulmonary infection with fever, cough, and purulent sputum.

• Abdominal type- Occurs following ruptured appendix or ulcer. It may spread throughout the peritoneum and involve multiple organs.

Actinomycosis can spread to bones and surrounding tissue and produce multiple discharging sinuses. Treatment requires prolonged administration of penicillin for 6–12 months. In penicillin-allergic patients, clindamycin or erythromycin is effective. Surgical excision and drainage may be required in some cases.

The given image shows cervicofacial actinomycosis.

The given image shows a patient with multiple cutaneous lesions caused by Actinomyces:

Sold by @itachibot

Solution to Question 19:

Both diphtheria toxin and pseudomonas exotoxin act by inhibiting elongation factor-2(EF-2).

Solution to Question 20:

The most common clinical form of actinomycosis is cervicofacial.

Actinomycosis is a chronic granulomatous infection characterized by indurated swellings, mainly in the connective tissue, suppuration, and discharge of sulphur granules. The most common causative agent is Actinomycetes israelli.

Various clinical forms of actinomycosis involves cervicofacial, thoracic, abdominal, and pelvic actinomycosis. Among them, actinomycosis most frequently occurs as oral-cervicofacial form presenting as a soft tissue swelling, abscess, or mass lesion involving the angle of jaw.

Sold by @itachibot

Clostridium and Bacillus

Question 1:

Which of the following organisms is capsulated?

a) Clostridium tetani

b) Clostridium difficile

c) Clostridium perfringens

d) Clostridium botulinum

Question 2:

A young man presented with complaints of severe abdominal pain and watery diarrhoea eight hours after consuming meat at a restaurant. A stool sample was sent for culture and the isolated organism showed this appearance on litmus milk fermentation. Which of the following is the most likely causative organism?



a) Bacillus anthracis

b) Clostridium tetani

c) Clostridium perfringens

d) Clostridium botulinum

If you bought this from someone else, it's been scammed. Sold by @itachibot

Sold by @itachibot

Question 3:

Which of the following reactions are positive for Clostridium perfringens?

a) M'fadyean reaction, CAMP test

b) M'fadyean reaction, Reverse CAMP test,

c) Target hemolysis, Reverse CAMP test

d) Stormy fermentation, Elek's test

Question 4:

A middle-aged diabetic man presented to the ER with severe abdominal pain, diarrhea and vomiting following consumption of pork and sweet potato curry. Following admission, the patient developed intra-abdominal sepsis and was taken into emergency surgery. Histopathology of the resected intestine showed mucosal necrosis. Which of the following toxins is likely implicated in his condition?

a) ° -toxin

b) β-toxin

c) θ-toxin

d) ε-toxin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

Which of the toxins produced by clostridium species is the basis of Nagler's reaction?

a) ° -toxin

b) β-toxin

c) ε-toxin

d) θ-toxin

Question 6:

A patient presents with a laceration over his leg following trauma with an unclean sharp metal rod 3 days ago. He also complains of difficulty in opening his mouth and abdominal cramps. Microscopy of the wound discharge revealed rod-shaped bacilli with terminal spherical spores. Which of the following is the organism most likely implicated here?

a) Clostridium perfringens

Sold by @itachibot

b) Clostridium botulinum

c) Clostridium tertium

d) Clostridium tetani

Question 7:

Which of the following is the first sign of generalized tetanus?

a) Fever

b) Increased respiratory and GI secretions

c) Trismus

d) Autonomic disturbance

Question 8:

Strychnine toxin acts by blocking the action of glycine at the post-synaptic receptors in the motor neurons of the anterior horn of the spinal cord. Which of the following has a similar mechanism of action but acts presynaptically?

a) ° -toxin of C. perfringens

b) Botulinum toxin

c) Tetanus toxin

d) Toxin A of C. difficile

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

An 18-year-old girl presents with a contaminated wound sustained 4 hours ago. She had received complete immunization against tetanus 7 years ago. Which of the following injections should be given to her?

a) Full course tetanus toxoid

b) Single dose tetanus toxoid

c) Human tetanus immunoglobulin

d) Human tetanus immunoglobulin and single dose of toxoid

Question 10:

Sold by @itachibot

An elderly diabetic man presented to the ER with a contaminated wound with necrotic material. As a first-year surgical resident, which of the following would be your first step in treating the conditon?

a) Anti-gas gangrene serum

b) Debridement

c) Antibiotics

d) Tetanus toxoid

Question 11:

An 18-year-old man presented with sudden onset descending paralysis and blurred vision 3 days after the intake of canned food. On examination, the patient has quadriparesis with areflexia. Both the pupils are nonreactive. What is the most probable diagnosis?

a) Poliomyelitis

b) Botulism

c) Gullian Barre Syndrome

d) Tetanus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Question 12:

Which of the following organisms is predominantly proteolytic in Robertson cooked meat medium?

a) Clostridium perfringens

b) Clostridium botulinum A

c) Clostridium difficile

d) Clostridium septicum

Question 13:

Which of the following organisms form the citron bodies?

a) Clostridium septicum

b) Clostridium histolyticum

c) Clostridium tetani

d) Clostridium perfringens

Sold by @itachibot

Question 14:

An elderly diabetic lady was brought to the hospital with complaints of diarrhoea and abdominal pain. She had recently been treated with intravenous clindamycin for cellulitis. She was febrile at the time of examination and the blood reports revealed a Hb level of 12g/dl; WBC count of 18000/mm3; and C-reactive protein 160mg/l. What is the most sensitive test to detect the causative organism?

a) Glutamate dehydrogenase immunoassay

b) PCR to detect toxin gene

c) Toxigenic culture

d) ELISA to detect toxin antigen

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Question 15:

A veterinarian developed an ulcer with central black eschar over the forearm as shown in the image. Which of the following is the most probable causative organism?

a) Pseudomonas aeruginosa

b) Yersinia pestis

c) Bacillus anthracis

d) Francisella tularensis

Sold by @itachibot

Question 16:

Which of the following bacilli characteristically shows the M'Fadyean's reaction?

a) Bacillus cereus

b) Clostridium

c) Bacillus anthracis

d) Corynebacterium diphthteriae

Question 17:

A dairy farmer presented to the dermatology OPD with an ulcer with central black eschar over the forearm. Which of the following media can be used as a selective medium for the growth of the likely etiological agent?

a) PLET medium

b) Buffered charcoal yeast extract medium

c) Cetrimide Agar

d) MYPA agar

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Question 18:

A swab was taken from an ulcer and sent to the laboratory for culture. The following image depicts a blood agar culture plate on which gamma phage suspension has been applied. Which of the following is the most likely organism implicated?



Sold by @itachibot

a) Erysipelothrix rhusiopathiae

b) Clostridium butyricum

c) Bacillus anthracis

d) Bacillus cereus

Question 19:

Which of the following tests is used for the diagnosis of anthrax?

a) Paul- Bunnell test

b) Kahn test

c) Ascoli's thermoprecipitation test

d) Rose-Waaler test

Question 20:

A young patient presented to the medicine OPD with complaints of nausea and vomiting. He gave a history of having fried rice from a local Chinese restaurant a few hours ago. What is the most likely causative organism?

a) Clostridium perfringens

b) Bacillus cereus

c) Staphylococcus aureus

d) Bacillus anthracis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

Which of the following can grow in an acidic pH?

a) Vibrio

b) Lactobacilli

c) Salmonella

d) Shigella

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	c
2	c
3	c
4	b
5	a
6	d
7	c
8	c
9	b
10	b
11	b
12	b
13	a
14	c
15	c
16	c
17	a
18	c
19	c
20	b
21	b

Detailed Explanations

Solution to Question 1:

Clostridium perfringens is the capsulated organism among the given options.

Most of the clostridial species are non-capsulated except C.perfringens and C.butyricum.

All clostridia are motile, with a peritrichate flagella, with the exception of:

• C. perfringens

• C. tetani type VI.

Solution to Question 2:

Sold by @itachibot

The given clinical history and the image showing stormy fermentation of lactose in litmus milk is suggestive of gastroenteritis caused by Clostridium perfingens type A.

The fermentation of lactose by this organism leads to the formation of acid which coagulates the casein in litmus milk. The clotted milk is disrupted due to vigorous gas production and this characteristic finding is obtained that is called stormy fermentation.

Solution to Question 3:

Target hemolysis and reverse CAMP test are positive for Clostridium perfringens.

A narrow zone of complete hemolysis due to θ-toxin and a wider zone of incomplete hemolysis due to ° -toxin results in development of target hemolysis on blood agar by C. perfringens.

In the reverse CAMP test, C.perfringens is streaked over the centre of the plate and Streptococcus agalactiae is streaked at right angles to it. A positive reverse CAMP test shows the presence of an arrow-shaped zone of enhanced hemolysis pointing towards C.perfringens. ° -toxin produced by C.perfringens interacts with the CAMP factor and produce synergistic hemolysis.

M'fadyean reaction is positive Bacillus anthracis.

The CAMP test is positive for Streptococcus agalactiae where it is inoculated on a plate streaked with Staphylococcus aureus.

Elek's test is positive for Corynebacterium diptheriae.

Solution to Question 4:

The condition described here is necrotising enteritis(pigbel) that is caused by β-toxin of Clostridium perfringens type-C.

This is a severe and often fatal enteritis caused by C.perfringens type C strains with heat-resistant spores that germinate in the intestine producing beta toxin, causing mucosal necrosis. The evocative name 'pigbel' is New Guinea pidgin for abdominal pain and diarrhea following unaccustomed feasting on pig meat along with trypsin inhibitors like sweet potatoes.

Immunisation with the type C toxoid has been shown to protect against this condition.

Solution to Question 5:

° -toxin produced by Clostridium perfringens forms the basis of Nagler's reaction.

The ° -toxin of C.perfringens has lecithinase activity. When C.perfringens is grown on a media containing egg yolk with the ° -antitoxin spread on one half of the plate, colonies on the other half without the antitoxin will be surrounded by a zone of opacity. There will be no opacity around the colonies on the half of the plate with the antitoxin due to the specific neutralization of ° -toxin. This is known as Nagler's reaction.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 6:

The above clinical vignette is strongly suggestive of tetanus caused by Clostridium tetani. It is a gram-positive, slender bacillus that has a spherical and terminal spore, giving it a characteristic drumstick appearance.

The genus clostridium consists of gram-positive, anaerobic, spore-forming bacilli.

Classification of Clostridium based on shape and position of spores:

- Central or equatorial: Clostridium bifermentans
- Subterminal: Clostridium perfringens, Clostridium novyi, and Clostridium botulinum
- Oval and terminal (tennis racket shaped): Clostridium tertium and Clostridium difficile
- Spherical and terminal (drumstick appearance): Clostridium tetani

The below image shows the characteristic drumstick appearance of C. tetani.

Sold by @itachibot



Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 7:

The first sign of generalized tetanus is increased tone of the masseter muscle, which results in trismus or lock-jaw.

Tetanus:

• Route of entry of organism- injury, unhygienic surgery, abortion/delivery, otitis media

• The incubation period is 6-10 days.

• First clinical sign- increased masseter tone (trismus/lock-jaw) followed by descending tetanus

• Deep tendon reflex is exaggerated.

• In neonates, difficulty in feeding is the usual presentation.

• Diagnosis is always clinical, microscopy is unreliable.

• Treatment includes cleaning the entry wound first in order to remove anaerobic foci of infection and prevent further toxin production. Metronidazole is preferred for antibiotic therapy. Antitoxin should be given early in an attempt to deactivate any circulating tetanus toxin. Spasms are controlled by heavy sedation with benzodiazepines. Symptomatic treatment is given for respiratory and cardiovascular derangements.

Solution to Question 8:

Tetanus toxin has similar mechanisms of action as strychnine but acts pre-synaptically.

Strychnine prevents the uptake of glycine at inhibitory synapses post-synaptically, while tetanospasmin blocks release of inhibitory neurotransmitters like glycine and gamma-aminobutyric acid (GABA) pre-synaptically. This leads to increased release of acetylcholine, thereby leading to the development of spastic paralysis.

Sold by @itachibot

The image below depicts the mechanism of action of tetanus toxin.



Solution to Question 9:

As she has received complete immunization against tetanus 7 years back, she should be given a single dose of tetanus toxoid for her contaminated wound.

Solution to Question 10:

Initial treatment for contaminated wounds with necrotic material is debridement.

A contaminated wound with necrotic material is suggestive of a higher risk of clostridial infection that may lead to gas gangrene. Emergent surgical exploration and thorough debridement are extremely important in the management of gas gangrene. Antibiotics (penicillin G plus clindamycin) and anti-gas gangrene serum have to be initiated after debridement. Hyperbaric oxygen therapy may also be considered.

Solution to Question 11:

The given clinical vignette of acute onset descending flaccid paralysis with a history of consumption of canned food is suggestive of botulism.

The incubation period from ingestion of contaminated food to the onset of symptoms in food-borne botulism is usually 8–36 h but can be as long as 10 days and is dose-dependent. The clinical syndrome of botulism consists of bilateral cranial-nerve palsies that may progress to respiratory compromise, a bilateral descending flaccid paralysis of voluntary muscles, and even death.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Cranial nerve deficits may manifest as some of the following:

• diplopia

• dysarthria

• dysphonia

• ptosis

• ophthalmoplegia

• facial paralysis and impaired gag reflex

The absence of cranial nerve palsies makes botulism highly unlikely, as does a lack of cranial nerve deficits at the onset of illness. Pupillary reflexes may be depressed and fixed or dilated pupils are sometimes noted.

The cornerstones of treatment for botulism are meticulous intensive care and administration of botulinum antitoxin.

Option A: In poliomyelitis, weakness is generally asymmetric, more proximal than distal, and may involve the legs (most commonly), the arms, or the abdominal, thoracic, or bulbar muscles.

Option C: Guillain–Barre syndrome is usually associated with ascending flaccid paralysis.

Option D: Tetanus causes spastic paralysis.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 12:

Clostridium botulinum A is predominantly proteolytic in Robertson cooked meat medium.

In Robertson cooked meat media, the saccharolytic species like C.perfringens, C.difficile, and C.septicum turn the meat pink, while the proteolytic species such as C.botulinum A turn the meat black.

Examples of predominantly proteolytic clostridia:

• C.bifermentans

• C.botulinum A, B, F

• C.histolyticum

C. tetani is slightly proteolytic but not saccharolytic.

Examples of predominantly saccharolytic clostridia:

• C.perfringens

• C.septicum

• C.novyi

• C.difficile

C.botulinum C, D and E are saccharolytic but not proteolytic.

Sold by @itachibot

Solution to Question 13:

Citron bodies are boat or leaf-shaped pleomorphic bacteria, usually associated with Clostridium septicum.

It is a gram-positive, spore-forming, obligate anaerobic bacterium.

Gram stain morphology of Clostridium septicum from CSF (a) and blood culture (b) in 1000× magnification with spore formation(arrow) is shown below.



Solution to Question 14:

The given clinical vignette of a patient having diarrhea following antibiotic administration is suggestive of pseudomembranous colitis due to Clostridium difficile. Toxigenic culture is the most sensitive method for the detection of Clostridium difficile infection.

The methods for diagnosis of Clostridium difficile infection are:

• Toxigenic culture- Anaerobic incubation of the sample in cycloserine-cefoxitin-fructose agar (CCFA) or its variant for several days. This is followed by identification of the organism in the growth and tests to confirm toxin production. This method has a sensitivity of 100%.

• Cell cytotoxicity neutralization assay (CCNA)- A stool filtrate is prepared and applied onto a monolayer of an appropriate cell line. The cells are observed for toxin-induced cytopathic effect (CPE). This is followed by a neutralization assay to confirm that the CPE is attributable to C.difficile toxins. Sensitivity is around 75-85% when compared to toxigenic culture (reference standard)

• Nucleic acid amplification tests (e.g.PCR)- More sensitive (93.3%) for C.difficile detection than toxin immunoassays but less sensitive than toxigenic culture.

• Toxin immunoassays- Enzyme immunoassays using monoclonal or polyclonal antibodies against C.difficile toxins. Sensitivity is lower than other tests.

if you purchased elsewhere, you may have been

Sold by @itachibot

• Glutamate dehydrogenase immunoassays- Detects glutamate dehydrogenase, a metabolic enzyme produced by all isolates of C.difficile (toxigenic and nontoxigenic). The test is highly sensitive but has low specificity.

Test	Substance detected	Sensitivity	Specificity
Toxigenic culture	C.difficilevegetative cells or spores	High	Low*
Nucleic acid amplification tests	C.difficilenucleic acid for toxin genes	High	Low/moderate
Glutamate dehydrogenase	C.difficilecommon antigen	High	Low*
CCNA	Free toxins	High	High
Toxin A and B immunoassays	Free toxins	Low	Moderate
*Must be combined with a toxin test to increase specificity			

Sold by @itachibot
If you purchased this from somewhere else
you may have been scammed

Solution to Question 15:

The given clinical scenario is suggestive of cutaneous anthrax which is caused by Bacillus anthracis.

Human anthrax presents in one of three ways:

• Cutaneous Anthrax(hide porter's disease)- Black eschar at the base, with a very prominent inflammatory ring around the eschar, is referred to as a malignant pustule.

• Pulmonary Anthrax(Wool sorter's disease)- Hemorrhagic pneumonia.

• Intestinal Anthrax- Hemorrhagic diarrhoea. It occurs mainly in primitive communities that eat the carcasses of animals dying of anthrax.

Robert Koch validated the germ theory of disease after discovering the anthrax bacillus as the cause of anthrax. He also discovered the causative agents of tuberculosis and cholera. Hence, he is known as the Father of Bacteriology.

Koch's postulates, which link specific microorganisms to specific diseases, state that the microorganism must be:

• Found in abundance in diseased organisms, but not in healthy organisms.

• Isolated from a diseased organism and grown in pure culture

• Able to cause disease when inoculated into a healthy organism

• Re-isolated from the diseased experimental host and identical to the original specific causative agent

Sold by @itachibot

Solution to Question 16:

M'Fadyean's reaction is shown by Bacillus anthracis.

When a blood film containing anthrax bacilli is stained with polychrome methylene blue for a few seconds and examined under the microscope, amorphous purplish material is noticed around the bacilli. This represents the capsular material and is called M'Fadyean's reaction.

The following micrograph shows M'Fadyean's reaction:



Solution to Question 17:

Based on his profession and clinical description of the lesion, the diagnosis is most likely to be cutaneous anthrax (most common form of anthrax) caused by Bacillus anthracis. The selective medium for this organism is PLET (polymyxin-lysozyme-EDTA-thallous acetate) medium.

Other options:

Option B: Buffered charcoal yeast extract medium is used for Legionella.

Option C: Cetrimide agar is a selective medium for Pseudomonas aeruginosa.

Option D: Mannitol, Egg Yolk, Phenol red, Polymyxin Agar (MYPA) medium is a selective medium for Bacillus cereus.

Solution to Question 18:

The given image with a circular spot of diminished growth in the culture plate from a swab from an ulcer depicts the gamma phage lysis assay. The most likely organism implicated is Bacillus Anthracis.

Sold by @itachibot

This test is performed by inoculating Bacillus Anthracis in sheep blood agar followed by the application of gamma phage suspension over one area of the culture plate. The gamma phage suspension inhibits the growth of Bacillus anthracis colonies resulting in an an area of diminished growth called a plaque.

Solution to Question 19:

Ascoli's thermoprecipitation test is used for the diagnosis of anthrax caused by Bacillus anthracis. It is an example of a ring precipitation test.

Option A: Paul-Bunnell test is used to diagnose infectious mononucleosis.

Option B: Kahn test is used to diagnose syphilis.

Option D: Rose-Waaler test is used to diagnose rheumatoid arthritis.

Solution to Question 20:

Based on the symptoms and a history of consuming fried rice 6 hours ago, we can diagnose this to be a case of food poisoning due to Bacillus cereus.

Food poisoning cause by B. cereus is of two types:

Bacillus cereus	Diarrheal type	Emetic type
Incubation period	8-16 hours	1-5 hours
Toxin	Secreted in intestine	Preformed
Heat	Heat labile	Heat stable
Food items	Meat	Rice
Clinical feature	Diarrhea, abdominal cramps	Vomiting, abdominal cramps

Solution to Question 21:

Among the given options, lactobacilli is the one that can grow in an acidic pH.

It is a gram-positive, facultatively anaerobic, non-motile, non–spore-forming bacterium. Lactobacillus derives lactic acid from glucose, creating an acidic environment, and can grow in an acidic medium. It is found in the vagina as part of its normal microbiota.

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Sold by @itachibot

Mycobacteria Tuberculosis

Question 1:

The following slide is given as a spotter in your microbiology practical examination. Identify the following organism.



a) Mycobacteria

b) Nocardia

c) Actinomyces

d) Corynebacterium

Question 2:

Which of the following is the component of mycobacteria which makes it acid fast?

a) Mycolic acid

b) Muramic acid

c) Teichoic acid

d) Talosaminuronic acid

Question 3:

Sold by @itachibot

A patient presents early in the morning to the DOTS center to provide sputum samples. Which of the following reagents are used in the digestion and decontamination of his samples before smear preparation?

a) NaOH and KOH

b) NaCl and N-acetyl cysteine

c) N-acetyl cysteine and NaOH

d) KOH and NaCl

Question 4:

A patient with suspected tuberculosis gives a sputum sample for microscopic examination. What is the minimum number of acid-fast bacilli that should be present in 1 mL of sputum for them to be detected by this method?

a) 100

b) 1000

c) 10000

d) 10

Question 5:

A patient has chronic cough with occasional blood in the sputum. He has a history of significant weight loss and night sweats. Which of the following media cannot be used to culture the organism causing his symptoms?

a) BACTEC 460

b) Middlebrook 7H10

c) Loefflers medium

d) McLeod's Medium

Question 6:

A microbiology resident is performing liquid automated culture on the sputum sample of a TB suspect patient. Which of the following culture media would he not use for the same?

a) BactiAlert

b) Versatrek

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

c) BACTEC MGIT 960

d) Tarshis

Question 7:

Which of the following tests can simultaneously detect TB and rifampicin resistance in <2 hours?

a) LAM detection by ELISA

b) Xpert MTB/RIF

c) BACTEC MGIT

d) Line probe assay

Question 8:

Which of the following is not a method of testing drug susceptibility in Mycobacterium tuberculosis?

a) Radiometric broth method

b) Molecular method

c) Disc diffusion method

d) Proportion Method

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

TB test done only on whole blood sample is _____.

a) TrueNat TB

b) Gene xpert

c) Interferon gamma release assay

d) All the above

Question 10:

A urine LAM (Lipoarabinomannan) assay was performed in an HIV +ve patient with a CD4+ count of 40/µL. Infection with which of the following organisms can be diagnosed using this test?

Sold by @itachibot

a) M. tuberculosis

b) M. leprae

c) Mycoplasma

d) Listeria monocyogenes

Question 11:

A middle-aged man came with a history of fever and night sweats for the past 2 weeks. He has had noticeable weight loss. Bronchoalveolar lavage was done, but Mycobacterium tuberculosis was not detected. Extra-pulmonary TB is suspected. Which site is most likely affected?

a) Genitourinary tract

b) Skeletal system

c) Gastrointestinal tract

d) Lymph nodes

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 12:

Your neighbour is worried about a scar on her 5-month-old daughter's arm that is shown below. She tells you the child has received all vaccines on time. Which of the following statements is true about the vaccine that caused this finding?

a) Distilled water is used as diluent for this vaccine

b) The dose administered is 0.5 ml

Sold by @itachibot

c) The route of administration is subcutaneous

d) WHO recommends Danish 1331 strain for vaccine

Question 13:

You are performing the pre-employment check up of a young man. You need to perform a tuberculin skin test as a part of this process. Which of the following tests would you not use for this purpose?

a) Pirquet test

b) Heaf test

c) QuantiFERON-TB gold

d) Tine test

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	c
4	c
5	d
6	d
7	b
8	c
9	c
10	a
11	d
12	d
13	c

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

The image shows pink non-branching acid-fast bacilli on a blue background. These findings are consistent with the staining characteristics of mycobacteria. The stain used is the Ziehl-Neelsen stain.

Other options:

Option B: The image below shows Nocardia stained using an acid-fast staining technique. Nocardia is partially acid-fast and appears as branching filaments.



Option C: The image given below is of a gram-stained specimen of Actinomyces israelii. A. israelii is non acid-fast



Option D: The images below show the staining characteristics of Corynebacterium diphtheriae. It is stained with Albert's stain which shows the needle appearance.

Sold by @itachibot

Corynebacterium diphtheriae-microscopy



Alberts stain



Gram stain

©MARROW

Sold by @itachi... from some... else, ...e been scammed.

Solution to Question 2:

Mycobacteria are acid-fast due to the presence of mycolic acid in their cell walls.

On being stained with carbol fuchsin or fluorescent dyes such as auramine-O or rhodamine, they resist decolourization by 20% sulphuric acid and absolute alcohol for 10 minutes. This property is referred to as acid and alcohol fastness and is routinely used to identify mycobacteria.

Mycobacterial cell wall structure



©MARROW

Solution to Question 3:

NaOH and N-acetylcysteine are used for the digestion and decontamination of sputum used in smear preparation for Mycobacterium tuberculosis.

Sold by @itachibot

Sputum and specimens from non-sterile sites need prior treatment for digestion (to liquefy the thick pus), decontamination (to inhibit the normal flora), and concentration (to increase the yield) before they can be used for testing.

Methods used for digestion and decontamination of sputum:

• Petroff's method- Uses 4% NaOH and HCl.

• N-acetyl cysteine (NALC) + 2% NaOH - NALC liquefies the sputum and NaOH kills the normal flora. This method is better suited for automated culture systems and is superior to Petroff's method.

Note: Formalin or hypochlorite can be used only if smear microscopy is being done. They are not useful for culture or animal pathogenicity testing.

If you p... d by @itachibot ...s from someone else, ...ye... ...scammed

Solution to Question 4:

At least 10,000 acid-fast bacilli should be present per milliliter of sputum for them to be readily demonstrable in direct smears.

Slide reporting as per NTEP guidelines:

No. of AFB	No. of oil immersionfields examined	Report
0	100	0
1-9	100	Scanty
10-99	100	1+
1-10	1	2+
>10	1	3+

Solution to Question 5:

This patient with chronic cough and hemoptysis with a concurrent history of significant weight loss and night sweats most likely has tuberculosis, caused by Mycobacterium tuberculosis. McLeod's medium is not used to culture M. tuberculosis. It is a selective medium for Corynebacterium diphtheriae.

Culture media used to cultivate Mycobacterium tuberculosis include:

• Solid- Loeffler's medium is a serum-containing solid medium used for the culture of M. tuberculosis. Other solid media used include Lowenstein- Jensen medium, Dorset medium, Tarshis medium, and Pawlowsky's medium.

Sold by @itachibot

• Liquid- Middlebrook's media are used for the preparation of antigens and vaccines and in sensitivity testing. Other examples of liquid mycobacterial culture media include Dubos' medium, Sula's medium and Sauton's medium.

• Automated culture methods:

•

• BACTEC 460- uses radioisotopes for detection

• BacT alert- colorimetric method of detection

• BACTEC MGIT- fluorometric detection method.

Sold by @itachibot

Sold by @itachi... if you purchased this from someone else, you may have been scammed.

Solution to Question 6:

Tarshis medium is a solid growth medium used for culturing mycobacterium species. Versatrek, BactiAlert, and BACTEC MGIT 960 are liquid automated cultures used for tuberculosis.

The diagnostic tools for TB are as follows:

1. Sputum microscopy for AFB

• Ziehl Neelson staining

• Fluorescence staining

2. Culture

• Solid media– Lowenstein-Jensen (most common), Petragnini and Dorset, Tarshis, Loeffler, Pawlowsky.

• Liquid automated culture – Versatrek, BactiAlert, BACTEC MGIT 960 (MGIT stands for Mycobacteria Growth Indicator Tube)

• Drug Sensitivity Testing / Proportionate Sensitivity Testing:

• Using MGIT, LJ medium

3. Rapid molecular diagnostic testing

• Line probe assay

• CB-NAAT- Nucleic acid amplification test using GeneXpert system

4. Other diagnostic tools are as follows:

• Chest X-ray

• Tuberculin sensitivity testing

Sold by @itachibot

Solution to Question 7:

Xpert MTB/RIF detects tuberculosis (TB) and rifampicin resistance simultaneously within 2 hours. It is a fully automated, real-time nucleic acid amplification technology, with high specificity and sensitivity.

The WHO recommends its use worldwide as the initial diagnostic test in adults and children who are presumed to have multi-drug resistant tuberculosis (MDR-TB) or HIV-associated TB. Xpert MTB/RIF should also be the first test applied to the cerebrospinal fluid (CSF) of patients in whom tubercular meningitis is suspected as well as a replacement test (over conventional microscopy, culture, and histopathology) for gastric lavage, pleural and other biopsies, and fine-needle aspiration specimens.

The image below shows the Xpert MTB/RIF system.



Solution to Question 8:

Disc diffusion method is used for drug susceptibility testing for most bacteria; however, it is not used for Mycobacterium tuberculosis.

The disc diffusion technique is not suitable for testing Mycobacterium tuberculosis because the organism grows so slowly that the drug has time to diffuse throughout the medium, and sharp zones of inhibition are not seen.

Sold by @itachibot



Methods used in the drug susceptibility testing of M.tb include:

For agar-based methods, 7H10 or 7H11 Middlebrook media are used and only the proportion method can be applied.

Radiometric techniques measure the metabolic activity of the bacteria. Based on changes in these values, drug susceptibility can be estimated.

Conventional drug-sensitivity testing	Radiometric Techniques	Molecular methods
Proportion methodAbsolute concentration methodResistance ratio method	BACTEC 460MB/BacTMycobacteria growth indicator tube (MGIT)VersaTREK	PCR-based assaysDNA probe-basedDNA microarrayGene Xpert

Sold by @itachibot If you found this from someone else, you have been scammed.

Solution to Question 9:

TB test done on a whole blood sample is the interferon-gamma release assay. It has been used for the diagnosis of latent tuberculosis in patients with HIV(who frequently have a negative Mantoux test).

Interferon-gamma release assays include:

• QuantiFERON-TB Gold

• T-SPOT.TB

QuantiFERON-TB Gold quantitates the amount of IFN-γ produced by CD4+ T cells in response to the antigens from Mycobacterium tuberculosis. QuantiFERON-TB Gold Plus (QFT- Plus) is an improved version of the QuantiFERON-TB Gold test. QuantiFERON-TB Gold detects only TB-specific CD4+ responses. However, QuantiFeron-TB Gold Plus detects both CD8+ and CD4+ T-cell responses to TB-specific antigens.

T-SPOT.TB determines the total number of individual effector T cells expressing IFN-γ.

Sold by @itachibot

The TrueNat TB test is a new molecular test that can diagnose TB in one hour as well as test for rifampicin resistance. This test uses a sputum sample taken from the patient. Only about 0.5 ml of the sample is required. Gene Xpert is also a molecular method of testing that cannot use blood as a sample. (Option A and B)

Solution to Question 10:

Urine lipoarabinomannan (LAM) assay is used for the diagnosis of tuberculosis.

Lipoarabinomannan (LAM) antigen is a lipopolysaccharide present in the mycobacterial cell wall, released from metabolically active cells or degenerating cells of patients having tuberculosis. Urine LAM is a useful tool in diagnosing tuberculosis, especially in patients with HIV and low CD4+ T cell count.

Most diagnostic tests for tuberculosis rely on sputum samples, which are difficult to obtain and have low sensitivity in immunocompromised patients, patients with disseminated TB, and children, delaying treatment initiation. Lipoarabinomannan (LAM) is the only WHO-endorsed TB biomarker that can be detected in urine, an easily collected sample.



Solution to Question 11:

Lymph nodes are the most common site of extrapulmonary TB.

The diagnosis is established by fine needle aspiration biopsy.

In descending order of frequency, the extrapulmonary sites most commonly involved in TB are:

• lymph nodes

• pleura

• genitourinary tract

Sold by @itachibot

• bones and joints

• meninges, peritoneum and pericardium.

In less than 50% of these cases, there is associated pulmonary disease.

Solution to Question 12:

The scar shown in the image results from the BCG (Bacillus Calmette-Guérin) vaccine. The WHO recommends Danish 1331 strain for BCG vaccine production.

Normal saline is recommended as the diluent for the BCG vaccine as distilled water is an irritant.

The dose administered is 0.1ml containing 0.1mg.

It is administered intradermally by using 26 gauge tuberculin syringe on the left deltoid.

Solution to Question 13:

QuantiFERON-TB gold is not a type of tuberculin skin test. It is a type of interferon-gamma release assays (IGRAs), a whole-blood test that can aid in diagnosing Mycobacterium tuberculosis infection.

There are 3 described methods for tuberculin skin testing (TST):

• Mantoux test, also known as the Pirquet test (preferred method)

• Heaf test

• Tine test

In contrast, IGRAs that have been approved by the FDA are:

• QuantiFERON-TB Gold In-Tube test (QFT-GIT)

• T-SPOT.TB test (T-Spot)

• QuantiFERON-TB Gold Plus (QFT- Plus)

IGRA is not used in India due to a higher probability of a false positive due to universal BCG vaccination. It can also lead to a diagnosis of active TB in a patient with latent TB and may lead to overtreatment. This test is banned in India after the WHO advised against it in low and middle-income countries.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Other Mycobacteria

Question 1:

Which of the following is a pigment-producing species of mycobacteria?

a) M. ulcerans

b) M. scrofulaceum

c) M. avium intracellulare complex

d) M. xenopi

Question 2:

A microbiologist working with mycobacteria notes moist, smooth, flat, white colonies growing on Lowenstein-Jensen media. Which species of mycobacteria is she likely studying?

a) Mycobacterium tuberculosis

b) Mycobacterium bovis

c) Mycobacterium kansasii

d) Mycobacterium avium-intracellulare

Question 3:

The following findings are obtained during biochemical testing of a mycobacterial species. Identify the organism.

a) M. bovis

b) M. tuberculosis

c) M. simiae

d) M. kansasii

Question 4:

A 54-year-old male presents with an ulcer on his hand. On swab culture, growth of M. ulcerans is seen. What is the diagnosis?

a) Hunner's ulcer

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) Meleney's ulcer

c) Buruli ulcer

d) Marjolin's ulcer

Question 5:

A 42-year-old man who works at an aquarium and handles the fish without protective equipment presents to the OPD with the following finding. Infection with which of the following bacteria led to his condition?

a) M. kansasi

b) M. marinum

c) M. paratuberculosis

d) M. gordonae

Question 6:

Which of the following statements is true about the microbiological diagnosis of the disease afflicting the patient shown below?

Sold by @itachibot

a) Acid fast staining is done by Ziehl-Neelsen technique, using 25% sulphuric acid

b) Sample should be collected from the centre of the lesion

c) The number of samples to be collected is 3-4

d) It is cultivated in nine banded armadillos and foot pads of mice

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 7:

What is the generation time of lepra bacilli?

a) 12-13 days

b) 12-13 hours

c) 15-20 hours

d) 1 month

Question 8:

What are the bacteria present in Virchow's cells?

a) M. avium intercellulare

b) M. tuberculosis

c) M. bovis

d) M. leprae

Sold by @itachibot

Question 9:

You counsel a patient with leprosy about lepra reactions before starting her on multidrug therapy. Which among these is correct regarding these reactions?

a) 1, 2 and 4

b) 3 and 5 only

c) 2, 4 and 5

d) 1 and 2 only

Question 10:

A patient diagnosed with leprosy was undergoing further evaluation for classification of type and immune response. Which of the following statements is false regarding the intradermal test used in such evaluation?

a) Negative in children <6 month age

b) It has no value in prognostication of leprosy

c) Used to classify leprosy

d) BCG vaccine may convert a negative test to positive

Question 11:

You are performing the test shown in the image on a patient. How many days later do you ask the patient to return so that you get an accurate read on the Mitsuda reaction?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Sold by @itachibot

a) 3 days

b) 10 days

c) 20 days

d) 30 days

Question 12:

A leprosy vaccine is being introduced on a pilot basis in India. Which of the following mycobacterial species is the basis for its development?

a) M. vaccae

b) M. bovis Danish 1331 strain

c) M.leprae murium

d) M. indicus pranii

Sold by @itachibot
If you bought this from someone else, you have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	c
5	b
6	d
7	a
8	d
9	c
10	b
11	c
12	d

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

Mycobacterium scrofulaceum is a pigment-producing species of mycobacteria.

M. scrofulaceum is a scotochromogen which produces yellow or orange colored pigments, even in the dark. It belongs to group II of Runyon's classification. It can cause scrofula or cervical adenitis in children.

M.ulcerans, M.avium intracellulare complex (MAC), and M. xenopi belong to group III-non photochromogens as they do not form pigment even on exposure to light.

Solution to Question 2:

The species of mycobacteria being studied here is Mycobacterium bovis. It forms moist, smooth, flat, white colonies on Lowenstein-Jensen (LJ) media that break-up easily when touched.

Mycobacterium bovis belongs to the Mycobacterium tuberculosis complex. It causes tuberculosis-like disease in humans.

Lowenstein–Jensen medium is a commonly used, egg-based, selective media for mycobacteria. It contains:

• Egg- Provides fatty acids and protein required for metabolism

• Malachite green- Prevents the growth of contaminants. Makes it a selective medium.

• Mineral salt solution- Buffer

• Glycerol- Source of carbon, favors growth of M. tuberculosis, but not M. bovis. LJ medium without glycerol, which contains pyruvate, is used for culture of M. bovis.

• L-asparagine- Nitrogen source

The image below shows M.bovis colonies on LJ media.



Mycobacterium tuberculosis colonies on LJ medium are dry, rough, raised and irregular with a wrinkled surface. They are creamy white or buff-colored, tenacious and difficult to emulsify(rough, tough, buff).

If you purchased this from someone else, Sold by @itachibot may have been scammed

Sold by @itachibot



Sold by @itachi... else, If you purchased this from some... may have been scammed.

Solution to Question 3:

The given biochemical findings are characteristic of Mycobacterium simiae. Eventhough atypical mycobacteria or non-tuberculous mycobacteria(NTM) are usually niacin test negative, some NTM like M. simiae and M. chelonae give a positive niacin test positive.

Mycobacteria can be identified based on their biochemical reactions as follows:

The first four tests are used to distinguish between M. tb complex and atypical mycobacteria while the last four tests are used to distinguish between M. tb and M. bovis.

Biochemical test	M. tb	M. bovis	NTM
Catalase	Weak+	Weak +	+++
Peroxidase	+	+	_
Arylsulfatse	-	-	+
Niacin	+	-	-
Nitrate reduction	+	-	N/A
Pyrazinamidase	+	-	N/A
TCH sensitivity	Resistant	Sensitive	N/A

Solution to Question 4:

Sold by @itachibot

Mycobacterium ulcerans causes Buruli ulcer.

M. ulcerans produces a toxin called mycolactone which leads to formation of Buruli ulcer. This is a chronic, debilitating, necrotizing disease of the skin and soft tissue which is characterized by the formation of ulcers that progressively enlarge over time if left untreated. The ulcers heal with disfiguring scars.

The image below shows a Buruli ulcer.

Option A: Hunner's ulcers are seen in a subset of patients with interstitial cystitis. These are areas of inflammation on the bladder wall.

The image below shows a cystoscopic image of a Hunner's ulcer.



Option B: Meleney's ulcer or post-operative synergistic bacterial gangrene occurs in immuno-compromised individuals following abdominal surgery. It is caused by a synergistic interaction between microaerophilic nonhemolytic Streptococcus and hemolytic aerobic Staphylococcus aureus.

Sold by @itachibot

Option D: Marjolin ulcer is a rare and aggressive form of squamous cell carcinoma that grows from burns, scars, or poorly healing wounds.

The image below shows a Marjolin's ulcer in a burn scar.

Solution to Question 5:

The image shows a fish tank granuloma that is caused by M. marinum.

The typical clinical presentation is as a single, small violet papule or nodule confined to one extremity, developing 2-3 weeks after trauma. The nodule may break down to form an indolent ulcer. M. marinarum is a natural pathogen of fish and amphibians, which spreads to humans who come in contact with contaminated swimming pools or fish tanks.

Other non-tuberculous mycobacteria causing disease in humans include:

• M. kansasii: Disease resembles pulmonary TB

• M. scrofulaceum: Cervical lymphadenitis

• M. avium complex: Pulmonary, disseminated TB, especially in AIDS patients

• M. fortuitum: Nail salon furunculosis

M. paratuberculosis (aka Johne's bacillus) causes Johne's disease in cattle. It is associated with Crohn's disease in humans.

Solution to Question 6:

The image depicts leonine facies in a patient suffering from leprosy or Hansen's disease caused by Mycobacterium leprae. It is cultivated in nine-banded armadillos and footpads of mice.

Sold by @itachibot
If you purchased this from elsewhere, you may have been scammed.

Sold by @itachibot

M. leprae is a Gram-positive, weakly acid fast bacillus. Thus, 5% sulphuric acid (option A) is used for decolourisation to demonstrate M.leprae's acid-fastness. On microscopy, the bacilli appear bound together in masses called globi, within which they are arranged in a 'cigar bundle' appearance.

For leprosy diagnosis, specimens are collected from the nasal mucosa, skin lesions and earlobes. Nasal smears are collected from the internal septum of the nose. Skin smears are collected from the edges of the lesions (option B) rather than from the centre. Slit skin smears are taken as well. The skin is pinched up tight to minimise bleeding and a cut is made with a scalpel, deep enough to get into the infiltrated layers. This is done over the buttocks, forehead, chin, cheek, and ears (at least 5-6 different areas) (option C).

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 7:

The generation time of lepra bacilli is 12-13 days.

Generation time, also known as doubling time, is the time it takes for the bacterial population to double. It can also be measured as the time between two cell divisions under optimum conditions. The generation time of different bacilli are given below:

- Lepra bacilli- 12-13 days average (may vary from 8-42 days)
- Tubercle bacilli- 15-20 hours
- Coliform bacilli- 20 minutes

Solution to Question 8:

The bacteria present in Virchow's cells is M. leprae.

Lepromatous leprosy lesions contain large aggregates of lipid laden macrophages (lepra cells/Virchow cells) which are often filled with globi (masses) of acid fast bacilli.

The image below shows a high-power Fite-Faraco stain of a skin lesion showing globi of M.leprae.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 9:

Statements 2, 4, and 5 regarding lepra reactions are correct.

Type 2 lepra reactions are unrelated to drug therapy, whereas type 1 reactions may occur within a year of starting treatment.

New lesions typically appear in type 2 lepra reactions. They are tender and erythematous subcutaneous and dermal nodules. Existing lesions show no changes.

Solution to Question 10:

The lepromin test has value as a prognostic test for leprosy.

The lepromin, or Mitsuda test, is a method to measure cellular immunity in leprosy. It is a skin test for delayed hypersensitivity to lepra antigens. A bacillary suspension of inactivated M. leprae is injected just under the skin and the induration reaction is measured 3–4 weeks after.

It is used to classify leprosy. A negative lepromin test is seen in lepromatous and borderline leprosy. It is negative in children <6 months of age, it may be positive at the end of 1st year. BCG vaccine is capable of converting a negative lepromin test to a positive.

Solution to Question 11:

Mitsuda's reaction is read in the 3rd week, i.e., on the 21st day or 20 days after the day of injection.

Lepromin test is a skin test for delayed hypersensitivity to lepra antigens. It is the only method for studying immunity in leprosy. A bacillary suspension of inactivated M. leprae is injected just

Sold by @itachibot

under the skin.

The first reaction to occur is the early reaction of Fernandez, which consists of erythema and induration developing in 24-48 hours. It usually remains for 3-5 days; it is usually poorly defined and carries less significance. The second and more meaningful result is the late reaction of Mitsuda, starting at 1-2 weeks (7-10 days), reaching a peak in three or four weeks, and gradually subsiding over the next few weeks. On the 21st day, if there is a nodule more than 5 mm in diameter, the reaction is said to be positive.

Note: The 21st day is 20 days after the day of injection which is counted as the 1st day.

Solution to Question 12:

Mycobacterium indicus pranii is the basis for the development of a leprosy vaccine.

The leprosy vaccine species was originally called Mycobacterium W (M.w) but was later renamed Mycobacterium indicus pranii. The vaccine is administered via intradermal route. The dose is 0.1 ml. It is divided into two equal doses and given on both arms.

Booster doses will be given to maintain the immunity level. These are given only in one arm.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Sold by @itachibot

Escherichia, Proteus and Klebsiella

Question 1:

An old woman presented to the OPD with dysuria and frequent micturition. Urinalysis showed pus cells, and culture on MacConkey agar showed the following. Which organism is involved here?



a) Escherichia

b) Salmonella

c) Shigella

d) Proteus

Question 2:

The stool culture of a patient with suspected food poisoning produced hemolytic colonies on blood agar. The isolated organism gave a positive indole test and a positive MUG test. You observe the following colonies on eosin methylene blue agar. Identify the organism.

Sold by @itachibot



a) Escherichia coli

b) Klebsiella pneumoniae

c) Proteus vulgaris

d) Klebsiella granulomatis

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 3:

What is the cut-off for bacteriuria beyond which a diagnosis of catheter-associated urinary tract infection (CAUTI) can be made?

a) >10^2 CFU/ml

b) >10^3 CFU/ml

c) >10^4 CFU/ml

d) >10^5 CFU/ml

Question 4:

A tourist came to the OPD with complaints of abdominal pain and watery diarrhea. Which of the following strains of E. coli is most commonly implicated in this case?

a) Enteropathogenic E. coli

b) Enterohemorrhagic E. coli

c) Enterotoxigenic E. coli

d) Enteroinvasive E. coli

Sold by @itachibot

Question 5:

Identify the organism below that causes effacement and attachment of intestinal cells.



a) ETEC

b) EIEC

c) EPEC

d) EAEC

Question 6:

You are investigating an outbreak of diarrhea. You find out that the etiological agent causes stacked-brick formation on hep-2 cells as shown below. What is the likely organism?

If you purchased this ... you may have been ... else, ... d.

Sold by @itachibot



a) Enteroaggregative E.coli

b) Enterohemorrhagic E.coli

c) Enterotoxigenic E.coli

d) Enteroinvasive E.coli

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 7:

A 29-year-old woman presented with bloody diarrhea, abdominal pain, hemolytic anemia, thrombocytopenia, and acute renal failure. She was diagnosed with hemolytic uremic syndrome and treated with plasmapheresis and dialysis. What is false about the infectious causative agent?

a) Serotype O157:H7 causes hemolytic-uremic syndrome.

b) It does not produce a Shiga like toxin

c) Fecal leukocytes may be seen

d) There is person to person transmission.

Question 8:

Sereny test was employed for laboratory diagnosis of which of the following organisms?

a) Enterotoxigenic E. coli

b) Enteropathogenic E. coli

c) Enterohemorrhagic E. coli

Sold by @itachibot

d) Enteroinvasive E. coli

Question 9:

What is the culture media used to differentiate E. coli O157:H7 from other E. coli?

a) Sorbitol-MacConkey agar

b) Mannitol salt agar

c) Cystine-lactose-electrolyte deficient medium

d) Thiosulphate citrate Bile salt sucrose medium

Question 10:

Match the following E.coli with their respective pathogenesis.

a) A-4,B-3,C-2,D-1

b) A-1,B-3,C-2,D-4

c) A-1,B-2,C-3,D-4

d) A-4,B-2,C-3.D-1

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 11:

Which of the following statements is false?

a) E.coli is an aerobe and facultative anaerobe.

b) Proteus infection increases risk of uric acid stones.

c) E.coli is motile by peritrichous flagella.

d) Proteus can metabolise phenylalanine to phenylpyruvic acid.

Question 12:

Which of the following shows Dienes phenomenon?

a) Proteus mirabilis

b) Klebsiella granulomatis

c) Providencia stuartii

d) Serratia marcescens

Sold by @itachibot

Question 13:

Blood culture of the sputum sample taken from a patient diagnosed with hospital-acquired pneumonia, showed a characteristic phenomenon as shown in the image. Identify the bacteria that exhibit this phenomenon.

a) Klebsiella

b) Enterobacter

c) Serratia

d) Proteus

If you purchased this document elsewhere, you may have been scammed.

Question 14:

A middle-aged man from north India presents with a deformity as shown in the image below. He gives a history of foul-smelling purulent nasal discharge before this nodule appeared. Which of the following organisms is responsible for his condition?

Sold by @itachibot

a) Rhinosporidium seeberi

b) Klebsiella rhinoscleromatis

c) Klebsiella ozaenae

d) Klebsiella oxytoca

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 15:

You notice a sputum sample of a patient suspected of having ventilator-associated pneumonia turns progressively redder over time. Microscopic examination of the sample reveals an absence of red blood cells. Which of the following organisms could have caused this?

a) Streptococcus pneumoniae

b) Proteus mirabilis

c) Klebsiella pneumonia

d) Serratia marcescens

Question 16:

A 19-year-old boy presented with a single painless papule on his penis with no enlarged lymph nodes. A smear was made from the lesion and stained with Wright- Giemsa stain which showed the finding as shown in the following picture. Identify the organism.

Sold by @itachibot



a) Treponema pallidum

b) Herpes simplex

c) Chlamydia trachomatis

d) Klebsiella granulomatis

Sold by @itachi... purchased this from some... else, If yo... may have been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	c
5	c
6	a
7	b
8	d
9	a
10	c
11	b
12	a
13	d
14	b

Sold by @itachibot

15	d
16	d

Detailed Explanations

Solution to Question 1:

This patient who presented with dysuria and frequent micturition with urinalysis showing pus cells, and growth of lactose-fermenting colonies on MacConkey agar suggests a diagnosis of urinary tract infection by Escherichia coli.

Enterobacteriaceae family is classified based on their action on lactose present in MacConkey's medium as:

• Lactose fermenters - Escherichia, Klebsiella

• Late lactose fermenters - Shigella sonnei

• Non-lactose fermenters- Salmonella species, Shigella species

The image below shows lactose fermenting and non-lactose fermenting colonies on MacConkey agar.



Mnemonic to remember non-lactose fermenters: (SPY PSM)

• Salmonella

• Proteus

• Yersinia

• Providencia- late fermenter

• Shigella (except S.sonnei which is a late lactose fermenter)

Sold by @itachibot

• Morganella

Solution to Question 2:

The given scenario of a patient with suspected food poisoning with stool culture growing hemolytic colony on blood agar with a positive indole test and MUG test with a iridescent green colony growth on eosin methylene blue agar is characteristic of Escherichia coli.

E. coli produces the enzyme β-glucuronidase which is tested using the MUG test. In it, the substrate MUG (4-methylumbelliferyl-β-glucuronide) is reduced by the enzyme to produce a fluorescent color which indicates a positive MUG test.

Solution to Question 3:

The cut-off value for bacteriuria to diagnose catheter-associated urinary tract infection (CAUTI) is >103 CFU/mL.

The minimum criterion for catheter-associated asymptomatic bacteriuria (CA-ASB) is >102 CFU/mL.

Solution to Question 4:

This patient with watery diarrhea most likely has traveller's diarrhea caused by Enterotoxigenic Escherichia coli (ETEC).

Enterotoxigenic E. coli (ETEC) is the most common organism that causes diarrhea in travelers. It produces the following toxins:

• LT (heat-labile toxin) - acts by ↑cAMP

• ST (heat-stable toxin) - two types of ST are known, ST-A (ST-I) and ST-B (ST-II)

• ST-A is soluble in methanol and acts by ↑ cGMP in the intestine.

• ST-B is insoluble in methanol, and the mode of action is not known.

Other options:

Option A: Enteropathogenic E. coli (EPEC) causes infantile diarrhea in developing countries. It adheres to the mucosal surface of the small intestine using E. coli adherent factor (EAF) and locus of enterocyte effacement (LEE) and causes effacement of microvilli.

Option B: Enterohemorrhagic E. coli (EHEC) produces verocytotoxin, which targets vascular endothelium. It causes hemorrhagic colitis and hemolytic uremic syndrome.

Option D: Enteroinvasive E.coli (EIEC) invades interstitial epithelial cells in the intestine and causes a condition resembling shigellosis. This ability is determined by a plasmid, which can be detected as a diagnostic test.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 5:

Attachment and effacement lesions on intestinal cells are caused by enteropathogenic Escherichia coli (EPEC).

E. coli is the second most common cause of diarrhea in the pediatric age group. Rotavirus is the most common cause. EPEC causes watery diarrhea which is self-limiting.

EPEC adheres to the mucosal cells of the small bowel and causes loss of microvilli (effacement). This leads to the formation of filamentous actin pedestals or cuplike structures and occasionally may cause entry of the EPEC into the mucosal cells.

Pathogenicity requires the presence of two important factors:

- Bundle-forming pilus encoded by plasmid containing EPEC adherence factor (EAF).
- Chromosomal locus of enterocyte effacement (LEE) pathogenicity island.



The image below shows an electron microscopy image of E Coli.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 6:

The stacked-brick pattern of adherence on hep-2 cells is caused by enteroaggregative E. coli (EAEC).

Enteroaggregative E. coli has been associated with persistent diarrhea. Intestinal colonization of EAEC is mediated by aggregative adherence fimbriae (AAF). EAST-1 toxin (enteroaggregative heat-stable toxin) is considered to be a major virulence factor for EAEC.

The given image shows the stacked-brick pattern of adherence on microscopy at 600x magnification.



Sold by @itachibot

Solution to Question 7:

Hemolytic-uremic syndrome (HUS) is caused by enterohemorrhagic E. coli (EHEC) serotype O157:H7. It produces a Shiga like toxin, therefore EHEC is also known as Shiga toxin-producing E. coli (STEC).

Other options:

Option A - Strains of E. coli producing verocytotoxin or Shiga toxin (stx2a-g) can cause diarrheal disease ranging from mild diarrhea to fatal hemorrhagic colitis and HUS. The most common EHEC belongs to serotype O157:H7.

Option C - In EHEC diarrhea, fecal leukocytes are commonly seen (70%) but although it is an infectious condition, fever is not seen. The absence of fever can lead to its misdiagnosis as a non-infectious condition such as intussusception or ischemic bowel disease.

Option D - The disease is transmitted through contaminated feces. Person-to-person transmission can occur if infected persons do not wash their hands after using the toilet or after changing diapers.

Note: Shiga toxin produced by S. dysenteriae is referred to as Stx.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 8:

The Sereny test is an animal pathogenicity test that used to be employed for laboratory diagnosis of enteroinvasive E. coli and Shigella.

Freshly isolated suspension of bacteria is instilled into the eyes of guinea pigs which leads to mucopurulent conjunctivitis and severe keratitis.

It is not in use anymore as we have better methods of detecting toxigenicity now like molecular methods.

Solution to Question 9:

Sorbitol-MacConkey agar is used to differentiate E. coli O157:H7 from other E. coli.

E. coli O157:H7 does not ferment sorbitol and produces pale colonies on Sorbitol-MacConkey agar. All other strains of E. coli ferment sorbitol. Thus they produce pink colonies on Sorbitol-MacConkey agar.

O157:H7 strains do not contain β-glucuronidase and so are unable to use the substrate 4-methylumbelliferyl-β-glucuronide(MUG). Thus, they give a negative MUG test.

Solution to Question 10:

The correctly matched pairs are as follows:

The various strains of Escherichia coli are:

Sold by @itachibot

• Enterotoxigenic E. coli (ETEC) is the most common organism that causes diarrhea in travelers. It produces the following toxins:

• LT (heat-labile toxin)- acts by ↑cAMP

• ST (heat-stable toxin)- two types of ST are known, ST-A (ST-I) and ST-B (ST-II)

• ST-A is soluble in methanol and acts by ↑cGMP in the intestine.

• ST-B is insoluble in methanol, and the mode of action is not known.

• Enteroinvasive E. coli (EIEC) invade interstitial epithelial cells in the intestine and cause a condition resembling shigellosis. This ability is determined by a plasmid, detection of which can also be a diagnostic test. Sereny's test was employed to diagnose EIEC but it is not in use anymore.

• Enterohemorrhagic E. coli (EHEC) produces verocytotoxin, which targets vascular endothelium. It causes hemorrhagic colitis and hemolytic uremic syndrome.

• Enteropathogenic E. coli (EPEC) causes infantile diarrhea in developing countries. It adheres to the mucosal surface of small intestine using EAF and LEE and causes effacement of microvilli. This leads to attachment and effacement lesions of intestinal cells.

• Enteroaggregative E. coli (EAEC) produces the stacked brick pattern of adherence on hep-2 cells. EAEC has been associated with persistent diarrhea. Intestinal colonization of EAEC is mediated by AAF (aggregative adherence fimbriae). EAEC produces EAST-1 toxin (enteroaggregative heat-stable toxin) which is considered to be a major virulence factor.

If you purchased this from someone else, you have been scammed. Sold by @itachibot

E. coli	Mechanism of action
1. ETEC	A. Produce a heat-stable toxin and cause activation of guanylyl cyclase
2. EIEC	B. Invades intestinal mucosal cells
3. EHEC	C. Produces a toxin against vascular endothelium cells
4. EPEC	D. Adhere to the mucosal surface and cause effacement of microvilli

Solution to Question 11:

Proteus increases the risk of struvite stones in the bladder as it leads to the alkalinisation of urine. Struvite stones are magnesium-ammonium phosphate stones that appear as rectangular/coffin-lid crystals shaped on urine microscopy. Uric acid stones are precipitated in acidic urine.

Proteus produces enzyme phenylalanine deaminase which converts phenylalanine to phenyl pyruvic acid. This property is the basis for the phenylalanine deaminase test.

Sold by @itachibot

The test is performed by inoculating the organism in phenylalanine agar. After incubation, 10% ferric chloride is added to the media. If phenyl pyruvic acid was produced, it will react with the ferric chloride and turn dark green. If the medium remains a straw colour, the organism is negative for phenylalanine deaminase production.

E. coli is an aerobe and facultative anaerobe. Its motility is due to the presence of peritrichous flagella. It is the most common cause of urinary tract infections.

Solution to Question 12:

Dienes phenomenon is exhibited by Proteus mirabilis.

When two identical strains of Proteus are inoculated at different places of the same culture plate, the resulting swarms of growth coalesce and no line is formed between swarming cultures of the same strain. However, if they are from two different strains, the spreading films of growth fail to coalesce and remain separated by a narrow but easily visible furrow (known as Dienes line). This is known as Dienes phenomenon.

The image below shows Dienes phenomenon.



Solution to Question 13:

The given image of the blood culture finding shows the characteristic bull's eye swarming pattern exhibited by Proteus.

Proteus species move very actively with the help of peritrichous flagella, resulting in swarming on solid media. It can be suppressed by the addition of:

• phenyl ethyl alcohol

• chloral hydrate

Sold by @itachibot

• sulphonamides

• sodium azide

• boric acid

• increasing the concentration of agar to 6%.

Swarming does not occur on MacConkey medium and cystine-lactose-electrolyte-deficient (CLED) medium.

Proteus is a gram-negative bacillus seen as a normal gastrointestinal flora. The two most common species causing infection in humans are P. mirabilis and P. vulgaris, which can be differentiated by spot indole test. P. mirabilis is indole negative while P. vulgaris is indole positive. Production of urease enzyme by Proteus, which hydrolyses urea to ammonia causes alkalinization of urine. Alkalinization of urine leads to the precipitation of organic and inorganic compounds which contributes to the formation of Struvite stones. P. mirabilis is associated with urinary tract infections while P. vulgaris is usually associated with wound and soft tissue infections.

Other organisms that show swarming include:

• Clostridium tetani

• Vibrio parahaemolyticus

• Serratia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 14:

The above image and clinical scenario are suggestive of rhinoscleroma and it is caused by Klebsiella rhinoscleromatis.

K. rhinoscleromatis is a gram-negative bacillus also known as Frisch bacillus. It is endemic in north India. Rhinoscleroma is characterized by chronic granulomatous hypertrophy of the nose. Biopsy of the lesion shows:

• Mikulicz cells- large foamy cells with a central nucleus and vacuolated, cytoplasm containing causative bacilli.

• Russell bodies- eosinophilic inclusion bodies found in the plasma cells. These are also observed in multiple myeloma.

Other options:

Option A: Rhinosporidium seeberi causes rhinosporidiosis.

Option C: K. ozaenae causes atrophic rhinitis.

Option D: K. oxytoca is associated with hemorrhagic colitis.

Solution to Question 15:

The given condition of sputum sample turning redder over time can be caused by Serratia marcescens.

Sold by @itachibot

Serratia marcescens are pleomorphic coccobacilli that can grow in sputum after collection and can mimic hemoptysis because they produce a red pigment called prodigiosin.

S. macescens mainly cause healthcare-associated infections such as ventilator-associated pneumonia, meningitis, endocarditis, septicemia and peritonitis. They can contaminate IV fluids, surgical instruments, and antiseptic solutions.

Solution to Question 16:

The given clinical vignette and the image showing Donovan bodies in Wright/Giemsa-stained impression smears indicate a diagnosis of donovanosis caused by Klebsiella granulomatis. They appear as rounded coccobacilli within cystic spaces in large mononuclear cells (called Pund cells).

Klebsiella granulomatis shows bipolar condensation of chromatin giving a safety pin appearance.

Bipolar staining (safety pin appearance) is shown by:

- H. ducreyi
- Vibrio parahaemolyticus
- Klebsiella granulomatis
- Yersinia pestis
- Burkholderia pseudomallei
- Burkholderia mallei

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

Shigella and Salmonella

Question 1:

A child presents with a high fever, acute abdominal pain, and malaise. The child passes small amounts of bloody mucopurulent stools with increasing tenesmus and cramps. What is the most likely causative organism?

a) Vibrio cholerae

b) Rotavirus

c) Shigella dysenteriae

d) Entamoeba histolytica

Question 2:

Which of the following toxins acts by inhibiting protein synthesis?

a) Cholera toxin

b) Shiga toxin

c) Pertussis toxin

d) Labile toxin of ETEC

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 4-year-old boy developed bloody diarrhea and abdominal cramps. The bacteria isolated from the stool specimen did not ferment lactose or mannitol. Which of the following organisms is implicated here?

a) Shigella dysenteriae

b) Shigella flexneri

c) Shigella boydii

d) Shigella sonnei

Question 4:

Which of the following salmonellae is non-motile?

Sold by @itachibot

a) Salmonella typhi

b) Salmonella enteridis

c) Salmonella gallinarum

d) Salmonella chester

Question 5:

A Greek patient with travel history to India is diagnosed with enteric fever. What is the minimum amount of pathogen required to cause this condition?

a) 1-10 bacilli

b) 100 bacilli

c) 10^3 -10^6 bacilli

d) 10^8 -10^10 bacilli

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

What is the incubation period of Salmonella typhi?

a) 2-5 days

b) 5-21 days

c) 14-25 days

d) 20-30 days

Question 7:

A 25-year-old man presented to the ER with fever, splenomegaly, and a rash as shown in the image below. After admission, he developed intestinal perforation and an exploratory laparotomy revealed several longitudinal ulcers in the small bowel. What is the probable diagnosis?

Sold by @itachibot

a) Enteric fever

b) Tuberculosis

c) Amoebiasis

d) Hemorrhagic colitis

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 8:

While evaluating a patient suspected to have enteric fever, you order a widely used tube agglutination test in the second week of his illness. Which of the following statements regarding this test is false?

a) H agglutinin titer > 160 and O agglutinin titer > 80 is considered significant

b) O antibody cannot differentiate between Salmonella types.

c) O antibody lasts longer and hence is not indicative of recent infection.

d) High titer value in a single test is not confirmative.

Question 9:

A young man with fever for 2 weeks has the following serology findings. Which of the given statements is true about the disease he has?

a) Intestinal perforation following infection with S.typhi occurs in the 1st week.

b) Faget's sign is seen in enteric fever.

c) In Widal test, Felix tube is used to detect H antigen.

Sold by @itachibot

d) Blood culture is the gold standard of diagnosis

Question 10:

A patient, suspected to have enteric fever, has been started on ceftriaxone by his local village healer 2 days ago. Which investigation will confirm his diagnosis at this stage?

a) Blood culture

b) Bone marrow culture

c) Stool culture

d) Urine culture

Question 11:

You are administering a Vi capsular polysaccharide vaccine to a child in Germany before his trip to India. Which of the following is false about the antigen used in this vaccine?

a) It is a capsular antigen

b) It is polysaccharide in nature

c) It is heat stable

d) It is alcohol stable

Question 12:

A man living in the UK is planning to travel to India. His doctor prescribes the necessary vaccines prior to travel including an oral typhoid vaccine. What is the schedule for this vaccine?

a) Day 1,3,5,7

b) Day 1,3,7,14

c) Day 1,7,14

d) Day 1,7,14,21

Question 13:

Which of the following drugs is not used for typhoid fever in adults?

a) Azithromycin

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) Ciprofloxacin

c) Ceftriaxone

d) Amikacin

Question 14:

A patient developed fever and diarrhoea after consumption of undercooked eggs. Which of the following statements is false about this condition?

a) Gastroenteritis due to non-typhoidal salmonellosis is usually self limited.

b) Stool cultures remain positive for 4-5 weeks after infection.

c) Most common isolate from stool culture is Salmonella enteritidis.

d) Non-typhoidal salmonellosis can be acquired from multiple animal reservoirs.

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	c
2	b
3	a
4	c
5	c
6	b
7	a
8	c
9	b
10	b
11	c
12	a
13	d
14	c

Detailed Explanations

Sold by @itachibot

Solution to Question 1:

Acute onset of symptoms (fever, abdominal cramps) with dysentery (bloody mucopurulent stools) and tenesmus is most likely due to Shigella dysenteriae.

Shigella dysenteriae is a Gram-negative bacillus causing bacillary dysentery in humans. It is spread through faeco-oral transmission.

Other options:

Option A: Vibrio cholerae results in watery diarrhea with characteristic rice watery stools.

Option B: Viruses such as rotavirus and norovirus causes watery diarrhea through mucosal irritation.

Option D: Entamoeba histolytica results in gradual onset of symptoms usually 1-3 weeks with dysentery.

Solution to Question 2:

Shiga toxin acts by inhibiting protein synthesis.

Shiga toxin or verotoxin is an exotoxin produced by Shigella species. It consists of two subunits A and B. The B subunit binds to globotriaosylceramide receptors on the target cell, and then, the A subunit gets internalized. The A subunit leads to inhibition of 60S ribosome, leading to shut down of protein synthesis. Its site of action is in the endothelial cells of small blood vessels of GIT.

Solution to Question 3:

Shigella dysenteriae does not ferment mannitol and lactose.

Shigella species cause bacillary dysentery (bloody diarrhea with abdominal cramps). They do not ferment lactose, except S.sonnei which is a late lactose fermenter. To differentiate between the various Shigella species, mannitol fermentation is noted.

Classification of pathogenic Shigella species:

Designation	Group	Mannitol Fermentation	Ornithine decarboxylase	LactoseFermentation
S. dysenteriae	A	-	-	-
S. flexneri	B	+	-	-
S. boydii	C	+	-	-
S. sonnei	D	+	+	Late lactose fermenter

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Solution to Question 4:

S. gallinarum is non-motile.

Salmonellae are aerobic or facultatively anaerobic Gram-negative bacilli. They are motile due to the presence of peritrichous flagella, except S. gallinarum and S. pullorum.

Solution to Question 5:

Enteric fever is caused by S. typhi or S. paratyphi. The minimum amount of S. typhi required to cause enteric fever is 103 to 106 organisms.

Enteric fever spreads through the feco-oral route. It presents with step-ladder fever and abdominal pain after an incubation period of 10–20 days. Over the next week, the patient can develop abdominal distension, diarrhea, splenomegaly and characteristic rose spots on the abdomen due to vasculitis.

The amount of pathogen required to cause an infection in the host is known as the infectious dose. The infectious doses of few bacteria are given below:

• Escherichia coli - 106 – 108 organisms

• Shigella - 10 – 100 organisms

• Escherichia coli O157:H7 - <10 organisms

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 6:

The incubation period for Salmonella typhi is from 5 to 21 days.

The incubation period for Salmonella typhi typically averages 10–14 days, but can range from 5 to 21 days. The duration is likely dependant on the inoculum size and the host's health and immune status.

Solution to Question 7:

The given image shows Rose spots. The combination of this with fever, splenomegaly, presence of longitudinal ulcers in the intestine and intestinal perforation is suggestive of enteric fever.

The term enteric fever includes typhoid fever caused by Salmonella typhi and paratyphoid fever caused by Salmonella paratyphi A, B and C.

Enteric fever presents with step ladder pyrexia. The fever is accompanied by relative bradycardia which is known as the Faget sign. Hepatosplenomegaly is a common finding on examination. Rose spots that fade on pressure appear on the skin during the second or third week.

Since bile is a good culture medium for the bacilli, they multiply abundantly in the gallbladder and are discharged continuously into the intestine. In the intestine, it causes inflammation and

Sold by @itachibot

necrosis leaving behind the characteristic typhoid ulcers which are oriented along the axis of the intestine (longitudinal ulcers). Ulceration of the bowel leads to two major complications- intestinal perforation and hemorrhage.

Option B: In intestinal tuberculosis, ulcers are perpendicular to the axis of the intestine (transverse ulcers).

Option C: Flask shaped ulcers are characteristic of amoebiasis.

Option D: Hemorrhagic colitis is caused by Shigella dysenteriae and some strains of E. coli. It is characterised by bloody diarrhea with tenesmus.

Solution to Question 8:

The tube agglutination test which is widely used for the diagnosis of enteric fever is the Widal test. The O antibodies detected via this test disappear within a few weeks of disease resolution and so are indicative of recent infection.

Widal test is a serological test for the measurement of O and H antigens of Salmonella typhi and Salmonella paratyphi present in the patient's serum. Four antigens are used- O antigens of S. typhi (TO), H antigens of S. typhi (TH), H antigens of S. paratyphi A (AH) and H antigens of S. paratyphi B (BH).

Paratyphoid O antigens cross-reacts with the typhoid O antigen due to the sharing of factor 12. It is not used in the test and thus, O antibody cannot differentiate between Salmonella types.

H agglutinin titer > 160 and O agglutinin titer > 80 is considered significant. However, a 4-fold rise in titers between 1st and 3rd week is more significant compared to a single high titer because of the possibility of an anamnestic reaction - a transient rise of titer due to unrelated infections (malaria, dengue) in persons who had a prior enteric fever. After one week interval, if the titer increases, it indicates true infection. If the titer decreases, it indicates an anamnestic reaction.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 9:

Widal results showing titers for O antibody >80 and H antibody for S.typhi to be >160 is suggestive of enteric fever. Faget's sign is seen in enteric fever.

Enteric fever, caused by Salmonella typhi and Salmonella paratyphi A, B and C, presents with step ladder pyrexia. The fever rises one day and falls the subsequent morning. It continues to form peaks and troughs. Hepatosplenomegaly is a common finding on examination. Rose spots that fade on pressure appear on the skin during the second or third week. Faget's sign is the unusual pairing of fever with bradycardia (relative bradycardia) instead of tachycardia. It is also observed in yellow fever, brucellosis, and tularemia. Intestinal perforation (option A) is a complication of enteric fever caused by Salmonella typhi. It occurs in the 3rd week following infection.

Widal test is a serological test used for the diagnosis of enteric fever during the 2nd week. It measures the levels of H and O antigens in the patient's serum.

Sold by @itachibot

Bone marrow culture is the gold standard (option D) for diagnosis of enteric fever. Blood culture is most commonly performed to confirm the diagnosis, however, bone marrow is more sensitive.

H antigen	O antigen
Flagellar antigen	Somatic antigen
Fluffy/woolly agglutination se en	Granular agglutination seen
Appear late and persist for lo nger duration	Appear early and persist for s horter duration
Dreyer's tube (conical bottom) used	Felix tube (round bottom) us ed

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

Bone marrow culture is preferred at this stage.

Bone marrow culture is the most sensitive test (>80% sensitivity) for diagnosis of enteric fever. It remains positive for upto 5 days after starting antibiotics, unlike blood culture. Hence, it is preferred in the patient above who has been on antibiotics for 2 days.

Blood culture has a sensitivity of about 60% for diagnosis of enteric fever. Its sensitivity drops with duration of illness, and is thus useful onlyduring the 1st week of enteric fever. It rapidly becomes negative on treatment with antibiotics. Stool culture has a sensitivity of 30% and only yields positive results from the 2nd or 3rd week onwards. Urine culture has low sensitivity and may be positive only after the 2nd week.

Note: The sensitivity of diagnostic methods for enteric fever is duodenal content culture > bone marrow culture > blood culture. However, duodenal content culture is rarely done in clinical practice.

Solution to Question 11:

Vi capsule polysaccharide antigen is a heat-labile antigen used in the Vi capsular polysaccharide vaccine (ViCPS) for typhoid.

Vi antigen is found in Salmonella typhi, S. paratyphi C and S. dublin. It is a capsular antigen and is a polysaccharide. It acts as a virulence factor by inhibiting phagocytosis and resisting complement activation and bacterial lysis. It envelopes the O antigen, and thus, many S. typhi strains fail to agglutinate with O antiserum. However, it can be destroyed by heat. It is alcohol stable. This antigen is poorly immunogenic.

The Vi capsular polysaccharide vaccine (Vi CPS) contains purified Vi polysaccharide. It is given as a single dose of 0.5ml intramuscularly to children over 2 years of age. It is recommended for travellers to areas endemic for Salmonella infections.

Sold by @itachibot

Solution to Question 12:

The oral typhoid vaccine is given on days 1, 3, 5, and 7.

Typhoral or Ty21a is a live attenuated oral typhoid vaccine. It is given in 4 doses given on days 1, 3, 5, and 7. No antibiotics should be given during this period. lt is indicated only in individuals over 6 years of age. Protective immunity starts after 7 days of the last dose and lasts for 4 years. Ty21a can also be administered to HIV-positive, asymptomatic individuals as long as their CD4 count is above 200/mm3. It is contraindicated in immunocompromised patients.

Solution to Question 13:

Amikacin is not used in the treatment of typhoid fever.

Amikacin is an aminoglycoside. Despite showing efficient in vitro killing of Salmonella, first- and second-generation cephalosporins and aminoglycosides have proven to be ineffective in actual clinical cases.

@itachibot someone else,

Antibiotic therapy for enteric fever in adults			
Indication	Agent	Dosage (Route)	Duration, D ays
Empirical Treatment			
	Ceftriaxone	2 g/d (IV)	10-14
Azithromycin	1 g/d (PO)	5	
Fully Susceptible			
Optimal treatment	Ciprofloxacin	500 mg bid (PO) or 400 mg q12 h (IV)	5-7
Azithromycin	1 g/d (PO)	5	
Alternative treatment	Amoxicillin	1 g tid (PO) or 2 g q6h (IV)	14
Chloramphenicol	25 mg/kg tid (PO or IV)	14-21	
Trimethoprim-sulfamethoxaz ole	160/800 mg bid (PO)	7-14	
Multidrug-resistant, fluroqui nolone-susceptible			
Optimal treatment	Ceftriaxone	2 g/d (IV)	10-14
Azithromycin	1 g/d (PO)	5	
Alternative treatment	Ciprofloxacin	500 mg bid (PO) or 400 mg q12h (IV)	5-14
Quinolone-resistant			
Optimal treatment	Ceftriaxone	2 g/d (IV)	10-14
Azithromycin	1 g/d (PO)	5	

Sold by @itachibot

Antibiotic therapy for enteric fever in adults			
Alternative treatment	High dose ciprofloxacin	750 mg bid (PO) or 400 mg q 12h (IV)	10-14
Ceftriaxone-resistant			
Optimal treatment	Meropenem	1 g q8h (IV)	10-14
Azithromycin	1 g/d (PO)	5	

Solution to Question 14:

The given case scenario is suggestive of non-typhoidal salmonellosis (NTS). The most common organism isolated in NTS from stool culture is Salmonella typhimurium.

Nontyphoidal salmonellosis refers to illnesses caused by all serotypes of Salmonella except for Salmonella typhi, Salmonella paratyphi A, B, and C. Transmission of nontyphoidal salmonellosis occurs usually through the consumption of food or water contaminated with animal feces. Transmission may also occur through food products, especially eggs, poultry, undercooked ground meat, and hence multiple animals act as reservoirs for non-typhoidal salmonella.

Gastroenteritis is the most common clinical presentation of NTS. The illness usually lasts 4–7 days, and most people recover without treatment. Culturing organisms continues to be the mainstay of clinical diagnostic testing for nontyphoidal salmonella infection. Approximately 90% of isolates are obtained from routine stool culture until 4-5 weeks after infection.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

Vibrio and Campylobacterales

Question 1:

A child presents with profuse, painless diarrhea. Stool microscopy shows the following finding. What is this characteristic appearance called?

a) School of fish appearance

b) Railroad appearance

c) Fish in stream appearance

d) Drumstick appearance.

Question 2:

While studying the microscopic features of Vibrio cholerae, you observe the following finding. What type of motility can you see here?

Sold by @itachibot



MARROW

a) Lashing motility

b) Swarming motility

c) Darting motility

d) Twitching motility

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 3:

On the basis of which antigen is Vibrio cholerae classified under the Sakazaki typing scheme?

a) O antigen

b) K antigen

c) H antigen

d) O and H antigen

Question 4:

Which of the following statements is false regarding non-agglutinable vibrios?

a) They belong to serogroup O2 – O200

b) They are mostly pathogenic

c) They are not agglutinated by O-1 antiserum

d) These are similar to V. cholera in most cultural and biochemical reactions

Sold by @itachibot

Question 5:

A microbiologist studying Vibrio cholera added sulphuric acid to a 24-hour peptone culture of the organism and a reddish-pink color developed. Which of the following is true about the test he performed?

a) It is known as methyl red test

b) A nitrosoindole compound is formed giving it the red colour

c) The red colour is obtained due to pigment production by the bacteria

d) The red colour is obtained due to acid & gas production by Vibrio

Question 6:

A 20-year-old woman presents with complaints of passing profuse loose stools which looked like rice water for the past 3 days. There's a history of similar symptoms in the family. Stool samples were sent to the lab for analysis and the test given in the image is performed. Identify this test.



a) Loop test

b) String test

c) Wire test

d) Duodenal string test

Question 7:

If you bought this, you were scammed by @itachibot. Please report this to Telegram if you got this from someone else.

Sold by @itachibot

Which of the following bacteria is not microaerophilic?

a) Helicobacter pylori

b) Campylobacter jejuni

c) Mycobacterium bovis

d) Vibrio cholera

Question 8:

The disruption of which of the following is responsible for watery diarrhea in cholera caused by Vibrio?

a) Intercalated discs

b) Macula densa

c) Zonula occludens

d) Hemidesmosomes

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 9:

A young child was brought to the ER with a history of loose stools. On examination, she is tachycardic, appears lethargic and skin goes back slowly after the skin pinch test. She was admitted and started on IV fluids and input-output charting was done. The given image shows stool output from the patient. Which of the following is the mechanism of action of the toxin responsible for the same?

Sold by @itachibot

a) Inhibit elongation factor -2

b) Toxin causes Increased guanyl cyclase activity

c) The toxin causes continued activation of adenylate cyclase

d) Inhibit 60S ribosomal subunit

Question 10:

Which of the following does not play a role in the virulence of Vibrio cholerae?

a) Lipopolysaccharide

b) Flagella

c) Pilum

d) Cholera toxin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 11:

An El Tor vibrio was identified to be the causative agent of a diarrhea epidemic. Which of the following is not a characteristic feature of this organism?

a) Resistant to El Tor phage V

b) Intrinsic resistance to polymyxin B

c) Voges-Proskauer test is positive

d) Not susceptible to group IV phage

Question 12:

You are a medical officer at a district hospital and receive 13 patients with rice-water diarrhea and dehydration. You suspect cholera and want to send the stool samples to the hospital lab. Which of the following media will you not use?

a) Venkatraman-Ramakrishnan medium

b) Cary-Blair medium

c) Monsur's GTTA medium

d) Nutrient agar

Question 13:

Sold by @itachibot

Which of the following bacteria shows Kanagawa phenomenon?

a) Vibrio cholerae

b) Vibrio parahaemolyticus

c) Vibrio vulnificus

d) Vibrio alginolyticus

Question 14:

An outbreak of food poisoning in a coastal city is determined to be due to contaminated seafood. The causative bacteria formed the following colonies on TCBS agar and showed hemolysis on Wagatsuma agar. Which of the following statements regarding the organism implicated here is false?



a) It is non-capsulated

b) It shows bipolar staining

c) It produces peritrichous flagella on solid medium

d) It gives a positive string test

Question 15:

Which of the following vibrios has the highest salt tolerance?

a) Vibrio mimicus

b) Vibrio parahaemolyticus

Sold by @itachibot

c) Vibrio vulnificus

d) Vibrio alginolyticus

Question 16:

A fisherman presented to the ER with a wound on his left leg. He tells you he had got a small cut a week back that got worse after his recent fishing trip. On examination, you note vesicles and bullae around the wound. Specimen taken from the bullae revealed lactose-fermenting colonies. What is the causative agent?

a) Vibrio cholerae

b) Vibrio parahaemolyticus

c) Vibrio vulnificus

d) Vibrio alginolyticus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

All of the following are correctly matched except:

a) Vibrio parahaemolyticus - Food poisoning

b) Vibrio alginolyticus - Infections of the eyes, ears and wounds

c) Vibrio vulnificus - Dysentery

d) Vibrio cholerae - Watery diarrhea

Question 18:

A 17-year-old presents to you with complaints of passing loose stools. Given below are the findings of his stool microscopy. Infection with which of the following bacteria would you consider in your differential diagnosis?

Sold by @itachibot



MARROW

a) Shigella dysenteriae

b) Campylobacter jejuni

c) Escherichia coli

d) Bacillus cereus

Sold by @Itachi... purchased this from some... else, you may have been scammed.

Question 19:

A young man presented with complaints of a tingling sensation in his hands and feet. He also complains of weakness in his legs that is slowly spreading to his upper body. He had an episode of bloody diarrhea 10 days ago. Which of the following organisms is the likely cause of this patient's condition?

a) Vibrio cholerae

b) Campylobacter jejuni

c) Leptospira interrogans

d) Helicobacter pylori

Question 20:

A young man presented with bloody diarrhoea, abdominal pain, and fever. Lab investigations showed leucocytosis and stool microscopy revealed RBCs and leucocytes. Darkfield microscopy showed a bacteria having darting motility. Which of the following is a selective culture medium for this organism?

a) Skirrow's media

Sold by @itachibot

b) PLET medium

c) BSK medium

d) BCYE medium

Question 21:

Identify the organism & the stain seen in the image given below.



a) Campylobacter jejuni; Gimenez stain

b) Helicobacter pylori; modified Giemsa stain

c) Treponema pallidum; Levaditi stain

d) Helicobacter pylori; Warthin-Starry stain

Question 22:

Which of the following organisms produce urease?

a) Helicobacter pylori

b) Salmonella

c) Escherichia coli

d) Shigella

Question 23:

Sold by @itachibot

You are a surgery resident evaluating a 45-year-old female patient with complaints of burning abdominal pain for the past 2 months that is relieved on taking food. You determine the disease to have a bacterial etiology. Which of the following is the most sensitive test to detect the causative organism here?

a) Culture

b) Serology

c) Urea breath test

d) Histology

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	b
5	b
6	b
7	d
8	c
9	c
10	a
11	a
12	d
13	b
14	a
15	d
16	c
17	c
18	b
19	b
20	a
21	d
22	a
23	c

If yo... ...chibot ...meone else,

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The given clinical vignette is suggestive of cholera caused by Vibrio cholerae. The image shows the characteristic fish in stream appearance of Vibrio.

Other options:

Options A and B: School of fish appearance and railroad appearance are shown by Haemophilus ducreyi.

Option D: Drumstick appearance is shown by Clostridium tetani due to its spherical and terminal spores.

Solution to Question 2:

The given microscopic finding shows the darting motility characteristic of Vibrio cholerae.

Vibrio cholerae is actively motile with a polar flagellum. The actively motile vibrios appear like a 'swarm of gnats'.

Solution to Question 3:

Vibrio cholerae are classified based on O antigen, under Sakazaki typing scheme.

Based on the presence or absence of O antigen, V. cholerae is classified as O1 which agglutinate with antisera to O1 group (agglutinable vibrios) and non-O1, which do not agglutinate with antisera to O1 group (non-agglutinable vibrios).

As per the Gardner and Venkatraman classification, Vibrios are classified into Group A (V. cholerae) and Group B (others) based on H antigen or flagellar antigen.

Gardner and Venkatraman classification:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 4:

Non-agglutinable vibrios are non-pathogenic (except O139) and hence, they are called non-cholera vibrios.

The vibrios which are not agglutinated by O-1 antiserum are called non-agglutinable (NAG) vibrios. They belong to the serogroups O2–O200 of which only serogroup O-139 is known to cause epidemics of cholera. They are similar to Vibrio cholerae in most biochemical reactions and culture characteristics, except that they are hemolytic to sheep erythrocytes and give a positive Voges-Proskauer reaction.

Solution to Question 5:

The test described here is the 'cholera red reaction' given by Vibrio cholera. When few drops of concentrated sulphuric acid are added to a 24-hour peptone water culture, a reddish-pink colour develops due to the formation of nitrosoindole. Vibrio cholerae reduce nitrates to nitrites and convert tryptophan to indole. These two properties contribute to the cholera red reaction.

Sold by @itachibot



Other options:

Option A: The methyl red (MR) test is used to detect the production of sufficient acid during the fermentation of glucose by bacteria. It is used to classify Enterobacteriacae.

Option C: Vibrio cholera does not produce any colored pigment.

Option D: Vibrio ferments carbohydrates leading to production of acid but no gas.

Solution to Question 6:

The given case scenario of passing profuse rice-water stools with similar complaints in the family suggests a probable diagnosis of cholera. The test shown in the above image is the string test and it is performed to identify Vibrio cholerae.

To perform the string test, a loopful of the bacterial culture is mixed with a drop of 0.5% sodium deoxycholate in saline on a slide. The bacterial cells of Vibrio cholerae are lysed by sodium deoxycholate, causing the suspension to lose its turbidity and become mucoid. A string is formed when the loop is drawn away from the suspension.

Note: The duodenal string test refers to the Entero test performed for the diagnosis of giardiasis.

Solution to Question 7:

Vibrio cholera is a facultative anaerobe and is not microaerophilic.

Solution to Question 8:

If you purchased this from some- else, you may have been scammed. Sold by @itachi

Sold by @itachibot

The disruption of zonula occludens is responsible for watery diarrhea in cholera caused by Vibrio cholerae.

Zonula occludens toxin (Zot) is an enterotoxin elaborated by Vibrio cholerae that increases intestinal permeability by interacting with a mammalian cell receptor with subsequent activation of intracellular signaling leading to the disassembly of the intercellular tight junctions.

Mechanism of action of cholera toxin:

• The cholera toxin molecule consists of 1 A and 5 B subunits. The B (binding) units attach to the GM1 ganglioside receptors on the surface of jejunal epithelial cells. The A (active) subunit, on being transported into the enterocyte, dissociates into A1 and A2. A2 fragment only links biologically active A1 to the B subunit.

• A1 causes prolonged activation of cellular adenylate cyclase and accumulation of cAMP, leading to outpouring into the small intestinal lumen, of large quantities of water and electrolytes. This leads to watery diarrhea which is isotonic with plasma but contains much more potassium and bicarbonate.

• The toxin also inhibits the intestinal absorption of sodium and chloride.

• All clinical manifestations and complications in cholera result from the massive water and electrolyte depletion thus caused.



Solution to Question 9:

The given clinical scenario of dehydration, profuse diarrhea, and rice water stools is suggestive of cholera. Cholera toxin causes continued activation of adenylate cyclase.

Mechanism of action of cholera toxin:

• The cholera toxin molecule consists of 1 A and 5 B subunits. The B (binding) units attach to the GM1 ganglioside receptors on the surface of jejunal epithelial cells. The A (active) subunit, on being transported into the enterocyte, dissociates into A1 and A2. A2 fragment only links biologically active A1 to the B subunit.

Sold by @itachibot

• A1 causes prolonged activation of cellular adenylate cyclase and accumulation of cAMP, leading to outpouring into the small intestinal lumen, of large quantities of water and electrolytes and the consequent watery diarrhea, that is isotonic with plasma but contains much more of potassium and bicarbonate.

• The toxin also inhibits the intestinal absorption of sodium and chloride.

• All clinical manifestations and complications in cholera result from the massive water and electrolyte depletion thus caused.



Solution to Question 10:

Vibrio cholerae possesses lipopolysaccharide (LPS) O antigen but it plays no role in the pathogenesis of cholera. It is responsible for the immunity induced by killed vaccines.

Factors involved in virulence of Vibrio cholerae:

• In the small intestine, vibrios can cross the protective layer of mucus and reach the epithelial cells by using mucinase and other proteolytic enzymes.

• Toxin co-regulated pilus(TCP), is essential for the adhesion to the epithelial surface and colonization of the bacteria in small intestine.

• Cholera toxin (also known as choleragen and cholera enterotoxin) causes watery diarrhea through cAMP-mediated pathway.

Solution to Question 11:

El Tor vibrios are susceptible to El Tor phage V.

Sold by @itachibot

Solution to Question 12:

Nutrient agar is not used to transport Vibrio cholerae. It is a simple media used to grow non-fastidious organisms, but not to transport them.

Transport media for Vibrio cholerae:

• Venkatraman-Ramakrishnan (VR) medium

• Cary-Blair medium

If the specimen can reach the laboratory in a few hours, it may also be transported in enrichment media. Examples include:

• Alkaline peptone water

• Monsur's medium.

If transport media are not available, strips of blotting paper may be soaked in the watery stool and sent to the laboratory packed in plastic envelopes.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 13:

Kanagawa phenomenon is shown by Vibrio parahaemolyticus. It refers to hemolysis occuring when the bacteria are inoculated on Wagatsuma agar.

Pathogenic strains of V. parahaemolyticus are hemolytic, while non-pathogenic strains are non-hemolytic. This phenomenon occurs due to a heat-stable hemolysin.

Solution to Question 14:

The given case scenario of an organism that caused food poisoning due to contaminated seafood in addition to the cultural characteristics is suggestive of food contamination by Vibrio parahaemolyticus. It is a capsulated organism.

V. parahaemolyticus is an enteropathogenic halophilic vibrio that is an important cause of food poisoning throughout the world. It inhabits the coastal seas, where it is found in shrimps, crabs, and oysters.

It forms opaque green colonies with a raised centre and flat translucent periphery on TCBS agar unlike V. cholera which forms yellow colonies. Pathogenic strains show hemolysis on Wagatsuma agar which is known as Kanagawa phenomenon. This can be used as a test for pathogenecity.

Other options:

Option B: It shows bipolar staining and has a tendency for pleomorphism, especially when grown on 3% salt agar.

Option C: Unlike other vibrios, it produces peritrichous flagella when grown on solid media. Polar flagella are formed in liquid cultures.

Option D: String test is positive like with V. cholera.

Sold by @itachibot

Solution to Question 15:

Vibrio with the highest salt tolerance of up to 10% NaCl is Vibrio alginolyticus.

The salt tolerances of vibrio species are as follows:

• V. cholerae optimally grows at 0.5–1% NaCl concentration. Its growth is inhibited by high concentration of >6% NaCl.

• V. mimicus grows best at low salt concentrations of 0.5-1% like V. cholerae.

• V. parahaemolyticus grows optimally salt concentration of 2-4 %. It has a salt tolerance up to 8%.

• V. vulnificus has a salt tolerance similar to V. parahaemolyticus.

• V. alginolyticus has the highest salt tolerance of up to 10% NaCl.

Solution to Question 16:

The given case scenario of contact with seawater of an existing wound leading to cellulitis by a lactose-fermenting bacteria suggests infection with Vibrio vulnificus.

V. vulnificus ferments lactose but not sucrose. It has a salt tolerance of up to 8%.

Infection with V. vulnificus can manifest in 2 ways:

• Soft-tissue infections can occur after contact with seawater and progress to cellulitis.

• Primary septicemia can occur in patients, typically with underlying liver disease, who have acquired the organism through the gastrointestinal tract after recent consumption of raw shellfish.

Note: V. cholerae, V. parahaemolyticus, and V. alginolyticus do not ferment lactose although lactose may be split very slowly by V. cholerae.

Solution to Question 17:

Vibrio vulnificus causes wound infection following contact of open wounds with sea water.

In immunocompromised hosts, and particularly those with liver disease, it penetrates the gut mucosa following ingestion. It may not cause any gastrointestinal manifestations and directly enters the bloodstream, leading to the rapid development of septicemia with high mortality.

Other options:

Option A: V. parahaemolyticus causes food poisoning associated with marine food. It also causes acute diarrhea, unassociated with food poisoning.

Option B: V. alginolyticus has been associated with infections of the eyes, ears, and wounds in human beings exposed to seawater.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option D: Cholera is caused by V. cholerae via its toxin but it does not cause dysentery which is characterized by bloody diarrhea. It is mainly associated with watery diarrhea.

Solution to Question 18:

The given clip shows darting motility which is seen in Vibrio cholerae and Campylobacter jejuni. Thus, among the given options, infection with C. jejuni should be considered as a differential diagnosis in this case.

Campylobacter jejuni possesses a single polar flagellum, hence it exhibits darting motility.

Shigella dysenteriae is a non-motile organism.

Solution to Question 19:

The given case scenario with ascending paralysis is suggestive of the development Guillain-Barré syndrome (GBS) following diarrhea. The organism that likely caused this patients condition is Campylobacter jejuni, which can cause diarrhea and trigger GBS.

Campylobacter is a slender, spirally curved, gram-negative bacillus. Growth occurs under microaerophilic conditions. The organism also shows darting motility. Asymptomatic campylobacter infection also may trigger Guillain-Barré syndrome.



Guillain-Barre syndrome is an acute, fulminant polyradiculoneuropathy that is autoimmune in nature. In C. jejuni infection, bacterial lipopolysaccharide triggers an autoimmune response against axons via molecular mimicry, hence causing GBS. Other organisms associated with GBS include cytomegalovirus, M. pneumonia, and Epstein-Barr virus.

If yo... Sold by @itachibot ...ed this from someone else, ...been scammed.

Sold by @itachibot

Solution to Question 20:

The given clinical scenario of fever, bloody diarrhea, abdominal pain and leucocytosis alongwith microscopy findings of RBCs and WBCs in fecal smear, and organism having darting motility are suggestive of infection with Campylobacter jejuni. The selective culture medium used for Campylobacter is Skirrow's media.

Other selective media for Campylobacter include:

• Butzler's medium

• Campy-BAP media

The transport medium used for Campylobacter is the Cary-Blair medium.

Solution to Question 21:

The given image shows Helicobacter pylori stained using Warthin-Starry stain.

H.pylori is a Gram-negative spiral rod that is motile by a unipolar tuft of lophotrichous flagella. The bacteria are present only in the mucus overlying the gastric mucosa and do not actually invade into the mucosa.

The given image shows Helicobacter pylori on gastric mucosa stained with Warthin-Starry silver stain.

Solution to Question 22:

Helicobacter pylori produces urease.

Urease-producing organisms: (Mnemonic-PUNCH)

• Proteus

Sold by @itachibot
If you bought this from someone else, it has been scanned.

Sold by @itachibot

- Ureaplasma urealyticum
- Nocardia
- Cryptococcus, Corynebacterium urealyticum
- H. pylori

Solution to Question 23:

The given case scenario is suggestive of a duodenal ulcer caused by Helicobacter pylori. The urea breath test is the most sensitive test for the diagnosis of Helicobacter pylori infection.

To perform this test, the patient drinks a urea solution labelled with radioactive carbon-14. The patient then blows into a tube. If H. pylori urease is present, the urea is hydrolyzed, and labelled carbon dioxide is detected in breath samples. The sensitivity of the urea breath test is >90%.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Pseudomonas and Burkholderiales

Question 1:

A 20 year old male presents with acute onset cough and thick sticky sputum. There is a history of repeated lung infections since childhood. Chest Xray shows evidence of bilateral upper lobe bronchiectasis. On evaluation, the chloride concentration in his sweat is >60 mEq/L. What is the organism majorly associated with this patient's respiratory disease?

a) Burkholderia pseudomallei

b) Pseudomonas putida

c) Pseudomonas aeruginosa

d) Burkholderia cepacia

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 2:

Which of the following organisms causes scombroid fish poisoning?

a) Salmonella

b) Staphylococcus

c) Pseudomonas aeruginosa

d) Peptostreptococcus

Question 3:

A patient comes to the dermatology OPD complaining of greenish discoloration of her nails. You diagnose her with green nail syndrome. Production of which of the following helps differentiate the organism implicated here from other members of its genus?

a) Pyocyanin

b) Pyoverdin

c) Pyorubin

d) Pyomelanin

Question 4:

Sold by @itachibot

The mechanism of action of exotoxin A of Pseudomonas resembles that of _____.

a) Cholera toxin

b) Diphtheria toxin

c) Tetanus toxin

d) Anthrax toxin

Question 5:

A patient with acute onset fever, cough and hemoptysis is diagnosed to have Whitmore's disease. Which of the following organisms causes this condition?

a) Burkholderia pseudomallei

b) Burkholderia mallei

c) Burkholderia cepacia

d) Paraburkholderia species

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A horse-groomer was diagnosed with ulcerative tracheobronchitis. Upon enquiry, it was discovered that one of his horses had recently died of pneumonia after developing a fever and subcutaneous nodules. Which of the following statements is false about the likely causative agent?

a) It is non-motile.

b) It is always oxidase negative

c) It is the causative agent for Glanders disease.

d) It induces Strauss reaction on inoculation into guinea pigs.

Question 7:

Which of the following is Burkholderia cepacia resistant to?

a) Ceftazidime

b) Trimethoprim-sulfamethoxazole

c) Temocillin

d) Cefotetan

Sold by @itachibot

Question 8:

An organism cultured in Bordet–Gengou medium shows smooth, refractile, glistening colonies resembling 'bisected pearls' or 'mercury drops'. The organism is most likely _____.

a) Bordetella parapertussis

b) Bordetella bronchiseptica

c) Bordetella pertussis

d) Bordetella petrii

Question 9:

An 8-year-old boy with incomplete vaccination history comes with vomiting after bouts of severe, violent cough. What is the mechanism of action of the toxin responsible for this child's disease?

a) Activation of phospholipase and degrades cell membrane

b) Inactivation of 60S ribosome

c) ADP ribosylation and lymphocytosis

d) Activation of guanylate cyclase

If you purchased this from someone else, you may have been scammed!
Sold by @itachibot

Question 10:

Which of the following statements is false about pertussis toxin?

a) It is expressed on the surface of the bacillus.

b) It cannot be toxoided.

c) "A" portion of A-B structure is the enzymatically active moiety.

d) It is similar to cholera toxin in its structure and function.

Question 11:

A 4-year-old unimmunized child presents with whooping cough. Which of the following constituents of the causative organism is responsible for facilitating its adhesion to the respiratory epithelium?

a) Adenylate cyclase

b) Lipopolysaccharide

Sold by @itachibot

c) Tracheal cytotoxin

d) Filamentous hemagglutinin

Question 12:

What is the incubation period of pertussis in humans?

a) <7 days

b) 3-5 days

c) 21-30 days

d) 7-14 days

Question 13:

You are evaluating a 6-year-old unimmunised girl with cough and abdominal pain. Her mother informs you that her daughter has bouts of violent coughing which frequently ends with a loud whoop and is accompanied by vomiting. She gives history of similar complaints in a classmate 2-3 weeks back. Which of the following drugs will you prescribe for this patient?

a) Erythromycin

b) Penicillin

c) Ciprofloxacin

d) Cotrimoxazole

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 14:

A young child with unknown immunization history is suspected to have whooping cough. Which of the following methods will you prefer to use to collect a specimen for culture?

a) Cough plate method

b) Per-oral swab

c) Per-nasal swab

d) Post-nasal swab

Question 15:

Sold by @itachibot

A pregnant woman comes to the ANC clinic for her 2nd dose of the Tdap vaccine. Which of the following constitutes the acellular component of this vaccine?

a) Pertactin, filamentous hemaglutinin, and cytotoxin

b) Endotoxin, fimbriae, and pili

c) Pertussis toxoid, cytotoxin, and pili

d) Pertussis toxoid, filamentous hemaglutinin, and fimbrial protein

Answer Key

Question No.	Correct Option
1	c
2	c
3	a
4	b
5	a
6	b
7	d
8	c
9	c
10	b
11	d
12	d
13	a
14	c
15	d

Detailed Explanations

Solution to Question 1:

Bilateral upper lobe bronchiectasis with a history of repeated lung infections since childhood and evelated sweat chloride concentration suggests a diagnosis of cystic fibrosis (CF). Pseudomonas aeruginosa is a common etiological agent associated with chronic airway infection in patients with cystic fibrosis (CF).

Sold by @itachibot

In cystic fibrosis, there is thick inspissated mucus and impaired mucociliary clearance. Due to these factors, lungs are often colonized in infancy and early childhood with Staphylococcus aureus and Haemophilus influenzae. In adulthood, they are replaced by P. aeruginosa. This results in chronic inflammation, which leads to airway modelling and bronchiectasis.

Note: Burkholderia cepacia complex is an opportunistic environmental pathogen. It causes fatal respiratory distress and septicemia in CF patients. This is termed as 'cepacia syndrome'.

Solution to Question 2:

Scombroid fish poisoning is associated with Pseudomonas aeruginosa.

It follows the consumption of both scombroid (mackerel-like) fish and nonscombroid fish that have not been refrigerated properly. The musculature of these dark- or red-fleshed fish undergoes decomposition by Morganella morganii, Escherichia coli, Proteus species, Klebsiella species, and Pseudomonas aeruginosa. There is consequent decarboxylation of the amino acid L-histidine to form histamine.

A second compound is thought to play a causative role in this intoxication because large doses of oral histamine do not reproduce the affliction. The toxin or toxins involved are heat stable and are not destroyed by cooking or freezing.

Symptoms develop within 15–90 min of ingestion. Most cases are mild, with tingling of lips and mouth, mild abdominal discomfort, and nausea. The more severe and commonly described presentation includes flushing, conjunctival hyperemia, pruritus, urticaria, and angioedema. Without treatment, the symptoms generally resolve within 8–12 h.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Green nail syndrome is caused by Pseudomonas aeruginosa. Production of pyocyanin helps differentiate it implicated here from other Pseudomonas species.

Pseudomonas aeruginosa is the only member of the Pseudomonas species that produces pyocyanin which is a bluish-green pigment. Other species of Pseudomonas also produce pigments but they cannot produce pyocyanin. Some of the other pigments produced by Pseudomonas species including P. aeruginosa are pyoverdin/flourescein (greenish-yellow pigment), pyorubin (red pigment) and pyomelanin (brown pigment).

Solution to Question 4:

The action of exotoxin A of Pseudomonas aeruginosa is similar to that of the diphtheria toxin.

Both exotoxin A and diphtheria toxin cause ADP ribosylation of EF-2 and inhibit protein synthesis. Exotoxin A is a single-chain polypeptide with A and B fragments. The B fragment binds the toxin while the A fragment inactivates EF-2 in the presence of nicotinamide adenine dinucleotide (NAD). This leads to the inhibition of polypeptide chain elongation and thus, protein

Sold by @itachibot

synthesis.



Solution to Question 5:

Burkholderia pseudomallei causes Whitmore's disease.

Melioidosis was first described by Whitmore and Krishnaswami. Whitmore isolated the bacillus, and hence, it is known as Whitmore bacillus and melioidosis is called Whitmore's disease. Melioidosis is similar in presentation to glanders disease (caused by Burkholderia mallei). Humans are infected tby contact with contaminated soil via skin abrasions.

Types of melioidosis:

• Acute - It causes acute pneumonia with upper lobe infiltrates or severe necrotizing disease. It can also mimic tuberculosis with fever, cough, hemoptysis. It may lead to septicemia.

• Chronic - It can lead to the development of multiple caseous or suppurative foci, with skin and subcutaneous abscesses. These can lead to lymphangitis and lymphadenopathy. It may involve bones and other internal organs.

The bacillus has a safety-pin appearance on methylene blue staining. Treatment is divided into two stages: an intensive 2 week phase of ceftazidime or a carbapenem followed by at least 12 weeks of oral cotrimoxazole to eradicate the organism and prevent relapse.

Solution to Question 6:

The development of ulcerative tracheobronchitis after contact with an infected horse is suggestive of glanders which is caused by Burkholderia mallei. It is an oxidase variable organism.

In this case, the horse likely had glanders, which presents with fever, pneumonia, and subcutaneous nodules called Farcy buds. Discharges from the respiratory tract and skin are highly infectious and can be transmitted to humans through close contact (occupational route). The

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

horse-groomer likely contracted the infection by the inhalational route and developed acute glanders, which is characterised by fever, mucopurulent nasal discharge and ulcerative necrosis of upper respiratory tract (ulcerative tracheobronchitis). It may lead to septicemia and has a high fatality rate. Percutaneous exposure causes local suppurative skin lesions with associated regional lymphadenopathy.

B. mallei is a non-motile Gram-negative bacillus causing Glanders disease. It is an aerobe and a facultative anaerobe. It is oxidase variable i.e. it is usually oxidase negative but may occasionally show a positive reaction. It is a biosafety level 3 pathogen and so must be handled only in specialised laboratories. When male guinea pigs are inoculated intraperitoneally with suspected infectious material, it causes Strauss reaction. B. mallei sets up a necrotizing inflammation in the scrotal sac within a few days and can be confirmed bacteriologically.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

Burkholderia cepacia is resistant to cefotetan.

B. cepacia complex strains are intrinsically resistant to aminoglycosides, polymixin, first and second-generation cephalosporins (e.g. cefotetan) and carboxypenicillins. This antimicrobial resistance is mediated by efflux pumps, antimicrobial degradation, modifying enzymes or altered membrane functions.

B. cepacia is a Gram-negative bacillus which can cause severe necrotizing pneumonia in patients with chronic granulomatous disease. It can lead to fatal septicemia in patients with cystic fibrosis.

Antimicrobial agents that are effective against B. cepacia complex include meropenem, ceftazidime, trimethoprim-sulfamethoxazole, piperacillin and temocillin.

Solution to Question 8:

The organism showing colonies resembling 'bisected pearls' or 'mercury drops' on Bordet-Gengou medium most likely is Bordetella pertussis.

Bordetella pertussis is a Gram-negative coccobacillus. It causes whooping cough or pertussis in children.

It can be isolated on the enriched medium Bordet-Genou medium (potato-blood-glycerol agar). After incubation for 48 to 72 hours, B. pertussis colonies are small, dome-shaped, smooth, opaque, viscid, greyish white, refractile, and glistening resembling 'bisected pearls' or 'mercury drops' and are surrounded by a hazy zone of hemolysis. If growth is confluent, an 'aluminium paint' appearance is seen. In culture films, Bordetella pertussis bacilli tend to be arranged in loose clumps, with clear spaces in between, giving a 'thumbprint' appearance.

B. parapertussis (option A) grows on nutrient agar with the production of a brown, diffusible pigment after two days. It also grows more rapidly than B.pertussis on charcoal blood agar.

Sold by @itachibot

Sold by @itachi... if you purchased this from some... else, you may have been scammed.

Solution to Question 9:

Post-tussive vomiting and bouts of cough in an unimmunized child points to a diagnosis of pertussis, caused by Bordetella pertussis. It secretes pertussis toxin (PT), which acts by ADP ribosylation and disrupts signal transduction of many cell types. This promotes lymphocytosis.

The pertussis toxin disrupts protein synthesis and causes necrosis of parts of the epithelium, leading to pseudomembrane formation. Pertussis toxin also promotes sensitization to histamine and enhances insulin secretion but these effects have not been demonstrable in patients. They are only seen in experimental animals.

B. pertussis also produces other toxins like:

• Adenylate cyclase toxin (ACT)- It acts by catalysing the production of cAMP by various types of cells and inhibiting phagocytosis.

• Heat labile toxin (HLT)- It is a cytoplasmic protein present in all bordetellae. It is dermonecrotic and lethal in mice. Its pathogenic role is not known.

• Tracheal cytotoxin (TCT)- It is low molecular weight peptidoglycan which induces ciliary damage in hamster trachea.

Solution to Question 10:

Pertussis toxin (PT) can be toxoided.

PT is one of the toxins produced by Bordetella pertussis. It is expressed on the surface of the bacillus and secreted into the surrounding medium. It has an A-B structure, the A portion being the enzymatically active moiety and B, the binding component. PT is similar to cholera toxin in its structure and function. PT has ADP-ribosylating activity and acts on inhibitory Gi ° subunit of G-protein coupled receptor (GPCR), whereas cholera toxin acts on stimulatory Gs ° subunit of

Sold by @itachibot

GPCR.

Pertussis toxoid is the major component of the acellular pertussis vaccine. A toxoid is a toxin that has lost its toxicity but retains its antigenicity.

Solution to Question 11:

Whooping cough is caused by Bordetella pertussis. Filamentous hemagglutinin (FHA) is responsible for facilitating the adhesion of Bordetella pertussis to the respiratory epithelium.

Bordetella pertussis produces 3 hemagglutinins: Pertussis toxin (PT), lipid factor and FHA. FHA is a surface protein and is readily shed. Along with fimbrae, it helps in the adhesion of B. pertussis to the cilia of the respiratory epithelium and to erythrocytes. FHA and PT also promote secondary infection by coating other bacteria such as Haemophilus influenzae and Pneumococci. They then assists their binding to respiratory epithelium. This phenomenon has been termed 'piracy of adhesins'.

Solution to Question 12:

The incubation period of pertussis in humans is 7-14 days.

Pertussis is caused by Bordetella pertussis. After the incubation period, the disease takes a protracted course comprising three stages:

• Catarrhal: It lasts 1-2 weeks and is characterized by coryza, lacrimation, mild cough, low-grade fever, and malaise.

• Paroxysmal: It lasts 2-4 weeks with the cough becoming more frequent and spasmodic. The cough comes in repetitive bursts of 5–10 in number, often within a single expiration. Post-tussive vomiting is frequent. The episode may be terminated by an audible whoop, which occurs upon rapid inspiration against a closed glottis at the end of a paroxysm.

• Convalescent: It lasts 1–3 months and is characterized by gradual resolution of coughing episodes.

Whooping cough is also known as '100-day cough'. For 6–12 months following the illness, intercurrent viral infections may be associated with a recrudescence of paroxysmal cough.

Solution to Question 13:

The given case scenario of whooping cough, post-tussive vomiting, and abdominal pain, along with positive history of exposure in an unimmunised child is suggestive of pertussis. Erythromycin or one of the newer macrolides such as azithromycin or clarithromycin is the drug of choice for the treatment of this condition.

Antibiotic therapy in pertussis is used to eradicate the bacteria from the nasopharynx. The treatment is beneficial if it is started within 10 days of the disease. Macrolides are the drugs of

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

choice. Trimethoprim-sulfamethoxazole (Cotrimoxazole) is recommended as an alternative for individuals allergic to macrolides. Macrolides may also be used in the chemoprophylaxis of pertussis. Post-exposure prophylaxis is given to household close contacts, and high-risk contacts (infants, immunocompromised, pregnancy).

Solution to Question 14:

Whooping cough is caused by Bordotella pertussis. The most effective method for collecting a specimen for the culture of this organism is the per-nasal or nasopharyngeal swab method.

Respiratory specimens may be collected for the culture (gold standard) of Bordetella pertussis by three different methods:

• Per-nasal swab: A Dacron or rayon tipped swab on a flexible nichrome wire is passed along the floor of the nasal cavity and material is collected from the pharyngeal wall. This specimen is the most effective and best suited for diagnostic procedures.

• Cough plate method: A culture plate is held about 10 to 15 cms in front of the patient's mouth during a bout of cough.

• Post-nasal (peroral) method: Secretions from the posterior pharyngeal wall are collected with a cotton swab on a bent wire passed through the mouth.

Other diagnostic tests for pertussis include direct fluorescent aantibody test and PCR on nasopharyngeal specimens. Enzyme immunoassays of antibodies after exposure to B. pertussis can also be used for patients between 2-4 weeks of illness.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 15:

Tdap vaccine is the tetanus, diphtheria and acellular pertussis vaccine. The component of the acellular pertussis vaccine is inactivated pertussis toxin (PT) and it may contain one or more bacterial components such as pertactin, filamentous hemagglutinin, and fimbrial protein.

Pertussis vaccine is usually administered in combination with the toxoids of diphtheria and tetanus (DTaP). Five doses of pertussis vaccine are recommended - at 2, 4, 6 and 15–18 months of age followed by a booster dose at 4–6 years of age.

Tdap is now also recommended for pregnant women in between 27 weeks and 36 weeks of gestation by the ACOG. This is because infants between the age of 0-2 months are vulnerable to pertussis as they do not begin their vaccine schedule before 6 weeks of age. Thus to protect these infants, placental transfer of maternal pertussis antibodies has proven to be the most effective approach. Children under 4 years who are contacts of patients should receive a booster even if they had been previously immunized. Non-immunized contacts should receive full series of vaccination and also chemoprophylaxis with erythromycin for 10 days after contact with the patient has ceased.

Sold by @itachibot

Haemophilus

Question 1:

A woman presents with fever with chills and a productive cough. Chest X-ray is suggestive of lobar pneumonia. She had undergone splenectomy 2 years back. Sputum culture on blood agar shows the following. What is the causative agent?

a) Hemophilus influenzae

b) Klebsiella granulomatis

c) Streptococcus pneumoniae

d) Neisseria meningitidis

Question 2:

A patient presented with fever, seizures, and altered sensorium. Gram stain of the CSF is given below. Satellitism was seen on blood agar. Which of the following statements is false about this disease?

Sold by @itachibot



a) This is the most serious disease produced by this organism.

b) The disease is more common in young adults.

c) The bacilli reach the meninges from the nasopharynx through the bloodstream.

d) Cefotaxime is the drug of choice

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following is not caused by non-typable H. influenzae?

a) Meningitis

b) Otitis media

c) Sinusitis

d) Puerperal sepsis

Question 4:

All of the following Hemophilus species require both Factors X and V except_____.

a) H. hemolyticus

b) H. influenzae

c) H. parahemolyticus

d) H. aegypticus

Sold by @itachibot

Question 5:

A 7-year-old girl presents to the emergency with acute stridor. The mother gives a history of fever, drooling, and breathing difficulty since that morning. An X-ray of her neck appears as given below. What is the most likely causative agent?



a) Hemophilus parainfluenzae

b) Streptococcus pneumoniae

c) Hemophilus influenzae

d) Parainfluenza virus

If you purchased this ... you may have been ... else, ... d.

Question 6:

Pneumonia due to a gram-negative pleomorphic bacillus growing on chocolate agar which is catalase and oxidase-positive is:

a) E.coli

b) Pseudomonas

c) Haemophilus influenzae

d) Actinomyces

Question 7:

A middle-aged man who is a known case of diabetes and chronic bronchitis presents to the OPD with low-grade fever and productive cough for 5 days. Which of the following is not a

Sold by @itachibot

virulence factor of the likely causative organism?

a) Outer membrane protein (OMP)

b) Lipooligosaccharide (LOS)

c) Capsular polysaccharide

d) Pili

Question 8:

Which of the following bacteria does not belong to the HACEK group?

a) Haemophilus parainfluenza

b) Aggregatibacter actinomycetemcomitans

c) Escherichia coli

d) Kingella kingae

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 9:

A 26-year-old man presented with painful, bleeding genital ulcers. On examination, tender inguinal lymphadenopathy is present. Scraping from the ulcer base was taken and the microscopic appearance of the Gram-stained smear is shown below. What is the causative organism?



a) Haemophilus ducreyi

b) Treponema pallidum

Sold by @itachibot

c) Chlamydia trachomatis serovar L1

d) Chlamydia trachomatis serovar L2

Question 10:

The following are true about chancroid:

a) C & D only

b) A, B,C & D

c) A B & C

d) A & B only

Question 11:

Which of the following is not involved in urethritis?

a) Trichomonas

b) Chlamydia

c) Gonococcus

d) H.ducreyi

Question 12:

A 5-year-old girl presents with conjunctivitis, fever, and hemorrhagic skin lesions. There's a history of travel to Brazil. Which of the following etiological agents would you suspect in this case?

a) Haemophilus influenzae type A

b) Haemophilus parahaemolyticus

c) Haemophilus aegypticus

d) Haemophilus parainfluenzae

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option

Sold by @itachibot

1	a
2	b
3	a
4	c
5	c
6	c
7	c
8	c
9	a
10	c
11	d
12	c

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Detailed Explanations

Solution to Question 1:

The given scenario of bacterial pneumonia in a splenectomised individual along with the image showing satellitism on blood agar points towards an infection due to Hemophilus influenzae.

Pneumonia can occur as a part of overwhelming post-splenectomy infection (OPSI). This is a fatal post-splenectomy complication occuring due to infections with encapsulated organisms such as S. pneumoniae, H. influenzae, and N. meningitidis. It can be prevented by vaccinating against these organisms within 2 weeks of undergoing splenectomy.

Satellitism is seen when S. aureus is streaked across a plate of blood agar on which H. influenzae has been inoculated. After overnight incubation, the colonies of H. influenzae will be large and well developed alongside the streak of Staphylococcus and smaller when they are farther away from the streak. This is due to the release of factor V by lysis of RBCs mediated by S. aureus. Factor V is required for the growth of H. influenzae.

The given image shows satellitism on blood agar

Sold by @itachibot

Sold by @itachi... from some... else,
If you purchased this ... you may have been scammed.

Solution to Question 2:

The given scenario and the image showing gram-negative coccobacilli in the CSF are suggestive of hemophilus meningitis caused by H.influenzae. Hemophilus meningitis is more common in children between 3 months and 3 years of age and not common in young adults.

Other options:

Option A: Meningitis is the most serious disease caused by H. influenzae. The case fatality rate is 90% if untreated.

Option C: H. influenze is transmitted via respiratory droplets. Thus, it is introduced into the nasopharynx of the contacts of an infected individual. It has to enter the bloodstream to reach the meninges.

Option D: Cefotaxime or ceftazidime are the drugs of choice for the treatment of hemophilus meningitis.

Solution to Question 3:

Meningitis is caused by typable H. influenzae.

Non-typable H. influenzae cause:

- Otitis media
- Sinusitis
- Puerperal sepsis
- Exacerbations of COPD.

Sold by @itachibot

H. influenzae is typed into 6 serotypes (a to f) based on the capsular polysaccharide. Serotype b (Hib) is the most virulent and accounts for most of the invasive infections.

The differences between type b strains and non-typable strains of H. influenzae are given in the table below:

Feature	Type B strains	Nontypable strains
Capsule	Made of Ribosyl-ribitol phos phate	Unencapsulated
Pathogenesis	Invasive infections due to he matogenous spread	Mucosal infections due to co ntagious spread
Clinical manifestations	Meningitis and invasive infec tions incompletely immunize d infants and childrenEpiglot titisCellulitisPneumonia	Otitis media in childrenExace rbation of COPD,Puerperal s epsisSinusitis
Vaccine	Highly effective conjugate va ccines	Not available

Sold by @ita... If you purchased this from so... you may have been scammed

Solution to Question 4:

Hemophilus parahemolyticus requires only factor V for its growth. It does not require factors X.

Hemophilus species requiring both factors X and V: (Mnemonic to remember: HAI)

- H. hemolyticus
- H. aegypticus
- H. influenzae

Factor V- It refers to the co-enzyme, NAD. It is heat-labile (destroyed at 120°C in a few minutes) and present inside RBC.

Factor X- It refers to Hemin. It is heat stable and is present in the blood.

Species	X factor requirement	V factor requirement
H. influenzae	+	+
H. aegypticus	+	+
H. ducreyi	+	-
H. parainfluenza	–	+
H. haemolyticus	+	+
H. parahaemolyticus	–	+

Sold by @itachibot

Solution to Question 5:

The given clinical vignette and 'thumb sign' on the X-ray are suggestive of acute epiglottitis caused by Hemophilus influenzae.

It is an acute inflammation of the epiglottis causing obstructive laryngitis and presents with high fever, severe sore throat, tachycardia, systemic toxicity, and drooling as patients are unable to swallow. The patient characteristically leans forward while sitting for ease of breathing.

The diagnosis often is made on clinical grounds. On direct fiberoptic laryngoscopy, the typical edematous cherry-red epiglottis is visualized. Neck radiographs reveal an enlarged edematous epiglottis (Thumb sign). The below image shows the Thumb sign.

In some countries, widespread vaccination against Hib has drastically decreased acute epiglottitis cases in children by over 90%. However, the incidence in adults has remained largely unchanged since the Hib vaccine was introduced. Despite vaccination efforts, many pediatric cases of epiglottitis today are still caused by Hib due to either lack of vaccination or vaccine ineffectiveness. In adults and increasingly in children, other bacterial pathogens, particularly group A Streptococcus, have been linked to cases of epiglottitis.

Treatment includes airway management followed by administration of antibiotics like ampicillin/sulbactam, cefotaxime, or ceftriaxone.



Solution to Question 6:

Pneumonia due to a gram-negative pleomorphic bacillus growing on chocolate agar which is catalase and oxidase-positive is most likely Haemophilus influenza.

Haemophilus pneumonia typically occurs in infants and is accompanied by empyema and sometimes meningitis as well. In older children and adults, the picture is of lobar pneumonia. Sputum is collected for the detection and isolation of Hemophilus. As the bacillus is very sensitive to low temperatures, specimens should never be refrigerated before inoculation. H.infiuenzae is a Gram-negative, non-motile, non-sporing bacillus, exhibiting considerable

Sold by @itachibot

pleomorphism. In sputum, it usually occurs as clusters of coccobacillary forms.

The bacillus has fastidious growth requirements. The accessory growth factors, named X and V, present in the blood are essential for growth. Catalase and

oxidase reactions are positive. Nitrates are reduced to nitrites. Biotyping has been done on the basis of indole production.

Cultures may be preserved for about a month on chocolate agar slopes. Chocolate agar is made by heating blood agar to 80-90°C for 15-20 minutes. This causes the release of V factor from within the erythrocytes.

Treatment- Amoxycillin-clavulanate or clarithromycin is more effective in treating respiratory infections by Haemophilus influenzae.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed

Solution to Question 7:

The given clinical scenario describes an acute exacerbation of COPD characterized by low-grade fever, increased cough, sputum production, and shortness of breath. The most common cause of these are non-typable H. influenzae which lack a capsule and so capsular antigen cannot be a virulence factor.

All H. influenzae have the following virulence factors:

• Outer membrane proteins (OMP)

• Lipooligosaccharide(LOS)

• Pili

In addition to the above factors, H. influenzae type b also has its capsular polysaccharide as a virulence factor.

The differences between Hib and non-typable strains are given below:

Feature	Type b strains	Nontypable strains
Capsule	Made of Ribosyl-ribitol phos phate	Unencapsulated
Pathogenesis	Invasive infections due to he matogenous spread	Mucosal infections due to co ntagious spread
Clinical manifestations	Meningitis and invasive infec tions incompletely immunize d infants and childrenEpiglot titisCellulitisPneumonia	Otitis media in childrenExace rbation of COPD,Puerperal s epsisSinusitis
Vaccine	Highly effective conjugate va ccines	Not available

Solution to Question 8:

Sold by @itachibot

Escherichia coli does not belong to the HACEK group.

The acronym HACEK refers to a group of fastidious slow-growing bacteria, normally residing in the mouth. These can sometimes cause severe infections, particularly culture-negative subacute bacterial endocarditis.

The group includes:

• Haemophilus species (H. parainfluenzae, H. aphrophilus, H. paraphrophilus)

• Aggregatibacter actinomycetemcomitans (previously called as Actinobacillus)

• Cardiobacterium hominis

• Eikenella corrodens

• Kingella kingae

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

The image shows the' school of fish' or 'railroad track' appearance of Haemophilus ducreyi. This appearance is seen when the bacilli are arranged in small groups, whorls or in chains. The given case scenario is suggestive of chancroid.

Chancroid is characterized by non-indurated, tender, irregular ulcers that may bleed on touch. The neighbouring lymph nodes are enlarged, painful, and tender.

Other options:

Option B - Treponema pallidum causes syphilis, characterised by chancre - a single painless ulcer with a clean base and a firm raised border, with painless lymphadenopathy. Persistent or multiple chancres may be seen in HIV-infected or other immunodeficient patients.

Options C and D - Chlamydia trachomatis serovars L1, L2 and L3 cause lymphogranuloma venereum, characterised by shallow, painless ulcers that progress to painful lymphadenopathy.

Solution to Question 10:

Chancroid is a painful genital ulcer caused by Haemophilus ducreyi, which is a gram-negative coccobacillus. Microscopic appearance of H. ducreyi is called the school of fish appearance or railroad appearance. Chancroid ulcers have a soft base that bleeds easily to touch. They also cause painful lymphadenopathy.

The image below shows the 'school-of-fish' appearance characteristic of H. ducreyi.

Sold by @itachibot



Note: Groove sign is a characteristic sign seen in lymphogranuloma venereum. It refers to the depression created by the inguinal ligament as it passes between the groups of inflamed inguinal lymph nodes.

Sold by @itachi... If you purchased this from someone else, you may have been scammed

Solution to Question 11:

Haemophilus ducreyi does not cause urethritis. It causes chancroid (soft chancre), which presents as a genital ulcer.

Urethritis is classified into two types:

- Gonococcal urethritis - caused by Neisseria gonorrhoea.
- Non-gonococcal urethritis - caused by:
- Chlamydia trachomatis (most common)
- Ureaplasma urealyticum
- Adenovirus
- Haemophilus vaginalis
- Mycoplasma genitalium
- Tricomonas vaginalis (rare)
- Herpes simplex virus (rare)

Solution to Question 12:

The given case scenario is suggestive of Brazilian purpuric fever (BPF) caused by Haemophilus aegypticus (Koch-Weeks bacillus).

Sold by @itachibot

Recent DNA studies have shown that the bacillus is identical to non-capsulated H.influenzae and it is now named H.influenzae biotype aegypticus.

It is worldwide in distribution and causes:

• A highly contagious form of conjunctivitis ('pink eye') which is especially common in the tropics and subtropics and may lead to epidemics. It responds to local sulphonamide or gentamicin administration.

• Brazilian purpuric fever characterized by fever, purpura, and conjunctivitis which proceeds to fulminant septicemia in infants and children. It has a high fatality rate.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Miscellaneous Bacteria - Yersinia, Brucella, Bartonella, Legionella

Question 1:

The milk ring test is found to be positive for a sample of milk. Which organism has it most likely been contaminated with?

a) Brucella

b) Bordetella

c) Bacillus

d) Burkholderia

Question 2:

Which of the following is an intracellular pathogen?

a) Brucella suis

b) Haemophilus influenzae

c) Pseudomonas aeruginosa

d) Staphylococcus aureus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A farmer was diagnosed with brucellosis. Which of the following patterns of fever will be seen in this disease?

a) Pel-Ebstein fever

b) Undulant fever

c) Relapsing fever

d) Saddleback fever

Question 4:

A 56-year-old dairy farmer presented with remittent fever and night sweats for 1 month. On examination, he was hypotensive and pallor was present. Rose Bengal card tests done on milk

Sold by @itachibot

samples from his farm were found to be positive. Which blood culture method with a biphasic medium will be used to isolate the organism?

a) BACTEC method

b) Castaneda method

c) Septi-check method

d) Oxoid signal method

Question 5:

A 50-year-old farmer presents with fever, weight loss, and lower backache for three weeks. Blood reports show predominant lymphocytosis. What are the drugs of choice for treating this condition?

a) Cotrimoxazole and rifampicin

b) Doxycycline and cotrimoxazole

c) Doxycycline and streptomycin

d) Streptomycin and rifampicin

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

A 34-year-old man presents to the hospital with fever, joint pain and a rash on his hands. He gives a history of a rat bite 1 week back. Which of the following can cause this disease?

a) 1 and 3.

b) 2 and 4.

c) 3 and 5.

d) 2 and 5.

Question 7:

While evaluating the Pap smear of a patient, you come across the following finding. What is the diagnosis?

Sold by @itachibot



a) Lymphogranuloma venereum

b) Bacterial vaginosis

c) Trichomoniasis

d) Gonorrhea

Sold by @itachi... purchased this from some... else, ...u may have been scammed.

Question 8:

A patient from Kolar, Karnataka presented with fever and hemoptysis. The findings of his peripheral smear examination are given below. Which of the following is the causative organism?



a) Yersinia pestis

Sold by @itachibot

b) Bacillus anthracis

c) Bordetella pertussis

d) Klebsiella pneumonia

Question 9:

A child from Uttarakhand suddenly developed fever with chills and malaise. He also developed a tender swelling which is shown in the image. There's a history of similar complaints in the community. What is false about the disease he's likely suffering from?

a) Wild rodents and their infected fleas are the primary reservoir.

b) The septicemic form cannot be acquired directly.

c) The pneumonic form is far more contagious than the bubonic form.

d) The bubonic form may progress to the pneumonic form.

Question 10:

A sudden epidemic of pneumonic plague has occurred in the Beed-Latur belt of Maharashtra. The primary contact of a suspected case came to you for preventive treatment. What will you prescribe for chemoprophylaxis in this patient?

a) Penicillin

b) Rifampicin

c) Erythromycin

d) Doxycycline

Sold by @itachibot

Question 11:

Which of the following organisms do not cause yersiniosis?

a) Yersinia enterocolotica

b) Yersinia pestis

c) Yersinia enterocolitica and Yersinia pseudotuberculosis

d) Yersinia pestis and Yersinia pseudotuberculosis

Question 12:

A 65-year-old patient presents to the OPD with high fever, chest pain, dyspnoea, and hemoptysis. The patient also complains of watery diarrhea. The direct fluorescent antibody stained film of his pleural fluid is given below. Identify the most likely etiologic agent.



a) Chlamydia pneumoniae

b) Legionella pneumophila

c) Hemophilus influenzae

d) Klebsiella pneumoniae

Question 13:

BCYE medium is used to culture:

Sold by @itachibot

a) Leptospira

b) Bacillus cereus

c) Campylobacter

d) Legionella

Question 14:

A 60-year-old man is diagnosed with Legionnaires' disease. He had recently attended a conference. What is the most likely way he acquired the infection?

a) From the air-conditioned rooms at the conference

b) From a chronic carrier of the disease

c) From sharing a towel with a fellow colleague in the conference

d) From drinking the water provided at the conference

Question 15:

A patient presented with fever and regional lymphadenopathy, following a scratch from his pet kitten. His lymph node biopsy specimen was examined with Warthin-Starry staining; the findings are shown in the given image. What is the causative organism of this condition?



a) Bartonella henselae

b) Bartonella bacilliformis

c) Bartonella quintana

If yo Sold by @itachibot rchased this from someone else, have been scammed.

Sold by @itachibot

d) Borrelia burgdorferi

Question 16:

Choose the correct statement regarding 5-day fever.

a) Transmitted by Phlebotomus sandflies.

b) Incubation period is 5 days

c) No animal reservoir is known

d) Treatment is necessary only in the severe cases

Question 17:

A young man presented to the ER with fever and suppurative lymphadenopathy of his left axilla a few days after he sustained a scratch from his pet cat. Which of the following will most likely be seen on biopsy of the lymph node?

a) Foreign body granuloma

b) Naked granuloma

c) Durck granuloma

d) Stellate granuloma

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A patient who is a known case of AIDS presents to the OPD with fever, chills, weight loss and the cutaneous lesions shown below. The microscopic examination of the lesions revealed predominant neutrophilic infiltration. What is the most likely diagnosis?

Sold by @itachibot

a) Kaposi sarcoma

b) Verruga peruana

c) Bacillary angiomatosis

d) Eczema herpeticum

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 19:

A patient has presented with lesions on the dorsum of the hand followed by axillary lymphadenopathy. What significant history will you ask for in this case?

a) History of exposure to rats

b) History of exposure to cats

c) History of similar complaint in family members

d) History of tick bite

Question 20:

Which of the following is incorrectly matched?

a) Bartonella henselae- Bacillary angiomatosis

b) Ehrlichia chaffeensis - Human monocytic ehrlichiosis

c) Ehrlichia phagocytophila - Human granulocytic ehrlichiosis

d) Bartonella bacilliformis - Parinaud's oculoglandular syndrome

Sold by @itachibot

Question 21:

A patient with fever and hemolytic anemia developed characteristic Peruvian warts 1 month later. What is the causative organism for this disease?

a) Bartonella hensalae

b) Bartonella quintana

c) Bartonella bacilliformis

d) Bartonella australis

If yo...rchibot ...meone else,

Answer Key

Question No.	Correct Option
1	a
2	a
3	b
4	b
5	c
6	d
7	b
8	a
9	b
10	d
11	b
12	b
13	d
14	a
15	a
16	c
17	d
18	c
19	b
20	d
21	c

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Milk ring test (MRT) is positive for Brucella. It is used for the detection of infected animals in dairy farms.

A sample of whole milk is mixed with a drop of Brucella antigen and incubated at 70 °C for 40–50 min. If antibodies are present in the milk, the bacilli are agglutinated and rise with the cream to form a blue ring at the top, leaving the rest of the milk unstained (MRT positive). If antibodies are absent, the solution remains uniformly blue (MRT negative).

Solution to Question 2:

Brucella suis is primarily an intracellular pathogen affecting reticuloendothelial system. This accounts for its refractoriness to chemotherapy.

Solution to Question 3:

Brucellosis is a bacterial zoonosis caused by Brucella sp which presents with undulant fever. It is referred to as undulant fever because of the rising and falling of fever like a wave (remittent fever). Patients may also have arthralgia, muscular pain, sweating, loss of weight, and appetite. CNS symptoms such as fatigue, irritability, and depression might also occur.

Other options

Option A: Pel-Ebstein fever is seen in Hodgkin's lymphoma. It is characterized by fever lasting 3–10 days followed by afebrile periods of 3–10 days.

Sold by @itachibot

Option C: Relapsing fever is seen in infection with Borrelia recurrentis in which days of fever are followed by a several-day afebrile period and then relapse into additional days of fever. A similar picture is also seen in malaria.

Option D: Saddleback fever is seen in dengue. Saddleback fever is defined as temperature >37.5°C with defervescence of at least one day, followed by a second peak lasting at least one day.

Solution to Question 4:

The given case scenario of remittent fever with anemia with a positive Rose Bengal card test is suggestive of brucellosis. Castaneda method for blood culture is used to isolate Brucella sp.

In the Castaneda method, blood is inoculated into a bottle of trypticase soy broth or brucella broth in a biphasic blood culture bottle (with liquid and solid media in the same bottle). This method minimizes materials and manipulation, thus reducing the chances of contamination and risk of infection to laboratory workers.

Blood cultures are positive in only 30–50% of cases even when repeated samples are tested. B.melitensis and B.suis are isolated more readily than B.abortus. The below image shows Castaneda's bottle.

Lab diagnosis of brucellosis is done using the following methods:

- Blood culture with
- Castaneda's method - recommended
- BACTEC - an automated method. It is faster than Castaneda's method.
- Serological tests:
- Standard agglutination test (SAT)
- ELISA for specific IgG/IgM

Sold by @itachibot

• Dipstick assays

• Detection in milk and infected cattle is carried out by:

• Microscopic demonstration by staining or immunofluorescence

• Milk ring test

• Rose Bengal card test

Solution to Question 5:

The given scenario of a farmer with fever, weightloss and back pain with a predominant lymphocytosis points towards a diagnosis of acute brucellosis. The drugs of choice for this condition in adults are intramuscular streptomycin with doxycycline.

The usual alternative regimen (and the current World Health Organization recommendation) is rifampin plus doxycycline. However, this is associated with a higher relapse rate, possibly because rifampicin increases clearance of doxycycline and reduces serum doxycycline levels.

In children, cotrimoxazole along with rifampicin or gentamycin may be used for the treatment of brucellosis.

Note: B. melitensis is endemic to the Middle East.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

The given case scenario is suggestive of rat-bite fever. This can be caused by Streptobacillus moniliformis and Spirillum minus.

Rat-bite fever (RBF) is characterized by fever, rash, and arthralgia which usually occurs days or weeks after a bite.

The two types of rat-bite fever are:

• Streptobacillary RBF develops 2–10 days after a rat bite, associated with sudden onset of fever, headache, and myalgia followed by petechial rash and arthritis. It is also called Haverhill fever or Erythema arthrititicum epidemicum.

• Spirillary RBF is similar to streptobacillary type but it is also associated with regional lymphadenopathy. It is also called Sodoku.

Diagnosis is based on demonstration of causative organisms in peripheral blood films. Serology can also be used for diagnosis.

RBF is treated with intravenous Penicillin G. Alternative drugs include ampicillin, cefuroxime, and cefotaxime.

Note: Pasteurella multocida causes cellulitis and lymphangitis following bites by cats and dogs.

Bartonella henselae causes cat sratch disease and bacillary angiomatosis.

Borellia burgdorferi causes Lyme disease.

Sold by @itachibot

Solution to Question 7:

The given image shows clue cells which are seen in bacterial vaginosis. They are epithelial cells with their surface covered by adherent bacilli seen in stained or unstained films.

Bacterial vaginosis is a polymicrobial infection characterized by a malodorous (fishy smell) vaginal discharge. Causative organisms include:

• Gardnerella vaginalis

• Mycoplasma hominis

• Prevotella

• Ureaplasma

• Peptostreptococcus.

Bacterial vaginosis is diagnosed using Amsel's criteria and Nugent's score:

1. Amsel's Criteria: (any 3 should be present)

• Thin, white/yellow homogenous vaginal discharge

• pH of Vaginal fluid>4.5

• Fishy odour of discharge accentuated on adding 10% KOH (Whiff test)

• Clue cells seen on microscopy

2. Nugent's score is obtained by counting the number of Gardnerella vaginalis, Mobilincus and lactobacillus in the gram-stained smear of vaginal discharge. A score of ≥7 is diagnostic

Treatment is oral metronidazole 500 mg BD for 7 days.

The images below show clue cells from microscopy.



If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot



Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Solution to Question 8:

The given case scenario and the image showing the characteristic "safety-pin appearance" in peripheral smear is suggestive of plague caused by Yersinia pestis.

On staining with Giemsa stain or methylene blue (Wayson's stain), the bacteria has bipolar staining with two ends densely stained and the central area clear. This is the characteristic 'safety-pin appearance' seen in Y. pestis.

Bipolar staining is also shown by the following organisms:

- Haemophilus ducreyi
- Vibrio parahemolyticus
- Calymmatobacterium granulomatis
- Burkholderia mallei
- Burkholderia pseudomallei.

Note: In India, 4 foci of plague are known:

- Kolar at the trijunction of Tamil Nadu, Andhra Pradesh, and Karnataka.
- Beed-Latur belt in Maharashtra
- Rhoru, Himachal Pradesh
- Uttarakhand

The image given below shows the safety-pin appearance of Yersinia pestis on Wayson's stain

Sold by @itachibot



Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

The given case scenario of sudden-onset fever with painful lymphadenopathy and a history of similar complaints in the community is suggestive of a diagnosis of plague caused by Yersinia pestis. The septicemic form of the disease may be acquired directly if the bacteria enter the bloodstream via infected blood transfusions.

Wild rodents and their infected fleas are the primary reservoirs of Yersinia pestis. Infection is transmitted to humans by the bite of infected fleas.

Human plague occurs in 3 clinical forms (in increasing order of severity): bubonic plague< pneumonic plague< septicemic plague

Bubonic plague- It is the most common variety of the disease. It is named after the buboes meaning swollen lymph nodes, which typically develop within a week after an infected flea bite. Other clinical features include sudden onset of fever and chills, headache, and fatigue.

Pneumonic plague- Bubonic plague can advance and spread to the lungs. It is the most virulent form of plague. Any person with pneumonic plague may transmit the disease via droplets to other humans. The symptoms include dyspnoea, chest pain, cough, fever, and hemoptysis.

Septicemic plague- When the bacteria enter the bloodstream directly and multiply there, it is known as septicemic plague. When left untreated, both bubonic and pneumonic plague can lead to septicemic plague.

Solution to Question 10:

The drug of choice for chemoprophylaxis of plague is doxycycline. It is given for 7 days to all close contacts of a patient having pneumonic plague.

Sold by @itachibot

Ciprofloxacin, levofloxacin, and moxifloxacin are the other first-line drugs used for antimicrobial prophylaxis.

The drug of choice for the treatment of plague is streptomycin. Streptomycin is nephrotoxic and ototoxic and should, therefore be cautiously used. Doxycycline or ciprofloxacin (and other fluoroquinolones) are considered as alternative drugs for the treatment of plague.

Solution to Question 11:

The term yersiniosis denotes infections with yersiniae other than Y.pestis.

Yersiniosis includes zoonotic infections by Y.pseudotuberculosis and Y.enterocolitica.

Solution to Question 12:

The given clinical vignette and the image showing fluorescent organisms in the direct fluorescent antibody (DFA) stained film of pleural fluid suggest the diagnosis of Legionnaire's disease, caused by Legionella pneumophila.

Two distinct clinical patterns of legionellosis have been identified:

• Legionnaires' disease presents with fever, non-productive cough, and dyspnea rapidly progressing to pneumonia if left untreated. Patients can also develop a hyponatremia. Diarrhea and encephalopathy are also common. It is often included in the differential diagnosis of atypical pneumonia. Atypical pneumonia is also caused by:

• Chlamydia pneumonia

• Chlamydia psittaci

• Mycoplasma pneumonia

• Coxiella burnetii

• Viruses such as influenza virus, adenovirus and respiratory syncytial virus

• Pontiac fever is a milder, non-fatal, influenza-like illness with fever, chills, myalgia, and headache.

Other options:

Option A: Chlamydia pneumonia causes pharyngitis, sinusitis, bronchitis, and pneumonia. It has also been associated with adult onset asthma. Outbreaks have been reported in closed communities. Real-time PCR is used for its diagnosis.

Option C: Haemophilus influenzae typically causes pneumonia in infants. While the primary infections occur due to capsulated strains, bronchopneumonia may occur as a secondary infection due to non-capsulated strains. Gram stain and culture are used for diagnosis.

Option D: Klebsiella pneumonia causes fatal pneumonia, which occurs in middle -aged or older persons with risk factors such as alcoholism, chronic bronchopulmonary disease or diabetes mellitus. Gram stain and culture are used for diagnosis.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 13:

BCYE medium (buffered charcoal yeast extract agar) is used to culture Legionella pneumophila.

Legionella pneumophila has fastidious growth requirements and grows on complex media such as BCYE medium with L-cysteine and antibiotic supplements (e.g., polymyxin B, cefazolin, vancomycin). The charcoal acts as a detoxifying agent. It grows slowly, and the colonies may take 3–6 days to appear.

Other options:

Option A: Leptospira is cultured on EMJH (Ellinghausen, McCullough, Johnson and Harris) medium

Option B: Bacillus cereus is cultured on PEMBA (Polymyxin pyruvate egg-yolk mannitol–bromothymol blue) agar

Option C: Campylobacter is cultured on Skirrow's, Campy BAP or Butzler's agar.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 14:

This patient most likely acquired infection with Legionella pneumophila from inhalation of aerosol in the air-conditioned rooms at the conference.

Legionella pneumophila can be found in natural, freshwater environments, but generally is not present in sufficient numbers to cause disease. Infection is typically by inhalation of aerosols produced by cooling towers, air conditioners, and showerheads which act as disseminators

No animal reservoirs exist. No evidence of human-to-human transmission has been found and no carrier state is established.

Solution to Question 15:

This patient presents with fever and lymphadenopathy after a cat scratch and the above image shows the black short rod-shaped bacilli arranged in chains or clumps, suggestive of cat scratch disease, This condition is caused by Bartonella henselae.

Cat scratch disease presents as a febrile illness with lymphadenopathy following a cat scratch. Patients present with a painless papule or pustule at the site of the scratch and lymph node enlargement and can be suppurative. The disease is usually self resolving. However, it can be treated with azithromycin only in patients who have severe complications.

Solution to Question 16:

5-day fever is an exclusively human disease and no animal reservoir is known.

Sold by @itachibot

5-day fever is also known as trench fever/quintan fever. It is caused by Bartonella quintana. It is characterized by sudden onset of fever accompanied by headache, malaise, restlessness, and shin pain.

Trench fever is transmitted by the body louse. (Option A). It has an incubation period ranging from 14-30 days. Symptoms coincide with the release of B. quintana in blood every 3-5 days with each episode lasting 5 days. Recrudescence and relapses are common. (Option B)

All cases are treated using doxycycline. Macrolides and fluoroquinolones can also be used in its treatment. (Option D)

Solution to Question 17:

The given scenario points to a diagnosis of cat-scratch disease caused by B. henselae. Stellate granulomas are characteristically present in this condition. They are granulomas containing central granular debris and neutrophils; giant cells are uncommon.

The image below shows a stellate granuloma

Other options:

Option A: Foreign body granulomas are incited by relatively inert foreign bodies, such as talc, sutures. Epithelioid cells and giant cells are apposed to the surface of the foreign body. The foreign material can usually be identified in the center of the granuloma.

The given image shows suture granuloma. The suture material is surrounded by lymphocytic infiltration

Sold by @itachibot



Option B: Naked granulomas are seen in sarcoidosis. They are non-caseating granulomas minimally surrounded by lymphocytes. The image below shows a naked granuloma



Option C: Durck granulomas are seen in cerebral malaria. The image below shows a Durck granuloma

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 18:

This clinical vignette of a patient with HIV infections and the violaceous lesions with a neutrophilic infiltrate is suggestive of bacillary angiomatosis.

Bacillary angiomatosis is seen in severely immunocompromised patients and is caused by B. henselae or B. quintana. It is characterized by neovascular proliferative lesions involving the skin. It consists of lobular proliferation of small blood vessels lined by enlarged endothelial cells interspersed with mixed infiltrates of neutrophils and lymphocytes, with a predominance of the former.

Both bacillary angiomatosis and Kaposi's sarcoma can have the clinical presentation mentioned in the question. However, the major differentiating factor is the microscopic examination:

• Bacillary angiomatosis predominantly has mixed infiltrate accompanied by neutrophils with clumps of bacilli.

• Kaposi's sarcoma predominantly has plasma cells and lymphocytic infiltrates.

Solution to Question 19:

This scenario of lesions on the hand followed by axillary lymphadenopathy is suggestive of cat scratch disease, hence we should ask about the history of exposure to cats. It is caused by Bartonella henselae.

Bartonella henselae is an intracellular, pleomorphic, gram-negative bacillus. They can be identified in lymph node biopsy with Warthin-Starry Silver stain. In addition to cat scratch disease, in immunocompromised individuals, it can also cause bacillary angiomatosis and peliosis hepatis.

Sold by @itachibot

In immunocompetent hosts, cat scratch disease is a self limiting illness. However in patients with severe disease, azithromycin can be used to treat the infection.



MARROW
Sold by @...
If you purchased this from ... else, you may have been scammed.

Solution to Question 20:

Parinaud's oculoglandular syndrome is an atypical form of cat scratch disease and is caused by Bartonella henselae . Bartonella henselae also causes bacillary angiomatosis in HIV-infected and other immunodeficient patients. . It presents as vascular nodules or tumours on the skin, mucosa, and other locations.

Bartonella bacilliformis causes Oroya fever also known as Carrion's disease.

Note: Ehrlichia phagocytophila has been renamed as Anaplasma phagocytophilum.

Solution to Question 21:

The above clinical scenario of fever, hemolytic anemia followed by verruga peruana lesions (Peruvian warts) is suggestive of Oroya fever. It is caused by Bartonella bacilliformis. The vector for Oroya fever is sandfly.

Oroya fever or Carrion's disease is associated with verruga peruana which are cutaneous vascular lesions carrying the bacteria.

It presents as fever with progressive anemia due to bacterial invasion of erythrocytes. A late sequela in survivors of Oroya fever or in those with asymptomatic infection is verruga peruana, which is characterized by vascular nodular skin lesions carrying the bacteria that occur in successive crops. An image of verruga peruana is given below.

Image shows verruga peruana lesions.

Sold by @itachibot

Bartonella quintana causes trench fever or 5 day fever.

Bartonella henselae causes cat scratch disease. It also causes bacillary angiomatosis and bacillary peliosis in HIV infected people.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Gram Negative Cocci

Question 1:

Which of the following is not a risk factor for Neisseria meningitidis infection?

a) Hypergammaglobulinemia

b) Mycoplasma infection

c) Hyposplenism

d) History of recent pilgrimage

Question 2:

A 22-year-old student presented with high-grade fever, and purpuric rash as shown in the image below. CSF sample showed gram-negative diplococci. Which of the following is the most likely aetiology?

a) Neisseria meningitidis

b) Pseudomonas aeruginosa

c) Streptococcus pneumoniae

d) E.coli

Question 3:

Sold by @itachibot

What is the most common manifestation of Neisseria meningitidis infection?

a) Meningitis

b) Septicaemia

c) Fever

d) Asymptomatic

Question 4:

A 5-year-old girl with thalassemia major was prophylactically immunized with a polysaccharide meningococcal vaccine before being taken up for elective splenectomy. Which group of Neisseria meningitidis would she be susceptible to even after immunization?

a) Group A

b) Group B

c) Group C

d) Group Y

Question 5:

A young man is brought to the ER with complaints of headache and fever. On examination, a maculopapular rash is seen over his trunk & extremities, and the following sign is elicited. Which of the following antibiotics should he be started on empirically?

a) Ceftriaxone

Sold by @itachibot
If you purchased this from someone else,
it may have been scammed.

Sold by @itachibot

b) Rifampicin

c) Ciprofloxacin

d) Cefazolin

Question 6:

You have to administer chemoprophylaxis to the contacts of a patient diagnosed with meningococcal meningitis. Which of the following combinations is correct?

a) 1, 2, 4 and 5

b) 2, 4 and 5

c) 1, 2, and 3

d) 1, 3 and 5

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 7:

Which of the following differentiates meningococcus from gonococcus?

a) It is oxidase positive

b) It ferments glucose

c) It ferments maltose

d) It reduces nitrates

Question 8:

A 30-year-old man has treatment-resistant gonorrhea. You plan to send a sample of his urethral discharge to the lab for culture and sensitivity testing. What transport media would you use for this purpose?

a) Cary-Blair medium

b) Thayer-Martin medium

c) Stuart's medium

d) Pike's medium

Question 9:

Sold by @itachibot

A 25-year old man presents with a history of urethral discharge for the last three days. A gram stain smear of the discharge is shown below. All of the following are true regarding the likely etiology except?



a) Catalase and oxidase positive

b) Pili is a virulence factor

c) Obligate intracellular organism

d) Shows twitching motility

If you purchased this from Sold by you may have been scammed. else,

Question 10:

Which of the following is the most common manifestation of gonorrhoea in males?

a) Asymptomatic

b) Acute urethritis

c) Discharging sinus

d) Genital ulcer

Question 11:

A patient with a history of sexual intercourse with a commercial sex worker 3 days back has developed genital discharge resembling 'flow of seed' as shown below. What medium should be used for the culture of the discharge?

Sold by @itachibot

a) Mannitol salt agar

b) Thayer-Martin media

c) Potassium tellurite agar

d) TCBS agar

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 12:

Which of the following is not a content of modified Thayer-Martin medium?

a) Vancomycin

b) Trimethoprim

c) Nystatin

d) Polymyxin B

Question 13:

An unresponsive middle-aged man with fever and petechial rashes was rushed into the ER. You successfully resuscitate him. Abdominal ultrasound shows bilateral adrenal hemorrhage. Which of the following statements is false regarding his condition?

a) Patients who recover may suffer from sloughing of the skin

b) The implicated organism can be isolated from the petechial rashes

c) The causative organism is Neisseria gonorrheae

d) Hypoplastic spleen is a risk factor for this condition

Sold by @itachibot

Question 14:

A 30-year-old woman with a history of recurrent pelvic inflammatory disease presents to the OPD with fever and right upper abdominal pain. A diagnostic laparoscopy was done and the findings are as given below. Which of the following organisms could be the causative agent?

a) Treponema pallidum

b) Haemophilus ducreyi

c) Neisseria meningitidis

d) Chlamydia trachomatis

Question 15:

Which is the true statement regarding gonococcal urethritis?

a) Symptoms are more severe in females than in males

b) Infection does not involve rectum and prostate

c) Dysuria is the most common presentation

d) Single dose of cefixime is the first line of treatment

Question 16:

Which of the following is not true about Neisseria gonorrhoeae?

Sold by @itachibot

a) It is exclusively a human pathogen

b) Some strains may cause disseminated disease

c) Vaccination has no role in prevention

d) Drug resistance is only due to acquisition of plasmids

Question 17:

Which of the following is true about Moraxella catarrhalis?

a) It is DNAse positive

b) An uncommon cause of COPD exacerbation

c) It ferments glucose

d) Causes angular conjunctivitis

Sold by @itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	a
2	a
3	d
4	b
5	a
6	a
7	c
8	c
9	c
10	b
11	b
12	d
13	c
14	d
15	c
16	d
17	a

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

Hypogammaglobulinemia is a risk factor for Neisseria meningitidis infection, not hypergammaglobulinemia.

Risk factors for meningococcal infection include:

• Overcrowding

• Cigarette smoking

• Mycoplasma infection

• Hyposplenism

• Hypogammaglobulinemia

• History of travel like pilgrimage

• Recent viral infection

• Terminal complement pathway deficiency (C5-C9 Deficiency) - increases the risk of disseminated gonococcemia

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

This patient has a high-grade fever and is a student, so he presumably spends a lot of time in crowded areas. The above image shows a purpuric rash. This patient's features are seen in meningococcemia caused by Neisseria meningitidis, which are gram-negative diplococci.

The organism enters the body through the nasopharynx by attaching to the epithelial cells with the help of pili. From the nasopharynx, they enter the bloodstream producing bacteremia, and the symptoms may resemble upper respiratory tract infection.

Fulminant meningococcemia is the severe form, where the patient presents with high-grade fever and hemorrhagic rash. There is thrombosis of several small blood vessels in various organs with perivascular infiltration and petechial hemorrhages. Some patients with fulminant meningococcemia may also develop disseminated intravascular coagulation, and circulatory collapse, a condition known as Waterhouse–Friderichsen syndrome.

Solution to Question 3:

Infection with Neisseria meningitidis is most commonly asymptomatic colonization in the nasopharynx of healthy individuals. Hence, a nasopharyngeal swab is used in the detection of carriers. In some patients, local inflammation ensues with rhinitis and pharyngitis.

Ceftriaxone as a single-dose IV/IM injection is highly (97%) effective in the eradication of carrier state and can be used at all ages and also in pregnancy. Ciprofloxacin is also highly effective and can be given orally, but it is not recommended in pregnancy and below 18 years of age.

Sold by @itachibot

Solution to Question 4:

Polysaccharide vaccines of Neisseria meningitidis do not offer protection against group B meningococci. This is because the group B capsule is made up of sialic acid residue which is also expressed on the surface of neural cells in the fetus. So, the B polysaccharide is perceived as 'self' and is poorly immunogenic.

To overcome this issue and provide protection against group B meningococci, a new vaccine was developed based on the surface protein i.e., non-capsular antigen of serogroup B meningococci. This is the multicomponent meningococcal vaccine - 4CMenB.

Meningococcal polysaccharide vaccines developed based on capsular antigen can be classified as:

- Plain polysaccharide vaccines:
- Bivalent (A and C),
- Trivalent (A, C and W-135)
- Tetravalent (A, C, Y, and W-135)
- Conjugated polysaccharide vaccines:
- Monovalent (A or C)
- Quadrivalent (A, C, Y, and W-135) MenQuadfi vaccine

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 5:

The clip given shows a positive Kernig's sign. This in combination with the other clinical features indicates a diagnosis of meningococcal meningitis. Third-generation cephalosporins such as ceftriaxone, cefotaxime, and ceftazidime are used empirically for the initiation of treatment for suspected meningococcal meningitis.

These drugs are administered as they also cover the various other potentially penicillin-resistant bacteria that may produce clinical features similar to that of meningococcal meningitis like penicillin-resistant pneumococcus. Also, some instances of reduced meningococcal sensitivity to penicillin have also been reported.

Both meningococcal meningitis and meningococcal septicemia are conventionally treated for 7 days, although courses of 3–5 days may be equally effective. Rifampicin can be used for prophylaxis for close contacts. Ceftriaxone can also be used for chemoprophylaxis, especially in pregnant individuals.

The image below shows petechial rashes in meningococcal infection

Sold by @itachibot

Sold by @itachibot
purchased this from someone else,
may have been scammed.

Solution to Question 6:

Ciprofloxacin, ceftriaxone, and rifampicin can be used for chemoprophylaxis of meningococcal meningitis.

Ciprofloxacin should not be given in pregnancy and to those below 18 years of age. Ceftriaxone is preferred in these cases as a single IM or IV dose. However, reduced susceptibility of isolates to ceftriaxone has occasionally been reported.

Recommended chemoprophylaxis for high-risk close contacts of meningococcal meningitis:

Age	Drug	Dose	Route and duration	Remarks
<1 month	Rifampicin	5 mg/kg	Oral every 12 hrs for 2 days	-
>1 month	Rifampicin	10 mg/kg	Oral every 12 hrs for 2 days	Not recommended in pregnancy
<15 years	Ceftriaxone	125mg	IM single dose	Can be given to children and is recommended in pregnancy
>15 years	Ceftriaxone	250 mg	IM single dose	Can be given to children and is recommended in pregnancy
>18 years	Ciprofloxacin	500 mg	Oral single dose	Not recommended in pregnancyPreferred for mass prophylaxis

Solution to Question 7:

Sold by @itachibot

Only meningococcus can ferment maltose, whereas both gonococcus and meningococcus can ferment glucose.

Both species are catalase and oxidase positive and neither of them can reduce nitrate. The only Neisseria species that can reduce nitrate is N. mucosa.

Solution to Question 8:

Stuart's medium is the transport media used for Neisseria gonorrheae and Neisseria meningitidis.

Stuart's medium is a semisolid, non-nutrient medium, which prevents microbial proliferation. It ensures that microorganisms present are able to survive for a sufficiently long period of time.

Solution to Question 9:

The given slide shows gram-negative diplococci within the polymorphs, characteristic of Neisseria gonorrhoeae infection. Neisseria is not an obligate intracellular organism, it can also occur extracellularly i.e, it is a facultative intracellular organism. The other neisseria that are found as normal inhabitants of the human respiratory tract occur extracellularly.

Neisseria gonorrhoeae is a gram-negative, non-sporulating, non-motile diplococcus that is aerobic but may grow anaerobically also. It is oxidase and catalase-positive.

Although Neisseria are non-motile organisms, they show twitching motility. This is a flagella-independent form of bacterial translocation over moist surfaces that occurs by the extension and retraction of polar type IV pili. These pili act as virulence factors by promoting attachment to host cells and inhibiting phagocytosis.

Virulence factors for gonococci:

• Outer Membrane Proteins (OMPs)

• IgA proteases

• Pili

• Lipo-oligosaccharide (LOS)

• Transferrin

• Lactoferrin-binding protein

Solution to Question 10:

Gonorrhea in males most commonly manifests as acute urethritis.

In males, the disease starts as acute urethritis. This infection may extend along the urethra to involve the prostate, seminal vesicles, and epididymis. Chronic urethritis may lead to stricture formation. The infection may also spread to the periurethral tissues causing abscesses and multiple discharging sinuses (watercan perineum).

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

In females, the initial infection involves the cervix, uteri and urethra which may extend to Bartholin glands, endometrium, and fallopian tubes. The vaginal mucosa is not usually affected in adults because the stratified squamous epithelium is resistant to infection by the cocci, and also because of the acidic pH of vaginal secretions.

Pelvic inflammatory disease and salpingitis may lead to sterility. Rarely peritonitis may develop with perihepatic inflammation (Fitz-Hugh-Curtis syndrome) which may also occur following Chlamydia trachomatis infection.

Solution to Question 11:

The clinical scenario is suggestive of gonococcal urethritis caused by Neisseria gonorrhoeae. Thayer-Martin and modified Thayer-Martin media are used for the culture of this organism. Gonorrhoea may manifest as mucopurulent genital discharge which can be described as resembling 'flow of seed' (semen).

Thayer-Martin medium contains vancomycin, colistin and nystatin.

Gonorrhea is a sexually transmitted disease caused by N. gonorrhea. The major symptoms include urethritis, vaginal discharge, dyspareunia, and dysuria. In the laboratory diagnosis of N. gonorrhea, the discharge on Gram's stain reveals gram■negative intracellular diplococci within phagocytes. It acidifies glucose but not maltose in serum sugars. Stuart's medium is used for transport while Thayer-Martin agar is used as a selective medium used to isolate N. gonorrhoeae. Cultures can be obtained from chocolate agar or Mueller-Hinton agar.

Stuart's medium is the transport media used for Neisseria gonorrhea and Neisseria meningitides. It is a semisolid, non-nutrient medium. Charcoal present in the medium neutralizes certain bacterial inhibitors while a reducing agent prevents oxidation.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 12:

Modified Thayer-Martin medium does not include polymyxin B. It includes polymyxin E, which is also known as colistin.

Modified Thayer–Martin medium is a selective media for Neisseria species. It includes:

• Vancomycin - to inhibit most gram-positive bacteria.

• Colistin (Polymyxin E) - to inhibit most gram-negative bacteria other than Neisseria.

• Nystatin or anisomycin - to inhibit yeast.

• Trimethoprim - to inhibit proteus.

The contents of the Thayer-Martin medium are the same as that of the modified Thayer-Martin medium except for the presence of trimethoprim in the modified version. The image below depicts Thayer-Martin medium.

Sold by @itachibot



Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Solution to Question 13:

The above scenario depicts Waterhouse-Friderichsen syndrome which is a complication seen in infection with meningococci. Waterhouse-Friderichsen syndrome is also known as hemorrhagic adrenalitis or fulminant meningococcemia and is characterized by bilateral adrenal gland hemorrhage.

Though Waterhouse-Freiderichsen syndrome is most often associated with meningococcal septicemia, it may complicate sepsis caused by other organisms like certain streptococcal species also. Hypoplastic spleen may also be associated with this condition and the implicated organism can be isolated from the petechial rashes.

Complications :

• Shock, extensive hemorrhage within the skin

• May fall into a coma

• Death, usually after a few hours, due to adrenal insufficiency

• Patients who recover may suffer from extensive sloughing of the skin and loss of digits due to gangrene.

Fulminant meningococcemia is a medical emergency and needs to be treated with adequate antibiotics as fast as possible. Ceftriaxone is an antibiotic commonly used nowadays.

Solution to Question 14:

The scenario depicted is Fitz-Hugh-Curtis syndrome which is a rare complication of pelvic inflammatory disease (PID) following infection with Chlamydia trachomatis or Neisseria gonorrhoeae.

Sold by @itachibot

It causes liver capsule inflammation leading to the creation of violin-string adhesions which can be seen on laparoscopy. On CT imaging it appears as perihepatic inflammatory stranding and hepatic capsular enhancement.

The major symptom and signs include right upper quadrant pain. It is also associated with minimal pelvic pain, vaginal discharge, or cervical motion tenderness.

Solution to Question 15:

Acute urethritis is the most common clinical manifestation of gonorrhea in males and major symptoms include urethral discharge and dysuria, usually without urinary frequency or urgency.

Other options:

Option A: The stratified squamous epithelium of the vagina is resistant to gonococcus, so infection is less severe in females.

Option B: In males, it can involve rectum, prostate, and epididymitis, but it never involves testes.

Option D: Rising minimum inhibitory concentration of cefixime worldwide has led the CDC to discontinue its recommendation of this agent as first-line treatment for uncomplicated gonorrhea.

Third-generation cephalosporins have remained highly effective as single-dose therapy for gonorrhoea. Ceftriaxone (250mg, IM) single dose + azithromycin 1 g PO, single dose or doxycycline 100 mg PO BD for 7 days is given. This also covers treatment for chlamydia (if chlamydial infection is not ruled out).

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 16:

Neisseria gonorrheae acquires resistance to antibiotics either by chromosomal mutations or by the acquisition of plasmids (R factors). These resistant strains are also known as:

• PPNG - penicillinase-producing N. gonorrhoeae (plasmid-borne resistance)

• TRNG - tetracycline-resistant N. gonorrhoeae (plasmid-borne resistance)

• CMRNG - chromosomally mediated resistance in N. gonorrhoeae (resistance to both penicillin and tetracycline)

• QRNG - quinolone resistant N. gonorrhoeae (due to alterations in DNA gyrase and topoisomerase IV)

Other options:

Option A: N. gonorrhea is a fastidious organism and is exclusively a human pathogen.

Option B: Some strains of N. gonorrhea can ascend up the genital tract and can lead to many disseminated diseases such as septic arthritis and pelvic inflammatory disease.

Option C: Vaccines are not effective against N. gonorrhea due to its antigenic variation and ability to invade neurophils.

Sold by @itachibot

Solution to Question 17:

Moraxella catarrhalis is a DNAse positive gram-negative diplococcus. It is the second most common bacterial cause of COPD exacerbations (after H. influenzae). It can also cause otitis media, sinusitis, tracheobronchitis, and pneumonia.

It differs from pathogenic Neisseria as:

• It does not ferment any carbohydrates - Asaccharolytic

• Gives a positive tributyrin hydrolysis test.

• Shows positive DNAse test.

Angular conjunctivitis is caused by Moraxella lacunata, which is a gram-negative rod-shaped bacteria.

Treatment is with oral amoxicillin and clauvulanic acid.

Sold by @Itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Rickettsia, Chlamydia and Mycoplasma

Question 1:

Transovarial transmission is a feature of which of the following diseases?

a) Scrub typhus

b) Epidemic typhus

c) Endemic typhus

d) Trench fever

Question 2:

A 36-year-old trekker presented with fever, headache, and rashes following a tick bite. He was later diagnosed to have Rocky Mountain spotted fever. Which of the following statements is false about the microbe implicated?

a) Obligate intracellular organism

b) Beta-lactams are ineffective

c) Reproduce by binary fission

d) Cell wall is absent

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following statements is true regarding endemic murine typhus?

a) Man to man transmission occurs

b) It is caused by R.prowazekii

c) Vector is Xenopsylla cheopis

d) Neill-Mooser reaction is negative

Question 4:

Multiple inmates at a prison are suspected to have epidemic typhus following an arthropod infestation. Which of the following vectors is implicated in the spread of this disease?

Sold by @itachibot



a) A

b) B

c) C

d) D

Sold by @itachi... purchased this from some... you may have been scammed. ...else,

Question 5:

A 45-year-old man from Kashmir presents with fever, chills and rash for 1 week. The rash appeared on the 5th day of the fever starting from the trunk and progressing to the limbs. Epidemic typhus is suspected. Which of the following statements is incorrect about the disease?

a) Transmitted by a louse

b) Brill-Zinsser disease is a recrudescent form

c) Rash is present on palms and soles

d) Transovarian transmission does not occur

Question 6:

What is the drug of choice for scrub typhus?

a) Doxycycline

b) Ciprofloxacin

c) Chloramphenicol

Sold by @itachibot

d) Azithromycin

Question 7:

A 25-year-old man came with complaints of fever and a skin lesion as shown below. He gives a history of hiking in scrub forests a week prior, where he suffered an insect bite. On examination, there is regional lymphadenopathy and a maculopapular rash. Which of the following arthropods is involved in the spread of this disease?

a) Gamasid mite

b) Trombiculid mite

c) Tick

d) Louse

Question 8:

A wildlife photographer presents with a blackish lesion at the site of an insect bite from a few days back. He also gives a history of fever. A positive Weil-Felix test with OX K agglutinins was observed. What is the diagnosis?

a) Rocky mountain spotted fever

b) Epidemic typhus

c) Endemic typhus

d) Scrub typhus

Sold by @itachibot

Question 9:

A veterinarian presents with new-onset mitral regurgitation and fever. An echo reveals valve vegetations and blood cultures show no growth. Q fever is suspected. Which of the following statements is false regarding the organism implicated?

a) Obligate intracellular

b) Gram-negative bacteria

c) Survives Holder's method of pasteurisation

d) Transmitted by mites

Question 10:

A patient presents with redness of the eye and foreign body sensation. On examination, there is follicular hypertrophy in the lower lids and moderately abundant exudates. Chlamydial infection is suspected. Which strains could have caused this condition?

a) Serotypes D to K

b) Serotypes A, B, Ba and C

c) Serotype L1, L2, L3

d) Serotype H- K

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 11:

A thick, yellowish discharge is noted from the eyes of a five-day-old neonate in the step-down ward during rounds. An iodine-stained smear of the exudates is positive for large cytoplasmic inclusions as shown in the image. Which of the following is false about the organism responsible for the above condition?

Sold by @itachibot



a) Filterable organisms

b) Obligate intracellular

c) Possess only DNA

d) Cannot grow on artificial media

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 12:

A poultry worker was admitted to ICU with severe pneumonia. Bronchoalveolar lavage was done; it showed Levinthal Cole Lillie bodies. Which of the following is the causative organism for the above condition?

a) Legionella pneumophila

b) Chlamydia psittaci

c) Mycoplasma pneumoniae

d) Moraxella catarrhalis

Question 13:

A patient presented with painful swellings on both sides of the groin as shown in the image. The clinician suspects lymphogranuloma venereum. Which is the most common serovar causing the above condition?

Sold by @itachibot

a) L1

b) L2

c) L3

d) L4

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 14:

What is the incubation period of lymphogranuloma venereum?

a) 1-5 days

b) 7-10 days

c) 10-30 days

d) 30-60 days

Question 15:

Lesion scrapings from a genital ulcer show numerous polymorphs, but no bacteria. Which of the following is the most appropriate culture medium for organisms responsible for the above condition?

a) McCoy cells

b) Thayer-Martin agar

c) Robertson's cooked meat

d) PPLO broth

Sold by @itachibot

Question 16:

A 28-year-old woman presented with a 3-day history of mucopurulent cervical discharge. Gram staining of the discharge revealed numerous pus cells, but no organisms. What is the investigation of choice for the diagnosis of this infection?

a) Saline mount

b) Serum antibody testing by immunofluorescence

c) Culture in Thayer Martin media

d) Nucleic acid amplification test

Question 17:

Which of the following microbes is also referred to as Eaton's agent ?

a) Klebsiella pneumoniae

b) Corynebacterium diphtheria

c) Chlamydia trachomatis

d) Mycoplasma pneumoniae

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Question 18:

A 20-year-old military recruit presented with fever, sore throat, and non-productive cough which showed no improvement with beta-lactam antibiotics. X-ray showed peribronchial pneumonia with thickened bronchial markings. Primary atypical pneumonia is suspected. Which of the following statements is false regarding the organism responsible for this condition?

a) Obligate intracellular

b) Cell wall is absent

c) Highly pleomorphic

d) Multiply by binary fission

Question 19:

Diene's method of staining is used for which of the following organisms?

Sold by @itachibot

a) Mycoplasma

b) Chlamydiae

c) Anthrax

d) Diphtheria

Question 20:

Mycoplasma is inherently resistant to which of the following antibiotics?

a) Beta lactam antibiotics

b) Macrolides

c) Tetracycline

d) Flouroquinolones

Sold by @itachibot
If this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	a
2	d
3	c
4	b
5	c
6	a
7	b
8	d
9	d
10	a
11	c
12	b
13	b
14	c
15	a
16	d
17	d
18	a

Sold by @itachibot

19	a
20	a

Detailed Explanations

Solution to Question 1:

Transovarial transmission is a feature of Orientia tsutsugamushi which causes scrub typhus.

Transovarial transmission refers to passing on the disease-causing bacteria from parent arthropod to offspring arthropod. Orientia tsutsugamushi is maintained by transovarial transmission in trombiculid mites. It is also seen in Rickettsia rickettsii, Rickettsia conorii, and Rickettsia akari.

The image below shows a trombiculid mite.



Trench fever (option D) is caused by Bartonella quintana.

Solution to Question 2:

Rocky Mountain spotted fever is caused by Rickettsia rickettsii. A cell wall is present in rickettsial organisms.

Rickettsiae are obligate intracellular, gram-negative pleomorphic coccobacilli. They are unable to grow in cell-free media and are thus cultured in yolk sacs of embryonated eggs. They are non-motile, non-capsulated organisms that contain both DNA and RNA. They reproduce by binary fission and are transmitted by arthropods like ticks and lice (except Coxiella burnetii and Bartonella henselae).

The rickettsiae are localized in endothelial cells of small blood vessels (except Ehrlichia and Anaplasma) in humans. They cause endothelial cell enlargement and degeneration, predisposing

Sold by @itachibot

to thrombus formation. Tetracyclines, macrolides, and fluoroquinolones are effective against this group but, beta-lactams are ineffective.

Solution to Question 3:

The vector for endemic murine typhus is the rat flea, Xenopsylla cheopis.

Endemic murine typhus is cuased by Rickettsia typhi (option B) while epidemic typhus is caused by Rickettsia prowazekii. The zoonotic reservoir for R. typhi is the rat. Humans are infected when flea feces contaminate pruritic bite lesions. Transmission can also occur via inhalation of aerosols of dried feces of rickettsemic rat flea. No man-to-man transmission occurs. Patients present with fever, chills, vomiting and a maculopapular rash that is more prominent on the trunk. Diagnosis is by serology and treatment is with doxycycline.

When male guinea pigs are inoculated intraperitoneally with blood from a case of endemic typhus, they develop fever and a characteristic scrotal inflammation. The scrotum becomes enlarged and the testes cannot be pushed into the abdomen. This is known as the Neil-Mooser reaction. It is positive with Rickettsia typhi (option D) and negative with Rickettsia prowazekii.

The image below shows the rat flea, Xenopsylla cheopis.

Solution to Question 4:

Louse (image B), Pediculus humanus corporis, transmits epidemic typhus.

Gaol fever/ Jail fever/ epidemic typhus is caused by Rickettsia prowazekii.

Sold by @itachibot



Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Solution to Question 5:

Epidemic typhus rash spares the palms and soles.

Epidemic typhus, caused by Rickettsia prowazekii, is transmitted by the louse Pediculus humanus corporis. There is no transovarial transmission of R. prowazekii. The disease spreads through infected louse feces deposited when the louse feeds on the human host. It is commonly seen in poor unhygienic or cold conditions.

Epidemic typhus starts with fever and chills. A characteristic rash appears on the fourth or fifth day, starting on the trunk and spreading over the limbs but sparing the face, palms, and soles. In some patients who recover from epidemic typhus, the rickettsiae may remain latent in the lymphoid tissues or organs for years. Such latent infection may, at times, be reactivated leading to recrudescent typhus (Brill-Zinsser disease).

Transovarial transmission refers to passing on the disease-causing bacteria from parent arthropod to offspring arthropod. It occurs with Orientia tsutsugamushi, Rickettsia rickettsii, R. conorii and R. akari.

Note: Rash on palms and soles is typically described in Rocky Mountain spotted fever, secondary syphilis and hand, foot, and mouth disease

The image below shows the rash of epidemic typhus.

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 6:

The drug of choice for scrub typhus is doxycycline.

To treat scrub typhus, oral doxycycline 100 mg is given twice daily for 7–15 days. In the case of pregnant women, oral azithromycin 500 mg for 3 days may be used as doxycycline is contraindicated in pregnancy.

Scrub typhus is a rickettsial infection caused by Orientia tsutsugamushi. The vector is a trombiculid mite. Humans are infected when they are bitten by the mite larvae known as chiggers. The incubation period is 1- 3 weeks. Patients develop a characteristic eschar at the site of the mite bite, along with regional lymphadenopathy and maculopapular rash. It can be associated with fever, headache, myalgia, meningoencephalitis, myocarditis, and pneumonia.

Serologic assays are used for diagnosis. Weil-Felix reaction is a heterophile agglutination reaction used for diagnosis, wherein the sera are tested for agglutinins to the O antigens of Proteus strains OX19, OX2, and OXK. OXK alone is agglutinated in scrub typhus.

Solution to Question 7:

A history of contact with scrub vegetation with fever, regional lymphadenopathy, maculopapular rash, and eschar at the bite site is suggestive of scrub typhus.

Scrub typhus is caused by Orientia tsutsugamushi and transmitted by trombiculid mite. Doxycycline 100mg BD for 7-14 days is the drug of choice for treatment of scrub typhus in adults. In case of pregnant women and children, azithromycin or chloramphenicol may be preferred where doxycycline is relatively contraindicated.

Sold by @itachibot

Solution to Question 8:

In the above clinical scenario, a history of insect bite followed by the development of a characteristic eschar at the site of the bite and a positive Weil-Felix test with OX K agglutinins suggests a diagnosis of scrub typhus.

Weil-Felix reaction is a heterophile agglutination reaction, wherein the sera are tested for agglutinins to the O antigens of certain Proteus strains OX 19, OX 2, and OX K. The basis of the test is sharing an alkali stable carbohydrate antigen by some Rickettsiae and by some strains of Proteus.

DISEASE	OX 19	OX 2	OX K
Epidemic typhus	+++	+	-
Endemic typhus	+++	+/-	-
Rocky Mountain spotted fever	++	++	-
Scrub typhus	-	-	+++
Rickettsial pox	-	-	-
Trench fever	-	-	-
Q fever	-	-	-

Solution to Question 9:

Coxiella burnetii is the causative organism of Q fever. It is transmitted by ixodid ticks in animals; however, it is not transmitted to humans via ticks.

Coxiella burnetii is a filterable, obligate intracellular pathogen. They are gram-negative but are better stained with Gimenez, Giemsa stains. They are heat resistant and hence survive Holder's method of pasteurization.

Q fever is a zoonosis and the primary source of human infection are infected cattle, sheep, and goats. It spreads through the inhalational route. Veterinarians are at higher risk due to increased exposure. Acute disease resembles influenza, nonbacterial (atypical) pneumonia, and hepatitis. Chronic disease presents as culture-negative bacterial endocarditis in patients with valvular heart disease.

Solution to Question 10:

Redness, foreign body sensation with follicular hypertrophy in the lower lids and moderately abundant exudates suggests inclusion conjunctivitis which is caused by Chlamydia trachomatis serotypes D to K.

Sold by @itachibot

Inclusion conjunctivitis is a type of acute follicular conjunctivitis caused by C. trachomatis serotypes D to K. The infection is transmitted to the eye from genitals, or contaminated swimming pools. Patients present with foreign body sensation, redness and mucopurulent discharge. On examination, there is acute follicular hypertrophy of the lower palpebral conjunctiva and fornix, non-tender pre-auricularlymphadenopathy and superficial keratitis in upper half of cornea.

C. trachomatis is a Gram-negative, obligate intracellular bacteria. It cannot produce ATP. It causes a variety of human diseases (ocular and genitourinary) and has two main biological variants:

- TRIC biovar which causes trachoma inclusion conjunctivitis (TRIC biovar). This is further divided into 12 serological variants:
- A, B, Ba, C serovars which cause blinding or endemic trachoma
- D to K servovars which cause inclusion conjunctivitis
- LGV biovar (L1, L2, L3 serotypes) which causes lymphogranuloma venereum

Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 11:

The given clinical scenario of a thick yellowish discharge from the eye showing iodine-stained inclusion bodies is suggestive of inclusion conjunctivitis caused by Chlamydia trachomatis. As chlamydia is a bacteria, it possesses both DNA and RNA.

Chlamydiae are obligate intracellular organisms that produce basophilic inclusion bodies. They are filterable like viruses. They are called energy parasites as they lack enzymes for ATP synthesis. They cannot grow on artificial media but can grow on cell lines like McCoy, HeLa, or the yolk sac of the chick embryo.

Chlamydiae occur in two forms, the elementary body and the reticulate body. The extracellular, infective form is called the elementary body. The intracellular, non-infective, growing, and replicative form is called the reticulate body. These are found inside a membrane-bound compartment called the inclusion body which can be visualized within infected epithelial cells by iodine or Giemsa stain. The cell wall of the reticulate body is fragile and pliable, leading to pleomorphism.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... you may have been scammed. ...else,

Solution to Question 12:

Pneumonia in a poultry worker with Levinthal-Cole-Lillie bodies on bronchoalveolar lavage specimens is suggestive of psittacosis.

Levinthal-Cole-Lillie bodies are inclusion bodies of Chlamydia psittaci seen in infected cells including alveolar macrophages from patients suffering from psittacosis. C. psittaci can be transmitted to humans by direct contact with infected birds or by inhalation of aerosols from avian feather dust. Psittacosis presents with fever, chills, muscular aches and pains, severe headache, hepatosplenomegaly and gastrointestinal symptoms. Severe pneumonia may develop. It is treated with tetracycline.

Solution to Question 13:

Lymphogranuloma venereum (LGV) is caused most commonly by Chlamydia trachomatis LGV serovar L2.

LGV is a sexually transmitted disease, characterized by suppurative inguinal adenitis.

Solution to Question 14:

The incubation period of Lymphogranuloma venereum ranges from 10-30 days.

Chlamydia trachomatis serovars L1, L2, and L3 cause lymphogranuloma venereum (LGV). The most common presentation is the inguinal syndrome, which is characterized by painful inguinal lymphadenopathy.

Sold by @itachibot

Solution to Question 15:

Clinical scrapings from a genital ulcer showing polymorphs in the absence of bacteria is suggestive of chlamydial infection and the most appropriate culture medium is McCoy cells (human synovial carcinoma cell line).

C. trachomatis is a small gram-negative bacterium and an obligate intracellular parasite that appears as an intracytoplasmic inclusion body. The gram stain of the cervical smear would show leucocytes but no organisms. Hence, selective culture media like the McCoy cells culture is used to isolate the organism.

Option B: Thayer–Martin agar is used for isolation of Neisseria spp.

Option C: Robertson's cooked meat is used for the culture of anaerobes, especially Clostridia

Option D: Pleuropneumonia-like organisms (PPLO) broth is used to culture Mycoplasma spp. and Ureaplasma spp.

Note: Abnormal urethral gram stain showing >10 PMNs/1000Å~ field in women with dysuria but without coliform bacteriuria supports the diagnosis of chlamydial urethritis.

Sold by @itachibot
If you purchased this from someone else you may have been scammed.

Solution to Question 16:

Mucopurulent cervical discharge for 3 days is suggestive of acute cervicitis. As gram stain does not show any organism, gonococcal infection is less likely. In this case, Chlamydia trachomatis is the most likely organism. For C. trachomatis, the investigation of choice is the nucleic acid amplification test (NAAT).

NAATs are the most sensitive tests to detect chlamydia. These tests also have a high specificity comparable to culture, and in contrast to culture, do not depend on viable pathogens. Therefore, NAATs are generally considered the test of choice for chlamydia and have replaced culture as the diagnostic gold standard.

Mucopurulent cervicitis is the inflammation of the columnar epithelium and subepithelium of the endocervix in women. Its commonly caused by Neiserria gonorrhoeae and Chlamydia trachomatis.

Solution to Question 17:

Eaton's agent refers to Mycoplasma pneumoniae.

Eaton was the first to isolate Mycoplasma pneumoniae. Because it was filterable, it was considered to be a virus previously.

It was formerly known as PPLO– Pleuro pneumonia-like organism.

Solution to Question 18:

Sold by @itachibot

The above clinical scenario of fever, non-productive cough, no improvement with beta-lactam antibiotics along with X-ray features showing peribronchial pneumonia with thickened bronchial markings is suggestive of Mycoplasma pneumonia, also known as primary atypical pneumonia or walking pneumonia.

Mycoplasmas are not obligate intracellular parasites. They are free-living bacteria devoid of cell walls. They are highly pleomorphic, with no fixed shape or size.

Because of their plasticity, they can pass through bacterial filters and have often been mistaken for viruses.

Mycoplasmas do not possess spores, flagella, or fimbria. Some species exhibit gliding motility. Mycoplasmas are gram-negative, but are better stained by Giemsa stain.

If yo... by @itachibot ... from someone else, ... sampled.

Solution to Question 19:

Diene's method is used for mycoplasma.

Mycoplasma shows a ‘fried-egg’ colony on Diene’s method of staining. The central dark bluish area represents the area of mycoplasmal down growth whereas the pale blue periphery indicates superficial growth of mycoplasma on the agar.



Solution to Question 20:

Mycoplasma is inherently resistant to beta-lactam antibiotics.

Mycoplasma lack cell wall, and therefore, are inherently resistant to beta-lactam antibiotics as these drugs act on the cell wall.

Sold by @itachibot

Spirochetes

Question 1:

Identify the organism.



a) Leptospira interrogans

b) Treponema pallidum

c) Borrelia burgdorferi

d) Helicobacter pylori

If you purchase this, you may have b... else,

Question 2:

A 30-year-old truck driver presented with a painless ulcer in the genital area. On examination, a single ulcer with an indurated base was found. An image of the tissue section is shown below. Identify the staining method.

Sold by @itachibot

a) Fontana's method

b) Levaditi's method

c) Castaneda's method

d) Albert's method

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 3:

What type of motility is shown by Treponema pallidum?

a) Tumbling motility

b) Gliding motility

c) Corkscrew motility

d) Darting motility

Question 4:

A 32-year-old pregnant lady tests VDRL positive at her 1st-trimester antenatal check-up. An FTA-ABS confirms the diagnosis. Which of the following is a false statement about the organism causing this disease?

a) It cannot be cultured in vitro

b) Endoflagella is in the periplasmic space.

c) Humans are the only known natural hosts

d) Reiter's strain is pathogenic

Sold by @itachibot

Question 5:

A patient had a solitary, painless penile ulcer 2 months back. Now, he presents with a generalized maculopapular rash over the body. What is the sensitivity of VDRL at this stage of the disease?

a) 25%

b) 50%

c) 75%

d) 100%

Question 6:

Which of the following is a false statement regarding the serological tests of syphilis?

a) Rapid plasma reagin test is a group specific treponemal test

b) VDRL test is the test of choice for response to therapy

c) VDRL is the test of choice for examining CSF

d) Treponemal tests are likely to remain reactive even after adequate therapy

Question 7:

A 55-year old man was brought to the OPD with complaints of rapid decline in cognition, impaired position, and vibratory sense. There is a history of sporadic short-term relationships with many promiscuous sexual partners. Romberg's sign is positive and bilateral Argyll Robertson pupils are seen on checking for pupillary reflex. Which of the following is the investigation of choice?

a) Treponema pallidum haemagglutination test

b) Venereal disease research laboratory (VDRL) test

c) Fluorescent Treponemal Antibody Absorption test

d) EIA-Enzyme-linked ImmunoAssay

Question 8:

You treated a case of secondary syphilis with a single dose of benzathine penicillin-G 2.4 million IU IM. What follow-up advice would you give for the assessment of the patient's

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Sold by @itachibot

response to treatment?

a) Follow up with VDRL after 6 months and 12 months

b) Follow up with VDRL after 3 months and 6 months

c) Follow up with TPHA after 6 months and 12 months

d) Follow up with TPHA after 3 months and 6 months

Question 9:

A pregnant woman at 10 weeks of gestational age tested VDRL positive but negative for specific treponemal tests. What is your interpretation?

a) Positive for syphilis

b) False negative

c) Positive for non pathogenic treponema

d) Biological false positive

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 10:

A 2-month-old infant presented with a runny nose and bullous lesions on the skin and legs. Her X-ray (given below) showed features suggestive of periostitis. Which of the following is the initial investigation of choice?



a) Venereal disease research laboratory (VDRL) test

b) Treponema pallidum hemagglutination (TPHA) assay

Sold by @itachibot

c) Fluorescent treponema antibody absorbed (FTA-ABS) test

d) Culture

Question 11:

What is the causative agent of yaws ?

a) T.pallidum

b) T.pertenue

c) T.carateum

d) T.endemicum

Question 12:

What is the causative organism and the corresponding vector for epidemic relapsing fever?

a) Borrelia burgdorferi - louse

b) Borrelia recurrentis - tick

c) Borrelia recurrentis - louse

d) Borrelia burgdorferi - tick

Question 13:

During a louse outbreak among jail inmates, a few of them developed fevers that subsided after 3-5 days. The fever relapsed after a week and subsided again. Many patients had tender splenomegaly on examination. Which culture medium is used to isolate the organism implicated here?

a) Thayer-Martin medium

b) Thiosulphate, citrate, bile salt agar

c) Barbour-Stoenner-Kelly broth medium

d) Mannitol, egg-yolk, phenol red, polymyxin agar

Question 14:

Which one of the following micro-organisms uses antigenic variation as a major means of evading the host immune system?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

a) Treponema pallidum

b) Borrelia recurrentis

c) Leptospira

d) Listeria

Question 15:

A 23 year old presented with fever, myalgia, and skin lesion as shown in the image. What is the causative organism for the above condition?

a) Orientia tsutsugamoshi

b) Ehrlichia chaffeensis

c) Borrelia burgdorferi

d) Coxiella burnetti

Question 16:

Following a flood, there is an outbreak of fever. Water samples from various sources are being studied for pathogens. The bacteria shown in the given image was isolated as a potential cause of disease. Which of the following animals is the most important reservoir of these organisms?

Sold by @itachibot



a) Rat

b) Cat

c) Fish

d) Dog

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 17:

A farmer who works in paddy fields presented with fever, intense myalgia, and hematochezia. Lab investigations showed sharply elevated creatinine levels and high liver enzymes. What is the causative organism?

a) Leptospira

b) Borrelia

c) Legionella

d) Listeria

Question 18:

At a relief camp in Karnataka, a few people developed fever, myalgia, and conjunctival suffusion. Rapid diagnostic IgM ELISA tests performed in the field showed leptospirosis. Which of the following culture media can be used to confirm this diagnosis?

a) EMJH

b) BCYE

c) TCBS

Sold by @itachibot

d) CCFA

Question 19:

A 16-year-old boy is admitted with fever, icterus, conjunctival suffusion, and hematuria for 15 days. Which serological test should be done for diagnosis?

a) Widal test

b) Microscopic agglutination test

c) Weil felix reaction

d) Paul Bunnell test

Question 20:

A 12-year-old boy presents with right upper quadrant pain, calf pain, conjunctival suffusion, icterus, and fever. Examination shows tender hepatomegaly. What is the most likely diagnosis?

a) Chikungunya

b) Leptospirosis

c) Dengue hemorrhagic fever

d) Encephalopathy caused by hepatitis A

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	d
5	d
6	a
7	b
8	a
9	d

Sold by @itachibot

10	a
11	b
12	c
13	c
14	b
15	c
16	a
17	a
18	a
19	b
20	b

Sold by @itachibot If you purchased this from someone else, have been scammed.

Detailed Explanations

Solution to Question 1:

.The given image shows Treponema pallidum demonstrated by dark ground microscopy.

Treponemes are short, slender with regular and sharp spirals.

Option A: Leptospira have a large number of closely wound spirals and characteristic hooked ends as shown in the image below



Sold by @itachibot



Option C: Borrelia are large with irregular and wide open coils.

Option D: Helicobacter pylori is a spiral rod (bacilli) that is not classified as a spirochete.

Solution to Question 2:

The given image shows Treponema pallidum demonstrated by Levaditi's method. Single painless ulcer with indurated base points towards the diagnosis of syphilis.

Syphilis is caused by Treponema pallidum, a spirochete bacillus. It is a sexually transmitted infection. Intrauterine transmission can lead to congenital syphilis.

It shows corkscrew motility which can be demonstrated by darkfield microscopy.

Treponema can be stained by silver impregnation methods.

• Fontana's method is used on smear

• Levaditi's method is used for tissue sections

The clinical manifestations of syphilis include:

Primary syphilis: The typical primary chancre usually begins as a single painless papule that rapidly becomes eroded and usually becomes indurated, with a characteristic cartilaginous consistency on palpation of the edge and base of the ulcer.

Inguinal lymphadenopathy is bilateral and may occur with anal as well as with genital chancres. The chancre generally heals within 4–6 weeks (range, 2–12 weeks), but lymphadenopathy may persist for months.

Secondary syphilis: The various features include mucocutaneous or cutaneous lesions and generalized nontender lymphadenopathy. macular, papular, papulosquamous, and occasionally pustular syphilides. Other features of secondary syphilis include sore throat, fever, weight loss, malaise, anorexia, headache, and meningismus. Ocular findings associated with secondary (or later/ unknown-stage) syphilis include pupillary abnormalities and optic neuritis as well as the

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

classic iritis or uveitis.

Latent syphilis: Early latent syphilis is limited to the first year after infection, whereas late latent syphilis is defined as that of 1 year's duration (or of unknown duration). T. pallidum may still seed the bloodstream intermittently during the latent stage, and latent syphilis in a pregnant woman may infect the fetus in utero.

Neurosyphilis: The major clinical categories of symptomatic neurosyphilis include meningeal, meningovascular, and parenchymatous syphilis.

Cardiovascular syphilis: Cardiovascular involvement results in uncomplicated aortitis, aortic regurgitation, saccular aneurysm (usually of the ascending aorta), or coronary ostial stenosis.

Congenital syphilis: The earliest manifestations of congenital syphilis include rhinitis, or "snuffles"; mucocutaneous lesions, bone changes, including periostitis detectable by x-ray examination of long bones; hepatosplenomegaly, lymphadenopathy, anemia, jaundice, thrombocytopenia, and leukocytosis.

Serologic tests for syphilis recommended by the Centers for Disease Control and Prevention (CDC):

• Screening or diagnosis (RPR or VDRL),

• Quantitative measurement of antibody to assess clinical syphilis activity or to monitor response to therapy (RPR or VDRL)

• Confirmation of a syphilis diagnosis in a patient with a reactive RPR or VDRL test (FTA-ABS, TPHA, EIA/CIA)

Treatment of syphilis is with 2.4 million units of intramuscular benzathine penicillin.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Treponema pallidum show corkscrew motility, i.e movement by rotation around the long axis.

The characteristic feature of spirochetes is the presence of endoflagella, situated between the outer membrane and cell wall, that facilitates this movement.

Sold by @itachibot



Sold by @itachi... from some... else, If you purchased this... you may have been scammed.

Solution to Question 4:

VDRL positive status is suggestive of syphilis, caused by Treponema pallidum. Fluorescent treponemal antibody absorption (FTA-ABS) is a confirmatory test.

Reiter's strain is a cultivable, non-pathogenic treponeme, showing morphological and antigenic similarities with T.pallidum. It is widely used as the antigen in group-specific treponemal tests for the diagnosis of syphilis.

Nichol's strain is virulent and non-cultivable in artificial media. However, it can be grown in cell culture through serial passage in live rabbit testes.

Solution to Question 5:

The sensitivity of VDRL is 100% in secondary syphilis. History of solitary painless ulcer with a generalized maculopapular rash is consistent with secondary syphilis.

The VDRL test becomes positive 4–5 weeks after exposure to T. pallidum and 1–2 weeks after the appearance of the chancre.

The VDRL test is a highly sensitive test. The sensitivity of the test depends on the stage of the disease:

• Primary syphilis- 60–75%

• Secondary syphilis- 100%

• High antibody titers may be found in patients developing gummatous lesions and neurovascular or cardiovascular complications.

• Tertiary syphilis- usually negative

• Negative test:

Sold by @itachibot

• Early primary syphilis

• Latent acquired syphilis

• Late congenital syphilis

Secondary syphilis is the most infective stage in the course of the illness. The patient may present with a roseolar or papular skin rash, condyloma lata, and mucous patches.

The image below shows lesions of secondary syphilis.

Solution to Question 6:

RPR is a non-specific test. It uses the VDRL antigen (cardiolipin) containing carbon particles. Hence, results can be seen by the naked eye, whereas VDRL requires a microscope for visualization of flocculation.

The test of choice for monitoring response to treatment and for examining CSF is VDRL

The test of choice for syphilis screening is RPR.

After treatment, in early syphilis,

• Non-treponemal tests- titers decline or become non-reactive

• Treponemal tests- remain reactive

For diagnosis of syphilis, a 2 test approach is used i.e., screening with an NTT (non-treponemal test) which includes either VDRL or RPR. Confirmation of syphilis is done by a treponemal test, most commonly by TPHA (Treponema pallidum hemagglutination test).

Follow-up is done using NTT (as treponemal tests remains positive for years after treatment).

Solution to Question 7:

Sold by @itachibot

The clinical scenario given above with a positive sexual history, Romberg test, and Argyll Robertson pupils is suggestive of neurosyphilis. The investigation of choice for neurosyphilis is VDRL.

The CSF VDRL test is highly specific and, when reactive, is considered diagnostic of neurosyphilis.

Involvement of the CNS in syphilis is detected by examination of CSF for:

• Mononuclear pleocytosis (>5 white blood cells/µL).

• Increased protein concentration (>45 mg/dL).

• CSF VDRL reactivity.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

The investigation used to assess response to treatment in syphilis is VDRL. In the case of secondary syphilis, follow-up VDRL should be done at 6 months and 12 months.

VDRL and RPR become negative after adequate treatment whereas specific treponemal tests remain positive indefinitely despite treatment.

Guidelines for follow up:

• Patients with primary or secondary syphilis should be examined 6 and 12 months after treatment.

• Patients with latent or late syphilis should be examined at 6, 12, and 24 months after treatment.

Solution to Question 9:

Pregnancy can give rise to false-positive VDRL.

Biological false-positive (BFP) reactions are defined as positive reactions obtained in non-specific treponemal tests (RPR, VDRL) with negative results in specific treponemal tests, in the absence of past or present treponemal infections and not caused by technical faults.

Solution to Question 10:

The newborn baby has characteristic signs and symptoms of congenital syphilis. Periostitis is a hallmark of skeletal manifestation. VDRL or RPR is the initial investigation of choice in congenital syphilis.

The earliest manifestations of congenital syphilis include rhinitis, or “snuffles”; mucocutaneous lesions, bone changes, including periostitis detectable by x-ray examination of long bones; hepatosplenomegaly, lymphadenopathy, anemia, jaundice, thrombocytopenia, and leukocytosis.

All neonates born to mothers who have reactive nontreponemal and treponemal test results should be evaluated with a quantitative nontreponemal serologic test (RPR or VDRL) performed

Sold by @itachibot

on the neonate's serum, because umbilical cord blood can become contaminated with maternal blood and yield a false-positive result.

FTA-ABS or TPHA is used for confirmation of syphilis in a patient with reactive RPR or VDRL test.

T. pallidum cannot be detected by culture.

Solution to Question 11:

Causative agent of yaws is Treponema pertenue.

It is one of the endemic treponematoses that spreads by direct contact.

The image shows ulcerative skin lesions seen in yaws.

Causative agent	Disease
T. pertenue	Yaws
T. pallidum	Venereal syphilis
T. carateum	Pinta
T. endemicum	Endemic syphilis

Solution to Question 12:

Epidemic relapsing fever is caused by Borrelia recurrentis and is transmitted by louse.

Sold by @itachibot

Endemic relapsing fever is caused by Borrelia duttoni, Borrelia hermsii, Borrelia parkeri, etc, and is transmitted by ticks.

Solution to Question 13:

The above clinical scenario is suggestive of louse-born relapsing fever (LBRF) caused by Borrelia recurrentis. Barbour-Stoenner-Kelly broth medium is used to isolate all Borrelia species except for B.miyaotoi.

Tender splenomegaly and relapsing fever in the background of a louse outbreak point towards the diagnosis of LBRF.

It was very common during wars and in the jails of former days but with improvements in hygiene and the discovery of insecticides, it has now become rare.

Solution to Question 14:

Borrelia recurrentis uses antigenic variation as a major means of evading the host immune system. It causes relapsing fever.

Borrelia readily undergoes antigenic variations in vivo and this is believed to be the reason for the occurrence of relapses in the disease. Antigenic variations occur because of DNA rearrangements in linear plasmids present in borrelia.

Solution to Question 15:

The image shows erythema migrans, seen in Lyme disease. It is caused by Borrelia Burgdoferi.

The lesion has a target center with partial central clearing and a bright red outer border. It usually begins as a red macule or papule at the site of the tick bite.

The tick implicated is the black-legged tick, Ixodes scapularis, shown in the image below.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

The most severe manifestation of Lyme disease is Lyme neuroborreliosis (LNB), reported in 10–15% of individuals with localized erythema migrans. The activation of the inflammatory response in LNB contributes to the pathogenesis of a broad spectrum of neurologic disorders.

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

The electron micrograph shows the spirochete, Leptospira interrogans, which causes the water-borne disease, leptospirosis. Rats are the most important reservoir.

Transmission of leptospires may follow direct contact with urine, blood, or tissue from an infected animal or exposure to environmental contamination.

Solution to Question 17:

The above clinical scenario suggests a diagnosis of Weil's disease which is caused by leptospira.

Long hours of working in a field with water (paddy field), is a high-risk factor for the development of leptospirosis.

Weil's syndrome is the severe form of leptospirosis that encompasses the triad of hemorrhage, jaundice, and acute kidney injury.

The treatment of choice is IV penicillin or IV ceftriaxone.

Solution to Question 18:

Ellinghausen, McCullough, Johnson, Harris (EMJH) media is the culture medium used for leptospira.

Sold by @itachibot

Other culture media that can be used are Korthof's, Stuart's, and Fletcher's media.

The image below shows EMJH medium, with Dinger's ring/zone. The growth of the leptospires is at a maximum density at this zone. It is identified as an area of increased turbidity, beneath the surface of the semi-solid medium. It is the zone of optimum oxygen tension for the growth of the bacteria.



Solu... if you purchased this ... you may have been ... else, ... d.

Solution to Question 19:

History of fever with liver and kidney involvement should arouse suspicion for leptospirosis (Weil's disease). A microscopic agglutination test is used for the diagnosis of leptospirosis.

Serotype-specific tests:

• Microscopic agglutination test

• Macroscopic agglutination test

Option A: Widal test is used to screen for typhoid fever

Option C: Weil-Felix reaction is used to diagnose rickettsial fevers

Option D: Paul-Bunnell test is the investigation of choice for infectious mononucleosis.

Solution to Question 20:

The most likely diagnosis based on the clinical vignette is leptospirosis.

Leptospirosis is a zoonosis caused by spirochetes of the genus Leptospira. Patients with mild disease may present with flu-like symptoms of sudden onset, fever with chills, headache, abdominal pain, conjunctival suffusion, and myalgia. The muscle pain usually affects the calves, abdomen, and back. Mild cases can be treated with oral doxycycline, ampicillin, or amoxicillin.

Sold by @itachibot

Severe leptospirosis is rapidly progressive and is associated with high fatality. The classic presentation of the severe disease called Weil's syndrome includes the triad of hemorrhage, jaundice, and acute kidney injury. Jaundice when present can be profound and physical examination may reveal an enlarged, tender liver. Temperature-pulse dissociation /relative bradycardia can be seen. Moderate to severe cases are treated with intravenous penicillin, ampicillin, or ceftriaxone.

The image below shows icterus and conjunctival suffusion in a patient with leptospirosis.



Leptospira are actively motile organisms and are best seen using a dark-field microscope. They are thin, tightly coiled, and flexible spirochetes with one end forming a hook. Diagnosis is by microscopic agglutination test.

Transmission occurs through cuts or breaks in the skin and mucous membranes (conjunctival and oral mucosa). Several outbreaks have been reported that resulted from exposure to floodwaters contaminated by urine from infected animals. The organism enters the bloodstream and lodges in the parenchymatous organs (liver and kidneys) leading to hemorrhage and necrosis of the tissue

Sold by @itachibot

causing dysfunction of the organs.

The lifecycle of Leptospira is shown below.



Illness is classically biphasic after an incubation period of up to 1 month. The stages are:

• Acute leptospiremic phase- the organism can be cultured from blood.

• Resolving immune phase- the organism can be cultured from urine.

The stages may not be clinically distinguishable.

Other options:

Option A - Chikungunya is a viral disease that presents with saddle-back fever, conjunctival suffusion, and photophobia. Migratory arthralgia, especially involving small joints may also be seen. A maculopapular or petechial rash is frequently seen on the trunk.

Option C - Dengue hemorrhagic fever is a form of severe dengue with bleeding manifestations. Evidence of plasma leakage and thrombocytopenia is also generally seen.

Option D - Encephalitis due to Hepatitis A is seen in fulminant hepatitis, a rare complication. Patients present with confusion, disorientation, ascites, and edema.

Sold by @ If you purchased this from you may have been scammed. else,

Sold by @itachibot

General Properties of Viruses

Question 1:

What is the protein coat covering the nucleic acid in a virion known as?

a) Capsid

b) Peplomer

c) Envelope

d) Tegument

Question 2:

Which of the following viruses has a single-stranded DNA?

a) Poxvirus

b) Parvovirus

c) Picornavirus

d) Reovirus

Question 3:

Which of the following is false regarding enveloped viruses?

a) Ether sensitive

b) Heat labile

c) Susceptible to bile salts

d) Lipid is viral coded

Question 4:

Which of the following is an enveloped virus?

a) Reovirus

b) Hepatitis A virus

c) Hepatitis C virus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

d) Adenovirus

Question 5:

A patient in the ICU requires fresh frozen plasma (FFP). The FFP that arrives from the blood bank is labeled as Solvent/Detergent plasma, which destroys lipid-enveloped viruses. Which of the following viral infections is the patient still susceptible to?

a) Human immunodeficiency virus

b) Hepatitis A virus

c) Hepatitis B virus

d) Hepatitis C virus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

In which of the following viruses is a segmented genome found?

a) Filovirus

b) Orthomyxovirus

c) Retrovirus

d) Poliovirus

Question 7:

A 6-year-old with HIV is brought with complaints of the rash shown below. Where does DNA replication occur in the viruses falling in the same family as the microbe causing this rash?

Sold by @itachibot

a) Host cytoplasm

b) Host nucleus

c) Viral cytoplasm

d) Viral nucleus

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

An epidemic involving an ss-RNA virus with 3'-5' RNA has broken out in West Africa. Which of the following enzymes do such viruses use in genome replication?

a) DNA-dependent RNA polymerase

b) RNA replicase

c) Reverse transcriptase

d) Taq polymerase

Question 9:

The virus shown below was suspected to be the cause of a deadly hemorrhagic fever outbreak in Guinea in March 2021. How is it transmitted?

Sold by @itachibot



a) Direct contact with infected body fluids

b) Mosquito-borne

c) Food-borne

d) Air-borne

Sold by @itachi... purchased this from some... else, you may have been scammed.

Question 10:

An elderly diabetic man presents with a complaints of fever, cough and difficulty in breathing for the past 3 days. He gives a history of international travel 10 days back. Laboratory reports show positive for COVID-19. What is the symmetry of the causative virus?

a) Naked helical symmetry

b) Icosahedral symmetry

c) Enveloped helical symmetry

d) Complex symmetry

Question 11:

Which of the following statements is false regarding symmetry of viruses ?

a) Viral symmetry is studied by heavy metals stains.

b) The most efficient closed shell sub-unit arrangement is icosahedral.

c) Icosahedral symmetry involves 12 faces and 20 vertices .

d) It is not possible for 'empty' helical particles to form

Sold by @itachibot

Question 12:

Which of the following viruses shows an enveloped icosahedral symmetry?

a) Rubella virus

b) Influenza virus

c) Measles virus

d) Poxvirus

Question 13:

An immunologist is working on growing the influenza virus for vaccine production at the National Institute of Virology, Pune. Which of the following sites is used for the given purpose in a fertilized egg?



a) 4

b) 1

c) 2

d) 3

Question 14:

Quantitative infectivity assay of viruses is done by _____.

Sold by @itachibot

a) Egg inoculation

b) Hemadsorption

c) Plaque assay

d) Electron microscopy

Question 15:

Which virus causes the cytopathic effect of syncytium formation?

a) SV 40

b) Enteroviruses

c) Measles virus

d) Adeno virus

Sold by @Itachibot
If you bought this from someone else, you've been scammed.

Answer Key

Question No.	Correct Option
1	a
2	b
3	d
4	c
5	b
6	b
7	a
8	b
9	a
10	c
11	c
12	a
13	b
14	c
15	c

Sold by @itachibot

Detailed Explanations

Solution to Question 1:

The protein coat covering the nucleic acid in a virion is called capsid.

The parts of a virus include:

• Nucleocapsid- capsid with enclosed nucleic acid

• Tegument- proteinaceous structure present between the envelope and capsid.

• Envelope- lipid-containing membrane that surrounds some virus particles. It is acquired during viral maturation by a budding process through a cellular membrane.

• Peplomers- virus-encoded glycoproteins, on the outer surface of the envelope as projections.



Characteristics of viruses:

• Do not have a cellular organization

• Have either DNA or RNA but never both

• Obligate intracellular parasites

• Lack enzymes necessary for protein and nucleic acid synthesis

• Filterable

• Cannot be grown on artificial media

• Unaffected by antibiotics.

Note : Martinus beijerinck proposed the theory of contagium vivum fluidum (living infectious fluid) while studying the cause of mosaic disease of tobacco plant. He is known as the father of virology.

Sold by @itachibot

Solution to Question 2:

Parvovirus has single-stranded DNA.

All DNA viruses are double-stranded, except parvovirus. (only single-stranded DNA virus)

All RNA viruses have only one copy of single-stranded RNA except:

• Reovirus- double-stranded RNA virus.

• Retroviruses- Two copies of single-stranded RNA.

Parvovirus is the smallest DNA virus that infect human beings.

	DNA	RNA
Largest virus	Poxvirus	Filovirus
Smallest virus	Parvovirus	Picornavirus

If you purchased this from someone else, you have been scammed. Sold by @itachibot

Solution to Question 3:

The lipid of enveloped viruses is largely of host-cell origin, whereas protein is virus coded.

The following are the differences between enveloped and non-enveloped viruses.

Enveloped viruses	Non-enveloped viruses
Ether sensitive	Ether resistant
Heat labile	Heat stable
Pleomorphic	Non-pleomorphic
Susceptible to bile salts	Resistant to bile salts

Solution to Question 4:

Hepatitis C virus is an enveloped virus.

It has a linear, single-stranded RNA genome, enclosed within a core and surrounded by an envelope.

Solution to Question 5:

The patient is still susceptible to hepatitis A infection. It is a non-enveloped virus, and hence it's not affected by SD plasma.

Sold by @itachibot

Solvent/Detergent plasma (SD plasma) is a form of blood plasma made from plasma collected from many people which is then processed with solvents in an attempt to remove viruses.

Only the viruses with an envelope are susceptible to SD plasma destruction. Hepatitis B, C, and Human immunodeficiency virus (HIV) are enveloped.

Solution to Question 6:

Segmented genome is found in orthomyxovirus.

Segmentation of viral genomes allows for the exchange of intact genes between related viruses during coinfection. This recombination is known as reassortment. There is a rearrangement and joining of nucleic acid sequences to form a unique chimeric genome. The new genome is usually less fit than the parent viruses, but occasionally, it results in an evolved virus with increased fitness.

Examples of segmented viruses (ARBO):

- Arenavirus (2 segments)
- Rotavirus (11 segments)
- Bunyavirus (3 segments)
- Orthomyxovirus (8 segments)

The image below shows the segmented genome seen in the influenza virus (Orthomyxoviridae)



Solution to Question 7:

The image shows pink, fleshy umbilicated papules suggesting molluscum contagiosum. It is caused by the Molluscum contagiosum virus belonging to Poxviridae. DNA replication in these viruses occurs in the host cytoplasm.

Sold by @itachibot this from someone else? Been scammed.

Sold by @itachibot

	DNA viruses	RNA viruses
Replication	Nucleus	Cytoplasm
Exception	Poxvirus- cytoplasm	Orthomyxovirus, retrovirus- nucleus

Solution to Question 8:

An ss-RNA with a 3'-5' strand is a negative-stranded RNA. These viruses use RNA replicase or RNA-dependent RNA polymerase during genome replication.

In positive-stranded RNA viruses, the viral RNA itself acts as mRNA.

In negative-stranded RNA viruses, the polarity is opposite that of mRNA. So the genome requires RNA-dependent RNA polymerase to synthesize their mRNA.

Examples of negative-stranded RNA viruses (Mnemonic: Always bring da polymerase or fail to replicate):

- Arenaviruses
- Bunyaviruses
- Deltaviruses
- Paramyxoviruses
- Orthomyxoviruses
- Filoviruses
- Rhabdoviruses

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

The image shown above is the Ebola virus. It is spread by direct contact with infected body fluids.

Ebola is a filamentous virus with a single-stranded RNA genome.

The shapes of some viruses are shown below.

Sold by @itachibot





Rotavirus: Wheel-shaped virus

Sold by @itachibot



Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Solution to Question 10:

The patient is suffering from coronavirus disease 2019 (COVID-19). The coronavirus implicated shows an enveloped helical symmetry.

Properties of Coronaviridae family:

• They are enveloped RNA viruses.

• They contain a genome of unsegmented single-stranded and positive-sense RNA

• They have helical nucleocapsid

Solution to Question 11:

Icosahedron has 20 faces (each an equilateral triangle) and 12 vertices.

Option A: Study of viral symmetry by standard electron microscopy requires the use of heavy metal stains (e.g., potassium phosphotungstate) to emphasize surface structure. It is a form of negative staining.

Option B: Symmetry observed with animal viruses is of the icosahedral pattern. It is the most efficient arrangement for subunits in a closed shell.

Option D: A regular, periodic interaction between capsid protein and nucleic acid is required to maintain a stable helical symmetry. Therefore, it is not possible for empty helical particles to form.

Given below is Icosahedron:

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 12:

Enveloped icosahedral symmetry is seen in the rubella virus.

Solution to Question 13:

Allantoic cavity inoculation (1) is used for the growth and replication of the influenza virus for vaccine production.

Embryonated eggs are used for the cultivation of viruses.



Sold by @itachibot

Note: Both allantoic and amniotic cavity inoculation are recommended for initial isolation of the influenza virus. However, it is simpler to inoculate the virus into the allantoic cavity, as compared to the amniotic cavity, thus allantoic inoculation is the best answer.

Solution to Question 14:

Quantitative infectivity assay of viruses is done by plaque assay.

Assay of infectivity of viruses

Physical methods	Biological methods
Can estimate total viral count	Can estimate total viral count
Cannot differentiate between infectious and noninfectious virus particles	Can detect infectious virus
Examples:Real-time PCRAntigen detection assayElectron microscopy	Examples:Qualitative- endpoint biological assayQuantitative- plaque and pock assay

The image below shows a plaque assay.



The image below shows pock assay.

Sold by @itachibot



Solution to Question 15:

The cytopathic effect of syncytium formation is caused by the measles virus.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

DNA Viruses

Question 1:

A 23-year-old lady presents with the lesions shown below. What is the causative organism?

a) Herpes simplex virus type 1

b) Epstein-Barr virus

c) Cytomegalovirus

d) Herpes simplex virus type 2

Question 2:

Cowdry Type A inclusions are seen in:

a) Herpes simplex

b) Rabies

c) Molluscum contagiosum

d) Adenovirus

Question 3:

Sold by @itachibot

A 13-year-old girl has had 4 episodes of meningitis over the last 4 years, with the disease lasting a few days at a time, leaving slowly resolving neurological symptoms such as visual and memory loss. A CSF cytology shows lymphocytic pleocytosis and a diagnosis of Mollaret's meningitis is made. Which organism is implicated here?

a) Measles virus

b) Mumps virus

c) Herpes simplex virus

d) Rubella virus

Question 4:

A dentist presents to the surgery OPD with the findings shown below. He is febrile. Microscopic examination of the tissue reveals Tzanck cells. What is the causative agent?

a) Pox virus

b) Adenovirus

c) Picornavirus

d) Herpes simplex virus

Question 5:

A young male presents to the STD clinic with painful lesions as shown in the image. What is the most likely microscopic finding of the lesion?

Sold by @itachibot

a) Tzank cells

b) Pund cells

c) Henderson-Patterson bodies

d) Guarnieri bodies

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 6:

Your neighbor's child complains of fever and develops a rash. On examination, the rashes are present all over the body, and look like 'dewdrops lying on the skin'. The child says that his friend in school had similar complaints a few days ago. Which of the given statements is true about this rash?

a) It is centripetal in distribution

b) The rash usually starts on the distal parts of limbs

c) All the lesions are of same age

d) Scabs are also infectious in chicken pox

Question 7:

While on rounds in the NICU, you notice a newborn baby with a zig-zag dermatomal cicatrix on the forehead and left side of the face. The limbs are small and atrophied. Microcephaly is noted. Which maternal infection are these findings characteristically associated with?

a) Toxoplasma

b) Cytomegalovirus

Sold by @itachibot

c) Varicella zoster virus

d) Herpes simplex virus type - 1

Question 8:

Which of the following is the predominant pathogen to infect renal transplant recipients one to four months following transplantation?

a) Cytomegalovirus

b) JC virus

c) BK virus

d) Epstein-Barr virus

Question 9:

An infant is brought by his worried parents with a rash and yellowish discoloration of the skin. On examination, the baby has petechiae, jaundice, hepatosplenomegaly, and microcephaly. The developmental assessment shows global developmental delay. An axial cut of the non-contrast CT of the brain is given below. What is the causative organism?

a) Rubella

b) Herpes simplex virus

c) Cytomegalovirus

d) Varicella

Sold by @itachibot
If you bought this from someone else, it was uncommed.

Sold by @itachibot

Question 10:

Choose the correct statement regarding CMV.

a) Ganciclovir resistance is due to mutation in UL97 phosphotransferase gene

b) CMV can be cultured in vitro in any mammalian cell

c) Commonest manifestation of CMV associated disease is watery diarrhea

d) UL54 mutated CMV strains do not show cross resistance to Foscarnet

Question 11:

Where are owl's eye inclusion bodies seen?

a) Yellow fever

b) Cytomegalovirus

c) Adenovirus

d) Herpes simplex virus

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

What does the Paul-Bunnell test diagnose?

a) Kissing disease

b) Hand-foot-and-mouth disease

c) Barlow's disease

d) Fifth disease

Question 13:

You are studying the association of human herpesvirus-4 with other diseases. Which of the following patients have diseases associated with this virus?

a) 1 and 5 only

b) 2, 3 and 4

c) 3 and 4 only

d) 1, 2 and 5

Sold by @itachibot

Question 14:

Which of the following viruses is known to cause hemorrhagic cystitis, diarrhea, and conjunctivitis?

a) RSV

b) Adenovirus

c) Rhinovirus

d) Rotavirus

Question 15:

Viral infection mimicking erythroblastosis fetalis is:

a) HSV

b) Epstein-Barr virus

c) Picornavirus

d) Parvovirus B19

Question 16:

Which organism causes the disease shown below?

a) Roseola virus

b) Parvovirus B19

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

c) Rubella virus

d) Rubeola virus

Question 17:

An 8-year-old child presents with skin lesions, suspected to be molluscum contagiosum. How is the diagnosis of this condition typically made?

a) Viral Culture

b) Skin biopsy

c) PCR for Viral DNA

d) Clinical appearance

If yo ld by @itachibot is from someone else, scammed.

Question 18:

What is the causative organism of the condition shown below?

a) Treponema pallidum

b) Human papillomavirus 6 and 11

c) Human papillomavirus 16 and 18

d) Hemophilus ducreyi

Question 19:

Sold by @itachibot

A patient presents with plantar warts as shown below. Which type of human papillomavirus is responsible for causing these?

a) 6

b) 11

c) 1

d) 7

Sold by @ If you purchased this from you may have been scammed. else,

Question 20:

As a CHC medical officer, you organize an outreach program for HPV vaccination and the prevention of cervical cancer. Which protein is used for antigenic response in this vaccine?

a) E6, E7

b) L1

c) E1, E2

d) L2

Question 21:

Nonavalent HPV vaccine works against which of the following strains of the virus?

a) 1, 6, 11, 16, 18, 31, 45, 52, 58

b) 6, 11, 16, 18, 31, 33, 45, 52, 58

c) 1, 16, 18, 21, 33, 45, 48, 52, 58

Sold by @itachibot

d) 6, 11, 16, 18, 33, 45, 48, 52, 58

Question 22:

An HIV-positive patient presents with the following skin lesions. Which of the following diseases is associated with the virus implicated here?

a) Post-transplant lymphoproliferative disease

b) Merkel cell carcinoma

c) Adult T-cell leukemia

d) Primary effusion lymphoma

Question 23:

Which of the following viruses is implicated in progressive multifocal leukoencephalopathy?

a) JC virus

b) BK virus

c) Measles virus

d) Herpes simplex virus

Answer Key

Sold by @itachibot

Question No.	Correct Option
1	a
2	a
3	c
4	d
5	a
6	a
7	c
8	a
9	c
10	a
11	b
12	a
13	d
14	b
15	d
16	b
17	d
18	b
19	c
20	b
21	b
22	d
23	a

Detailed Explanations

Solution to Question 1:

The lesions shown are cold sores, also known as herpes labialis caused by Herpes simplex virus type 1 (HSV-1). Herpes labialis primarily affects the lip and typically present as thin-walled, umbilicated vesicles, the roof of which breaks down with superficial ulcers. They generally heal without scarring.

HSV-1 can be isolated from lesions in and around the mouth and is transmitted by direct contact or droplet spread from cases or carriers. HSV-2 is responsible for the majority of genital herpes infections and is commonly transmitted venereally.

The images below show lesions of herpes labialis.

Sold by @itachibot

Solution to Question 2:

Cowdry type A inclusions are seen in the vesicles of Herpes Simplex infection

The classic Cowdry type A inclusion appears as a light purple, homogeneous intranuclear structure surrounded by a clear halo. The inclusions stain with antibodies to HSV, permitting a rapid, specific diagnosis of HSV infection.

Sold by @itachibot

HSV inclusion bodies



©MARROW

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 3:

Mollaret's meningitis or benign recurrent lymphocytic meningitis is most commonly caused by the herpes simplex virus (HSV). Demonstration of HSV antibodies in CSF or persistence of HSV DNA in CSF can establish the diagnosis.

Solution to Question 4:

The image suggests herpetic whitlow, confirmed by the presence of Tzanck cells on microscopy. It is caused by HSV-1.

It is often contracted by health care workers that come in contact with the virus when exposed to oral secretions.

On Tzanck smear, which is a rapid, diagnostic method, Tzanck cells are seen. These are multinucleated giant cells with faceted nuclei and homogenously stained ground-glass chromatin. Tzanck smear is also used in the diagnosis of Hailey-Hailey disease, Darier's disease, hand-foot-and-mouth disease, etc.

Sold by @itachibot

Diagnosis of herpes simplex virus:

• Most sensitive test- HSV DNA- PCR (can differentiate between HSV 1 and 2)

• Most definitive test- virus isolation using human diploid fibroblast in which cytopathic changes can be observed as early as 24 - 48 hours.

• Microscopy - Tzanck smear is prepared by Wright or Giemsa stain. Intranuclear type A inclusion bodies may be seen in Giemsa-stained smears.

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 5:

Clusters of vesicles are seen on an erythematous base along with painful and pruritic lymphadenopathy suggest a diagnosis of genital herpes. Tzanck cells or multinucleated giant cells are present on microscopy.

The Tzanck smear is a rapid, sensitive, and inexpensive diagnostic method. Smears are prepared from the lesions, from the base of vesicles, and stained with a 1% aqueous solution of O-toluidine blue for 15 seconds. Multinucleated giant cells with faceted nuclei and homogeneously stained ground glass chromatin - Tzanck cells are seen.

The typical herpes lesions are thin-walled, umbilicated vesicles, the roof of which breaks down, leaving tiny superficial ulcers. They heal without scarring. The most common site of cutaneous lesions in the face - on the cheeks, chin, around the mouth, or on the forehead.

Sold by @itachibot

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 6:

The condition described here is chickenpox. It is caused by the varicella-zoster virus. The following are the characteristics of rash seen in chicken pox :

• The rash is centripetal in distribution.

• It evolves rapidly, progressing through macule, papule, vesicle, pustule, and finally scab.

• It is a pleomorphic rash and appears in crops, hence varying stages of lesions can be noticed at a given time.

• The lesions resemble a dewdrop lying on the skin, as they are superficial and do not involve the deeper layers of the skin.

Sold by @itachibot

Solution to Question 7:

The given clinical scenario is suggestive of congenital varicella syndrome. Hence, an antenatal history of varicella zoster virus (VZV) infection would be significant in the mother of this baby.

Fetal varicella infection occurs if the mother gets infected during the first half of pregnancy.

Clinical features include:

- Cicatrix, a zigzag scarring in a dermatomal distribution
- Microphthalmia
- Cataracts
- Optic atrophy
- Chorioretinitis
- Microcephaly
- Limb hypoplasia
- Motor and sensory deficits

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 8:

The predominant pathogen to infect renal transplant recipients in the period around 1 to 4 months following the transplantation is cytomegalovirus (CMV).

The most common CMV presentation in kidney transplant patients are fever (common), leukopenia, hepatosplenomegaly, myalgia, and arthralgia.

JC virus and BK virus are common infections after solid organ transplantation based on the site of infection. This means BK and JC viruses are the most common organisms to infect the kidney after any solid organ transplant. But CMV is the predominant organism to cause infection in a person who has undergone a renal transplant. (Not the most common organism to affect the kidney after any solid organ transplant).

Because of the frequency and severity of CMV disease, antiviral prophylaxis with ganciclovir or valganciclovir is being used in renal transplant recipients whenever indicated to prevent CMV disease.

Solution to Question 9:

The clinical features and imaging suggest a diagnosis of congenital CMV infection. The non-contrast CT head shows periventricular calcifications and ventriculomegaly.

Features of congenital CMV infection:

- Petechial rash
- Jaundice

Sold by @itachibot

• Hepatosplenomegaly
• Microcephaly
• Deafness
• Chorioretinitis
• IUGR
• Periventricular calcification and ventriculomegaly



Option A: Congenital rubella syndrome is a triad of cardiac defects, cataracts, and congenital hearing loss.

Option B: Neonatal herpes presents as skin lesions with eye involvement, encephalitis, and disseminated disease.

Option D: Fetal varicella syndrome is manifested as cicatrizing/scarring skin lesions, chorioretinitis, hypoplastic limbs, low birth weight, and CNS defects due to cerebral atrophy.

Note: In 15-20% of infants with congenital CMV infection, microcephaly is not detected at birth. It may develop as the child grows and is detected at follow-up visits for other complaints.

Solution to Question 10:

Resistance to ganciclovir most commonly occurs due to mutations in the UL97 phosphotransferase gene.

Option B: Human CMV replicates in vitro only in human fibroblasts, although the virus is often isolated from epithelial cells of the host.

Option C: CMV infection is mostly subclinical.

Option D: UL54 mutations can confer resistance to ganciclovir, cidofovir, and foscarnet.

Sold by @itachibot

Solution to Question 11:

Owl's eye inclusion bodies are intranuclear basophilic inclusion bodies seen in CMV infection.



Solution to Question 12:

Paul-Bunnell test is used to diagnose kissing disease (infectious mononucleosis) caused by Epstein–Barr virus (EBV).

It is a heterophile agglutination test used to identify the heterophile antibodies that agglutinate sheep erythrocytes. Patients suffering from infectious mononucleosis due to Epstein–Barr virus (EBV) will produce antibodies that will react and agglutinate sheep erythrocytes.

Solution to Question 13:

Human herpesvirus-4 or Epstein-Barr virus causes infectious mononucleosis, Burkitt's lymphoma, and is associated with oral hairy leukoplakia in HIV patients.

EBV may cause the following malignancies:

• Burkitt's lymphoma

• Nasopharyngeal carcinoma (in people of Chinese origin),

• Lymphomas in immunodeficient persons such as AIDS or transplant recipients.

Endemic or African Burkitt's lymphoma has got nearly 100% association with EBV, as compared to non-endemic and HIV-associated Burkitt's lymphoma.

Oral hairy leukoplakia is triggered by EBV in patients with poor immune status, most commonly with HIV.

Oral submucous fibrosis is associated with the human papillomavirus (HPV).

Sold by @itachibot
If you purchased this from, you may have been scam, else,

Sold by @itachibot

Adult T cell leukemia is associated with the human T-cell leukemia virus.

Solution to Question 14:

Adenovirus is known to cause hemorrhagic cystitis, diarrhea, and conjunctivitis.

Acute hemorrhagic cystitis in children is caused by adenoviruses types 11 and 21.

The following table shows various syndromes caused by adenoviruses and associated serotypes:

Syndrome	Adenovirus serotypes
Respiratory disease in children	1, 2, 5, 6
Sore throat, febrile cold, pneumonia	3, 4, 7, 14, 21
ARD in military recruits	4, 7, 21
Follicular (swimming pool) conjunctivitis/ Pharyngoconjunctival fever	3, 7
Epidemic keratoconjunctivitis (shipyard eye)	8, 19, 37
Diarrhea	40, 41

Solution to Question 15:

Parvovirus B19 infection mimics erythroblastosis fetalis.

Parvovirus B19 is associated with:

• Erythema infectiosum (fifth disease)

• Transient aplastic crisis

• Pure red cell aplasia

• Erythroblastosis fetalis (most commonly in early pregnancy)

Picornavirus, Epstein-Barr virus, HSV infection do not mimic erythroblastosis fetalis.

Solution to Question 16:

The image showed a rash on the cheeks and face, with a slapped cheek appearance, characteristic of erythema infectiosum/ fifth disease. It is caused by Parvovirus B19.

Sold by @itachibot

Parvovirus B19 usually presents as a respiratory infection, with an erythematous maculopapular rash and arthralgia. The rash starts as prominent erythema of the cheek, popularly known as slapped-cheek appearance. The rash further spreads to the trunk and limbs followed by lymphadenopathy and arthralgia.

It induces aplastic crisis in sickle cell anemia patients.

Infection during pregnancy leads to non-immune fetal hydrops (highest risk in the first half of gestational period).

Solution to Question 17:

Diagnosis of molluscum contagiosum is based on clinical appearance.

Molluscum contagiosum manifests as pink pearly wart-like lesions with a characteristic dimple at the center (umbilicated), not associated with inflammation and necrosis.

Option B: Skin biopsy or smear of expressed material shows characteristic inclusion bodies but is necessary only when the diagnosis is uncertain.

Options A and C: Viral culture and PCR for viral DNA are not used for diagnosis.

Solution to Question 18:

The image shows condyloma acuminata. It is most commonly caused by Human papillomavirus (HPV) 6 and 11.

Condyloma acuminata is a benign, genital wart which is a moist, soft, pedunculated wart found on the external genitalia.

Microscopically, the characteristic features include:

Sold by @itachibot

• A branching villous structure with connective tissue stroma

• Epidermis showing hyperkeratosis and acanthosis (thickening)

• Cytoplasmic vacuolization of the squamous cells (koilocytosis) is characteristic of HPV infection.

• No evident dysplasia



Option A: Treponema pallidum can cause condylomata lata in secondary syphilis. It is smoother and flatter and may be associated with induration and ulceration. (image below)

Option C: Human papillomavirus 16 and 18 are associated with carcinoma of the cervix and anogenital region.

Option D: Hemophilus ducreyi causes chancroid. Given below is the image of chancroid.

Sold by @itachibot

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 19:

HPV types 1a, 57, and 27 accounts for most plantar warts.

HPV 7 causes hand warts in butchers (butcher's warts).

HPV 6 and 11 cause condyloma acuminata, sinonasal papilloma, and squamous papilloma of the larynx.

Solution to Question 20:

The HPV vaccines use virus-like particles (VLPs) that consist of the HPV L1 major capsid protein. The L1 protein self-assembles into VLPs when expressed in eukaryotic cells.

These VLPs contain the same epitopes as the HPV virion. But they do not contain genetic material and cannot transmit infection. The immunogenicity of the HPV vaccines relies on the development of conformational neutralizing antibodies to epitopes displayed on viral capsids.

The nonavalent HPV vaccine works against strains 6, 11, 16, 18, 31, 33, 45, 52 and 58. It is usually commercialized in the name of Gardasil-9. It has received expanded indication which now includes the prevention of oropharyngeal and other head and neck cancers.

HPV Vaccine	Types
Quadrivalent (Gardasil)	6, 11, 16, 18
Bivalent (Cervarix)	16, 18
Nonavalent (Gardasil-9)	6, 11, 16, 18, 31, 33, 45, 52, 58

Sold by @itachibot

Solution to Question 21:

The nonavalent HPV vaccine works against strains 6, 11, 16, 18, 31, 33, 45, 52 and 58. It is usually commercialized in the name of Gardasil-9. It has received expanded indication which now includes the prevention of oropharyngeal and other head and neck cancers.

HPV Vaccine	Types
Quadrivalent (Gardasil)	6, 11, 16, 18
Bivalent (Cervarix)	16, 18
Nonavalent (Gardasil-9)	6, 11, 16, 18, 31, 33, 45, 52, 58

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

The above image showing red-purple macules and plaques in an HIV-positive patient is suggestive of Kaposi sarcoma, caused by human herpesvirus 8. It also causes primary effusion lymphoma. It is one of seven currently known human cancer viruses or oncoviruses.

HHV-8 or Kaposi's sarcoma-associated herpesvirus (KSHV) is known to cause Kaposi's sarcoma, in immunocompromised persons. Cancer patients receiving chemotherapy, AIDS patients, and organ transplant recipients are all at a high risk of showing signs of infection.

KSHV is a known causative agent of four diseases:

- Kaposi's sarcoma
- KSHV-associated multicentric Castleman's disease (MCD)
- Primary effusion lymphoma
- KSHV inflammatory cytokine syndrome

Solution to Question 23:

Progressive multifocal leukoencephalopathy is caused by the JC virus. JC virus-induced progressive multifocal leukoencephalopathy is one of the neurologic diseases associated with HIV infection.

Option B: BK virus is associated with nephropathy in transplant recipients.

Option C: The long-term sequelae associated with measles infection is subacute sclerosing panencephalitis.

Option D: Herpes simplex virus causes temporal lobe encephalitis.

Sold by @itachibot

Hepatitis

Question 1:

Which of the following is a non-enveloped virus that causes hepatitis?

a) Hepatitis C

b) Hepatitis B

c) Hepatitis A

d) Hepatitis D

Question 2:

A 25-year-old man hailing from Mumbai presents to you with complaints of abdominal pain and malaise. On examination, he is icteric and the liver is palpable 6 cm below the costal margin. On further questioning, he informs you that a few weeks back the drains surrounding his house were overflowing due to the monsoon flooding. Which of the following is the most likely etiology?

a) Hepatitis A or D

b) Hepatitis B or E

c) Hepatitis A or B

d) Hepatitis A or E

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

A 18-year-old man presents with fever, diarrhea, and jaundice. He gives a history of consuming food at a roadside stall. Which of the following statements is incorrect about the etiological agent?

a) It belongs to family Hepatoviridae

b) Majority of infections are symptomatic

c) It is transmitted by feco-oral route

d) A formalin inactivated vaccine is used for prevention

Question 4:

Sold by @itachibot

What is the average incubation period for Hepatitis A virus?

a) <1 week

b) 30 days

c) 90 days

d) 120 days

Question 5:

Which of the following is called as Australia antigen?

a) HBsAg

b) HBeAg

c) HBcAg

d) HBV-DNA

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Which of the following is also referred to as Dane particle?

a) HAV

b) HBV

c) HCV

d) HDV

Question 7:

A 35-year-old male patient with chronic hepatitis B has high levels of serum HBeAg. Which of the following viral genes is involved in coding the same?

a) C gene

b) S gene

c) P gene

d) X gene

Question 8:

Sold by @itachibot

While drawing blood from a patient with hepatitis B, it splashes into the eyes of a nurse. Which of the following markers will appear first in her blood if she gets infected?

a) HBeAg

b) HBsAg

c) Anti HBs

d) Anti HBe

Question 9:

A patient with hepatitis B presents to OPD for follow-up. Which of the following serological markers will not be detected in his blood?

a) HBsAg

b) Anti-HBs

c) HBcAg

d) Anti-HBc

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

There are 3 to 5% healthy hepatitis B carriers in India who are asymptomatic. They are at the risk of developing hepatocellular carcinoma in the future because _____

a) They are unable to mount inflammation against the virus

b) Virus can integrate with host DNA and form complementary DNA

c) There is a risk of elevation of transaminases

d) Liver parenchymal cells are in a state of high proliferation

Question 11:

The given life cycle is suggestive of which of the following viruses?

Sold by @itachibot



a) HIV

b) Hepatitis B

c) Influenza

d) Herpes simplex

Question 12:

A patient is diagnosed with Hepatitis B infection. Which of the following markers will always be present in his serum, even during recurrences, or if he develops chronic hepatitis?

a) HBc Ag

b) Anti HBs Ab

c) Anti HBc Ab

d) HBe Ag

Question 13:

Which of the following is not synthesized by pre-core mutants of hepatitis B virus?

a) HBsAg

b) HBxAg

c) HBcAg

d) HBeAg

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Question 14:

The serological profile of a patient is given below. What is your diagnosis?

a) Acute HBV infection with high infectivity

b) HBV vaccination

c) Chronic HBV infection with high infectivity

d) Acute HBV infection with low infectivity

Question 15:

An HBsAg positive primigravida presents to you for routine antenatal care. She asks you about the risk of transmission of HBV to her baby. Which of the following is not related to the same?

a) Intrauterine transmission

b) Normal vaginal delivery

c) Forceps vaginal delivery

d) Breastfeeding

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

An intern takes a hepatitis B vaccine booster dose before his ward postings begin. Which of the following serological markers will indicate immunity against HBV in this case?

a) HBsAg

b) Anti-HBs

c) Anti-HBc

d) Anti-HBe

Question 17:

Which of the following statements is false regarding HBV vaccination?

a) The current vaccine is prepared by cloning 'S gene' of HBV in Baker's yeast

b) A single dose of HBV vaccine is recommended for high-risk groups for pre-exposure prophylaxis

c) For high-risk mothers who are seronegative, hepatitis B vaccine can be given during pregnancy.

Sold by @itachibot

d) Hepatitis B vaccine should be administered intramuscularly in the deltoid muscle.

Question 18:

A known case of hepatitis B going into hepatic failure is found to have a high viral load. Which of the following serological markers will not contribute to the viral load?

a) HBeAg

b) HBV DNA

c) HBV DNA polymerase

d) Anti-HBs

Question 19:

A 30-year-old man with heroin addiction presents with fatigue, palpable purpura, joint pains, and abdominal pain. Urine analysis shows massive proteinuria. Which of the following viruses should be tested for in this patient?

a) Hepatitis A

b) Hepatitis C

c) Hepatitis D

d) Hepatitis E

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 20:

Which hepatitis virus shows maximum chronicity in adults?

a) Hepatitis A

b) Hepatitis E

c) Hepatitis C

d) Hepatitis D

Question 21:

A patient with fulminant hepatitis is found to have coinfection with hepatitis B and hepatitis D viruses. Which of the following structures of HBV aid in the survival of HDV?

a) Reverse transcriptase

Sold by @itachibot

b) Envelope protein

c) Transcriptase to transcribe mRNA

d) Replicase enzyme

Question 22:

In the year 1955, following sewage contamination of drinking water in New Delhi, an outbreak of jaundice ensued. Electron microscopy of fecal samples from the affected population revealed a virus resembling caliciviruses. Which of the following was the etiological agent?

a) Hepatitis C

b) Hepatitis A

c) Hepatitis B

d) Hepatitis E

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Question 23:

Which of the following is false about viral hepatitis?

a) Anti LKM-3 antibody is associated with hepatitis B infection

b) Anti-LKM1 antibodies are associated with hepatitis C infection

c) IgM anti-HAV antibody appears during the late incubation period

d) Enterically transmitted non-A, non-B hepatitis is caused by hepatitis E virus

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	b
5	a
6	b
7	a

Sold by @itachibot

8	b
9	c
10	b
11	b
12	c
13	d
14	a
15	d
16	b
17	b
18	d
19	b
20	c
21	b
22	d
23	a

Sold to ... If you purchased this ... you may have been scammed. ... else,

Detailed Explanations

Solution to Question 1:

Hepatitis A and E are non-enveloped viruses, while Hepatitis B, C, and D are enveloped viruses.

Solution to Question 2:

The given case scenario with icterus, hepatomegaly, and environmental history is suggestive of acute viral hepatitis, likely to be caused by hepatitis A or E viruses.

Hepatitis A and E are transmitted by feco-oral route. Overcrowding, poor hygiene, and fecal contamination of water supplies such as after monsoon flooding can lead to spreading of the disease.

Hepatitis B, C, and D are transmitted by parenteral routes, e.g. blood transfusions.

Solution to Question 3:

The given clinical scenario is suggestive of viral hepatitis which is most likely to be Hepatitis A. Hepatitis A virus belongs to the family Picornaviridae. It is transmitted by feco-oral route and most of the infections are symptomatic.

Sold by @itachibot

There is no definitive treatment, thus, only supportive therapy is given.

Solution to Question 4:

The average incubation period of the Hepatitis A virus (HAV) is around 30 days. It ranges from 15-50 days. The clinical disease has a prodromal stage and an icteric stage.

Solution to Question 5:

HBsAg (Hepatitis B surface antigen) is also known as the Australia antigen. It is called so because it was first isolated from the serum of an Australian person.

Sera from hepatitis B patients when observed under an electron microscope show three types of particles:

• A spherical particle: 22nm, most abundant, contains only HBsAg.

• Filamentous or tubular particle: 22nm, contains only HBsAg.

• A double-walled spherical structure: 42nm diameter, is the complete Hepatitis B virus. It was described by Dane and is called Dane particle.



Solution to Question 6:

Dane particle refers to Hepatitis B.

Sera from hepatitis B patients when observed under an electron microscope show three types of particles:

• Spherical particle: 22nm, most abundant, contains only HBsAg.

Sold by @itachibot

• Filamentous or tubular particle: 22nm, contains only HBsAg.

• A double-walled spherical structure: 42nm diameter, is the complete Hepatitis B virus. It was described by Dane and is called Dane particle.



Sold by
If you purchased this fro
you may have been scam
else,
d.

Solution to Question 7:

HBeAg is coded by the C gene. The presence of high levels of HBeAg (and HBV DNA) suggests high infectivity.

HBV Genome has four overlapping genes. They code for:

• S gene-It codes for surface antigen.

• C gene-

• When the C region alone is translated: HBcAg is formed.

• When translation begins from Pre C region, HBeAg is formed.

• P gene- Codes for DNA polymerase.

• X gene- Codes for a non-particulate protein HBxAg.

Sold by @itachibot



©MARROW

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

HBsAg is the first marker to appear in the blood after infection.

It can be detected even before the elevation of serum transaminases and onset of clinical illness. Usually, it disappears within 2 months of onset of clinical disease but may last for even 6 months and beyond. If it persists beyond 6 months, it indicates chronic hepatitis.

Anti-HBs is the protective antibody, which appears soon after HBsAg, disappears and remains in serum for a long time.

Solution to Question 9:

Hepatitis B core antigen (HBcAg) is not present in blood because it is enclosed within the HBsAg coat. However, antibodies against HBc (Anti-HBc) are the earliest antibody markers to be seen in the blood after 1-2 weeks.

Note:

• HBsAg: Earliest marker to be detected in the serum of Hepatitis B-infected persons.

• Anti HBc: Earliest antibody to be detected in the serum of Hepatitis B-infected persons.

The given image shows hepatitis B virus and its antigens

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 10:

In most tumors, HBV DNA is integrated into the cell genome, and in most cases, the virus DNA has undergone rearrangements, including deletions. After integration, it promotes the overexpression of cell genes that can also contribute to oncogenesis.

The only viral gene product that is consistently present in the tumor cells is the X protein.

Solution to Question 11:

The given life cycle is of the hepatitis B virus.

Steps:

• The infectious virion attaches to cells and uncoating occurs.

• Cell enzymes in the hepatocytes convert dsDNA to covalently closed circular double-stranded DNA (cccDNA)

• HBV mRNA and pre-genome RNA is synthesized using cccDNA as a template.

• pre-genome RNA is encapsidated by packaging signal into new core particles.

• These new core particles are templates for HBV reverse transcriptase.

• Negative-strand DNA is synthesized by viral polymerase by reverse transcription.

• The cores bud from the pre-Golgi membranes, acquiring HBsAg-containing envelopes.

• Once viral particles are assembled, they either exit the cell or cause reinfection of the same cell.

Sold by @itachibot

Solution to Question 12:

A patient who has had exposure to HBV will have anti-HBc antibody even when he has moved into the chronic or recurrent stages. IgM anti-HBc reappears during acute recurrences while IgG anti-HBc is found in chronic states.

The HBsAg antigen is present in chronic hepatitis B, while the antibody anti-HBs is not.

Solution to Question 13:

Pre-core mutants of hepatitis B virus are unable to synthesize HBeAg.

In the HBV genome, the C gene consists of 2 regions:

• Pre-C region- When translation begins from the Pre-C region, HBeAg is synthesized.

• C-region- When it begins from the C-region, HBcAg is synthesized.

In the HBeAg sequence of pre-core mutants, there is a G to A switch at the 28th amino acid, causing a stop codon. Hence, pre-core mutants are unable to synthesize HBeAg.

The image below shows the HBV genome :



Solution to Question 14:

HBsAg and IgM Anti-HBc are positive, indicating an acute hepatitis B infection. HBeAg positive indicates that the patient is highly infectious.

Solution to Question 15:

Sold by @itachibot

Breastfeeding is not related to the vertical transmission of hepatitis B virus. Breastfeeding is not contraindicated in HBsAg-positive mothers.

Mother-to-child transmission of HBV can occur via three modalities:

• Intrauterine transmission- Transplacental transmission is uncommon. Another route is via a breach in the placental barrier exposing maternal blood to fetal blood e.g. - threatened preterm labor.

• Transmission during delivery- It is the most frequent method of vertical transmission. It is mostly due to newborn contact with the mother's infected secretions or blood at the time of delivery (normal/instrumental).

• Postpartum transmission- Though HBV DNA is present in breast milk, it is not transmitted to the fetus. However, cracked nipples expose the infant to maternal blood, and can transmit the virus to the infant.

Immunoprophylaxis with HBIg is indicated for babies born to HBsAg-positive mothers followed by a three-dose hepatitis B recombinant vaccine.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 16:

Presence of anti-HBs with the absence of all other serological virus markers indicates immunity following HBV vaccination.

Solution to Question 17:

Pre-exposure prophylaxis consists of administration of a 3–4 dose series of hepatitis B vaccine given over a 6-month period.

The pre-exposure prophylaxis of Hepatitis B vaccine schedule is as follows:

The vaccine is genetically engineered by cloning 'S gene' in baker's yeast. It has non-glycosylated HBsAg particles given along with an alum adjuvant.

Pregnancy is not a contraindication for hepatitis B vaccination. The efficacy has been shown to be similar to that for non-pregnant adults, with overall seroconversion rates approaching 95 percent after three doses which can be taken during routine prenatal care.

The vaccine is given intramuscularly, in the deltoid muscle and can be co-administered with other vaccines.

Group	Vaccine dose in mL (Active in gredient in µg/mL)	Schedule
Newborn	0.5 mL (10µg)	<24 hours, 6 weeks, 14 weeks (if administration of the first dose within 24 hours of birt h is not possible, then 6, 10, and 14 weeks)

Sold by @itachibot

Group	Vaccine dose in mL (Active in gredient in µg/mL)	Schedule
Healthy persons aged ≤19 yea rs	0.5 mL (10 µg)	0, 1, 2 / 6 months
Healthy persons aged ≥20 ye ars	1.0 mL (20 µg)	0, 1, 2 / 6 months
Persons on hemodialysis or i mmunocompromised person s	2.0 mL (40 µg)	0, 1, 2, 6 months

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 18:

Anti-HBs will not contribute to the high viral load, as it is not a marker of active viral replication.

HBeAg, HBV DNA, and HBV DNA polymerase are markers of active viral replication and will contribute to the high viral load.

HBeAg is a qualitative marker while HBV DNA is a quantitative marker of the replicative phase.

Solution to Question 19:

The given clinical scenario is suggestive of mixed cryoglobulinemia, which is strongly associated with hepatitis C. Hence, the patient should be tested for hepatitis C.

Solution to Question 20:

Maximum chronicity is seen in HCV infection, followed by HBV infection.



Sold by @itachibot

5-10% of adults infected with HBV develop chronic infection. The majority of neonates affected with hepatitis B develop chronic hepatitis.

HEV is associated with chronic infection only in immunocompromised individuals. HAV is not associated with chronicity.

Note: The most common cause of post-transfusion hepatitis in developed countries is hepatitis C

Solution to Question 21:

Hepatitis D virus (HDV) is an incomplete/defective virus. It requires HBV, to provide the envelope protein (HbsAg) to aid in its survival. Hepatitis D is called the delta virus.

Two types of infection may occur:

• Co-infection - acute HDV and HBV infections occur simultaneously

• Super-infection - Delta viral infection occurs in a person already harboring HBV. This most often leads to chronic hepatitis

Coinfection and superinfection can cause fulminant hepatitis. HBV vaccination is protective against HDV.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

Hepatitis E virus (HEV) resembles caliciviruses in morphology and physical characteristics. Hepatitis E is a non-enveloped virus with ssRNA genome

Note: The largest HEV epidemic occurred in New Delhi in 1955-1956.

Solution to Question 23:

The anti-LKM-3 antibody is associated with hepatitis D infection, not hepatitis B.

Anti–liver-kidney microsomal antibodies (anti-LKM) are many of several autoantibodies that are detected in the serum of patients with different types of acute or chronic liver disease. These antibodies are targeted against antigens of the cytochrome P450 system.

They are also associated with chronic hepatitis C and autoimmune hepatitis.

Option C: IgM anti-HAV antibody indicates current/recent infection. It appears during the late incubation period, peaks in 2-3 weeks, and disappears after 3-4 months. IgG denotes remote infection or immunity. It appears around the same time, peaks at 3-4 months, and persists for a long time.

Option D: Hepatitis E infection was previously known as an epidemic or enterically transmitted non-A, non-B hepatitis. It is transmitted by feco-oral route.

Sold by @itachibot

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

HIV

Question 1:

Which of the following subtypes of HIV is most common in India?

a) Subtype A

b) Subtype B

c) Subtype C

d) Subtype D

Question 2:

Which of the following statements is incorrect about HIV?

a) Cross blood-brain barrier

b) Has 2 single strands of RNA

c) Inhibited by 0.3 % H_2O_2

d) Is a thermostable virus

Question 3:

Which of the following is a nonstructural gene of HIV?

a) gag gene

b) tat gene

c) pol gene

d) env gene

Question 4:

Which of the following represents the correct sequence of HIV replication?

a) RNA-DNA-RNA

b) RNA-DNA

c) DNA-RNA

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) DNA-RNA-DNA

Question 5:

As part of developing the public health policy on HIV in India, which of the following modes of transmission would you focus on the most?

a) Parent to child transmission

b) Heterosexual transmission

c) Blood transfusion

d) Homosexual transmission

Question 6:

A 35-year-old bisexual man goes to an ICTC centre for counselling about HIV. Which of the following is false about HIV transmission?

a) Male-to-female HIV transmission has more risk of causing infection than female-to-male transmission

b) The risk of acquiring HIV with unprotected receptive anal intercourse is lower than that with penile-vaginal intercourse

c) The quantity of HIV-1 in plasma is a primary determinant of the risk of HIV-1 transmission

d) Male circumcision is associated with a lower risk of HIV acquisition for heterosexual men

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 7:

Specific binding of the HIV to the CD4 receptor occurs via which of these?

a) gp120

b) gp41

c) CCR5

d) CXCR4

Question 8:

A patient presents to you with a history of sharing needles during IV drug abuse with an HIV-positive partner. HIV-antibody spot test was negative, however p24 antigen assay was positive. You explain to him that he is currently in the window period for HIV. Which of the

Sold by @itachibot

following statements would you use to explain the concept to him?

a) It is the time period between infection and onset of first symptoms

b) It is the time period between infection and detection of antibodies

c) It is the time period between exposure and onset of viral multiplication

d) It is the time period between exposure and appearance of symptoms

Question 9:

A 41-year-old man was admitted for hernioplasty and initial investigations revealed that he is HIV-positive. He gives no history of contact and has no HIV-associated symptoms. Which of the following statements is incorrect about this phase of HIV infection?

a) Viral replication is active

b) Sequestered in lymphoid tissue

c) Infective

d) Rapid decline of CD4 count is common

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 10:

A 40-year-old man presents to the clinic requesting an HIV test. He had unprotected sexual contact with another man 3 weeks ago. Antibodies against which of the following will be detected first by ELISA or Western blot?

a) gp120

b) gp 41

c) p24

d) p55

Question 11:

You are an officer with the Center for Disease Control and are screening a high-risk population for HIV. Which of the following methods will you use as a screening tool?

a) Western blot

b) ELISA

c) DNA PCR

d) p24 antigen assay

Sold by @itachibot

Answer Key

Question No.	Correct Option
1	c
2	d
3	b
4	a
5	b
6	b
7	a
8	b
9	d
10	b
11	b

If you purchased this, you may have been else,

Detailed Explanations

Solution to Question 1:

Subtype C of HIV is the most common in India. It causes more than 95% of HIV infections.

Subtypes A and C are more readily transmitted by heterosexual contact than subtype B, which is mainly transmitted by homosexual contact and injections.

Solution to Question 2:

HIV is a thermolabile virus. It is readily inactivated by heating to 60 °C for 10 minutes and at 100 °C in seconds. At room temperature (20–25 °C) in the dried blood, it may survive for 1 week.

Option A: HIV crosses the blood-brain barrier. It may cause encephalopathy leading to loss of higher functions, progressing to dementia.

Option B: HIV is a spherical, enveloped virus. Its genome is composed of two copies of identical single-stranded positive-sense RNA.

Option C: HIV is inactivated by treatment with 0.3% hydrogen peroxide, 50% ethanol, 35% isopropanol or 10% household bleach. A 2% solution of glutaraldehyde is effective for the disinfection of medical instruments.

Sold by @itachibot

Solution to Question 3:

The tat gene (trans-activating gene) is a non-structural gene.

HIV has structural and non-structural genes:

• Structural genes- Genes coding for structural protiens:

• gag- Codes for core and shell protein p55, which is cleaved to p15, p18 and p24

• env- Codes for envelope glycoprotein gp160 which is cleaved to gp120 and gp41

• pol- Codes for enzymes: Polymerase, reverse transcriptase, protease and endonuclease

• Non-structural and regulatory genes:

• tat- Trans-activating gene

• nef- Negative factor gene

• rev- Regulator of virus gene

• vif- Viral infectivity factor gene

• vpu- (only in HIV-1) and vpx (only in HIV-2)

• vpr- Stimulates only the promoter region of the virus

• LTR- Long terminal repeat sequences

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 4:

HIV replicates by the sequence RNA-DNA-RNA. The provirus initiates viral replication by directing the synthesis of viral RNA and other components. Hence, the sequence is RNA–ssDNA–dsDNA–RNA.

HIV genome is composed of 2 single-stranded positive-sense RNA. When a virus infects a cell, the viral RNA is transcribed by the enzyme reverse transcriptase, first into single-stranded DNA and then to double-stranded DNA which is integrated into the host chromosome.

Note: If the options include RNA-DNA-RNA and RNA-DNA-DNA-RNA then the latter would be the choice.

The given image represents HIV replication

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

Heterosexual transmission is the most common mode of transmission of HIV infection in developing countries such as India and African countries. In many Western countries, male-to-male sexual transmission is the dominant mode of transmission.

Blood transfusion carries the highest risk per 10000 exposures of getting HIV from an infected person. However, it is not the most common mode of transmission.

Solution to Question 6:

The risk of HIV acquisition associated with unprotected receptive anal intercourse is relatively higher than that seen in penile-vaginal intercourse. This is because the rectal mucosal membrane is thin and fragile and anal intercourse is associated with microtrauma (rectal tears).

The differences in transmission rates between men and women are because the vagina and cervix are exposed to semen for a longer duration, increasing the risk of transmission. However, the penis and urethral orifice are exposed relatively briefly to infected vaginal fluid, lowering the risk.

Solution to Question 7:

The specific binding of the HIV to the CD4 receptor is by the envelope glycoprotein gp120.

Option B - Cell fusion occurs via binding to gp41. Cell fusion is essential for infection to take place.

Option C, D - Binding to CD4 receptor requires the participation of a co-receptor CXCR 4 for T cell tropic HIV strains and CCR 5 for macrophage-tropic strains.

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else, it may have been scammed.

Solution to Question 8:

In the given scenario, the patient has experienced a high-risk exposure to HIV due to sharing needles during IV drug abuse with an HIV-positive partner.

The period between infection and detection of antibodies against HIV is known as the window period.

The window period may last anywhere between 2–8 weeks to a few months. During this period, the individual may be highly infectious. HIV infection can be detected during this period by p24 antigen assay and PCR.



Sold by @itachibot

Solution to Question 9:

The given patient is in the latent period of HIV infection. During the latent phase of HIV infection, the CD4+ T cell count declines steadily and not rapidly.

All patients infected with HIV pass through a phase of clinical latency during which they are asymptomatic. This can last for years. During the latent phase, patients show positive HIV antibody tests and they are infectious. There will be varying degrees of viral multiplication.

Throughout the course of untreated infection and even during the stage of clinical latency, HIV is actively replicating or is sequestered in lymphoid tissues.



Solution to Question 10:

Antibodies against gp41 will be detected first by ELISA or Western blot.

Antibodies to HIV usually appear within 3–6 weeks of primary infection. The appearance of HIV-binding antibodies occurs prior to the appearance of neutralizing antibodies. The only viral proteins that elicit neutralizing antibodies are the envelope proteins gp120 and gp41.

The order of appearance of antibodies is as follows:

Antibody to envelope gp41 > Antibodies to gag protein (p24 and p55) > Antibodies to envelope gp120 > Antibodies to gag protein p17 > Antibodies to pol gene products (p31 and p66).

Note: The earliest antigen to appear in the blood following HIV infection is p24.

Solution to Question 11:

ELISA is most commonly used as a screening test for HIV.

Sold by @itachibot

Screening tests possess high sensitivity but are not highly specific. ELISA is a simple, relatively inexpensive screening test.

The most commonly used confirmatory test is Western blot. Western blot demonstrating antibodies to products of all three of the major genes of HIV (gag, pol, and env) is conclusive evidence of infection with HIV. It is considered positive if it shows antibodies against at least two of the following gene products: p24, gp41, gp120/160.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Myxoviruses and Rhabdoviruses

Question 1:

Which of the following RNA viruses has an 8 segmented genome with its site of replication being the nucleus?

a) Influenza virus

b) Measles virus

c) Rubella virus

d) Mumps virus

Question 2:

A microbiologist working on influenza virus in a lab, observes reversal of hemagglutination. What is this phenomenon known as?

a) Hemolysis

b) Precipitation

c) Elution

d) Complement fixation

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following statements is true regarding antigenic drift?

a) There is a gradual sequential change in antigenic structure

b) It doesn't account for the periodic epidemics of influenza

c) The new antigens thus formed are totally unrelated to the previous antigens

d) Antibodies to the predecessor viruses do not neutralise the new variants

Question 4:

Which of the following organisms is represented by the given form of replication?

Sold by @itachibot



a) Hepatitis virus

b) Influenza virus

c) HIV

d) Herpes virus

Question 5:

You receive multiple patients from a poultry farm presenting with fever, dyspnea, cold, and cough. You suspect an outbreak of Avian flu. Which of the following virus strains causes the same?

a) H1N1

b) H5N1

c) H3N2

d) H2N2

Question 6:

A 4-year-old girl with a history of fever, rhinorrhea, sore throat, and myalgia presented with multiple episodes of vomiting and seizures. She was diagnosed with Reye's syndrome. Which of the following is incorrect about the most likely etiological agent?

a) Virus shedding is present before symptoms begins

b) 5-12 years is a high risk group

c) Aquatic birds act as reservoir

Sold by @itachibot If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) Secondary attack rate is 5-15%

Question 7:

A 12-year-old boy is brought to the OPD with complaints of headache and drowsiness since 2 days. He had a history of fever, ear pain, and difficulty in swallowing for 1 week. On examination, you see the following. Brudzinki's sign is positive. What is the most likely diagnosis?

a) Croup

b) Mumps

c) Infectious mononucleosis

d) Herpangina

Question 8:

A 9-month-old child is brought to the OPD for MMR vaccine. Which strain of mumps virus is used in the same?

a) Edmonston-Zagreb strain

b) Nakayama strain

c) Jeryl-Lynn strain

d) RA 27/3

Sold by @itachibot

Question 9:

A mother brings her 5-month-old baby to the casualty with complaints of fast breathing and poor feeding. She also gives a history of fever, cold, and cough for 3 days. On examination, nasal flaring and expiratory wheeze are present. Which of the following statements is false about the viral etiology of the same?

a) The cytopathic effect is syncytium formation

b) Ribavirin is the drug of choice

c) Omalizumab is given for prophylaxis

d) It is the most important cause of bronchiolitis

Question 10:

An unimmunized man presents with fever, malaise, and an erythematous rash starting from behind the ears and progressing to the face and trunk. Given below is an image of lymphoid tissue taken from this patient. What pathological finding do you see here?



a) Warthin-Finkeldey giant cells

b) Warthin-Starry stain

c) Langhans giant cells

d) Aschoff giant cells

Question 11:

If you found this scanned by @itachibot from someone else, it is from scanned.

Sold by @itachibot

A 3-year-old boy with no prior immunization history presents with fever, cough, and coryza. On examination of the buccal mucosa, you see the following. Which stage of the disease is this finding seen in?



a) Prodromal stage

b) Post measles stage

c) Eruptive stage

d) All the stages

Sold by
If you purchased this fro
you may have been scam

Question 12:

Which of the following disease has the highest mortality once clinically manifested?

a) Nipah

b) Rabies

c) Ebola

d) Zika

Question 13:

A 26-year-old pet-groomer was brought to the casualty with a history of breathlessness, excessive agitation, and restlessness. His bystanders give a history of progressive behavioral change and he had a fever, sore throat, and generalized weakness 1 month back. On examination, pulse was 100 bpm, SpO2 was 98% on room air. Which of the following samples will not be preferred to detect the etiological agent?

Sold by @itachibot

a) Corneal impression

b) Cerebrospinal fluid

c) Saliva

d) Skin biopsies including hair follicles

Question 14:

An officer of the Indian Forest Services is bitten by a rabid fox while on duty. Which of the following statements is true about the etiological agent?

a) The incubation period is 4 to 5 days

b) It doesn't produce Negri bodies

c) It causes disease

d) It doesn't affect salivary gland

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 15:

A 23-year-old man was brought to you with a history of fever, behavioral changes, and seizures. He is foaming at the mouth and is unable to ingest liquids. The patient was admitted to the ICU, however he died soon after. Post mortem brain samples showed the following finding. What is the diagnosis?



a) Yellow fever virus

b) Rabies virus

c) CMV

Sold by @itachibot

d) Molluscum contagiosum

Question 16:

Babes nodules are seen in which of the following conditions?

a) Polio

b) Rabies

c) Diphtheria

d) Herpes

Question 17:

Which of the following statements is false regarding rabies vaccination?

a) Bitten dog should be observed for a minimum of 10 days.

b) Four-dose regimen is given as 0.1ml intramuscularly on days 0, 3, 7 and 28.

c) Pre-exposure prophylaxis regimen consists of 3 doses.

d) The dose of human anti rabies immunoglobulin is 20 IU/kg.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	b
5	b
6	b
7	b
8	c
9	c
10	a
11	a
12	b

Sold by @itachibot

13	a
14	c
15	b
16	b
17	b

Detailed Explanations

Solution to Question 1:

Influenza virus has an 8 segmented genome. Its site of replication is the nucleus, unlike other RNA viruses. Influenza virus belongs to the Orthomyxoviridae family.

Most DNA viruses replicate in the nucleus, except pox. Most RNA viruses replicate in the cytoplasm, except retroviruses and orthomyxoviruses.

Examples of segmented viruses (ARBO):

• Arenavirus (2 segments)

• Rotavirus (11 segments)

• Bunyavirus (3 segments)

• Orthomyxovirus (8 segments)

The image below shows the segmented genome seen in the influenza virus (Orthomyxoviridae)



Solution to Question 2:

Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Sold by @itachibot

Elution is the process of reversal of hemagglutination and the release of the virus from the red cell surface.

Influenza virus contains two surface antigens:

• Hemagglutinin (HA) - causes RBC agglutination; facilitates viral entry into cells

• Neuraminidase (NA) - an enzyme that destroys the receptor HA uses to bind to RBCs causing elution/reversal of hemagglutination. Hence, it is also called a receptor destroying enzyme (RDE).



Solution to Question 3:

Antigenic drift causes a gradual sequential change in antigenic structure.

It accounts for periodic epidemics. This property forms the basis for the development of annual influenza vaccines.

Antigenic variation is a property of influenza viruses by which it alters the antigenic structure on its surface and thus avoids a host immune response. It is of two types - Antigenic shift and antigenic drift.

	Antigenic shift	Antigenic drift
Antigenic structure	Abrupt, drastic change	Gradual, sequential change
Periodicity	10 - 20 years	2 - 3 years
Epidemiological effect	Pandemics and major epide mics	Periodic epidemic and spora dic cases
Genetic cause	Large genetic re-assortment	Point mutation
Antibodies to predecessor str ains	Do not neutralize the new va riant	Neutralizes the new variant t o varying degrees

Sold by @itachibot

	Antigenic shift	Antigenic drift
Clinical relevance	Animal-origin viruses infecting humans e.g. Swine flu, COVID-19	Annual influenza vaccine developmentCommon cold - People can get flu more than once

Solution to Question 4:

The given image shows the replication of influenza virus.

Stages of replication:

• Attachment, penetration, uncoating- the virus is internalized by receptor-mediated endocytosis

• Transcription and translation- Viral transcription occurs in the nucleus. The mRNAs are produced from viral nucleocapsids.

• Viral RNA replication- Viral genome replication is accomplished by the same virus-encoded polymerase proteins involved in transcription.

• Maturation- The virus matures by budding from the surface of the cell.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 5:

H5N1 is a strain of avian influenza.

Transmission occurs only from birds (e.g. poultry). Recombination occurs in pigs. Human-to-human transmission is very rare.

It causes flu-like symptoms and can cause severe respiratory illness.

Option A: H1N1 causes swine flu

Option C: H3N2 causes Hong Kong flu

Option D: H2N2 caused a pandemic in 1957

Solution to Question 6:

The most likely etiological agent in the given case of Reye's syndrome is the influenza virus. Children <5-years, especially those <2-years of age are at the highest risk of serious complications related to influenza infection. Other high-risk groups for influenza include adults ≥60 years of age, persons with chronic diseases, persons who are immunocompromised.

Option A: The virus is shed in respiratory secretions a day before the onset of illness and for 3-4 days thereafter.

Option C: Birds, especially aquatic birds act as reservoirs of influenza viruses.

Option D: Secondary attack rate is 5-15%.

Sold by @itachibot

Note: Reye's syndrome is precipitated by the interaction of a viral infection (such as influenza) with salicylates. Therefore, aspirin is not given to pediatric patients with influenza and influenza-like illness.

Solution to Question 7:

The given clinical scenario with fever, parotitis, headache, drowsiness, and positive Brudzinki's sign is suggestive of meningitis secondary to mumps.

Mumps is caused by the mumps virus and is characterized by fever, bilateral or unilateral parotid swelling which may be tender. Parotitis can present with ear pain and difficulty in swallowing. It usually resolves in a week.

Meningoencephalitis/aseptic meningitis can occur before, along with, or after parotitis. Most commonly, it manifests 5 days after parotitis. It can present with headache, neck stiffness, and drowsiness.

Complications of mumps:

• Aseptic meningitis or meningoencephalitis

• Epididymo-orchitis- Usually unilateral. Seen in post-pubertal males

• Oophoritis

• Other complications include pancreatitis and myocarditis

Note: Non-mumps parotitis can be caused by:

• Epstein-Barr virus - causes infectious mononucleosis

• Coxsackievirus - causes herpangina

• Parainfluenza virus - causes croup

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 8:

The strain of mumps virus used in the MMR vaccine (Measles, Mumps, Rubella) is the Jeryl-Lynn strain.

The virus is attenuated by serial passage in embryonated eggs and grown in chick embryo fibroblast culture.

Note: The schedule of MMR vaccine as per the National Immunisation Schedule is from 9 completed months to 12 months. (can be given till 5 years of age).

Solution to Question 9:

The given clinical scenario is suggestive of bronchiolitis caused by the respiratory syncytial virus (RSV). Palivizumab is given for RSV prophylaxis in high-risk infants such as premature babies,

Sold by @itachibot

immunocompromised infants, etc.

Option A - The characteristic cytopathic effect is syncytium formation (multinucleated giant cell).

Option B - Inhaled ribavirin is the drug of choice.

Option D - RSV causes upper and lower respiratory tract infections in infants, and is the most common cause of bronchiolitis from ages 6 weeks to 6 months. It can cause pneumonia and tracheobronchitis and is common during winters and monsoon.

Solution to Question 10:

Warthin-Finkeldey cells are seen here. These are multinucleate giant cells, which are found in the lymphoid tissues of patients with measles.

The cytopathic effect of measles consists of multinucleate syncytium formation with numerous acidophilic nuclear and cytoplasmic inclusions, as shown in the image below.

Option B: Warthin-Starry stain is used to detect spirochetes and to stain H. pylori and B. henselae

Option C: Langhans giant cells are seen in tuberculosis

Option D: Aschoff giant cells are seen in rheumatic fever



Solution to Question 11:

The given image shows Koplik's spots present in measles. They are seen during the prodromal phase of the illness.

Koplik's spots are small bluish-white ulcerations on the buccal mucosa opposite to the first and second lower molars and are pathognomonic of measles. They contain giant cells, cytoplasmic and intranuclear inclusions, and virion components. They can involve the entire buccal mucosa and fade as the maculopapular rash develops.

Sold by @itachibot

Stages of measles:

The natural history of measles has 3 stages as follows:

• Prodromal stage - It begins 10 days after infection and lasts until day 14. It consists of cough, coryza, redness of eyes, lacrimation, photophobia, vomiting, and diarrhea. Koplik's spots appear in this stage.

• Eruptive stage - A dusky, red maculopapular rash appears. It starts behind the ears spreads downwards to the trunk and lower extremities. Antibodies appear in this stage and are detectable by IgM ELISA. The symptoms such as fever and rash usually resolve spontaneously or go on to develop complications.

• Post-measles stage - Weakness and weight loss may persist. The infection may become chronic and cause secondary infections such as cancrum oris, candidiasis etc.

The image below shows Koplik's spots



The image below shows the characteristic maculopapular rash in measles

Sold by @itachibot

Solution to Question 12:

Rabies is a vaccine-preventable, zoonotic, viral disease. Once clinical symptoms appear, rabies is virtually 100% fatal.

Spread of rabies:

• Centripetal spread -

• The virus enters the peripheral neurons, it moves via retrograde fast axonal transport towards the spinal cord or brainstem a.k.a the central nervous system (CNS) at a speed of up to 250 mm/day.

• Once within CNS, it spreads quickly to other regions in the CNS through fast axonal transport. It infects neurons primarily, but may rarely infect astrocytes.

• Centrifugal spread - After CNS infection is established, it spreads along the sensory and autonomic nerves to:

• Salivary glands

• Adrenal glands

• Skin

Rabies virus replicates in acinar cells of the salivary glands and is secreted in the saliva of rabid animals that serve as vectors.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Solution to Question 13:

The given case scenario is highly suggestive of rabies. Corneal impression smears are not performed, as they are of low diagnostic yield.

Laboratory diagnosis of rabies is performed by the following methods:

• Antigen detection - Direct immunofluorescence (IF) from specimens of :

• Skin biopsy from the nape of the neck including hair follicles. Rabies antigen is present in cutaneous nerves at the base of hair follicles. It is the most sensitive and specific test.

• Corneal smears were previously used. They are not used anymore as they are of low diagnostic yield.

• Viral isolation- Intracerebral inoculation in mice can be attempted from the brain, CSF, and saliva.

• Antibody demonstration from serum and CSF.

• Viral RNA detection by RT-PCR .

• Postmortem examination of brain tissue samples- demonstration of Negri bodies.

Sold by @itachibot

Solution to Question 14:

The given scenario is suggestive of infection with the street form of rabies virus. Street rabies causes disease.

All mammals are susceptible to rabies. Cattle, cats and foxes are highly susceptible, whereas dogs and humans have intermediate susceptibility.

Differences between street rabies and fixed rabies:

Street rabies	Fixed rabies (produced by repeated passages)
Causes disease	Doesn't cause disease
Produces Negri body	Doesn't produce Negri body
Affects salivary glands	Doesn't affect salivary glands
Incubation period- 3 months	Incubation period- 4-5 days

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 15:

The given image showing Negri bodies and the case scenario of fever, neurological symptoms like seizures and altered sensorium, hydrophobia (due to painful muscle contractions during swallowing liquids), and hypersalivation are suggestive of encephalitic rabies.

Negri bodies are eosinophilic cytoplasmic inclusions in neurons that are composed of rabies virus proteins and viral RNA. These are commonly observed in Purkinje cells of the cerebellum and in pyramidal neurons of the hippocampus. Their presence is a characteristic pathologic finding in rabies.

The image below shows Negri bodies:



Sold by @itachibot

Solution to Question 16:

Babes nodules are seen in rabies. Mononuclear inflammatory infiltration in the leptomeninges, including microglial nodules are called Babes nodules.



The most characteristic pathologic finding in rabies is the Negri body. These are eosinophilic cytoplasmic inclusions in brain neurons that are composed of rabies virus proteins and viral RNA. The image given below shows Negri bodies.



Solution to Question 17:

Sold by @itachibot

According to the Updated Thai Red Cross Regimen (2-2-2-0-2) for post exposure prophylaxis, reconstituted rabies vaccine is given as follows:

• 0.1 ml intradermally

• Sites - two sites at the same visit. It is given in both the deltoids

• Schedule - Days 0, 3, 7, and 28

Healthy dogs, cats, or ferrets may be confined and observed for 10 days.

The dosage of equine rabies immunoglobulin (eRIG) is 40 IU/kg while that of human anti-rabies immunoglobulin (hRIG) is 20 IU/kg body weight.

PrEP regimens for individuals of all ages are as follows:

• Three intramuscular or intradermal vaccine doses on days 0, 7, and 21 or 28.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Arboviruses and Picorna Viruses

Question 1:

Which family does yellow fever virus belong to?

a) Arboviridae

b) Flaviviridae

c) Bunyaviridae

d) Picornaviridae

Question 2:

A 10-year-old boy from Uttar Pradesh was rushed to the emergency department with a headache, fever, and blurred vision followed shortly by unresponsiveness to verbal commands. He is suspected to have viral encephalitis. Which of the following transmits this endemic disease?

a) Aedes aegypti mosquito

b) Culex mosquito

c) Anopheles mosquito

d) Mansonia mosquito

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

Which of the following is not true about Japanese encephalitis?

a) Transmitted by Culex vishnui

b) Pig is the amplifier host

c) Man acts as the reservoir

d) Vaccinating pigs is helpful in disease control

Question 4:

A mother brings her 9-month-old child for vaccination against Japanese encephalitis. Which of the following statements is false regarding this vaccine?

Sold by @itachibot

a) Two doses are recommended at age 9 months and 18 months

b) Inactivated vero cell culture is given 4 weeks apart

c) Live vaccine SA 14-14-2 is administered intramuscularly

d) The live attenuated vaccine can be given to adolescents

Question 5:

A patient presents with fever, arthralgia, and pain behind the eyes for 3 days. On examination, she has the following rash. Lab reports show thrombocytopenia and leucopenia. What is the most likely diagnosis?

a) Dengue fever

b) Malaria

c) Typhoid

d) Yellow fever

Question 6:

A 35-year-old wildlife photographer presents with sudden onset of fever, headache, and myalgia. On examination, you notice gingival bleeds and conjunctival hemorrhages. He recently visited a forest in Karnataka that has a large population of monkeys. Which of these is the vector for the suspected disease?

a) Fleas

b) Mite

Sold by @itachibot

c) Ticks

d) Mosquito

Question 7:

A middle-aged man presents with delirium, fever, jaundice, and a hemorrhagic diathesis. The patient is agitated and has had no urine output in the last 6 hours. On investigation, there is azotemia and elevated liver enzymes. A viral disease with the vector shown below is suspected. Which of the following inclusion bodies may be seen?

a) Henderson-Patterson bodies

b) Torres bodies

c) Levinthal Cole Lille bodies

d) Paschen bodies

Question 8:

What is the duration of validity of 17D vaccination?

a) 1 year

b) 5 years

c) Lifetime

d) 10 years

Sold by @itachibot

Question 9:

A pest control technician with a history of fever, vomiting, hematuria, and decreased urine output was diagnosed with hemorrhagic fever with renal syndrome. Which of the following statements is false about the causative virus?

a) Hantaviruses are the natural pathogens of rodents

b) Transmission primarily occurs by Culex mosquitoes

c) Hantavirus pulmonary syndrome presents with pulmonary edema

d) Hantaviruses cause epidemic hemorrhagic fever

Question 10:

An epidemic of paralytic poliomyelitis occurs in a non-immune population. Which among the following strains of poliovirus is the most common cause of this?

a) Type 1

b) Type 2

c) Type 3

d) Type 4

Question 11:

A 5-year-old girl is brought to the pediatrics OPD with a history of lower-limb paralysis and the following deformity. Which of the following will be the most common form of presentation of this illness?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed!

Sold by @itachibot

a) Inapparent infection

b) Abortive form

c) Paralytic form

d) Nonparalytic form

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

A 20-year-old unimmunized patient presents to you with complaints of weakness of lower limbs, inability to pass urine, and fever, for 3 days. His stool samples were sent to the district lab to confirm the diagnosis. Which region of the spinal cord will primarily be affected in these cases?

a) Anterior horn cells

b) Posterior horn cells

c) Lateral horn cells

d) Intermediate column

Question 13:

Which of the following statements is false regarding vaccination against polio?

a) Sabin type has the live attenuated virus

b) OPV should not be administered to immunocompromised persons

c) Salk vaccine is associated with mutations that lead to reversal of virulence

Sold by @itachibot

d) Both OPV and IPV induce humoral IgG

Question 14:

Which of the following strains of poliovirus have been eradicated worldwide?

a) Type 2

b) Types 2 and 3

c) Types 1 and 3

d) Type 3

Question 15:

A 68-year-old woman presented with acute chest pain which was stabbing type in nature. The pain is spasmodic and associated with fever spikes. There's an outbreak of coxsackievirus B in her community. What is the most likely diagnosis?

a) Intercostal neuralgia

b) Bornholm disease

c) Herpangina

d) Pericarditis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 16:

A 5-year-old boy presents with headache, fever, and vomiting. On examination, you observe the following. He is diagnosed with herpangina. Which group of viruses causes the same?

Sold by @itachibot



a) Enteroviruses

b) Rhinoviruses

c) Myxoviruses

d) Rhabdoviruses

Sold by @itachi... else, purchased this from some... may have been scammed.

Question 17:

Which of the following organisms causes the following disease?

a) Coxsackievirus

b) Human herpes virus 7

Sold by @itachibot

c) Pox virus

d) Molluscum contagiosum

Question 18:

A 60-year-old presented with eye pain, watering of eyes, and increased sensitivity to light. On examination, you see the following. Which of the following viruses can cause this condition?



a) Rhinovirus and Coxsackie A24

b) Rhinovirus and Enterovirus-70

c) Enterovirus 70 and Adenovirus 3

d) Enterovirus-70 and Coxsackie A24

Question 19:

A 42-year-old woman presented with fever, joint pains, and headache. Chik sign (image given below) is present on the patient's nose. Which of the following vectors transmits the etiological agent?

Sold by @itachibot



a) Culex tritaeniorhynchus

b) Anopheles culicifacies

c) Aedes aegypti

d) Aedes albopictus

Sold by @itachi... purchased this from some... else, If you... may have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	c
4	c
5	a
6	c
7	b
8	c
9	b
10	a
11	a
12	a
13	c
14	b

Sold by @itachibot

15	b
16	a
17	a
18	d
19	c

Detailed Explanations

Solution to Question 1:

Yellow fever virus belongs to the family flaviviridae, which contains the genus Flavivirus.

The important species under genus Flavivirus include:

- Yellow fever virus
- Dengue virus
- Japanese encephalitis virus
- Kyasanur forest disease virus
- West Nile encephalitis
- Zikavirus

Note: Arbovirus is a term used to refer to any viruses that are transmitted by arthropod vectors. There is no family or order by the name arbovirus. The word arbovirus stands for ARthropod-BOrne virus.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

Endemic viral encephalitis is suggestive of Japanese encephalitis. It is transmitted by Culex tritaeniorhynchus and Culex vishnuii.

It is the leading cause of childhood neurological viral infection in India. In endemic areas such as North India, it typically affects children <15 years of age as by early adulthood, the majority of the population has protective immunity following natural exposure to the virus.

Ardeid birds like herons and egrets are reservoir hosts, while the amplifier hosts are pigs. It is transmitted by Culex mosquitoes, and human infection is a tangential dead end.

Solution to Question 3:

Humans are the dead-end hosts for the Japanese encephalitis virus.

Sold by @itachibot

Option A: The disease is transmitted by Culex tritaeniorhynchus and Culex vishnui (common in India) species.

Option B: Herons act as reservoir hosts and pigs as amplifier hosts.

Option D: The control measures include mosquito control, locating piggeries away from the human dwellings and vaccinating pigs.

The image below shows the life cycle of the Japanese B encephalitis virus.



LIFE CYCLE OF JAPANESE ENCEPHALITIS

If you purchased this, you may have been ripped off. else,

Solution to Question 4:

Live SA 14-14-2 strain of Japanese encephalitis (JE) vaccine is administered subcutaneously.

Types of JE Vaccines:

	SA 14-14-2 cell culture derived vaccine	Inactivated vero cell culture derived Kolar (Jenvac 82156 XY)	
Live attenuated	Inactivated		
Route of administration	Subcutaneous	Intramuscular	Intramuscular
Dosage	2 doses	2 doses	2 doses
Duration between doses	At ages 9months and 16-18 months	4 weeks apart	4 weeks apart
Ages	9 months, 16-18 months, adolescents and non-immune adults	up to 15 years	up to 18 years

Sold by @itachibot

Solution to Question 5:

The given clinical scenario with the symptoms and the rash suggests dengue fever.

It presents with sudden-onset fever, retro-orbital pain, myalgia, arthralgia, and a rash. The rash in dengue appears as white islands on a red sea due to erythema with petechiae. The alternative name for dengue, breakbone fever, comes from the associated muscle and joint pains.

Suspected cases of dengue are confirmed by the presence of NS1 antigen or IgM antibodies.

The treatment of dengue is supportive with hydration taking top priority.

Solution to Question 6:

The given case scenario with exposure to monkeys, fever, myalgia, headache, conjunctival hemorrhage and gingival bleeds is suggestive of Kyasanur forest disease (KFD), which is transmitted by ticks (Haemophysalis spinigera). It is endemic to the forests of Karnataka and Kerala.

The clinical features include:

• Fever

• Headache

• Conjunctivitis

• Myalgia

• Meningoencephalitis

• Hemorrhagic symptoms- hemorrhages in skin, mucosa and gastrointestinal bleeds.

The case fatality rate is about 5 percent.

Monkeys are amplifier hosts, rodents are reservoirs and humans are incidental hosts.

A killed KFD virus vaccine was used in a small field trial and appeared to provide some degree of protection.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 7:

The image shows the Aedes aegypti mosquito and the clinical scenario with hepatic and renal dysfunction with delirium points towards a diagnosis of yellow fever. Torres bodies are amorphous eosinophilic nuclear inclusions associated with yellow fever, a disease caused by the Flavivirus.

Inclusion bodies are structures with distinct sizes and shapes that are seen in virus-infected cells under light microscopy.

Note: Dengue hemorrhagic fever can be a differential diagnosis, but there are no inclusion bodies seen in dengue.

Sold by @itachibot

Solution to Question 8:

17D vaccine (yellow fever vaccine) is valid for a lifetime, beginning 10 days after vaccination. According to the amendment in 2016 to International Health Regulations 2005, the validity of the related international certificate of vaccination against yellow fever has been changed from ten (10) years to the lifespan of the person (traveler) vaccinated.

It is a live-attenuated vaccine derived from the 'Asibi strain', administered subcutaneously. Immunity after a single dose is long-lasting and provides lifetime protection. It is thermolabile.

Vaccination is compulsory for travel to and from endemic areas.

In India, 17D is manufactured at the Central Research Institute, Kasauli.

Solution to Question 9:

Hemorrhagic fever with renal syndrome (HFRS) is caused by Hantaviruses. They are rodent-borne viruses that are transmitted by inhalation of the virus contained in dried excreta of rodents. They are the natural pathogens of rodents like mice, rats, and voles.

HFRS presents with acute kidney injury, hemorrhages, fever, and circulatory collapse. Hantaviruses can also cause Hantavirus pulmonary syndrome.

Note: In Aug 2016, The International Committee on Taxonomy of Viruses (ICTV), has renamed and moved the genus Hantavirus to Orthohantavirus.

Solution to Question 10:

Type 1 poliovirus is the most common cause of epidemics in populations.

Poliovirus strains have been classified into 3 types:

• Type 1- most common and causes most of the epidemics.

• Type 2- endemic infections.

• Type 3- epidemics but rarer than type 1.

Immunity is type-specific.

Note :- Type 2 is declared eradicated in September, 2015 and Type 3 is declared eradicated in October, 2019

Solution to Question 11:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

The given case scenario with a history of lower limb paralysis and deformity is suggestive of poliomyelitis. The given image shows deformity of the right lower limb, namely hyperextension at the knee joint and muscle wasting.

Inapparent infection is the most common (90-95%) presentation of polio. Only seroconversion occurs without any clinical manifestations.

Option B: Abortive poliomyelitis is associated with primary viremia and presents as fever, headache, sore throat, and malaise lasting for 1-5 days.

Option C: Paralytic poliomyelitis occurs following minor illness, characterized by biphasic fever with headache, stiff neck, and flaccid paralysis.

Option D: In non-paralytic poliomyelitis, the disease does not progress beyond the stage of aseptic meningitis.

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 12:

The given case scenario of fever, lower limb paralysis, urinary retention, and stool samples being sent for diagnosis, is suggestive of paralytic polio. The anterior horn cells of the spinal cord are primarily affected, causing acute flaccid paralysis. Posterior horns and intermediate columns/lateral horns may also be involved to some extent.

In the central nervous system, the virus multiplies selectively in the neurons and destroys them. The earliest change is the degeneration of Nissl bodies (chromatolysis). Nuclear changes follow.

Note: Fecal samples of all cases of acute flaccid paralysis are obtained (2 samples) and transported to the laboratory within 72 hours of collection at 4–8°C or frozen at -20°C. This is known as the reverse cold chain and is done as a part of gold standard polio surveillance/acute flaccid paralysis surveillance (AFP surveillance).

Solution to Question 13:

Salk type is the killed vaccine, hence there is no risk of mutation that can lead to a reversal of virulence.

There are two types of vaccines for polio - IPV (inactivated poliovirus vaccine) and OPV (oral poliovirus vaccine).

Solution to Question 14:

Type 2 and 3 have been eradicated worldwide.

Type 2: In September 2015, WHO declared the eradication of the wild type 2 poliovirus strain. In India, the last type 2 infection was detected in 1999.

Type 3: It has not been detected anywhere in the world since November 2012. On World Polio Day in October 2019, the type 3 virus was declared to be eradicated worldwide.

Sold by @itachibot

The only circulating strain is type 1.

Solution to Question 15:

The given case scenario with stabbing type chest pain which is spasmodic in nature, associated with fever spikes, and history of an outbreak of coxsackievirus B is suggestive of Bornholm disease or epidemic pleurodynia.

Symptoms include paroxysms of acute pleuritic chest pain associated with fever spikes. Involved muscles are tender. Chest X-ray is usually normal. Most cases occur during epidemics of coxsackievirus B. The disease is named after the Danish island of Bornholm where early cases occurred.

The treatment is non-steroidal anti-inflammatory agents or hot fomentation of the involved muscles.

Other diseases associated with coxsackievirus B:

• Neonatal myocarditis and pericarditis

• Juvenile diabetes mellitus

• Orchitis

• Severe disseminated congenital infection- hepatitis, meningoencephalitis, adrenocortical involvement

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 16:

The given image shows palatal ulcers in herpangina. It is caused by the coxsackievirus A, a large subgroup of the enteroviruses.

Herpangina or vesicular pharyngitis is characterized by grayish- white papulovesicular lesions on an erythematous base that ulcerate. These are present on:

• Soft palate

• Anterior pillars of tonsils

• Uvula

• Posterior oropharynx

It is a common clinical manifestation of coxsackievirus A infection in children, but it can also be caused by coxsackievirus B, echoviruses, or enterovirus 71.

Treatment is usually only supportive, as the disease is self-limiting and usually runs its course in less than a week.

Solution to Question 17:

Sold by @itachibot

The image shows vesicles on the palms, soles, and oral mucosa, consistent with hand-foot-and-mouth disease. It is caused by coxsackievirus.

Hand-foot-and-mouth disease (HFMD) is an exanthematous fever affecting mainly young children, characterized by oral and pharyngeal ulcerations and a vesicular rash of the palms and soles that may spread to the arms and legs. Vesicles heal without crusting, which clinically differentiates them from the vesicles of herpesviruses and poxviruses. It is usually a benign illness resolving in 1-2 weeks.

Coxsackie A16, A9, and B 1-3 are common causative agents.

Other diseases caused by Coxsackie A virus:

• Aseptic meningitis associated with a maculopapular rash.

• Herpangina

• Minor respiratory infections

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 18:

The given case scenario is suggestive of acute hemorrhagic conjunctivitis (AHC) caused mainly by Enterovirus 70 and Coxsackievirus A24 variant (CA24v).

It is characterized by sudden swelling, congestion, and watering of the eyes. Subconjunctival hemorrhages are a characteristic feature and corneal involvement may occur transiently.

There is no treatment currently available. The disease is self limiting, and resolves within a five to seven-day period.

Note: Acute follicular conjunctivitis is caused by Adenovirus serotypes 3, 4, 7 and 11.

Solution to Question 19:

The given case scenario with fever, joint pain, headache, and Chik sign is suggestive of chikungunya which is transmitted by Aedes aegypti mosquitoes.

Chikungunya presents with sudden onset of fever, crippling joint pain, lymphadenopathy, and conjunctivitis. The fever is typically biphasic with a period of remission after 1–6 days. Nose pigmentation is striking in cases of Chikungunya fever and is termed as Chik sign

No animal reservoir has been identified.

Note: Although Chikungunya is transmitted by both Aedes aegypti and Aedes albopictus, A.Aegypti is considered primary vector of transmission for Chikungunya.

Sold by @itachibot

Miscellaneous Viruses - Rubella, Coronaviruses, Prions, Rotavirus, Filovirus, Zika Virus and Nipah Virus

Question 1:

A newborn baby has a characteristic rash as shown below. On examination, bilateral pearly white nuclear cataracts are noted, and a continuous murmur is auscultated. What family does the virus implicated in this syndrome belong to?

a) Rhabdoviridae

b) Filoviridae

c) Matonaviridae

d) Calciviridae

Question 2:

A brain biopsy done at the time of autopsy shows spongiform encephalopathy. Which of the following is unlikely to be the diagnosis?

Sold by @itachibot



a) Creutzfeldt Jakob Disease

b) Kuru

c) Mad cow disease

d) Sub acute sclerosing pan-encephalitis

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 3:

Which strain of the Ebola virus has the highest case fatality rate?

a) Zaire strain

b) Sudan strain

c) Reston strain

d) Taï Forest strain

Question 4:

Which of the following is a leading cause of diarrheal death in children?

a) Calcivirus

b) Adenovirus

c) Rotavirus

d) Norwalk virus

Sold by @itachibot

Question 5:

An 18-year-old went on an exchange program to Singapore. An adventurous eater, she tried clams and oysters on a cruise ship. 2 days later, she developed fever, nausea, vomiting, and diarrhea. What is the causative agent?

a) Adenovirus

b) Rotavirus

c) Norwalk virus

d) Enterovirus

@itachibot someone else,

Question 6:

A 70-year-old lady with recent onset, progressive dementia comes with myoclonus. The brain imaging is shown below. What biochemical component is responsible for this disease?



a) DNA

b) RNA

c) Protein

d) Polysaccharide

Question 7:

A 32-year-old lady is admitted to the ICU with COVID-19. Which of the following statements is false regarding the organism causing it?

Sold by @itachibot

a) These viruses have club-shaped peplomers on their surface

b) It spreads by direct contact

c) SARS virus is highly mutable

d) It acts by binding to ACE1 and TMPRSS4 receptors

Question 8:

A person who comes in contact with a case of coronavirus (COVID- 19) needs to be followed up for _____.

a) 7 days

b) 21 days

c) 14 days

d) 42 days

Question 9:

In a patient undergoing testing for SARS-CoV, a FELUDA test is done. Which of the following statements is false about this technology?

a) It is the acronym for FNCAS9 Editor Linked Uniform Detection Assay

b) Based on CRISPR gene editing technology

c) It is slower than RT PCR in detecting SARS-CoV2

d) It uses CAS-9 protein to bind to the viral signature

Question 10:

A confirmed case of which of the following viral haemorrhagic fevers was found in India?

a) Ebola

b) Yellow fever

c) Crimean congo hemorrhagic fever

d) Marburg

Question 11:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

A woman presents with a positive urine pregnancy test, 1 month after taking the MMR vaccine. At the time of receiving the vaccine, she had been advised to avoid conception. What advice do you give her now?

a) Wait and watch

b) Termination of pregnancy mandatory

c) Low risk; termination of pregnancy may be considered

d) High risk of anomalies; serious consideration for termination

Question 12:

While working at a WHO outreach post in Africa, you conduct an educational program about the Zika virus. What family does this virus belong to?

a) Togaviridae

b) Flaviviridae

c) Bunyaviridae

d) Reoviridae

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 13:

Zika virus is transmitted by:

a) Aedes aegypti

b) Culex

c) Anopheles

d) Phlebotomus papatasii

Question 14:

Which of the following features do not support a diagnosis of Zika fever?

a) Petechial rash

b) Guillan barre syndrome

c) Fever with arthralgia

d) Conjunctivitis

Sold by @itachibot

Question 15:

The transplacental transmission of the organism given in the image which causes microcephaly is-



a) Nipah

b) Zika virus

c) CMV

d) Rubella

Question 16:

Tropical spastic paraparesis is caused by:

a) Polyomavirus SV40

b) Epstein-Barr virus

c) Human immunodeficiency virus

d) Human T-lymphotropic virus

Question 17:

What family does the Nipah virus belong to?

a) Paramyxoviridae

b) Togaviridae

If you purchased this you may have been sold else.

Sold by @itachibot

c) Flaviviridae

d) Bunyaviridae

Answer Key

Question No.	Correct Option
1	c
2	d
3	a
4	c
5	c
6	c
7	d
8	c
9	c
10	c
11	a
12	b
13	a
14	a
15	b
16	d
17	a

Detailed Explanations

Solution to Question 1:

The clinical scenario of bilateral nuclear cataracts, blueberry muffin rash (image), and patent ductus arteriosus(murmur) suggest congenital rubella syndrome. Rubella virus belongs to the family Matonaviridae.

Rubella virus previously belonged to family Togaviridae, but has been recently changed to family Matonaviridae by the International Committee on the Taxonomy of viruses (ICTV), making it the sole genus under the family Matonaviridae.

Option A: Rhabdoviridae includes Rabies virus.

Sold by @itachibot

Option B: Filoviridae includes Marburg virus and Ebola virus.

Option D: Calciviridae includes Norwalk agent.

Solution to Question 2:

Spongiform encephalopathy is a feature of prion disease. Subacute sclerosing panencephalitis (SSPE) is not a prion disease.

SSPE is a rare and chronic form of progressive brain inflammation caused by a persistent infection with the measles virus.

Slow virus diseases are characterized by a long asymptomatic period, often months or years in duration, between the introduction of the infectious agent and the appearance of clinical illness. Examples: AIDS, Kuru, CJD, SSPE, PML

CJD, Kuru, and mad cow disease are classified under slow virus diseases which are caused by prions.



Solution to Question 3:

The Zaire strain has a case fatality rate of 90%

There are five identified Ebola virus strains, four of which are known to cause disease in humans:

• Ebola virus (Zaire ebolavirus)

• Sudan virus (Sudan ebolavirus)

• Taï Forest virus (Taï Forest ebolavirus)

• Bundibugyo virus (Bundibugyo ebolavirus).

Sold by @itachibot

The fifth, Reston virus (Reston ebolavirus), has caused disease in nonhuman primates, but not in humans.

The Bundibugyo ebolavirus, Zaire ebolavirus, and Sudan ebolavirus have been associated with large outbreaks in Africa. The strain that caused the 2014–2016 West African outbreak belongs to the Zaire ebolavirus species.

Solution to Question 4:

Rotavirus is a leading cause of diarrheal death among children in the developing world.

Rotavirus infections occur in most children below the age of 5 years and are most frequent between 6-24 months of age. Neonatal infections are common but are often asymptomatic or mild.

Vaccines for rotavirus: RotaTeq and Rotarix, given orally.

Note: Viruses that can cause diarrhoea in children include:

- Rotavirus (most common)
- Norwalk virus
- Adenovirus- types 40 and 41
- Astrovirus
- Coronavirus
- Sapovirus

The image below shows an electron micrograph image of the Rotavirus.



Sold by @itachibot
If you purchased this from someone else, you have been scammed.

Solution to Question 5:

Sold by @itachibot

The clinical scenario describes a case of Norwalk infection. It is the most common cause of diarrhea in older children and adults.

They occur in group settings such as banquet halls, cruise ships, dormitories, and campgrounds, where outbreak sources are usually contaminated food and/or water. Potential sources of contaminated food include shellfish such as clams and oysters, which are contaminated by water.

Clinical features are nausea, vomiting, watery diarrhea, abdominal pain. Symptoms of gastroenteritis develop 12 to 48 hours after being exposed to norovirus.

The image below is an electron micrograph of Norovirus (Norwalk virus).



If you purcha... you may have... else,

Solution to Question 6:

The clinical features and imaging suggest a diagnosis of Creutzfeldt-Jakob disease(CJD), caused by prions. Prions are composed of protein.

These are infectious protein particles that lack any nucleic acid. They are filterable like viruses but are resistant to agents of sterilization.

Prion proteins have two isoforms:

• PrPSc is the prion protein that causes disease.

• PrPc is the normal prion protein, found in human cells.

Prions cause diseases such as CJD, Kuru, Alzheimer's, Parkinson's.

Laboratory diagnosis of prion diseases:

• Measurement of PrPSc by immunoassay

• Brain MRI

• Brain biopsy

• Stress protein 14-3-3

Sold by @itachibot

• Sequencing PRNP gene

• Abnormal EEG



Sold by @ If you purchased this from you may have been scamm... else, d.

Solution to Question 7:

The coronavirus spike protein binds to the ACE2 and TMPRSS2 receptors to enter the human cells.

Coronaviruses are a group of spherical or pleomorphic enveloped RNA viruses having club or petal-shaped peplomers. Coronaviruses are named for the crown-like spikes on their surface.

It spreads through coughing, sneezing, and close personal contact such as touching the mouth, nose.

The virus is highly mutable and so vaccine prophylaxis for all strains is difficult.

Coronaviruses cause upper respiratory tract infections, severe acute respiratory syndrome (SARS), and Middle Eastern respiratory syndrome (MERS). They spread by inhalation of the virus present in droplets or aerosols of respiratory secretions of the patients.

Basic protective measures against the new coronavirus: (SARS-CoV-2)

• Wash your hands frequently for at least 20 seconds.

• Maintain social distancing- Maintain at least 1 meter distance between yourself and anyone who is coughing or sneezing

• Avoid touching eyes, nose, and mouth

• Practice respiratory hygiene- cover your mouth and nose with your bent elbow or tissue when you cough or sneeze, then dispose of the used tissue immediately.

• If you have fever, cough, and difficulty breathing, seek medical care early

• Stay informed and follow the advice given by your healthcare provider

Sold by @itachibot

Solution to Question 8:

A person who comes in contact with a coronavirus(COVID-19) case (confirmed or probable) needs to be followed up for apperance of symptoms of coronavirus for the maximum incubation period of 14 days.

Treatment: There is no specific antiviral treatment recommended for COVID-19. People with COVID-19 should receive supportive care to help relieve symptoms. For severe cases, treatment should include care to support vital organ functions.

Basic protective measures against the new coronavirus: (SARS CoV 2)

• Wash your hands frequently for atleast 20 seconds.

• Maintain social distancing- Maintain at least 1 metre distance between yourself and anyone who is coughing or sneezing

• Avoid touching eyes, nose and mouth

• Practice respiratory hygiene- cover your mouth and nose with your bent elbow or tissue when you cough or sneeze, then dispose of the used tissue immediately.

• If you have fever, cough and difficulty breathing, seek medical care early

• Stay informed and follow advice given by your healthcare provider

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

FELUDA is faster than RT-PCR in detecting SARS-CoV2.

FELUDA (FNCAS9 Editor Linked Uniform Detection Assay) is a newly approved method based on CRISPR gene-editing technology to identify and target the genetic material of SARS-CoV2.

The CRISPR technology uses a highly specific CAS9 protein to find and bind to the target Covid signature. This is then coupled with paper-strip chemistry to elicit a visual readout on a paper strip.

It is a very affordable, sensitive (96%), and specific (98%) strip test which can detect viral genetic material within 45 minutes.

Solution to Question 10:

Crimean Congo hemorrhagic fever is a severe form of viral hemorrhagic fever, is caused by Nairovirus species.

It is zoonotic in origin with ticks as vectors. The outbreak was caused due to person-to-person transmission in hospitals.

Sold by @itachibot

Solution to Question 11:

In this case scenario, the doctor should advise the patient to wait and watch.

The vaccine should not be administered during pregnancy. If pregnancy occurs within 28 days of immunization, the patient should be counseled on the theoretical risks for the fetus.

Studies of over 200 women who had been inadvertently immunized with rubella vaccine during pregnancy showed that none of their offspring developed congenital rubella syndrome. Therefore, termination of pregnancy is probably not warranted.

Solution to Question 12:

Zika virus belongs to the family Flaviviridae.

It is classified by the International Committee on Taxonomy of Viruses (ICTV):

• Family: Flaviviridae

• Genus: Flavivirus

• Species: Zika virus

Zika virus is an RNA virus that is transmitted by Aedes aegypti mosquito bite and by sexual contact. It causes microcephaly and is associated with Guillain-Barré syndrome.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 13:

Zika virus may be transmitted by the bite of infected Aedes aegypti and Aedes albopictus.

The symptoms of Zika virus disease include fever, rash, conjunctivitis, muscle and joint pain, malaise, or headache. These symptoms are mild and usually last for 2-7 days.

Zika virus infection during pregnancy may cause lead to microcephaly or congenital Zika syndrome.

The complications of Zika virus infection include Guillain-Barré syndrome, neuropathy, and myelitis.

Solution to Question 14:

A petechial rash is not a primary feature of Zika virus infection.

Though petechiae may rarely appear in Zika virus disease it is however not included in the CDC case definition of Zika virus disease. Petechial rash points towards a diagnosis of dengue fever while a maculopapular rash suggests a Zika virus disease.

Zika virus infection is characterized by influenza-like clinical signs including fever, headache, and malaise. A maculopapular rash, conjunctivitis, myalgia, and arthralgia usually accompany or follow those manifestations.

Sold by @itachibot

The complications of Zika virus are:

• Guillain-Barre syndrome

• Fetal microcephaly in pregnant women.

Solution to Question 15:

The transplacental transmission of Zika virus given in the image causes microcephaly.

The condition shown in the image is microcephaly. It is caused by Zika virus transmission from an infected pregnant mother to her child. Zika virus is a mosquito-borne Flavivirus mainly transmitted through Aedes aegypti.

Symptoms are mild and usually last for 2-7 days.

• Fever

• Rash

• Conjunctivitis

• Muscle and joint pain

• Malaise

• Headache

• In pregnancy- congenital Zika syndrome or microcephaly

The complications of Zika virus infection include Guillain-Barré syndrome, neuropathy, and myelitis.

NCDC Delhi and the National Institute of Virology Pune, are the only institutes that have the capacity to provide the diagnosis in an acute febrile stage. RT-PCR is the standard test available for the diagnosis of Zika virus infection.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Solution to Question 16:

is the causative agent of the neurodegenerative disorder HTLV-1-associated myelopathy or tropical spastic paraparesis (HAM/TSP).

The condition resembles multiple sclerosis and is associated with progressive weakness of legs and lower body. The patients develop spastic paraparesis or paraplegia, hyperreflexia, ankle clonus, and extensor plantar response. The disease progresses slowly and takes the form of thoracic myelopathy.

Other conditions caused by HTLV-1 include:

• Adult T-cell leukemias and lymphomas (ATL)

• Uveitis

• Infective dermatitis

Sold by @itachibot

- Arthritis
- Sjo■gren's syndrome

Solution to Question 17:

Nipah virus belongs to the family Paramyxoviridae.

Togaviridae contains chikungunya virus.

Flaviviridae contains yellow fever virus, dengue virus, Japanese encephalitis virus, hepatitis C virus, Zika virus

Bunyaviridae contains sandfly fever virus, hantaviruses.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Superficial and Systemic Mycoses

Question 1:

Which of the following is a component of the fungal cell wall?

a) Ergosterol

b) Lipoteichoic acid

c) Mycolic acid

d) Chitin

Question 2:

Which of the following is a yeast?

a) Candida

b) Cryptococcus

c) Fusarium

d) Talaromyces

Question 3:

Which among the following is not a sexual spore?

a) Zygospore

b) Sporangiospore

c) Basidiospore

d) Ascospore

Question 4:

Which of the following fungal spores are formed by condensation of the hyphal elements?

a) Arthrospores

b) Conidiospore

c) Basidiospores

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

d) Ascospores

Question 5:

A young patient presents to the dermatology OPD with a history of itchy hypopigmented patches over her chest for 6 months. A KOH mount is prepared with the skin scraping sample, which on microscopy reveals a characteristic appearance as shown in the image. What is the most probable diagnosis?



a) Pityriasis alba

b) Candidiasis

c) Tinea corporis

d) Pityriasis versicolor

Question 6:

Which of the following statements about dermatophytes is true?

a) Epidermophyton infects skin and hair

b) Microsporum infects skin and hair

c) Epidermophyton infects nails and hair

d) Microsporum infects nails and hair

Question 7:

Sold by @itachibot

A 23-year-old-lady came to the dermatology clinic with complaints of hair loss and the presence of firmly adherent, black, gritty, hard nodules along the hair shaft as shown. There is no associated pruritis of the scalp. Which of the following could be the most probable causative agent?



a) Piedraia hortae

b) Trichosporon beigelii

c) Hortaea werneckii

d) Malassezia furfur

Sold
If you purchased this
you may have been sc
else,
d.

Question 8:

A 45-year-old diabetic patient presented with complaints of discoloration of his toenail for the past 4 months as shown in the image below. Which pathogen most commonly causes this condition?

Sold by @itachibot



a) Trichophyton rubrum

b) Trichophyton schonleinii

c) Epidermophyton floccosum

d) Trichophyton mentagrophytes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 9:

Which of the following organisms causes disease following traumatic inoculation into the skin?

a) Sporothrix schenckii

b) Blastomyces dermatitidis

c) Coccidioides immitis

d) Paracoccidioides brasiliensis

Question 10:

Which of the following is false about the most commonly used culture media for growing fungi?

a) Contains taurocholate

b) Is a specific media

c) Contains cycloheximide

d) Has an acidic pH

Sold by @itachibot

Question 11:

A gardener from the sub-Himalayan region presented with multiple ulcerative eruptions along the lymphatics of his hand. A tissue biopsy was taken and sent for culture. Microscopic examination of the organism cultured is shown in the image. Identify the organism.



a) Sporothrix schenckii

b) Histoplasma capsulatum

c) Phialophora richardsiae

d) Exophiala jeanselmei

Question 12:

A farmer developed multiple ulcerative nodules along the lymphatics following a rose thorn prick a week ago. Microscopic finding of the skin biopsy is given below. What is the likely diagnosis?

Sold by @itachibot



a) Cryptococcosis

b) Sporotrichosis

c) Histoplasmosis

d) Aspergillosis

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 13:

A farmer from the sub-Himalayan belt had a thorn prick injury a year back and developed a warty, verrucous growth at the site of prick. A skin biopsy sample was taken and sent for histopathological examination. The characteristic microscopic body as shown in the image was seen. Which of the following could be the most likely diagnosis?



Sold by @itachibot

a) Blastomycosis

b) Pheohyphomycosis

c) Chromoblastomycosis

d) Sporotrichosis

Question 14:

A middle-aged man from south India presented to the OPD with a swelling in his left foot that is unresponsive to antibiotics. On examination, multiple discharging sinuses were noted. The discharge contained brown granules that were soft in consistency. Which of the following is likely the causative agent?

a) Actinomadura madurae

b) Sporothrix schenckii

c) Fonsecaea pedrosoi

d) Madurella mycetomatis

Question 15:

Which of the following is not a dimorphic fungus?

a) Sporothrix schenckii

b) Coccidioides immitis

c) Pneumocystis jirovecii

Sold by @itachibot

d) Talaromyces marneffei

Question 16:

A 54-year-old man had presented to the clinic with a history of chronic cough. His chest X-ray showed a mass in the left lung with hilar lymphadenopathy. Biopsy of the mass was done and findings are shown below. Which of the following is the most likely diagnosis?



a) Gilchrist disease

b) Darling's disease

c) Desert rheumatism

d) Jorge Lobo's disease

Question 17:

A 7-year-old boy with a recent history of travel to South America was brought to the hospital with fever, enlarged lymph nodes and multiple skin lesions on his face. Tissue sample from the lesion shows the following findings under microscopy. Which of the following is the most likely causative organism?

Sold by @itachibot

a) Coccidioides immitis

b) Paracoccidioides brasiliensis

c) Histoplasma capsulatum

d) Blastomyces dermatitidis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 18:

A 50-year-old male presents with low-grade fever and a skin lesion as shown below. He had a history of recent travel to North America. Chest X-Ray reveals consolidation in the right lung with hilar lymphadenopathy. Culture on Sabouraud dextrose agar at 37°C shows thick-walled, large (7-20 μ), round yeast cells with a single broad-based bud. What is the most probable diagnosis?

Sold by @itachibot

a) Blastomycosis

b) Histoplasmosis

c) Coccidioidomycosis

d) Paracoccidioidomycosis

Question 19:

A middle-aged man residing in a rural area presents with psychosis, ataxia and convulsions. On further inquiry, you find out there is history of similar complaints in the family members who recently consumed food prepared with grains as shown in the image. What is the poisoning is most likely due to?

a) Claviceps purpurea

b) Aspergillus flavus

c) Amanita phalloides

d) Aspergillus fumigatus

Question 20:

The below given image shows result of hair perforation test. The organism likely to show below given appearance is

Sold by @itachibot



a) Trichophyton rubrum

b) Trichophyton violaceum

c) Trichophyton schoenleinii

d) Trichophyton mentagrophytes

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	a
5	d
6	b
7	a
8	a
9	a
10	a
11	a
12	b
13	c
14	d

Sold by @itachibot

15	c
16	b
17	b
18	a
19	a
20	d

Detailed Explanations

Solution to Question 1:

Chitin is a component of the fungal cell wall.

General characteristics of Fungi:

• They are unicellular/multicellular eukaryotes.

• The cell wall contains chitin, glucan, mannan and other polysaccharides.

• The cytoplasmic membrane contains ergosterol.

• They possess true nuclei with the nuclear membrane and paired chromosomes.

• They divide sexually, asexually or by both.

Raymond Sabouraud is known as the Father of Mycology.



Solution to Question 2:

Cryptococcus is a yeast.

Sold by @itachibot
If you have purchased this from someone else, it has been scammed.

Sold by @itachibot

Option A: Candida is a yeast-like fungus.

Option C: Fusarium is a mold/filamentous fungus.

Option D: Talaromyces is a dimorphic fungus.

Solution to Question 3:

Sporangiospore is not a sexual spore. It is an asexual spore.

Fungi can be classified into 4 classes based on sexual spore formation:

Fungi imperfecti (Deuteromycetes): These are the fungi whose sexual phases are yet to be identified. Most of the fungi of medical importance belong to this group.

Note: The earlier classification of Phycomycetes, under which Zygomycetes, Oomycetes, and other sub-phyla were present, is now obsolete. Phylum Glomerulomycota has replaced it.

Class	Sexual spore	Asexual spore
Glomerulomycetes	Oospores, Zygospores	Sporangiospores
Ascomycetes	Ascospores	Conidia
Basidiomycetes	Basidiospores	Conidia
Fungi imperfecti (Deuteromycetes)	Yet to be identified	Conidia

Solution to Question 4:

Arthrospores are asexual spores that are formed along the mycelium/hyphae by segmentation and condensation of hyphae.

Arthrospores are commonly formed by dermatophytes, especially those belonging to the genus Trichophyton. They are also formed by dimorphic fungi like Coccidiodes sp.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 5:

The given clinical picture and the KOH mount preparation showing the spaghetti and meatball appearance suggest pityriasis versicolor.

Pityriasis versicolor is a chronic superficial infection of the stratum corneum caused by Malassezia globosa and Malassezia furfur. The most common site involves the neck, chest, back, and upper arms. The lesions often appear as hypopigmented areas.

The KOH mount reveals fungal hyphae that are short and thick (like spaghetti or banana) and many variously sized spores (like meatballs or grapes) referred to as spaghetti and meatballs or bananas and grapes appearance.

Treatment includes topical azoles or terbinafine for 2 weeks.

The image below shows the hypopigmented lesions of P. versicolour over the chest.

Sold by @itachibot

Electron microscopic image of Malassezia furfur



Sold by @... If you purchased this from ... else, you may have been scammed.

Solution to Question 6:

Microsporum infects only skin and hair, not nails.

Dermatophytes include 3 genera:

• Microsporum: It infects only hair and skin (not nails). The species included in this genus are-

• M. gypseum

• M. canis

• M. nanum.

• Epidermophyton: It infects only skin and nails (not hair). E. floccosum is the only species in this genus.

• Trichophyton: It infects skin, hair, and nails. The species falling under this genus are-

• T. rubrum

• T. mentagrophytes

• T. tonsurans

• T. schoenleinii

• T. violaceum

Solution to Question 7:

Sold by @itachibot

The presence firmly adherent, black, gritty, hard nodules along the hair shaft is suggestive of black piedra. The causative agent of this condition is Piedraia hortae.

Piedra is a superficial fungal infection of the hair shaft. It is of two types:

• Black piedra: It is characterized by firmly adherent, black, gritty, hard nodules along the hair shaft. Scalp hair is most often affected. Hair becomes brittle and breaks easily leading to complains of hair loss. It is caused by Piedraia hortae.

• White piedra: It is characterized by soft, white or light brown transparent nodules that are easily detached from the hair. It affects pubic, axillary, facial hair more commonly than scalp hair. It is caused by Trichosporon beigelii.

Management involves shaving or cutting off the affected hair along with topical antifungals.

Black piedra



White piedra

Solution to Question 8:

Sold by @itachibot

The given clinical scenario points to a diagnosis of onychomycosis and the pathogen most commonly causing this condition is Trichophyton rubrum. Diabetes increases the risk of onychomycosis.

Onychomycosis refers to nail infections caused by fungi. It can be due to dermatophytes, non-dermatophytic moulds, and candida. Tinea unguium refers to nail infections caused by dermatophytes. This condition may affect toenails or fingernails, but toenail infections are particularly common.

The following photomicrograph of the nail segment in KOH shows T. rubrum fungal hyphae invading nail cells.



The following image is of a lactophenol cotton blue(LPCB) mount showing pyriform shaped microconidia on the hyphae of T. rubrum.



Sold by @itachibot

Solution to Question 9:

Sporothrix schenckii causes infection followed by traumatic inoculation into the skin.

It is a dimorphic fungus that lives on vegetation such as grasses, trees, rose bushes, etc. Following traumatic inoculation into the skin as in thorn prick, S schenckii causes sporotrichosis, a chronic granulomatous infection. This is usually further followed by spread along the draining lymphatics and lymph nodes as shown in the image below.



Subcutaneous mycoses are usually caused by traumatic inoculation of the causative fungus. These include:

Subcutaneous mycoses	Causative agent/s
Sporotrichosis	Sporothrix schenckii

Sold by @itachibot

Subcutaneous mycoses	Causative agent/s
Chromoblastomycosis	Phialophora verrucosaFonsecaea pedrosoiFonsecaea compactaRhinocladiella aquaspersaCladophialophora carrionii
Phaeohyphomycosis	Exophiala jeanselmeiPhialophora richardsiaeBipolaris spiciferaWangiella dermatitidis
Mycetoma	Pseudallescheria boydii (Anamorph Scedosporium apiospermum).Madurella mycetomatisMadurella griseaE jeanselmeiAcremonium falciforme

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 10:

Sabouraud dextrose agar (SDA) is the most commonly used culture media for growing fungi. It does not contain taurocholate.

It consists of antibiotics like gentamicin or chloramphenicol to inhibit the contaminant bacteria and also cycloheximide to inhibit the growth of saprophytic fungi. It has an acidic pH of 5.4.

Media used in mycology can be classified as:

• Specific media-

• SDA agar

• Czapek-Dox medium

• Birdseed agar

• Cornmeal agar

• Non-specific media-

• Blood agar

• Brain heart infusion agar

Sold by @itachibot



Sold by @itachibot
If you purchased this from someone else,
you can have been scammed.

Solution to Question 11:

The presence of multiple ulcerative eruptions along the lymphatics in a gardener from the sub-Himalayan region and the microscopic finding of flower-like clusters of conidia are suggestive of sporotrichosis, caused by Sporothrix schenckii. It is a thermally dimorphic fungus that lives on vegetation.

The lesions occur following traumatic inoculation of the fungus. A granulomatous nodule is first formed at the site of inoculation that may progress to form a necrotic or ulcerative lesion. The infection can spread via lymphatic channels from the skin and subcutaneous tissue to the lymph nodes resulting in formation of secondary ulcers on the lymph nodes. This condition is known as lymphocutaneous sporotrichosis. This disease is commonly encountered among gardeners, forest workers and manual labourers.

Clinical presentation of lymphocutaneous sporotrichosis:

Sold by @itachibot

Solution to Question 12:

The image shows an asteroid body which is characteristically seen in a case of sporotrichosis.

Asteroid bodies are formed due to the deposition of the immune complexes around the yeast cell. It consists of a central rounded or oval basophilic, cigar-shaped yeast cell surrounded peripherally by eosinophilic radiations. Formation of such structures is called the Splendore-Hoeppli phenomenon.

Note: Asteroid bodies are also seen in sarcoidosis. Those are irregular, star-shaped inclusions seen in multinucleate giant cells.

Solution to Question 13:

The history of a slowly spreading warty, verrucous lesion following a thorn prick injury and the histopathological image showing medlar bodies is suggestive of chromoblastomycosis.

Chromoblastomycosis (chromomycosis) is a subcutaneous mycotic infection that is usually caused by traumatic inoculation. Over months to years, the primary lesion becomes verrucous and wart-like with extension along the draining lymphatics.

Histologically, the lesions are granulomatous and the dark sclerotic bodies may be seen within leukocytes or giant cells called medlar/copper penny bodies.

Subcutaneous mycoses	Causative agent/s
Sporotrichosis	Sporothrix schenckii
Chromoblastomycosis	Phialophora verrucosaFonsecaea pedrosoiFonsecaea compactaRhinocladiella aquasperaCladophialophora carrionii
Phaeohyphomycosis	Exophiala jeanselmeiPhialophora richardsiaeBipolaris spiciferaWangiella dermatitidis
Mycetoma	Pseudallescheria boydii (Anamorph Scedosporium apiospermum).Madurella mycetomatisMadurella griseaE jeanselmeiAcremonium falciforme

Solution to Question 14:

If you purchased this from someone else, it may have been scanned.
Sold by @itachibot

Sold by @itachibot

A swelling with multiple discharging sinuses with brown granules that is unresponsive to the antibiotics is suggestive of eumycetoma caused by Madurella mycetomatis.

Mycetoma is a chronic, granulomatous infection of the skin, subcutaneous tissues, fascia and bone, which most often affects the foot or the hand. It can be divided into three types- eumycetomas, actinomycetomas and botryomycosis.

Infection follows traumatic inoculation of the organism into the subcutaneous tissue from soil or vegetable sources. Within host tissues, the organisms develop to form compacted colonies (grains) 0.5–2 mm in diameter, the color of which depends on the organism responsible. It results in tumefactions, deformities and draining sinuses discharging these grains or granules.

Laboratory Diagnosis:

• A. Direct Examination: The presence of grains in pus collected from draining sinuses or in biopsy material is diagnostic.

• B. Culture Samples should also be cultured, at both 25–30°C and 37°C, on brain-heart infusion agar or blood agar for actinomycetes and on Sabouraud agar (without cycloheximide) for fungi.

Treatment:

The management of eumycetoma is difficult, involving surgical debridement or excision and chemotherapy. Actinomycetoma on the other hand, responds well to rifampicin in combination with sulfonamides or cotrimoxazole.

Option A: Actinomadura madurae causes actinomycetoma which would respond to antibiotics.

Option B: Sporothrix schenckii causes sporotrichosis which is characterised by lesions along the route of lymphatics.

Option C: Fonsecaea pedrosoi causes chromoblastomycosis which would not have discharging sinuses.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 15:

Pneumocystis jirovecii is a yeast-like fungus and not a dimorphic fungus.

Dimorphic fungi exist in the yeast form in body tissues and cultures maintained at 37°C, and in the filamentous form in cultures at room temperature(25°C).

Examples of dimorphic fungi include:

• Blastomyces dermatitidis

• Coccidioides immitis

• Paracoccidioides brasiliensis

• Talaromyces marneffei

• Sporothrix schenckii

• Histoplasma capsulatum

• Coccidioides posadasii

Sold by @itachibot

Solution to Question 16:

The given clinical history and the presence of tiny yeast forms within macrophages are suggestive of Histoplasmosis which is also called Darling's disease. Other names of this condition include Cave's disease and Ohio disease.

The causative agent is a dimorphic fungus called Histoplasma capsulatum. Transmission occurs via inhalation of conidia.

After inhalation, the conidia develop into yeast cells and are engulfed by alveolar macrophages. Within macrophages, the yeasts may disseminate to reticuloendothelial tissues, such as the liver, spleen, bone marrow, and lymph nodes.

Biopsy reveals tiny (2-4 microns) oval-shaped yeast cells inside macrophages.

On culture and microscopy, tuberculate macroconidia and hyaline septate mold with microconidia are seen.

It is treated using azole antifungals.

Solution to Question 17:

The given clinical history and the characteristic pilot's wheel appearance on culture(image) is suggestive of paracoccidioidomycosis caused by Paracoccidioides brasiliensis. It is also called Lutz mycosis/ South American blastomycosis.

Paracoccidioides brasiliensis is a thermally dimorphic fungus that causes a chronic granulomatous disease affecting the lungs, skin, mucosa, lymph nodes. On culture, yeast form is seen at 37° C and the mycelial form at 25° C.

Microscopy of tissue sample reveals thick walled, round, yeast cells with numerous buds attached to each other circumferentially giving the classic mickey mouse or pilot wheel appearance.

Most infections are asymptomatic and few individuals with acute disease present with hepatosplenomegaly, lymphadenopathy and skin lesions.

The below image shows skin lesions in a child with paracoccidioidomycosis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Treatment:

• Mild/moderate disease: Itraconazole

• Severe disease: Amphotericin B for 2-3weeks followed by Itraconazole.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 18:

The given clinical scenario with pulmonary and cutaneous involvement along with a history of travel to North America and the characteristic culture findings suggests the diagnosis of Blastomycosis caused by Blastomyces dermitidis.

Blastomycosis is a chronic infection, characterized by the formation of suppurative and granulomatous lesions, especially, in the lungs and skin.

It is most common in North America and so is also called North Amerian Blastomycosis.

The causative organism is a thermally dimorphic fungus called Blastomyces dermatitidis. Humans are infected by the inhalation of spores in the soil.

Blastomycosis can present in three forms:

• Pulmonary disease:

• Presents with cough, sputum production, chest pain, and fever

• Hilar lymphadenopathy may be present

• Focal or diffuse consolidation, miliary lesions, or abscess formation may occur.

• Resembles TB and Histoplasmosis

• Cutaneous disease:

• Involvement of face and exposed parts of the skin

• Skin lesion: ulcerated verrucous granulomas with an advancing border and central scarring. The border is filled with microabscesses and has a sharp, sloping edge. (image in stem)

Sold by @itachibot

• Disseminated disease:

• Multiple skin lesions are the most common presentation

• Highly fatal

• Genitourinary tract (especially prostate) is frequently involved

• Bone involvement resembles chronic osteomyelitis.

It can be diagnosed based on culture findings. Characterictic findings include the presence of thick-walled round yeast cell with a single broad-based bud when cultured at 37°C. (figure of 8 appearance)

Treatment:

• Localised mild/moderate disease - Itraconazole (6 months)

• Disseminated/ severe disease - Amphotericin B

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 19:

The given clinical vignette and the presence of dark coloured ergot growing on the wheat as shown in the image are suggestive of ergotism due to Claviceps purpurea

Mycotic poisoning is of two types: Mycetism and Mycotoxicosis

Mycetism:

The fungus itself causes toxic effects.

Examples:

• Claviceps → Ergot poisoning

• Coprine → coprine poisoning

• Inocybe → muscarine poisoning

Mycotoxicosis:

Fungal toxins contaminate food and cause toxic effects.

Examples:

• Aflatoxins: These are a type of mycotoxins produced by Aspergillus species of fungi, such as Aspergillus flavus and Aspergillus parasiticus.

• Fusarium toxins

• Citrinin toxin was first isolated from Penicillium citrinum.

Solution to Question 20:

Image shows surface erosion of shaft caused by Trichophyton mentagrophytes.

Hair perforation test:

Sold by @itachibot

• Done to differentiate Trichophyton rubrum from mentagrophytes.

• Few strands of human hair are placed in a petridish with 20 ml of distilled water and autoclaved.

• Few drops of sterile 10% yeast extract and a few fragments of the test fungus are added to the hair strands.

• Incubated at 250C for 2-3 weeks.

• Hair is removed and examined microscopically in a lactophenol cotton blue stained wet preparation.

• T.mentagrophytes causes surface erosion of the shaft resulting in wedge shaped appearance of the shaft.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Opportunistic Mycoses

Question 1:

What is the stain used for identifying cryptococcus?

a) Giemsa stain

b) Ziehl-Nielsen stain

c) Gram staining

d) Mucicarmine

Question 2:

Which of the following is a false statement regarding Cryptococcus neoformans?

a) It is laccase negative

b) It most commonly causes meningitis in AIDS patients

c) There are 3 serological types based on capsular polysaccharide

d) Niger seed agar is used to demonstrate melanin production

Question 3:

Which of the following organisms cannot be grown on any culture media?

a) 1 & 4

b) 1, 3 & 5

c) 1 , 2 & 5

d) 2 & 4

Question 4:

A patient with HIV presented complaining of cough and progressively worsening breathing difficulty. Chest X-ray showed bilateral interstitial infiltrates. The findings of microscopic examination of his bronchoalveolar lavage is given below. Which is the most probable causative organism?

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot



a) Pneumocystis jirovecii

b) Histoplasma capsulatum

c) Cryptococcus neoformans

d) Rhinosporidium seeberi

Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Question 5:

A middle-aged immunocompromised patient presents with complaints of pain during swallowing. She describes it as a burning sensation that radiates to her back every time she swallows. Oral examination reveals creamy white patches on the oral mucosa. The wet mount microscopy of the scraping obtained from the patches is given below. Identify the organism.



Sold by @itachibot

a) Candida albicans

b) Cryptococcus neoformans

c) Aspergillus fumigatus

d) Histoplasma capsulatum

Question 6:

The following test is positive for identification of _____



a) Candida albicans

b) Candida glabrata

c) Candida parapsilosis

d) Candida tropicalis

Question 7:

A 35-year-old HIV positive■ patient presents with mucosal lesions in the mouth as shown in the image below. On microscopy, budding yeasts and pseudohyphae are seen. What is the likely diagnosis?

Sold by @itachibot



a) Oral candidiasis

b) Hairy leukoplakia

c) Lichen planus

d) Diphtheria

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 8:

Galactomannan in serum is seen in:

a) 1 & 2 only

b) 1 & 3 only

c) 1 only

d) 1, 2 & 3 only

Question 9:

Identify the organism in the given slide stained with Gomori methenamine silver stain.

Sold by @itachibot



a) Obtuse angled mucor

b) Acute angled penicillium

c) Acute angled branching aspergillus

d) Obtuse angled rhizopus

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 10:

A young lady complains of recurrent rhinitis, nasal discharge and bilateral nasal blockage for two years. She has a history of allergic rhinitis. On rhinoscopy, multiple ethmoidal polyps with mucosal thickening and impacted secretions are seen in both the nasal cavities. A tissue biopsy is taken and sent for culture, which on microscopic examination shows the following appearance. What is the most probable organism implicated here?



Sold by @itachibot

a) Aspergillus fumigatus

b) Aspergillus flavus

c) Aspergillus niger

d) Penicillium chrysogenum

Question 11:

What is the drug of choice for invasive pulmonary aspergillosis?

a) Itraconazole

b) Voriconazole

c) Lyophylised Amphotericin B

d) Posaconazole

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

β 1-3 glucan assay is used to diagnose all except?

a) Candidiasis

b) Aspergillosis

c) Pneumocystis pneumonia

d) Mucormycosis

Question 13:

Which fungus is most commonly associated with orbital cellulitis in patients with diabetic ketoacidosis ?

a) Candida

b) Rhizopus

c) Aspergillus

d) Trichophyton

Question 14:

Sold by @itachibot

A 68-year-old post-COVID-19 patient presented with a black, foul-smelling nasal discharge and blurred vision. On examination, a blackish necrotic mass eroding the septum and hard palate was visualised. Lactophenol cotton blue mount from the culture is given below. Which of the following statements is false regarding the condition?



a) It is a fatal condition

b) Iron administration to patients with active infection should be avoided

c) Posaconazole can be used to treat it

d) Pulmonary form is the most common

If you purchased this ... you may have been sc... else, ...d.

Answer Key

Question No.	Correct Option
1	d
2	a
3	d
4	a
5	a
6	a
7	a
8	c
9	c
10	a

Sold by @itachibot

11	b
12	d
13	b
14	d

Detailed Explanations

Solution to Question 1:

Mucicarmine stain is used in the identification of cryptococcus.

Mucicarmine stained specimen of lung tissue containing cryptococci:



Solution to Question 2:

Pathogenic varieties of cryptococcus including C. neoformans and C. gattii are laccase positive. They also have ability to grow at 37° C and these factors are used to differentiate them from non-pathogenic cryptococci.

Cryptococci have been classified into 5 serotypes (A, B, C, D, and AD) based on their capsular polysaccharide:

• Three serotypes fall under Cryptococcus neoformans - A(var. grubii), D(var. neoformans), and AD(a hybrid of serotypes A and D). Although serotype AD is very similar to serotypes A and D, the isolates still comprise a discrete group that can be distinguished.

• The serotypes B and C are now considered a different species - Cryptococcus gattii.

Niger seed agar is a solid medium used for selective and differential isolation of Cryptococcus neoformans from clinical specimens. C. neoformans and C. gattii produces brown

Sold by @itachibot

colonies on this medium due to the formation of melanin. Melanin is produced by laccase using caffeic acid as the substrate.

The following image shows pigmented colonies of C. gattii growing on niger seed agar.



Cryptococcal meningitis caused by Cryptococcus neoformans, is an opportunistic infection seen in individuals with AIDS. Flucytosine and amphotericin B are used in the treatment of this condition.

Sold by @itachibot
If you purchased this from elsewhere, you may have been scammed.

Solution to Question 3:

Rhinosporidium seeberi and Pneumocystis jirovecii are non-cultivable organisms and cannot be grown on any culture media.

Another example of a non-cultivable fungus is Lacazia loboi. It causes Jorge Lobo's disease which is a chronic subcutaneous mycosis.

Rhizopus oryzae is a mold which is easily cultivable on SDA. Dimorphic fungi like Histoplasma capsulatum and Paracoccidiodes brasiliensis are identified based on their colony morphology on SDA.

Note: Recently, molecular biological techniques have demonstrated that Rhinosporidium is an aquatic protistan parasite. Infections due to Rhinosporidium have been classified under infections due to Protozoa under the ICD 11.

Solution to Question 4:

The given clinical picture and the GMS stained smear showing the black-colored cysts resembling crushed ping pong balls are suggestive of pneumocystis pneumonia (PCP) caused by Pneumocystis jirovecii.

Sold by @itachibot

Patients with PCP generally present with fever and cough that is usually non-productive. Chest X-ray is either a normal film if the disease is suspected early, or with diffuse bilateral interstitial infiltrates. The classic finding of dense perihilar infiltrate is unusual in patients with AIDS.

Treatment of choice is trimethoprim-sulfamethoxazole (cotrimoxazole) and pentamidine isethionate. Cotrimoxazole can also be used for prophylaxis in patients having AIDS.

The following X-ray image shows a cyst (arrow) and bilateral lung infiltrates in a patient with PCP:





Solution to Question 5:

The given clinical scenario of odynophagia, creamy white patches on the oral mucosa and the wet mount microscopy showing budding yeasts with pseudohyphae are suggestive of oral candidiasis caused by Candida albicans.

Sold by @itachibot

It is an opportunistic infection commonly seen in immunocompromised states like AIDS, diabetes etc.

Treatment includes topical azoles or nystatin.

Solution to Question 6:

The image shows the germ tube test used for identifying Candida albicans.

Unlike other species of candida, yeast cells of Candida albicans will begin to form germ tubes after incubation in serum maintained at 37°C for about 90 minutes. This is known as the Reynolds-Braude phenomenon. This test is positive for Candia dubliniensis as well.

Solution to Question 7:

The most likely diagnosis of the given clinical scenario is oral candidiasis.

Oral candidiasis or thrush is a patchy, pseudomembranous lesion that develops over the tongue, lips, gum or palate. It is composed of epithelial cells, yeasts, and pseudohyphae. The causative agent is usually Candida albicans.

It occurs in almost all individuals with AIDS. Other risk factors include hyperglycemia, cellular immunodeficiency, corticosteroids or antibiotics.

It is treated with topical nystatin or oral ketoconazole or fluconazole.

Option B: Hairy leukoplakia is a wart-like growth on the tongue, caused by Epstein Barr virus. It is commonly seen as a white lesion on the lateral side of the tongue.

Option C: Lichen planus is a pruritic disease characterized by papules commonly seen over the flexor surfaces and trunk. Oral and vulvovaginal lesions may also occur. Oral lesions appear as lacy, white, and raised lesions.

Option D: Diphtheria is characterized by the presence of a grey pseudomembrane covering the tonsils and pharynx. It is caused by Corynebacterium diphtheriae.

Solution to Question 8:

The galactomannan assay is used for the diagnosis of invasive (deep-seated) mycoses caused by Aspergillus species.

Galactomannan is a component of the cell wall of Aspergillus. It can be detected in serum or bronchoalveolar lavage by ELISA for the diagnosis of invasive aspergillosis in adults and children. This test is especially useful in hematopoietic stem cell transplant recipients or patients with hematologic malignancies.

Other diagnostic features of invasive aspergillosis are:

• A positive culture from the affected organ

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

• Aspergillus antigen detection of galactomannan

• Narrow, septate hyphae with branching at 45o and conidiophores

• Halo sign on CT scan

Solution to Question 9:

The given slide stained with GMS shows acute-angled (45o) uniform, branching septate hyphae which is suggestive of Aspergillus species.

Aspergillus species grow rapidly, producing aerial hyphae that bear characteristic conidial structures. Long aseptate conidiophores are seen with terminal vesicles on which phialides are arranged in basipetal chains.

The following image shows the lactophenol cotton blue solution (LPCB) mount of Aspergillus fumigatus showing the long conidiophores with terminal vesicles.



Option A: Mucor produces nonseptate, broad hyphae with aerial sporangiophores which end in a sporangium.

Option B: Penicillium produces septate hyphae with branched, septate and brush-like conidiophores.

Option D: Rhizopus appears similar to Mucor but it has rhizoids with sporangiophores arising in groups directly above the rhizoids.

Solution to Question 10:

The given clinical picture and the microscopic appearance is suggestive of fungal rhinosinusitis. The most probable causative organism is Aspergillus fumigatus. It is commonly involved in paranasal sinus mycoses (fungal sinusitis).

Sold by @itachibot

The disease has been described to have four types:

• allergic

• non-invasive

• invasive

• fulminant

A possible fifth type called non-invasive destructive type may also exist.

Zygomycetes like mucor and rhizopus are the common agents in fulminant type of fungal sinusitis.

The characteristic microscopic appearance of Aspergillus fumigatus:

• The vesicle is conical.

• Phialides are arranged in a single row.

• Conidia arise from the upper third of the vesicle and are hyaline.

Option B: Aspergillus flavus appears as follows:



• The vesicle is globular.

• Phialides are arranged in one or two rows.

• Conidia arise from the entire vesicle and are hyaline.

Option C: Aspergillus niger appear as follows:

If yo... itachibot ... someone else, ...ed.

Sold by @itachibot



• The vesicle is globular shaped.

• Phialides are in two rows.

• Conidia arise from the entire vesicle and are black.

Option D: Penicillium chrysogenum appear as follows:



• Septate hyphae with branched conidiophores bearing chains of spores(brush or broom appearance).

Solution to Question 11:

Voriconazole is the drug of choice in the treatment of invasive pulmonary aspergillosis. It can also be used against candida species (including glabrata and krusei), scedosporium and fusarium.

If yo... sold by @itachibot ...this from some... else, ...n scammed.

Sold by @itachibot

The other options can be used as alternatives for the treatment of invasive pulmonary aspergillosis if voriconazole is unavailable.

Solution to Question 12:

The β 1-3 glucan assay cannot be used to diagnose mucormycosis. This condition is usually diagnosed based on clinical features and culture.

The test detects the presence of (1-3)-β-d-glucan (BDG) which is an abundant cell wall polysaccharide found in most fungi.

It is a chromogenic, quantitative enzyme immunoassay (EIA) designed to detect beta D Glucan by using purified, lysed horseshoe crab (Limulus polyphemus) amebocytes. 1,3 Beta D Glucan level of ≥80 pg/mL is considered a positive result.

Fungi that do not contain BDG in their cell walls include

• Cryptococcus neoformans

• Zygomycetes like rhizopus and mucor

• Blastomyces dermatitidis

Mucormycosis is an opportunistic mycosis caused by zygomycetes, which do not contain (1-3)-β-d-glucan, and hence, the β 1-3 glucan assay cannot be used for its diagnosis.

Solution to Question 13:

Rhizopus is the most common fungus associated with orbital cellulitis in patients with diabetic ketoacidosis.

Diabetic patients are immunocompromised and so prone to develop opportunistic infections. Orbital cellulitis with Rhizopus may turn fatal in these patients as it is an angioinvasive fungus and could lead to septicemia and thromboses.

Solution to Question 14:

The given clinical scenario and the LPCB mount showing aseptate hyphae branching at right-angles is strongly suggestive of mucormycosis. Rhino-orbito-cerebral mucormycosis is the most common form of mucormycosis. Pulmonary mucormycosis is the second most common form.

Mucormycosis is an invasive disease caused principally by species of Rhizopus, Mucor, and Lichtheimia. The disease is often characterized by hyphae growing in and around the blood vessels. It can turn fatal on cerebral involvement and development of speticemia.

Risk factors for the infection include:

• Diabetes (Diabetic ketoacidosis).

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Sold by @itachibot

• Elevated levels of free iron.

• Defects in phagocytic function.

• Organ or hematopoietic stem transplantation.

Treatment of choice is Amphotericin-B. Posaconazole can be used as an alternative drug.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

General Parasitology

Question 1:

A 9-year-old child infected with Toxocara canis is diagnosed to have visceral larva migrans. It is an example of_____.

a) Obligate parasite

b) Facultative parasite

c) Accidental parasite

d) Aberrant parasite

Question 2:

Which of the following refers to a host in which the larval stage of the parasite remains viable without further development?

a) Definitive host

b) Intermediate host

c) Paratenic host

d) Reservoir host

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 3:

A 46-year-old farmer came to the clinic with cough and chest pain for one week and fatigue for a month. He was diagnosed to have a hydatid cyst in his right lung. Which of the following classes does the causative organism of this disease belong to?

a) Cestodes

b) Nematodes

c) Trematodes

d) Zoonoses

Question 4:

Which of the following is the smallest cestode infecting humans?

Sold by @itachibot

a) Diphyllobothrium latum

b) Hymenolepis nana

c) Heterophyes heterophyes

d) Strongyloides stercoralis

Question 5:

Which of the following parasites has a direct life cycle?

a) Enterobius vermicularis

b) Taenia solium

c) Toxoplasma gondii

d) Trichinella spiralis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Question 6:

Match the following:

a) 1-B, 2-D, 3-C, 4-A

b) 1-B, 2-D, 3-A, 4-C

c) 1-D, 2-B, 3-A, 4-C

d) 1-D, 2-A, 3-B, 4-C

Question 7:

An entomologist who had recently visited Mexico presented to the OPD with fever, fatigue, and an alleged history of contact with the insect given in the image. Which of the following diseases is it a biological vector for?

Sold by @itachibot

a) Sleeping sickness

b) Leishmaniasis

c) Babesiosis

d) Chagas disease

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 8:

Autoinfection is not seen with which of the following parasites?

a) Strongyloides stercoralis

b) Taenia solium

c) Hymenolepis nana

d) Ascaris lumbricoides

Question 9:

A microbiologist is doing research about antigenic variations in parasites. Which of the following parasites will exhibit this phenomenon?

a) Trypanosoma brucei

b) Hymenolepis nana

c) Taenia solium

d) Enterobius vermicularis

Sold by @itachibot

Question 10:

You order a peripheral blood film examination while evaluating a patient with pyrexia of unknown origin. Which of the following organisms cannot be detected by this investigation?

a) Giardia

b) Trypanosoma

c) Babesia

d) Leishmania

Question 11:

A Giemsa stain of a thin peripheral smear is prepared. Which of the following cannot be diagnosed?

a) Coxiella burnetii

b) Toxoplasma gondii

c) Bartonella henselae

d) Ehrlichia chaffeensis

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 12:

A patient has presented to the PHC with fever and chills. You order a rapid diagnostic test to detect HRP-2 antigen. Which of the following diseases is it used to diagnose?

a) Malaria

b) Filaria

c) Amoebiasis

d) Giardiasis

Question 13:

In a patient with a positive Casoni's skin test, which of the following infections will you suspect?

a) Schistosomiasis

b) Kala-azar

c) Hydatid disease

Sold by @itachibot

d) Toxoplasmosis

Question 14:

A patient presented to the OPD with cough, chest pain, and expectoration of rusty colored sputum. He is diagnosed to have an oriental lung fluke infection. Which of the following parasites is implicated here?

a) Clonorchis sinensis

b) Paragonimus Westermani

c) Fasciolopsis buski

d) Metagonimus yokogawai

Question 15:

A college student from Mangalore comes to the OPD with intermittent fever, headache, and chills for 1 week. She gives a history of malaria 1 year back for which she had received treatment. Which of the following species you would expect to see in the peripheral smear of this patient?

a) Plasmodium vivax and falciparum

b) Plasmodium ovale and falciparum

c) Plasmodium vivax and ovale

d) Plasmodium vivax and malariae

Question 16:

A middle-aged woman comes to the clinic with abdominal pain, diarrhea, fatigue, and weight loss. Stool examination revealed bile-stained eggs. Which of the following organisms may not be implicated in this case?

a) Ascaris lumbricoides

b) Trichuris trichuria

c) Enterobius vermicularis

d) Taenia solium

Question 17:

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

You are examining a wet mount stool sample of a patient with diarrhea. As the eggs are not visible, you concentrate the sample using the saturated salt solution technique. Which of the following eggs will not float in this technique?

a) Trichuris tricuhria

b) Hymenolopis nana

c) Strongyloides stercoralis

d) Enterobius vermicularis

If yo ibot one else,

Answer Key

Question No.	Correct Option
1	d
2	c
3	a
4	b
5	a
6	b
7	d
8	d
9	a
10	a
11	a
12	a
13	c
14	b
15	c
16	c
17	c

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

Toxocara canis (dog roundworm) infecting a human is an example of an aberrant parasite. It is defined as a parasite that infects a host where they cannot develop further.

Parasites can be classified as:

• Ectoparasites - inhabit only the body surface of the host without penetrating the tissue. The term infestation is often used for parasitization. e.g., lice, ticks, and mites.

• Endoparasite - lives within the body of the host and causes infection.

Endoparasites are further classified as follows:

• Obligate parasites - cannot exist without a host. e.g., Toxoplasma gondii and Plasmodium.

• Facultative parasites - exist as either parasitic or free-living forms. e.g., Naegleria fowleri.

• Accidental parasites - infects an unusual host. e.g., Echinococcus granulosus infects man

• Aberrant/Wandering parasites - infects a host where they cannot develop further. e.g., Toxocara canis (dog roundworm) infecting humans.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 2:

A host in which the larval stage of the parasite remains viable without further development is referred to as the paratenic host.

For example, in the lifecycle of Diphyllobothrium latum (fish tapeworm), humans are the definitive host. The first intermediate host is cyclops and the second is small freshwater fish. If the small freshwater fish with the plerocercoid larva get eaten by a larger fish, the larva survives in the larger fish. This makes the larger fish a paratenic host.

Host is an organism that harbors the parasite and provides nourishment and shelter to it. It is relatively larger than the parasite.

Types of hosts:

• Definitive host - harbors stages of sexual reproduction of the parasite.

• Intermediate host - harbors stages of asexual multiplication of the parasite.

• Paratenic host - harbors only the larval stage of the parasite which remains viable without further development.

• Reservoir host - harbors the parasite and acts as an important source of infection to other susceptible hosts, especially in endemic areas.

• Accidental host - harbors a parasite that is not usually found in the host.

Solution to Question 3:

The causative organism of hydatid disease is Echinococcus granulosus and it belongs to the class cestodes.

It is a tape-like, segmented worm. It infects man accidentally, giving rise to hydatid cysts.

Sold by @itachibot

The image below shows the classification of parasites.



Note: Zoonoses refers to diseases caused by zoonotic organisms. It is not a class of organisms.

Solution to Question 4:

The smallest cestode infecting humans is Hymenolepis nana.

Option A: Diphyllobothrium latum is the longest cestode infecting man.

Option C: Heterophyes heterophyes is the smallest trematode infecting man.

Option D: Strongyloides stercoralis is the smallest nematode infecting man.

Also, Fasciolopsis buski is the largest trematode infecting humans.

Solution to Question 5:

Enterobius vermicularis has a direct life cycle.

Parasite requiring a single host to complete its development is said to have a direct life cycle.

Parasites having a direct life cycle are:

Parasite requiring two or more hosts to complete its development is said to have indirect life cycle, e.g. Plasmodium spp.

Sold by @itachibot
If you purchased this from so... else, you may have been scammed.

Sold by @itachibot

Protozoa	Helminths
Entamoeba histolyticaGiardia lambliaTrichomonas vaginalisBalantidium coliCryptosporidium parvumCyclospora cayetanensisIsospora belliMicrosporidia.	Ascaris lumbricoidesEnterobius vermicularisTrichuris trichuraAncylostoma duodenale Necator americanusHymenolepsis nana.

Solution to Question 6:

The correct answer is option B.

A boy from Kolkata with fever and massive splenomegaly is likely suffering from Kala-azar caused by Leishmania donovani. Its intermediate host is sandfly.

A lady from Kerala with bilateral non-pitting pedal edema is likely suffering from lymphatic filariasis caused by Wuchereria bancrofti. Its intermediate host is the culex mosquito.

Sold by @itachibot

A man presenting with new-onset seizures and MRI brain showing cyst with a dot sign is likely suffering from neurocysticercosis caused by ingestion of Taenia solium eggs (Cysticercus cellulose). The intermediate host of Taenia Solium is the pig.



A woman having fever with her peripheral smear showing Maltese cross appearance is likely suffering from babesiosis caused by Babesia microti. Its intermediate host is human.

Sold by @itachibot



Sold by @itachi... else,
If you purchased this from some...
you may have been scammed.

Solution to Question 7:

The image shows reduviid bug which is the biological vector for the transmission of Chagas disease. Acute Chagas disease can present as fever with fatigue and a subcutaneous swelling known as a chagoma. It may lead to fatal myocarditis.

Biological vectors not only help in the transfer of parasites but the parasites undergo development or multiplication in their body as well. They are also called true vectors.

Examples of biological vectors are:

• Mosquito - Malaria, filariasis

• Sandfly - Kala-azar

• Tsetse fly - Sleeping sickness

• Reduviid bugs - Chagas disease

• Ticks - Babesiosis

Solution to Question 8:

Autoinfection is not seen in infection with Ascaris lumbricoides.

Autoinfection refers to either of the following:

• Reinfection by microbes or parasitic organisms that are present on or within the body. It is also known as internal reinfection. e.g., Strongyloides.

• Self-infection by direct contact with a contagious agent, as with parasite eggs in the infectious state transmitted by fingernails. e.g., Pinworm.

The following are the parasites causing autoinfection (mnemonic: CHEST.C)

Sold by @itachibot

• Cryptosporidium parvum

• Hymenolepis nana

• Enterobius vermicularis

• Strongyloides stercoralis

• Taenia solium

• Capillaria philippinensis.

Solution to Question 9:

Trypanosoma brucei exhibits antigenic variation within the host. This genetic switch protects them from antibodies.

These trypanosomes exhibit unique antigenic variations of their glycoproteins. There is a cyclical fluctuation in the trypanosomes occurring in the blood of infected vertebrates after every 7–10 days.

Parasites exhibiting antigenic variation include:

• Trypanosoma brucei gambiense

• Trypanosome brucei rhodesiense

• Plasmodium spp.

• Giardia lamblia

The following image shows Trypanosoma brucei.



The following image shows Plasmodium spp.

Sold by @itachibot
If you purchased this from someone else, you have been scammed

Sold by @itachibot



The following image shows the trophozoite of Giardia lamblia.

Solution to Question 10:

Stool examination is useful for the detection of Giardia lamblia, not peripheral blood film examination.

Examination of blood is of vital importance for demonstrating parasites that circulate in blood vessels.

Examples of parasites found in the peripheral blood film:

Sold by @itachibot

Protozoa	Nematodes
Plasmodium spp.Babesia spp. Trypanosoma spp.Leishmania spp.	WuchereriabancroftiBrugia malayiLoa loaMansonella spp.

Solution to Question 11:

The diagnosis of Coxiella burnetii cannot be done using a Giemsa stain of a thin peripheral smear.

It is done mainly by serology, which shows a rise in the titer of specific antibodies to C. burnetii.

Microorganisms such as Histoplasma, Leishmania, Toxoplasma, and Pneumocystis can also be detected with Giemsa, and in gastric tissues, Helicobacter pylori (H. pylori) appear thin and distinctly blue. Giemsa's stain is frequently used for diagnostic purposes in haematology to differentiate the nuclear and cytoplasmic morphology of platelets, RBCs, WBCs, and parasites.

If you purchased this from someone else Sold by @itachibot they have been scammed.

Solution to Question 12:

HRP-2 antigen detection is used in the diagnosis of malaria.

Malaria antigens detected by the rapid immunochromatographic test are:

• pLDH - positivity is seen in all Plasmodium species

• HRP-2 - positivity is specific for P. falciparum

The below image shows rapid diagnostic test (RDT) kit for malaria.

Solution to Question 13:

Sold by @itachibot

Casoni's skin test is done in patients with hydatid disease.

It is an immediate hypersensitivity reaction (type I).

This test is abandoned now due to non-specificity and has been supplemented by serological tests.

Solution to Question 14:

Paragonimus westermani is commonly known as oriental lung fluke or Japanese lung fluke. It causes a pulmonary syndrome characterised by cough, chest pain and hemoptysis.

Paragonimus westermani is a trematode. The transmission to humans occurs by eating raw or undercooked crab or crayfish containing metacercariae. The metacercariae excyst in the human gut and young worms migrate to the lungs, where they become encapsulated in lung tissue.

The patient complains of cough, chest pain, and hemoptysis. The viscous sputum is speckled with golden-brown eggs.

Treatment: Praziquantel (25 mg/ kg TDS for 1-2 days) is the drug of choice.

The following image shows the life cycle of Paragonimus westermani, the oriental lung fluke.



Solution to Question 15:

The given clinical scenario is suggestive of malaria relapse and it is caused by Plasmodium vivax and Plasmodium ovale species.

In case of P.vivax and P.ovale, two kinds of sporozoites are seen. Some multiply inside hepatic cells to form schizonts while others persist and remain dormant as hypnozoites (resting forms). These dormant forms may get activated and release merozoites, which go on infecting RBCs producing true clinical relapse. It usually occurs after 6 to 12 months of complete cure from the disease.

Sold by @itachibot

In case of P.falciparum and P.malariae, the initial tissue phase disappears completely and no hypnozoites are found, hence relapse is not seen.

In Plasmodium falciparum, recrudescence can be seen which is defined as the recurrence of asexual parasitemia. It usually occurs within 8 weeks of treatment and is due to incomplete treatment or drug resistance.

Solution to Question 16:

Enterobius vermicularis eggs are non-bile stained and the others are bile stained.

Enterobiasis commonly occurs in young children and is characterised by anal pruritus. The worm may migrate up to the uterus, fallopian tubes and into the peritoneum in adult women. This can lead to:

• Chronic salpingitis

• Cervicitis

• Peritonitis

• Recurrent urinary tract infections.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 17:

The eggs of Strongyloides stercoralis will not float in a saturated salt solution.

Concentration methods are used when the parasites in stool are scanty and difficult to identify in routine stool examination. It is of 2 types:

• Floatation methods- the feces are suspended in a solution of high specific gravity and eggs will float up. e.g. saturated salt solution technique, zinc sulfate centrifugal floatation technique.

• Sedimentation Methods - the feces are suspended in a solution with low specific gravity and eggs will sediment at the bottom. e.g. formol-ether sedimentation technique.

Sold by @itachibot

Protozoology - Amoebae, Ciliates & Flagellates

Question 1:

The life cycle given below is of which of the following:



a) E. vermicularis

b) A. duodenal

c) B. coli

d) E. histolytica

Question 2:

A 40-year-old man is diagnosed to have amoebic dysentery. Which of the following is the infective form of the protozoan causing this disease?

a) Trophozoite

b) Mature quadrinucleate cyst

c) Metacyclic trophozoite

d) Precyst

Question 3:

Sold by @itachibot

A 25-year-old woman presented with complaints of abdominal pain and 6 episodes of passing foul-smelling dark stool mixed with mucus and blood streaks. There is no history of fever and tenesmus. On examination, vitals are stable and localized tenderness is present in the right iliac fossa. Which of the following is most likely to cause this condition?

a) Giardia lamblia

b) Entamoeba histolytica

c) Campylobacter jejuni

d) Shigella dysenteriae

Question 4:

You received a stool sample of a patient suspected to have dysentery in the microbiology lab. On microscopy, you observe the structures shown in the image. What other findings you will expect to see in this stool sample?



a) RBCs as discrete or in rouleaux

b) Charcot-Leyden crystals

c) Abundant cellular exudates

d) Absent eosinophils

Question 5:

A patient with abdominal pain and mucus in stool is diagnosed to have intestinal amoebiasis. Which of the following is the most common extraintestinal complication of this disease?

Sold by @itachibot

a) Hepatic amoebiasis

b) Pulmonary amoebiasis

c) Amoebic appendicitis

d) Splenic abscess

Question 6:

A 40-year-old man was brought to the casualty with complaints of nausea, vomiting and disorientation. His brother tells you they had recently gone on a camping trip where they had swum in a freshwater lake. On examination, Kernig's sign is positive. A spinal tap is performed and the findings of CSF wet mount microscopy are shown below. Which of the following is the most likely causative organism?



a) Naegleria fowleri

b) Acanthamoeba spp.

c) Angiostrongylus cantonensis

d) Balamuthia mandrillaris

Question 7:

A college student presents with ocular redness, foreign body sensation, and corneal ulcers following the use of contact lens. The saline mount preparation of the corneal scraping shows polygonal cysts. What would be the most probable cause?

a) Acanthamoeba

Sold by @itachibot

b) Naegleria

c) Entamoeba

d) Onchocerca

Question 8:

Which of the following protozoan parasites is found in the lumen of the human small intestine?



a) 1

b) 2

c) 3

d) 4

Question 9:

A 30-year-old HIV-positive patient presented with a chronic history of watery diarrhea. The histopathology of the ■duodenum is shown in the image below. What is the likely diagnosis?

Sold by @itachibot



a) Giardiasis

b) Echinococcosis

c) Isosporiasis

d) Cryptosopridiasis

Sold by @itachi... purchased this from some... else, may have been scammed.

Question 10:

A patient is undergoing the test depicted in the image given below. Which of the following diseases does it help to diagnose?



a) Amoebiasis

Sold by @itachibot

b) Trichomoniasis

c) Giardiasis

d) Echinococcosis

Question 11:

The findings of the microscopic examination of the wet mount preparation of stool of a patient with diarrhoea is shown below. Which of the following organisms will exhibit this type of motility?



a) Vibrio cholera

b) Giardia lamblia

c) Campylobacter jejuni

d) Listeria monocytogenes

Question 12:

A patient came to the gynecology OPD with complaints of foul-smelling frothy vaginal discharge and intense itching. The per speculum finding is given below. Which of the following statements is false regarding the causative organism of this disease?

Sold by @itachibot



a) It exists in cyst and trophozoite form

b) It is an obligate parasite

c) The gold standard for diagnosis is culture on CPLM or PEM

d) It can cause male infertility

Question 13:

A 12-year-old child presented to you with fever, malaise, and weight loss. On examination, you notice the finding shown in the image. In which of the following conditions is this seen?

a) West African sleeping sickness

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

b) Chagas disease

c) Leishmaniasis

d) Babesiosis

Question 14:

Which of the following statements is false regarding Trypanosoma?

a) Tsetse fly is the vector for both T. rhodesiense and T. gambiense

b) T. gambiense exhibits antigenic variation

c) T. rhodesiense can cause myocarditis

d) Humans are the reservoir for T. rhodesiense

Question 15:

A middle-aged woman presents to the OPD with fever and malaise for the past 4 months. On examination, the following ocular finding is noted along with hepatosplenomegaly and lymphadenopathy. What is the most probable diagnosis?

a) West African Sleeping Sickness

b) Chagas disease

c) Toxoplasmosis

d) East African Sleeping Sickness

Sold by @itachibot

Sold by @itachibot

Question 16:

Which of the following is used for the culture of Trypanosoma?

a) NNN medium

b) Boeck and Drbohlav medium

c) Craig's medium

d) NIH polygenic medium

Question 17:

The given life cycle belongs to which of the following organisms?



a) Trypanosoma brucei rhodesiense

b) Trypanosoma cruzi

c) Trypanosoma brucei gambiense

d) Trypanosoma rangeli

Question 18:

The given life cycle belongs to which of the following organisms?

Sold by @itachibot



a) Trypanosoma brucei gambiense

b) Trypanosoma brucei rhodesiense

c) Leishmania donovani

d) Babesia microti

Sold by @itachi... If you purchased this from someone else, may have been scammed.

Question 19:

You are performing the following diagnostic procedure in a patient having fever for the past 2 weeks to find the causative organism. Which disease are you most likely testing for?



a) Kala-azar

Sold by @itachibot

b) Schistosomiasis

c) Toxoplasmosis

d) Chagas disease

Question 20:

What is the vector for the condition diagnosed by the image shown below?



a) Anopheles mosquito

b) Ixodes Tick

c) Sandfly

d) Tsetse fly

Question 21:

A 15-year-old boy from Delhi presented to the OPD with the finding seen in the image. It is diagnosed to be an oriental sore. Which of the following causes this?

Sold by @itachibot

a) Leishmania mexicana complex

b) Leishmania braziliensis complex

c) Leishmania donovani

d) Leishmania tropica complex

Question 22:

A medicine resident is evaluating a patient with abdominal pain and diarrhoea. The stool wet mount finding is given below. What is the most likely causative organism?



a) Balantidium coli

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) Isospora belli

c) Cryptosporidium parvum

d) Enteromonas hominis

Question 23:

What is the investigation of choice for the diagnosis of kala-azar?

a) Rapid dipstick test

b) Aldehyde test of Napier

c) ELISA

d) Indirect fluorescent antibody test

If you bought this from someone else, you've been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	d
2	b
3	b
4	b
5	a
6	a
7	a
8	b
9	a
10	c
11	b
12	a
13	a
14	d
15	b
16	a
17	b
18	c
19	a

Sold by @itachibot

20	c
21	d
22	a
23	a

Detailed Explanations

Solution to Question 1:

The given life cycle is of Entamoeba histolytica.

Its life cycle includes 2 stages - the motile trophozoite and the non-motile cyst, both of which are seen in the image shown above. The transmission is primarily by the fecal-oral route. There is no animal reservoir.

The ingested cysts differentiate into trophozoites in the ileum which invade the colonic epithelium. The trophozoites penetrate to the submucosal layer and form abscesses which break down to form flask-shaped ulcer.

The trophozoites multiply by binary fission and produce cysts, and both stages are passed in the feces. Trophozoites are rapidly destroyed outside the body. Cysts survive in the external environment and are responsible for transmission.

The most frequent site of systemic disease is the liver.

Note: The genus Entamoeba has four morphologically identical species - E. histolytica, E. dispar, E. bangladeshi, and E. moshkovskii. E. moshkovskii which was initially thought to be non-pathogenic has recently been shown to be a causative agent of non-invasive diarrhea.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 2:

The mature quadrinucleate cyst is the infective form of Entamoeba histolytica causing amoebic dysentery. It is passed in feces and can remain viable under moist conditions for about 10 days.

The following are the characteristic features of various life stages of Entamoeba histolytica:

The trophozoite is found only in tissues. Its cytoplasm has a ground-glass appearance. The nucleus has a central karyosome with radiating fibrils called linin network giving a cartwheel appearance. Pseudopodia are present which exhibit gliding motility.

The below image shows trophozoite form.

Sold by @itachibot



The precystic stage has a large glycogen vacuole and 2 chromatid bars. It secretes a refractile cyst wall to become a cyst.

The cystic stage has a mature cyst with 4 nuclei and no glycogen mass. Chromatid bars disappear.

The below image shows cystic form.



Solution to Question 3:

The given clinical scenario is suggestive of amoebic dysentery and the causative agent is Entamoeba histolytica.

Clinical features of intestinal amebiasis:

• Acute intestinal amebiasis- dysentery, lower abdominal discomfort, and flatulence.

Sold by @itachibot

• Chronic amebiasis- low-grade symptoms such as occasional diarrhea, weight loss, and fatigue.

• The amoebic ulcer which is flask-shaped is the typical lesion seen in intestinal amoebiasis. The ulcers are multiple and are confined to the colon. Sometimes, in secondary intestinal lesions, the ulcers are numerous in the cecum, followed by the sigmoido-rectal region.

A granulomatous lesion called an ameboma may form in the cecal or rectosigmoid areas of the colon which can resemble an adenocarcinoma of the colon.

The most appropriate treatment is intravenous metronidazole.

Option A - Giardia is often asymptomatic, but in some cases can lead to fat malabsorption, dull epigastric pain, and flatulence.

Option C - Campylobacter presents with fever, pain, and watery diarrhea. The stool contains leucocytes and blood.

Option D - Shigella causes bacillary dysentery which is characterised by fever and generalised abdominal tenderness.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 4:

The given image is showing erythrophagocytosis (trophozoite with ingested RBCs) and this in the stool sample of a patient with features of dysentery is indicative of amoebic dysentery. The other stool microscopic finding seen in amoebic dysentery is Charcot-Leyden crystals.

The ingested erythrocytes appear as dark inclusions. Erythrophagocytosis is the only characteristic that can be used to morphologically differentiate E. histolytica from the nonpathogenic entamoeba.

Amoebic dysentery:

Clinically, it is insidious in onset with localized abdominal tenderness. Tenesmus, toxicity, and fever are absent.

Under microscopy, RBCs are clumped, yellowish-brown in color. Charcot-Leyden crystals and eosinophils are present.

Charcot-Leyden crystals are diamond-shaped, clear, and refractile structures that may be seen in stool specimens under microscopy and represent breakdown products of eosinophils.

The below image shows Charcot-Leyden crystals:

Sold by @itachibot

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 5:

The most common extraintestinal complication of amoebiasis is hepatic amoebiasis.

In intestinal amoebiasis, most of the trophozoites reach the liver. However, only a small portion of them actually lodge and multiply in the liver cells. Patients usually present with an enlarged, tender liver without detectable impairment of liver function called amoebic hepatitis.

Few patients go on to develop amoebic liver abscess. The center of the abscess contains thick chocolate brown pus (anchovy sauce pus), which is made up of liquefied necrotic liver tissue. It is bacteriologically sterile and free of amoeba. The periphery is made up of nearly normal liver tissue containing invading amoeba.

Complications and sequelae of intestinal amoebiasis:

- Fulminant amoebic colitis
- Toxic megacolon
- Perianal ulceration
- Amoeboma
- Extraintestinal amoebiasis
- Hepatic amoebiasis (most common)- Amoebic hepatitis and Amoebic liver abscess
- Amoebic appendicitis and peritonitis
- Pulmonary amoebiasis
- Cerebral amoebiasis
- Splenic abscess
- Cutaneous amoebiasis
- Genitourinary amoebiasis

Sold by @itachibot



Sold by @itachi... if you purchased this from some... you may have been scammed. ...else,

Solution to Question 6:

The given clinical scenario and microscopy showing amoeboid trophozoite form with pseudopodia is suggestive of primary amoebic meningoencephalitis. It is caused by Naegleria fowleri.

It is also known as brain-eating amoeba. It occurs in 3 forms: cyst, amoeboid trophozoite and flagellate trophozoite forms.

The infection usually follows swimming in freshwater rivers or ponds. The amoebae invade the nasal mucosa and pass through the olfactory nerve branches in the cribriform plate into the meninges. Incubation period is 2 days to 2 weeks.

Treatment is with intravenous amphotericin B.

Options B & D: Both Acanthamoeba and B.mandrillaris cause granulomatous amoebic meningoencephalitis which is a subacute condition.

Option C: A. cantonensis is the most common cause of eosinophilic meningitis in Southeast Asia. It is a nematode(roundworm).

Solution to Question 7:

The given clinical history and the demonstration of polygonal cysts are suggestive of amoebic keratitis caused by acanthamoeba.

Acanthamoeba culbertsoni is a free-living amoeba. It exists as trophozoite form and resistant cystic form. Both these forms are infective. The trophozoite is large and characterized by spine-like pseudopodia (acanthopodia).

Human beings acquire the infection by inhalation of cysts or trophozoites, ingestion of cysts, or directly through traumatized skin or eyes.

Sold by @itachibot

It presents as two chronic conditions- acanthamoeba keratitis and granulomatous amoebic encephalitis.

Diagnosis of amoebic keratitis is made by the demonstration of the cyst in corneal scrapings by wet mount, histology, and culture.

The below image shows the ring lesion seen in acanthamoeba keratitis.



The below image shows cysts of acanthamoeba in lactophenol cotton blue staining.



Solution to Question 8:

Image 2 shows the trophozoite of Giardia lamblia, which is the only protozoan parasite found in the lumen of the human small intestine. It lives in the duodenum and upper jejunum.

Sold by @itachibot

Other protozoan parasites found in the human small intestine do not occur in the lumen but are present in the epithelial cells lining the small intestine. These include:

• Isospora belli

• Cyclospora cayetanensis

• Cryptosporidium parvum

• Sarcocystis hominis

Options A and C: Entamoeba histolytica and Balantidium coli are found in the human large intestine.

Option D: Trichomonas vaginalis is found in the genital tract.

The image given below shows the trophozoite of Giardia lamblia.

The image given below shows the trophozoite of Entamoeba histolytica.



Sold by @itachibot

The image given below shows the trophozoite of Balantidium coli.



The image given below shows the trophozoite of Trichomonas vaginalis.

Solution to Question 9:

The given clinical scenario with the histopathology showing trophozoites of Giardia lamblia in the duodenal wall is suggestive of giardiasis.

Giardia lamblia, also called Giardia duodenalis, is the causative agent of giardiasis. It is the only common pathogenic protozoan found in the duodenum and jejunum of humans.

Infection occurs by the ingestion of mature cysts in contaminated water and food and is transmitted by the faeco-oral route.

Sold by @itachibot

It is weakly pathogenic in humans. Secretory IgA and IL-6 are responsible for clearing the infection. Hence, giardiasis usually occurs in immunosuppressed, agammaglobulinemic, or malnourished individuals.

Clinical features:

• Diarrhea

• Fat malabsorption

• Dull epigastric pain

• Belching and flatulence

• Iron deficiency and growth stunting.

• Stool will have excess mucus and fat but no blood.

Detection of antigen by ELISA is more sensitive and specific than microscopy. Entero test is also used in the diagnosis.

Histopathology of the duodenum reveals several heart-shaped or pear-shaped trophozoites with two equally sized nuclei. The small intestinal morphology is usually normal. However, in heavy infections, there may be villous blunting and intraepithelial infiltrates.

Metronidazole/tinidazole/nitazoxanide are the first-line agents in the treatment.

The image given below shows a magnified view of the pear-shaped trophozoites of Giardia lamblia.



Sold by @itachibot
If yo
this from someone else,
en scammed.

Solution to Question 10:

The given image depicts a string test or entero test and is used in the diagnosis of giardiasis.

It is a useful method for obtaining the duodenal specimen in giardiasis.

Sold by @itachibot

In this test, a coiled thread inside a small weighted gelatin capsule is swallowed by the patient, after attaching the free end of the thread in the cheek. The capsule passes through the stomach to the duodenum. After 2 hours, the thread is withdrawn, placed in saline, and is shaken. The centrifuged deposit of the saline is then examined for giardia.

Solution to Question 11:

The given wet mount preparation of stool shows organisms having falling leaf motility. It is exhibited by Giardia lamblia.

The trophozoite is motile with a slow oscillation along its long axis resembling a falling leaf.

Solution to Question 12:

The given clinical scenario with the image showing strawberry cervix is suggestive of Trichomonas vaginalis infection. It exists only in trophozoite form.

Trichomonas vaginalis is the only protozoan with one stage in its life cycle ie., trophozoite form (no cyst stage is seen). It is an obligate parasite and cannot live without close association with vaginal, urethral or prostatic tissues.

The common presentation of trichomoniasis is foul-smelling frothy vaginal discharge, itching, and dysuria. The gold standard for diagnosis is culture on CPLM (cysteine-peptone-liver-maltose) or PEM (plastic envelope medium). The treatment of choice is metronidazole.

T. vaginalis infection in women may lead to infertility, preterm birth and low-birth-weight infants. In men, trichomoniasis has been also associated with infertility through inflammatory damage to the genitourinary tract or interference with sperm function.

The below image shows the trophozoite form of Trichomonas vaginalis.



Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Solution to Question 13:

The given image shows posterior cervical lymphadenopathy which is called Winterbottom's sign. It is seen in West African sleeping sickness.

It is seen in stage I of West African sleeping sickness caused by Trypanosoma brucei gambiense.

The table below shows diseases caused by various Trypanosoma species.

Protozoa	Disease
Trypanosoma brucei gambiense	West African sleeping sickness
Trypanosoma brucei rhodesiense	East African sleeping sickness
Trypanosoma cruzi	Chagas' disease/South American trypanosomiasis

Solution to Question 14:

Humans are the reservoir for T. gambiense, not for T. rhodesiense.

The table below shows characteristic differences between West African and East African trypanosomiasis.

Parasites causing myocarditis include:

- Trypanosoma brucei rhodesiense
- Trypanosoma cruzi
- Toxoplasma gondii
- Echinococcus granulosus
- Trichinella spiralis

Characteristics	West African trypanosomiasis	East African trypanosomiasis
Organism	T. brucei gambiense	T. brucei rhodesiense
Vector	Tsetse fly	Tsetse fly
Reservoir	Humans	Wild and domestic animals
Course of disease	Chronic	Acute
Parasitemia	Low	High
Virulence	Less	More

Sold by @... If you purchased this from ... you may have been scammed. ... else,

Sold by @itachibot

Characteristics	West African trypanosomiasis	East African trypanosomiasis
Mortality	Low	High

Solution to Question 15:

The given clinical scenario with the image showing Romana's sign is suggestive of Chagas disease.

It is caused by T. cruzi. Reduviid bugs are vectors for Chagas disease.

Inoculation of T. cruzi in conjunctiva causes unilateral, painless edema of periocular tissues in the eye is called Romana's sign. This is classically seen in acute Chagas disease that lasts for 1 to 4 months.

The following image shows a patient with Romana's sign.

Acute Chagas disease is also characterised by the presence of chagomas which are subcutaneous lesions occurring at the site of inoculation.

Chronic Chagas disease is characterized by megaesophagus, megacolon, and myopathy.

Shown below is an image of the Reduviid bug.

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

NNN (Novy, McNeal, and Nicolle) medium is used for the culture of Trypanosoma brucie and cruzie. It is also used for the culture of Leishmania.

The other media used in the culture of Trypanosoma are Weinman's medium and Tobie's medium.

Options B, C & D: Boeck and Drbohlav medium, Craig's medium and NIH polygenic medium are used for the culture of E.histolytica.

Solution to Question 17:

The given life cycle belongs to Trypanosoma cruzi.

T. cruzi is the causative organism of Chagas disease or South American trypanosomiasis. They are intracellular parasites.

It passes its life cycle in two hosts- the definitive host is the man and the intermediate host (vector) is reduviid bug or triatomine bugs.

Metacyclic trypomastigotes are the infective forms to the man that are found in feces of reduviid bugs. Transmission occurs when mucus membranes, conjunctiva, or wound on the surface of the skin are contaminated by feces of the bug.

Inside the definitive host, the trypomastigotes differentiate into intracellular amastigotes. The amastigotes multiply by binary fission and differentiate into trypomastigotes, which are then released into the bloodstream. These are the infective stage for triatomine bug. No multiplication occurs in this stage.

Inside the triatomine bug, the trypomastigotes are transformed into epimastigotes and then into metacyclic trypomastigotes in the hindgut which are excreted in feces.

Sold by @itachibot

Note: Amastigote forms resemble the Leishmania spp. and hence they are frequently known as Leishmanial forms. They are found in muscular tissue, nervous tissue, and reticuloendothelial system.

Solution to Question 18:

The given life cycle belongs to Leishmania donovani.

The definitive hosts are man, dog, and other mammals and the vector/intermediate host is female sandfly. Promastigotes are infective forms to man.

L. donovani exists in 2 forms:

• Amastigote form- in humans and other mammals

• Promastigote form- in the sandfly and artificial culture

When the infected sandfly bites a person, infectious promastigotes are released into the host dermis along with the sandfly saliva, which potentiates parasite infectivity.

Amastigote form (as shown in the image below) is also known as Leishman-Donovan (LD) body. It is an intracellular form found inside macrophages, monocytes and reticuloendothelial cells. It contains a kinetoplast which consists of a parabasal body and a blepharoplast. Flagellum is absent.



Promastigote form (as shown in the below image) is present in the midgut of the female sandfly. The kinetoplast lies transversely near the anterior end. A single flagellum is seen. It is the infective form of Leishmania donovani for humans.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 19:

The given image shows bone marrow aspiration and it is most commonly done to diagnose Kala-azar.

The microscopic demonstration of amastigotes from bone marrow is the gold standard for the diagnosis of Kala-azar

Generally, the sternal marrow is aspirated by puncturing the sternum at the level of the 2nd or 3rd intercostal space, using a sternal puncture needle.

Other samples collected are peripheral blood, splenic aspirate, and enlarged lymph node.

In kala-azar:

• Most common specimen collected for microscopy- bone marrow aspirate.

• Most sensitive method- splenic aspirate.

Option B: Schistosomiasis is typically diagnosed by stool examination for parasite eggs.

Option C: Toxoplasmosis is usually diagnosed based on serologic testing.

Option D: Chagas disease is diagnosed by peripheral blood smear examination, culture and examination of aspirates from lymph nodes and chagoma.

Solution to Question 20:

The image shown is of the Leishman-Donovan bodies whose presence are diagnostic of Kala Azar. The vector of Kala Azar is a sandfly.

Leishman-Donovan bodies are amastigotes with a double dot appearance containing a nucleus and small rod-like kinetoplast. They are present inside the macrophage.

Sold by @itachibot

Solution to Question 21:

The given image shows cutaneous leishmaniasis and is caused by Leishmania tropica complex. It is also called oriental sore/Delhi boil/Bagdad boil/Aleppo button. It is seen mainly among children in endemic areas.

Old world Leishmania tropica complex includes 3 species:

• L. tropica

• L. major

• L. aethiopica

New world Leishmania tropica complex includes 3 species:

• L. mexicana complex

• L. braziliensis complex

It begins as a raised papule that grows into a nodule which then ulcerates over some weeks. The margins of the ulcer are raised and indurated. The ulcer is usually painless unless secondary bacterial infection occurs.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 22:

The given image shows the trophozoite form of the Balantidium coli. Living trophozoites/cysts are yellowish or greenish in color on wet mount.

Balantidium coli occurs in two stages:

• Trophozoites are oval in shape and contain two nuclei. There is the presence of cilia over the entire surface and are larger around the mouth. Its anterior end is narrow and the posterior end is broad. At the anterior end, there is a groove leading to the mouth (cytostome). Posteriorly, there is a small anal pore (cytopyge).

• Cysts are smaller, round with a micronucleus, a macronucleus and many contractile vacuoles.

Solution to Question 23:

The rapid dipstick test is the method of choice for the diagnosis of kala-azar.

The following tests are used for the laboratory diagnosis of kala-azar:

• Rapid diagnostic tests - rK39 and dipstick test strip.

• Parasitological diagnosis - demonstration of LD bodies on microscopy is confirmatory

• Surveillance purposes - Aldehyde test of Napier

Sold by @itachibot

• Serological diagnosis - Rapid dipstick test, direct agglutination test, ELISA, indirect fluorescent antibody test

• Assessing immune status - Leishmanin (Montenegro) test

• Induration of ≥ 5 mm is considered positive

• Usually becomes positive 4-6 weeks after onset of cutaneous leishmaniasis

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Protozoology - Sporozoa

Question 1:

A 55-year-old man comes with complaints of fever and increasing fatigue for the past week. On examination, he is febrile, hypotensive and tachycardic. The findings of microscopic examination of his peripheral blood smear is given below. What is the causative agent?



a) Plasmodium vivax

b) Plasmodium falciparum

c) Plasmodium ovale

d) Plasmodium malariae

Question 2:

During your rural posting, a patient presents with a high fever and chills. Malaria is endemic in the area so you order a peripheral smear. While examining the peripheral blood smear, you note that the schizont stage is not seen. Which of the following organisms is implicated here?

a) Plasmodium vivax

b) Plasmodium malariae

c) Plasmodium falciparum

d) Plasmodium ovale

Sold by @itachibot

Question 3:

Which of the following Plasmodium spp. has the longest incubation period?

a) Plasmodium vivax

b) Plasmodium falciparum

c) Plasmodium malariae

d) Plasmodium ovale

Question 4:

Which of the following Plasmodium spp. mainly affects the old RBCs?

a) Plasmodium vivax

b) Plasmodium falciparum

c) Plasmodium malariae

d) Plasmodium ovale

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 5:

A microbiologist is examining the slide shown in the following image under the microscope. Which of the following is false regarding this?



©MARROW

a) 'B' is used for determining the species

b) 'B' is more sensitive than 'A'

Sold by @itachibot

c) 'A' is air dried rapidly and stained with one of the Romanowsky stains

d) For 'B' to be considered negative, at least 200 oil immersion fields should be examined

Question 6:

A 15-year-old boy presents to the OPD with high-grade fever, chills, and rigors for the past 3 days. A peripheral blood smear revealed the following. Which of the following is false regarding the diagnostic methods used to detect the condition he is likely suffering from?



a) Jaswant Singh Bhattacharya stain is used for peripheral smear examination

b) HRP-2 antigen test is done for detection of P.vivax

c) Acridine orange dye is used in quantitative buffy coat method

d) Rapid diagnostic tests have lesser sensitivity than microscopy

Question 7:

A 50-year-old man with a 5-day history of fever is brought to the casualty by his son after an episode of seizures. He is diagnosed with malignant tertian malaria. Which of the following complications will you not expect to see in this patient?

a) Hypoglycemia

b) Acute renal failure

c) Pulmonary edema

d) Metabolic alkalosis

Sold by @itachibot

Question 8:

A 24-year-old patient presents with a high-grade fever, headache, and weakness for 5 days. He gives a history of blood transfusion 4 months back. The microscopic examination of the thin blood smear is given below. What is the most probable causative agent?



a) Plasmodium knowlesi

b) Babesia microti

c) Plasmodium falciparum

d) Plasmodium malariae

If you purchased this, you may have been

Question 9:

Sabin-Feldman dye test is used for diagnosis of which of the following condition?

a) Paragonimiasis

b) Hydatid cyst

c) Toxoplasmosis

d) Cryptococcus

Question 10:

Which of the following statements is true regarding Toxoplasma gondii?

a) Humans are the definitive host

Sold by @itachibot

b) Oocysts are formed within the intermediate host

c) The tachyzoites inside the tissue cyst can be reactivated on immunosuppression

d) Human toxoplasmosis is a zoonosis

Question 11:

An HIV positive pregnant woman was brought to the ER after an episode of generalized tonic-clonic seizures. CECT was done which showed multiple enhancing lesions. A stereotactic biopsy of the cerebral lesions was done and the image is given below. Which of the following is the most appropriate drug for her treatment?



a) Trimethoprim

b) Sulfadadiazine

c) Spiramycin

d) Albendazole

Question 12:

A Kinyoun-positive oocyst (25 µm) isolated from an HIV patient with diarrhea is shown in the image. What is the likely organism implicated here?

Sold by @itachibot



a) Cystoisospora

b) Cyclospora

c) Cryptosporidium

d) Microspora

Sold by @itachi... purchased this from some... else, ...u may have been scammed.

Question 13:

A patient who underwent a renal transplant 3 months ago came with complaints of diarrhea for the last 9 days. The stool sample subjected to Kinyoun staining revealed 4-6 µm structures as shown below. Which of the following is true about the causative organism?



a) It does not shows autoinfection

Sold by @itachibot

b) It is an obligate intracellular organism

c) Oocysts are immediately infective after being excreted in stool

d) Metronidazole is the drug of choice

Question 14:

Match the following:

a) A-2, B-1, C-3, D-4

b) A-3, B-2, C-1, D-4

c) A-4, B-1, C-2, D-3

d) A-1, B-2, C-4, D-3

If you bought this from someone else, you've been scammed. Sold by @itachibot

Answer Key

Question No.	Correct Option
1	b
2	c
3	c
4	c
5	a
6	b
7	d
8	b
9	c
10	d
11	c
12	a
13	c
14	b

Detailed Explanations

Solution to Question 1:

Sold by @itachibot

The given clinical history and the blood smear showing multiple ring forms in RBCs are suggestive of malaria caused by Plasmodium falciparum.

Falciparum malaria infects both young and mature erythrocytes. Binucleate rings (double chromatin) seen in RBCs are common and they resemble stereo headphones in appearance. The rings are often attached to the margins of the red cells and are described as form applique or accole. There may be several rings present within a single erythrocyte.

Hypnozoites are not present in P. falciparum infection.

The peripheral smear finding of P. falciparum malaria are:

• Multiple ring form

• Accole form

• Absent schizont

• Banana shaped gametocytes

• Maurer's dots

The below image shows banana shaped gametocyte:



If yo...itachibot...someone else,...ed.

Solution to Question 2:

The schizont stage is usually not seen in the peripheral smear of individuals having Plasmodium falciparum malaria.

A mature schizont contains merozoites and hemozoin. This will burst to release merozoites into circulation. The merozoites invade fresh erythrocytes, in which they go through the same process of development called erythrocytic schizogony or merogony.

After the ring form is formed in peripheral blood, the remaining RBC cycle of falciparum occurs in the capillaries of the brain and other internal organs. Hence only the ring form is seen in the peripheral smear.

Sold by @itachibot

However, for other Plasmodium species, all erythrocytic stages occur in peripheral blood vessels.

The below image shows the life cycle of Plasmodium spp.



Solution to Question 3:

Plasmodium malariae has the longest incubation period of 28 days.

The incubation period is the interval between the infective mosquito bite and the first appearance of clinical symptoms. The average incubation periods for Plasmodium species are:

• P. vivax: 8-31 days (14 days)

• P. falciparum: 8-14 days (12 days)

• P. ovale: 8-31days (14 days)

• P. malariae: 18-40 days (28 days)

Note: The pre-patent period is the interval between the entry of the parasites into the host and the time when they first become detectable in blood.

Solution to Question 4:

Plasmodium malaria mainly affects the old RBCs.

Options A and D- P. vivax and P. ovale infect young RBCs and reticulocytes.

Option B- P. falciparum infects RBCs of all ages.

The degree of parasitization is not heavy for P. vivax and P. malariae while very high intensity of parasitization is seen in infection with P. falciparum.

Sold by @itachibot
If you purchased this from elsewhere, you may have been scammed.

Sold by @itachibot

Solution to Question 5:

In the given image 'A' is a thin smear and 'B' is a thick smear. The thin smear is used for determining the species, not the thick smear.

The two types of smears are prepared from the peripheral blood are:

• Thin smear– It is used for detecting the parasites and determining the species. They are air-dried rapidly and stained with Romanowsky stains.

• Thick smear– It is more sensitive and examined under oil immersion microscope. It is more suitable for the rapid detection of malarial parasites. Species identification is difficult. It is recommended that 200 oil immersion fields should be examined before a thick film is declared negative.

The image below shows the thin and thick smears.



Solution to Question 6:

The given clinical scenario and the peripheral smear showing ring forms are suggestive of P. falciparum malaria. HRP-2 antigen detection used in rapid diagnostic tests for malaria is specific for P. falciparum.

Jaswant Singh Bhattacharya (JSB) stain is routinely used in India to stain peripheral blood smears for the diagnosis of malaria.

Acridine orange dye is used in the quantitative buffy coat (QBC) method of diagnosis which provides better yield compared to peripheral smears.

Rapid diagnostic tests are based on the detection of antigens using immunochromatographic methods. They are less sensitive than microscopy and also cannot differentiate between species except P. falciparum. The 2 antigens detected are:

• pLDH (parasite lactate dehydrogenase)- pan malarial antigen

Sold by @itachibot

• HRP-2 (histidine-rich protein 2)- specific for P.falciparum.

Solution to Question 7:

Metabolic acidosis (and not alkalosis) is a complication of malignant tertain malaria caused by Plasmodium falciparum.

Complications of falciparum malaria include:

• Metabolic acidosis
• Cerebral malaria
• Algid malaria
• Septicemic malaria
• Blackwater fever
• Pulmonary edema
• Acute renal failure
• Hypoglycemia
• Hyperpyrexia
• Severe anemia
• Shock

If you purchased this from someone else, you may have been scammed.
Sold by @Itachibot

Solution to Question 8:

The given clinical history and the microscopy showing Maltese crosses or tetrads are suggestive of babesiosis caused by Babesia microti.

The definitive host for babesia are ixodid ticks. Man is an incidental, dead-end host.

On microscopic examination, babesia trophozoites appear as intraerythrocytic round or pyriform or ring form. The ring form is the most common and it lacks the central hemozoin deposit which differentiates it from P. falciparum.

Other distinguishing features are the absence of schizonts and gametocytes and the presence of tetrads/Maltese crosses, which are pathognomic of B. microti or B. duncani.

The below image shows Maltese cross appearance of Babesia microti.

Sold by @itachibot



If you purchased this from some- else, you may have been scammed. Sold by @itachi

Solution to Question 9:

Sabin-Feldman dye test is used for the diagnosis of toxoplasmosis.

It is based on the principle that the presence of anti-toxoplasma antibodies in the patient's serum inhibits the staining of trophozoites by alkaline methylene blue dye.

Toxoplasma trophozoites and complement are added to the patient's serum which is then incubated for an hour at 37°C. After incubation, methylene blue dye is added.

If anti-toxoplasma antibodies are present in the serum, these are activated by complement and the parasite membrane is lysed. Thus, the trophozoites are not stained (positive result).

If there are no antibodies, trophozoites will take up the stain and appear blue under the microscope. This is considered to be a negative result.

The dilution of the test serum at which 50% of the tachyzoites are not stained is reported as the antibody titer of the test serum. The test is highly sensitive and specific.

It cannot differentiate between recent or past infections as it detects only IgG antibodies.

Solution to Question 10:

Human toxoplasmosis is a zoonosis. It is transmitted by cats.

Toxoplasma gondii has two types of life cycles:

• Enteric cycle

• Exoenteric cycle

The enteric cycle occurs in definitive hosts such as cats. Exoenteric cycle occurs in intermediate hosts such as humans, mice, rats, etc.

Sold by @itachibot

Both asexual reproduction (schizogony) and sexual reproduction (gametogony) occur within the mucosal epithelial cells of the small intestine of the cat. The tachyzoites and tissue cyst represent stages in asexual multiplication, while the oocyst is formed by sexual reproduction.

All three forms, tachyzoites, tissue cyst, and oocyst occur in the definitive host. Only the asexual forms, tachyzoites, and tissue cysts are present in the intermediate host.

Tachyzoites continue to multiply and spread locally by the lymphatic system and blood, while the bradyzoites multiply slowly inside the tissue cyst. The dormant bradyzoites inside the cyst may be reactivated on immune suppression causing renewed infection in the host.

The below image shows the life cycle of Toxoplasma gondii.



Solution to Question 11:

Sold by @itachibot

The given clinical scenario and the image showing lymphocytes, macrophages, and eosinophils surrounding the toxoplasma bradyzoites which are encased in cysts are suggestive of cerebral toxoplasmosis. Spiramycin is the drug of choice during pregnancy.

Cerebral toxoplasmosis is an opportunistic infection commonly found in HIV +ve patients and usually occurs when CD4+ T cell counts <100/µL.

It appears as ring enhancing lesions on contrast enhanced CT. Calcification may also be seen in old lesions.

Sabin-Feldman dye test is a serological test done for the detection of antibodies against toxoplasma. IIF (indirect immunofluorescence), IHA (indirect heamagglutination), and CFT (complement fixation test) are also used for the diagnosis of toxoplasmosis. Dye test remains positive throughout life, while CFT becomes negative soon after acute infection.

ELISA is most commonly used and IgM and IgG antibodies are estimated separately.

Standard treatment is sulfadiazine/pyrimethamine and leucovorin for a minimum of 4–6 weeks. The drug of choice for toxoplasmosis in pregnancy is spiramycin.

Sold by @itachibot
purchased this from someone else,
may have been scammed.

Solution to Question 12:

The given clinical scenario with diarrhea in an HIV patient is suggestive of a coccidian parasitic infestation. Kinyoun-positive oocysts with the size of 25 µm in the image are formed by Cystoisospora.

Comparison of coccidian parasites causing diarrhea in an immunocompromised host:

The below image shows oocysts of Cyclospora cayetanensis in the modified acid-fast stain.

Property	Cryptosporidium	Cyclospora	Cystoisospora
Sporulated oocyst	4-6 µm, round, contains four sporozoites	8-12 µm, round, contains 2 sporocysts, each having two sporozoites	23-36 µm, oval, contain two sporocysts, each having four sporozoites
Acid-fast detection	Uniformly acid fast	Variable acid fast	Uniformly acid fast
Sporulation of the oocyst	Occurs inside the host cell	Occurs in soil	Occurs in soil
Diagnostic form	Sporulated cyst	Unsporulated cyst	Unsporulated oocyst

Sold by @itachibot



The below image shows oocysts of Cryptosporidium parvum in the modified acid-fast stain.



Solution to Question 13:

The given clinical scenario with Kinyoun stain positive oocysts of size 4-6 µm is suggestive of cryptosporidiosis. The oocysts are immediately infective after being excreted in the stool.

Cryptosporidium is a coccidian parasite and implicated in causing traveler's diarrhea in immunocompromised patients. The two cryptosporidium species responsible are C. hominis and C. parvum.

The infective form is the oocyst excreted in the patient's feces, which can be acquired through drinking or during recreational water activities. The oocysts are immediately infectious (unlike isospora oocysts, which require maturation), and may cause autoinfection or be implicated in

Sold by @itachibot

person-to-person transmission. Because of their small size (4-5micron), they is difficult to see using routine staining. Hence acid-fast/Kinyoun's stain is used.

Once ingested, the 4 sporozoites are released from the cyst and attach to the small intestinal mucosa. It causes crampy abdominal pain and non-inflammatory watery diarrhea.

Treatment is mainly supportive, though nitazoxanide has been FDA approved for use in cryptosporidium infection.

Comparative features of coccidian parasites:

Property	Cryptosporidium	Cyclospora	Cystoisospora
Sporulated oocyst	4-6 µm, round, contains four sporozoites	8-12 µm, round, contains 2 sporocysts each having two sporozoites	23-36 µm, oval, contain two sporocysts, each having four sporozoites.
Acid-fast detection	Uniformly acid-fast	Variable acid-fast	Uniformly acid-fast
Sporulation of the oocyst	Occurs inside the host cell	Occurs in soil	Occurs in soil
Diagnostic form	Sporulated cyst	Unsporulated cyst	Unsporulated oocyst

Sold by @ If you purchased this from ... have been scam...

Solution to Question 14:

The below image shows trophozoites of Trichomonas vaginalis with flagella.

Parasite	Organ of locomotion
Balamuthia mandrillaris	Pseudopod
Balantidium coli	Cilia
Trichomonas vaginalis	Flagella
Toxoplasma gondi	No Particular organ

Sold by @itachibot

The below image shows the trophozoite of Balantidium coli with cilia.



The below image shows the trophozoite of Balamuthia mandrillaris with pseudopodia.

Sold by @itachibot



The below image shows tachyzoites of Toxoplasma gondi. They have no specialised structure for locomotion.



Sold by @itachibot

Helminthology - Cestodes & Trematodes

Question 1:

Which of the following cestodes requires only one host?

a) Hymenolepis nana

b) Echinococcus granulosus

c) Diphyllobothrium latum

d) Multiceps multiceps

Question 2:

A 10-year-old child is brought to your clinic with complaints of diarrhea and abdominal pain for the last 4 days. Stool microscopy is shown below. The eggs present in the stool float in a saturated solution of salt. Which of the following can you use to treat her?



a) Ivermectin

b) Mebendazole

c) Praziquantel

d) Paromomycin

Question 3:

If yo... sold by @itachibot ...this from someone else, ...scammed.

Sold by @itachibot

A patient is brought to the hospital, with complaints of seizures and the MRI image of the brain is given below. The disease is most probably caused by?



a) Cysticercosis cellulosae

b) T. saginata

c) Cysticerocosis bovis

d) T. asiatica

Sold by @ ... If you purchased this from ... you may have been scamm... else, ...d.

Question 4:

A 42-year-old man presented to the ER with complains of abdominal discomfort, indigestion, nausea and diarrhea after consuming measly pork. Stool examination reveals round eggs as shown in the image below. Which of the following statements is true regarding the causative organism?

Sold by @itachibot

a) Eggs are not infective to man

b) Hooks are absent

c) The eggs do not float in saturated salt solution

d) Intermediate host is cow

Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Question 5:

The adult form of echinococcus is seen in which of the following?

a) Dog

b) Cat

c) Swine

d) Sheep

Question 6:

Which of the following organisms cause multiple alveolar-like lesions in the liver?

a) Echinococcus multilocularis

b) Echinococcus granulosus

c) Entamoeba histolytica

d) Cysticercus cellulosae

Sold by @itachibot

Question 7:

While examining a patient with right upper quadrant pain, you note the abdomen is soft with localised tenderness present. CT abdomen findings are given below. Based on these findings, you post the patient for a PAIR procedure. Which of the following is not used in this procedure?



a) Cetrimide

b) 95% Alcohol

c) Sodium hypochlorite

d) Hypotonic saline

Question 8:

A 24-year-old woman presented to the OPD with a history of progressive breathlessness and cough for 3 months. A chest X-ray was done which showed a cavitary lesion in the lower lobe of the right lung. Wedge resection of the lesion was done and the histopathology image is given below. What is the most probable causative agent?

If you purchased elsewhere, you may have been defrauded.

Sold by @itachibot



a) Strongyloides stercoralis

b) Fasciola hepatica

c) Echinococcus granulosus

d) Paragonimus westermani

Sold by @itachi... purchased this from someone else, If yo... may have been scammed.

Question 9:

Which of the following organisms does the given life cycle belong to?



a) Hymenolepis diminuta

b) Echinococcus granulosus

Sold by @itachibot

c) Hymenolepis nana

d) Diphyllobothrium latum

Question 10:

A 35-year-old woman presented to you with complaints of tingling and numbness in her legs. On examination, her hands appear as shown in the image, ankle reflex was absent and Romberg's test was positive. Which of the following is most likely implicated in this condition?

a) Echinococcus granulosis

b) Diphyllobothrium latum

c) Hymenolepis nana

d) Taenia solium

Question 11:

Which of the following organisms most commonly causes Katayama fever?

a) Schistosoma haematobium

b) Fasciola hepatica

c) Schistosoma mansoni

d) Clonorchis sinensis

Sold by @itachibot

Question 12:

A 20-year-old male patient presents to the OPD with complaints of fever, intermittent abdominal pain, and burning micturition. He gives a history of swimming in a freshwater lake 10 days back. Urine microscopy shows the following. Which of the following is the causative agent?

a) Schistosoma hematobium

b) Schistosoma mansoni

c) Schistosoma japonicum

d) Schistosoma species cannot be determined based on egg

Question 13:

A 50-year-old fisherman presented with complaints of blood in the urine. A diagnostic cystoscopy was done and the microscopic view of the biopsied specimen is given below. Which of the following is most likely to be associated with this condition?

Sold by @itachibot



a) Schistosoma hematobium

b) Fasciolopsis buski

c) Clonorchis sinensis

d) Schistosoma mansoni

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 14:

Identify the parasite from the given life cycle.



a) Fasciola hepatica

b) Fasciolopsis buski

Sold by @itachibot

c) Pargaonimus westermani

d) Clonorchis sinensus

Question 15:

Which of the following organisms does the given life cycle belong to?



a) Diphyllobothrium latum

b) Paragonimus westermani

c) Fasciola hepatica

d) Clonorchis sinensis

Question 16:

A wet mount examination of the stool sample of a patient with abdominal pain and diarrhea is given below. Which of the following parasites is he likely infected with?

Sold by @itachibot

a) Fasciola hepatica

b) Ancyclostoma duodenale

c) Schistosoma japonicum

d) Enterobius vermicularis

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 17:

A 23-year-old patient has a history of travel to China and the consumption of pickled crab. He now complains of rusty sputum and cough. What is the most likely causative organism?

a) Fasciola buski

b) Fasciola hepatica

c) Paragonimus westermani

d) Schistosoma japonicum

Answer Key

Question No.	Correct Option
1	a
2	c
3	a
4	c

Sold by @itachibot

5	a
6	a
7	d
8	c
9	d
10	b
11	c
12	a
13	a
14	a
15	d
16	a
17	c

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Detailed Explanations

Solution to Question 1:

Hymenolepis nana requires only one host for its development. Man serves as the host for H. nana.

Usually, cestodes complete their life cycle in 2 different hosts except:

• Hymenolepis requires only 1 host, man.

• Diphyllobothrium requires 3 hosts

Parasite requiring a single host to complete its development is said to have a direct life cycle.

Parasites having a direct life cycle are:

Protozoa	Helminths
Entamoeba histolyticaGiardia lambliaTrichomonas vaginalisBalantidium coliCryptosporidium parvumCyclospora cayetanensisIsospora belliMicrosporidia.	Ascaris lumbricoidesEnterobius vermicularisTrichuris trichuraAncylostoma duodenale Necator americanusHymenolepsis nana.

Solution to Question 2:

The given clinical history and the egg seen on stool microscopy are suggestive of infection with Hymenolepis nana. A single dose of praziquantel (25 mg/kg) is the drug of choice.

Sold by @itachibot

The Hymenolepis nana eggs have a thin colorless outer membrane and inner embryophore enclosing the hexacanth oncosphere. The space between two membranes contains yolk granules and 4-8 thread-like polar filaments arising from two knobs on the embryophore. The eggs float in saturated salt solution and are non-bile stained.

Hymenolepiasis occurs more commonly in children and is usually asymptomatic. In heavy infections, they present with nausea, anorexia, abdominal pain, diarrhea, and irritability. Sometimes pruritus may occur due to an allergic response.

The image given below shows the egg of Hymenolepis nana.



Solution to Question 3:

The given clinical scenario and MRI brain showing scolex within multiple cystic lesions are suggestive of neurocysticercosis caused by the larval form of T. solium called Cysticercosis cellulosae.

Neurocysticercosis is the most common central nervous system parasitic infection.

The clinical features of neurocysticercosis include:

- Epilepsy
- Increased intracranial tension
- Hydrocephalus
- Psychiatric disturbances
- Meningoencephalitis
- Transient paresis
- Behavioral disorders
- Aphasia

Sold by @itachibot

• Visual disturbances.

The CT and MRI findings seen in neurocysticercosis are:

• Scolex within a cyst

• Cystic lesions with or without ring enhancement

• Nodular calcifications

• Focal enhancing lesions

• Multilobulated cystic lesions in the subarachnoid space.

Note: About 70% of adult-onset epilepsy is due to neurocysticercosis.

Albendazole is the drug of choice for neurocysticercosis. Glucocorticoid therapy is usually begun before initiating albendazole (antiparasitic drug). This is continued for several days after the institution of therapy to reduce the incidence of side effects from inflammatory reactions to dead and dying cysticerci of Taenia solium.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 4:

Development of the given symptoms following the consumption of measly pork is suggestive of intestinal taeniasis caused by Taenia solium (pork tapeworm). The eggs of Taenia solium do not float in the saturated salt solution.

The eggs of T. solium are infective to man and pigs. The eggs of T. saginata are infective only to cattle and not to man.

The adult worm of Taenia solium is usually 2-3 meters long and the scolex has dagger-shaped hooks.

Man is the definitive host. The intermediate host is usually pig, but occasionally it can be man.

The below image shows the life cycle of T. solium.



Sold by @itachibot

Solution to Question 5:

The adult form of echinococcus is seen in dogs and other canids which are its definitive hosts.

The larval stage is passed in intermediate hosts such as sheep, cattle, goat, swine, horse, camel, and man.

Man acts as an accidental intermediate host (dead end).

Human infection follows the ingestion of eggs due to the intimate handling of dogs or by eating raw vegetables or other food contaminated with dog feces.

Note: Humans can become accidental definitive hosts, second intermediate hosts, or paratenic hosts for spirometra. These are cestodes belonging to the Diphyllobothriidae family. They cause a condition called sparagnosis.

The image below shows the life cycle of Echinococcus granulosus.



Solution to Question 6:

Echinococcus multilocularis causes multiple alveolar-like lesions in the liver.

E. multilocularis leads to formation of a multilocular hydatid cyst, most commonly affecting the liver. It presents as multiple small cysts in the liver.

Note: Echinococcus granulosus causes a hollow fluid-filled unilocular cyst in the liver.

The CT below shows water lily sign, which is seen in the hydatid disease of the lung. It is caused by Echinococcus granulosus.

Sold by @itachibot



The water-lily sign occurs due to detachment of the endocyst membrane from the cyst wall giving the apparance of floating membranes within the pericyst. This mimics the appearance of a water lily.

It is classically visualised on plain radiographs (mainly chest X-ray) when the collapsed membranes are calcified but may be seen on ultrasound, CT, or MRI.

Sold by @itachi... If you purchased this from someone else, you may have been scammed

Solution to Question 7:

The given clinical scenario and CT findings are suggestive of hydatid disease of the liver. Hypotonic saline is not used as a scolicidal agent for the PAIR procedure (Puncture, Aspiration, Injection, and Re-aspiration) which is the treatment of choice in the early stages of this condition.

Computed tomography (CT) scan is the best imaging modality for hydatid cyst. It shows the presence of a space-occupying lesion with a smooth outline with septa.

In PAIR the steps involved are:

• USG or CT guided puncture of the cyst

• Aspiration of the cyst fluid

• Injection of scolicidal agent

• Re-aspiration of fluid after 5 minutes

The following agents can be used as scolicidal agents:

The below image depicts the PAIR procedure.

Scolicidal agent	Complications associated wit h it
Cetrimide	Acidosis
95% Alcohol	Cholangitis

Sold by @itachibot

Scolicidal agent	Complications associated wit h it
Hypertonic saline	Hypernatremia
Sodium hypochlor ite	Hypernatremia
Hydrogen peroxid e	



Solution to Question 8:

The given clinical scenario and the histological slide showing the hydatid cyst are suggestive of infection with Echinococcus granulosus.

The hydatid cyst wall consists of 3 layers:

• Pericyst- the outer dense fibrous capsule that is formed as a result of host inflammatory response.

• Ectocyst- the intermediate layer consisting of acellular, chitinous, laminated hyaline material.

• Endocyst- the inner germinal layer that is cellular and consists of a number of nuclei embedded in protoplasmic mass and is extremely thin. It is the vital layer of the cyst and it is the site of asexual reproduction giving rise to brood capsules with scolices.

The clinical features of hydatid disease of lung are breathlessness, cough, hemoptysis and chest pain. Treatment includes wedge resection or lobectomy and administration of albendazole.

The below image shows the histology of hydatid cyst.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 9:

This is the life cycle of Diphyllobothrium latum. It is also known as fish tapeworm or broad tapeworm.

The 3 hosts of Diphyllobothrium latum are:

• Definitive host- man, dog, and cat.

• First intermediate host- freshwater crustacean

• Second intermediate host- freshwater fish.

The infective form to humans is the third stage plerocercoid larvae. It possesses a rudimentary scolex.

The adult worm lives inside the small intestine and lays operculated eggs. These freshly passed eggs contain an immature embryo surrounded by yolk granules. This hexacanth embryo with 6 hooklets inside the egg is called oncosphere.

Solution to Question 10:

The given clinical scenario with cutaneous hyperpigmentation and features of subacute combined degeneration are suggestive of vitamin B12 deficiency. This can be caused by Diphyllobothrium latum.

The anemia develops because the tapeworm absorbs a large quantity of vitamin B12 and interferes with its ileal absorption leading to vitamin B12 deficiency. This can also lead to megaloblastic anemia (Bothriocephalus anemia).

In severe cases, patients may exhibit neurological features of vitamin B12 deficiency like:

• Tingling and numbness

Sold by @itachibot

• Bilateral lower limb weakness

• Bilateral hypertonia

• Sensory ataxia (Romberg's positive)

• Absent deep tendon reflexes.

It is treated with parenteral vitamin B12 supplementation.

Solution to Question 11:

Among the given options, Katayama fever is most commonly caused by Schistosoma mansoni.

It is an immune complex mediated disease caused by antibodies to the schistosomulae, adult worms, and eggs. This acute illness consists of fever, abdominal pain, diarrhea, and allergic manifestations.

The diagnosis is by the demonstration of the eggs in feces.

Praziquantel is effective in the treatment. Glucocorticoids are given to suppress hypersensitivity reactions.

Instances of Katayama fever are rarer in infections with Schistosoma haematobium compared to S. mansoni.

*Note: Katayama fever is most commonly caused by S. japonicum and S. mansoni. But, among the given options, S. mansoni is the most common cause of Katayama fever.

Solution to Question 12:

The given clinical picture and image showing an egg that is non-operculated with the terminal spine at one pole are suggestive of infection with Schistosoma hematobium.

Species differentiation of schistosoma can be done based on the location of the spine as follows:

• Terminal spine- Schistosoma hematobium

• Lateral spine- Schistosoma mansoni

• Rudimentary lateral spine- Schistosoma japonicum

The image below shows egg of Schistosoma mansoni. Adult worms of S. mansoni live in the inferior mesenteric vein.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Adult worms of S. japonicum live in the superior mesenteric vein draining the ileocecal region. They are also seen in the intrahepatic portal venules and hemorrhoidal plexus of veins.

The image below shows an egg of Schistosoma japonicum.



Adult worms of S. hematobium live in the vesical and pelvic plexus of veins.

The below image shows the life cycle of S.hematobium:

Sold by @itachibot



Sold by @itachi... if you purchased this from some... you may have been scammed. ...else,

Solution to Question 13:

The given clinical scenario and the image showing keratin pearls are suggestive of squamous cell carcinoma of the urinary bladder. It is associated with Schistosoma hematobium infection.

Chronic schistosomiasis presents as painless hematuria, dysuria, and proteinuria. Fibrosis may cause obstructive uropathies like hydronephrosis and hydroureter. It has been associated with squamous cell carcinoma of the bladder. Such malignancy is detected in a younger age group as compared to transitional cell carcinoma of the bladder.

The following parasites cause malignancies:

• Schistosoma hematobium- bladder carcinoma

• Clonorchis sinensis and Opisthorchis viverrini- bile duct carcinoma

Solution to Question 14:

The given life cycle belongs to Fasciola hepatica.

Fasciola hepatica (sheep liver fluke) causes disease primarily in sheep and other domestic animals.

Humans are infected by eating watercress (or other aquatic plants) contaminated by larvae (metacercariae) that excyst in the duodenum, penetrate the gut wall, and reach the liver, where they mature into adults.

The adult flukes in the bile ducts produce eggs, which are excreted in the feces. The eggs hatch in freshwater and miracidia enter the snails. Miracidia develop into cercariae, which then encyst on aquatic vegetation. Sheep and humans eat the plants, thus completing the life cycle.

Sold by @itachibot

Most infections are asymptomatic. Halzoun is a painful pharyngitis caused by the presence of adult flukes on the posterior pharyngeal wall after eating raw sheep liver. Obstructive jaundice can occur months or years later.

Diagnosis is made by identification of the eggs in the feces.

Solution to Question 15:

The given life cycle belongs to Clonorchis sinensis. It causes clonorchiasis (Asian liver fluke infection).

The hosts for Clonorchis sinensis are:

• Definitive host- humans

• First intermediate host- snail

• Second intermediate host- fish

Humans are infected by eating raw or undercooked fish containing the encysted larvae (metacercariae). In the duodenum, metacercaria will undergo excystation and migrate to bile ducts to develop into adult worms. The eggs containing miracidia are excreted in human feces which are then ingested by snails. Miracidium hatches out in the midgut of the snail. Inside the snail, sporocyst develops into 1st and 2nd generation redia and then to cercariae. The cercariae now penetrate the fish and develop into metacercaria (infective form).

Chronic clonorchiasis is strongly associated with cholangiocarcinoma.

If you purchased this from someone else, you may have been scammed.
Sold by @itachibot

Solution to Question 16:

The given image showing an operculated egg (egg having a lid-like structure on its end) in the stool sample is suggestive of Fasciola hepatica infection.

The eggs of Fasciola hepatica are large, ovoid, operculated, and bile stained. They contain an immature larva (miracidium) and do not float in the saturated salt solution.

The following are the parasites with operculated eggs:

• Fasciola hepatica

• Fasciolopsis buski

• Fasciola gigantica

• Paragonimus westermani

• Opisthorchis felineus

• Clonorchis sinensis

• Gastrodiscoides hominis

• Diphyllobothrium latum

Sold by @itachibot

Eggs of Schistosoma japonicum, Enterobius vermicularis, and Ancylostoma duodenale are non-operculated.

Solution to Question 17:

The given clinical scenario is suggestive of infection with Paragonimus westermani.

Paragonimus westermani is a trematode. The transmission to humans occurs by eating raw or undercooked crab or crayfish containing metacercariae. The metacercariae excyst in the human gut and young worms migrate to the lungs, where they become encapsulated in lung tissue.

The patient complains of cough, chest pain, and hemoptysis. The viscous sputum is speckled with golden-brown eggs.

Praziquantel (25 mg/ kg TDS for 1-2 days) is the drug of choice.

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Helminthology - Nematodes

Question 1:

Which of the following parasites is ovoviviparous?

a) Enterobius vermicularis

b) Trichuris trichiura

c) Strongyloides stercoralis

d) Trichinella spiralis

If yo @itachibot rom someone else, mmed.

Question 2:

Identify the helminth from the life cycle given below.



a) Ancylostoma duodenale

b) Strongyloides stercoralis

c) Ascaris lumbricoides

d) Enterobius vermicularis

Question 3:

Sold by @itachibot

A 24-year-old patient presented with complaints of diarrhea, vomiting, and abdominal pain for 1 week. The stool microscopy revealed the following. What is the appropriate treatment for this condition?

a) Ivermectin

b) Tinidazole

c) Albendazole

d) Praziquantel

Sold by
If you purchased this fro
you may have been scam
else,
d.

Question 4:

An 8-year-old child was brought with complaints of perianal itching which is worse in the night. The scotch tape method demonstrates eggs as shown in the image given below. What is the most likely causative agent?

Sold by @itachibot



a) Hookworm

b) Whipworm

c) Pinworm

d) Roundworm

Question 5:

Which of the following helminthic infections that is commonly transmitted by soil, is characterized by■ the egg in stool as shown in the image below?



a) Trichuris trichiura

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

b) Ancyclostoma duodenale

c) Necator americanus

d) Ascaris lumbricoides

Question 6:

A college student came to the OPD with fever and muscle aches. He tells you he had consumed smoked pork at a party 2 weeks back. Following this, he had developed abdominal cramps and diarrhea which had resolved on their own. CBC revealed eosinophilia and the muscle biopsy showed the following finding. What is the causative organism?

a) Trichuris trichiura

b) Trichinella spiralis

c) Ascaris lumbricoides

d) Enterobius vermicularis

Question 7:

In a patient with a positive Bachman intradermal test, which of the following infections will you suspect?

a) Schistosomiasis

b) Kala-azar

c) Hydatid disease

d) Trichinellosis

Sold by @itachibot

Question 8:

Identify the organism shown in this image.



a) Strongyloides stercoralis

b) Enterobius vermicularis

c) Ancylostoma duodenale

d) Ascaris lumbricoides

Question 9:

A 9- year-old boy was brought to the casualty with fever and colicky abdominal pain. An erect X-ray abdomen showed multiple air-fluid levels. Stool examination revealed the following egg. Which of the following is false about the organism causing this condition?

If you purchased this ... you may have been ... else, ... d.

Sold by @itachibot

a) Human is the only host

b) Can cause pancreatitis

c) Can cause Loeffler's syndrome

d) Egg containing L1 larva is infective

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 10:

What is the most common cause for visceral larva migrans?

a) Toxocara canis

b) Ancylostoma braziliense

c) Ascaris lumbricoides

d) Strongyloides stercoralis

Question 11:

A middle-aged woman comes to the OPD with complaints of weakness for the past 3 months. She is diagnosed to have microcytic hypochromic anemia. The stool microscopy reveals the following. What is the most probable causative agent?

Sold by @itachibot



a) Ancylostoma duodenale

b) Strongyloides stercoralis

c) Enterobius vermicularis

d) Ascaris lumbricoides

Sold by @itachi... else, if you purchased this from some... you may have been scammed.

Question 12:

A patient presented to the OPD with dry cough, low-grade fever and wheezing. He is diagnosed with Loeffler's syndrome. Which of the following parasites does not cause this?

a) Ascaris lumbricoides

b) Necator americanus

c) Strongyloides stercoralis

d) Enterobius vermicularis

Question 13:

A 27-year-old woman had gone on a trek in the Amazon rainforest. A month later she presented to the OPD with the following lesion which was painful and pruritic. Her systemic examination was normal. Which of the following organisms is most likely to cause this condition?

Sold by @itachibot

a) Strongyloides stercoralis

b) Ancylostoma braziliense

c) Toxocara canis

d) Necator americanus

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 14:

Which the following filarial worms are non-pathogenic to humans?

a) Onchocerca volvulus

b) Brugia malayi

c) Wuchereria bancrofti

d) Mansonella ozzardi

Question 15:

A 55-year-old man came to the hospital with complaints of progressive swelling of his legs. On examination, you note bilateral non-pitting edema of the lower limbs with thickening of the overlying skin. What is the best next step in the management of this patient?

Sold by @itachibot

a) PCR

b) Lymphoscintigraphy

c) Night blood test for worms

d) Lymphangiography

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 16:

A child from Bihar is brought to the AIIMS ER with a high-grade fever. On examination, bilaterally enlarged, tender inguinal lymph nodes are present. The peripheral smear examination finding is shown in the image. What is the most likely causative organism?

a) Wuchereria bancrofti

Sold by @itachibot

b) Brugia malayi

c) Loa loa

d) Onchocerca volvulus

Question 17:

A patient from Kerala has presented to your OPD with fever. A magnified view of peripheral blood smear is shown below. Which of the following is false regarding the organism seen?

a) DEC provocation test is used to demonstrate microfilaria during the day

b) Second stage filariform larva is the infective form

c) Microfilaria is the major diagnostic form

d) Filarial stage of adult worms are responsible for disease

Question 18:

A 54-year-old farmer from Assam presented with the finding shown in the image. He had also developed a painless swelling in his elbow 2 years back which has gradually been increasing in size. Which of the following is the drug of choice to treat this condition?

Sold by @itachibot

a) Eflornithine

b) Suramin

c) Ivermectin

d) Diethylcarbamazine

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 19:

A middle-aged woman presents to your clinic with photosensitivity, redness, and watering of the eyes for the last 2 days. She shows you this photo of a swelling in her hand that had arisen 2 weeks back but disappeared in just 3 days. What is the most likely diagnosis?

a) Loiasis

Sold by @itachibot

b) River blindness

c) Tropical pulmonary eosinophilia

d) Occult filariasis

Question 20:

Which of the following is also known as Weingarten syndrome?

a) Tropical pulmonary eosinophilia

b) Loasis

c) Onchocerciasis

d) Tropical basophilia

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 21:

Which of the following is known as serpent worm?

a) Gnathostoma spinigerum

b) Angiostrongylus cantonensis

c) Dracunculus medinensis

d) Capillaria philippinensis

Question 22:

Match the following parasites with their respective hosts:

a) A-3, B-2, C-4, D-1

b) A-2, B-1 ,C-4 ,D-3

c) A-1, B-2, C-3, D-4

d) A-4, B-3, C-1, D-2

Answer Key

Question No.	Correct Option
1	c

Sold by @itachibot

2	b
3	a
4	c
5	a
6	b
7	d
8	d
9	d
10	a
11	a
12	d
13	b
14	d
15	c
16	b
17	b
18	c
19	a
20	a
21	c
22	a

If you pu... you may ... else, ...d.

Detailed Explanations

Solution to Question 1:

Strongyloides stercoralis is an ovoviviparous parasite.

Nematodes are classified as follows:

• Oviparous - lay eggs:

• Unsegmented eggs: Ascaris lumbricoides, Trichuris trichuria

• Segmented eggs: Ancylostoma duodenale, Necator americanus

• Eggs containing larva: Enterobius vermicularis

• Viviparous - produce larvae. e.g., Trichinella spiralis, Wuchereria bancrofti, Brugia malayi

• Ovoviviparous – lay eggs containing fully formed larvae that hatch immediately. e.g., Strongyloides stercoralis

Sold by @itachibot

Solution to Question 2:

The image shows the life cycle of Strongyloides stercoralis.

The infective form of strongyloides is the filariform larva. Infection occurs through penetration of the skin by larvae in the soil. Then, they migrate to the small intestine.

The larvae molt twice and become adult female worms in the small intestine. The eggs laid in the mucosa of the intestine hatch immediately releasing the rhabditiform larva (L1 stage). These rhabditiform larvae in the bowel can develop directly into filariform larvae that penetrate the colonic wall or perianal skin and enter the circulation to repeat the migration. This establishes ongoing internal reinfection.

The rhabditiform larva can be seen on microscopic examination of stool.

Solution to Question 3:

The given clinical scenario and the stool microscopy showing the rhabditiform larva of S. stercoralis are suggestive of strongyloidiasis. The most appropriate treatment for this condition would be ivermectin.

Strongyloidiasis is caused by Strongyloides stercoralis. It is generally a benign and asymptomatic condition. In some cases, there may be cutaneous, pulmonary, and intestinal manifestations.

Solution to Question 4:

The given clinical scenario of perianal itching and the scotch tape microscopy showing the eggs are suggestive of infestation with pinworm (Enterobius vermicularis).

The eggs of pinworm are plano-convex in shape, colorless, non-bile stained, and float in the saturated salt solution. Eggshell is double layered and relatively thick, though transparent.

Since pinworms are not released in feces, the diagnosis cannot be made by conventional fecal ova and parasite tests. NIH swab method and the scotch tape method are used for the collection of specimens.

The NIH swab (National Institutes of Health, USA) consists of a glass rod, at one end of which a piece of transparent cellophane is attached with a rubber band. The cellophane part is used for swabbing by rolling over the perianal area. It is returned to the test tube and sent to the laboratory. The cellophane piece is detached, spread over a glass side, and examined microscopically.

The image given below shows the NIH swab.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



In scotch tape method, the tape is firmly pressed against the anal margin, covering all sides. The tape is transferred to a glass slide with sticky side down. Then a drop of toluene is added for clearing and examined under the microscope.

Scotch tape method



The below image shows the life cycle of Enterobius vermicularis.

Sold by @itachibot



Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 5:

The given image shows the egg of Trichuris trichiura (whipworm).

The characteristic features of whipworm eggs are:

• Eggs are barrel-shaped in the middle with a projecting mucus plug at each pole containing an unsegmented ovum. The plugs are colorless.

• They float in the saturated salt solution.

• It has a triple shell, the outermost layer of which is stained brown.

They are commonly transmitted by soil. The eggs passed in the stool are unembryonated. Microscopic identification of the eggs in a stool specimen is diagnostic of whipworm infection.

In moderate infections of 20 adult worms or less, the worms are confined to the caecum and ascending colon. The symptoms include epigastric and lower abdominal pain, diarrhoea (rarely bloody), vomiting, flatulence and distention, weight loss, and increased blood and iron loss.

In heavy infections of 200 adult worms or more (trichuriasis dysentery syndrome), the worms are in the distal colon and rectum. The symptoms include bloody diarrhoea with profuse mucus, abdominal pain, tenesmus, weight loss leading to cachexia, severe anemia, and rectal prolapse. On examination, clubbing may be present. Blood investigations may show moderate eosinophilia.

Chronic infection leads to growth retardation and nutritional deficiencies. Around 0.005 ml of blood is lost per day per worm. In heavy infections, as much as 4 to 5 ml can be lost per day per worm.

Treatment is mebendazole or albendazole.

Solution to Question 6:

Sold by @itachibot

The given clinical scenario and muscle biopsy showing larvae are suggestive of trichinellosis. It is caused by Trichinella spiralis.

Trichinella spiralis is known as the pork worm since the infection is acquired by consuming raw or undercooked pork. The initial infection presents with diarrhea and abdominal pain. In the later phase of the infection, the larvae move from the intestines into the circulation and migrate into muscles where they are encysted. This causes symptoms like fever and myalgia.

These larvae are calcified in about 5-6 months. Eosinophilia caused by parasitic infection also comes down after 6 months.

Detection of spiral larva in muscle tissue is done by performing a muscle biopsy. Deltoid, biceps, gastrocnemius, or pectoralis muscles are usually selected for biopsy.

Other parasites that can be demonstrated in a muscle biopsy are:

- T. solium
- T. cruzi (amastigote)

The image below shows Trichinella spiralis larvae within the muscle.



If yo...tachibot...someone else,...d.

Solution to Question 7:

Bachman intradermal test is used for the diagnosis of trichinellosis.

In this test, 1:5,000 or 1:10,000 dilution of the trichinella larval antigen is injected intradermally. An erythematous wheal appears in positive cases within 15-20 minutes. The test remains positive for years after infection.

Solution to Question 8:

Sold by @itachibot

The given image shows the Ascaris lumbricoides (roundworm). It is the most common of human helminths and is distributed worldwide.

Adult worms live in the small intestine (85% in the jejunum and 15% in the ileum). The roundworm, Ascaris lumbricoides is the largest nematode parasite in the human intestine.

The adult worms are large cylindrical worms, with tapering ends, the anterior end being more pointed than the posterior.

Man is the natural host. There is no intermediate host.

Infection occurs through food or water contaminated with nightsoil or human feces containing eggs of ascaris. The eggs contain the infective rhabditiform larva.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Solution to Question 9:

The given clinical scenario and the image are suggestive of ascariasis caused by Ascaris lumbricoides. Embryonated egg containing the L3 larva is the infective form.

Man is the only natural host. There is no intermediate host. The infection is acquired via the consumption of food or water contaminated with eggs. The eggs hatch after reaching the duodenum to release the larvae. These penetrate the intestinal mucosa and enter the portal vessels. From the liver, larvae reach the systemic circulation, lungs and then reach the small intestine to produce symptoms.

Clinical features:

Due to the migrating larvae:

- Fever
- Dry cough
- Asthmatic wheezing
- Loeffler's syndrome

Due to the adult worm:

- Asymptomatic
- Nutritional effects - protein-energy malnutrition, Vitamin A deficiency
- Toxic effects - urticaria, angioedema due to hypersensitivity
- Mechanical effects - colicky abdominal pain, volvulus, and intestinal obstruction
- Ectopic ascariasis - acute biliary obstruction, pancreatitis, and obstructive appendicitis.

The life cycle of Ascaris lumbricoides is given below.

Sold by @itachibot



Sold by @itachi... If you purchased this from someone else, you may have been scammed.

Solution to Question 10:

The most common cause for visceral larva migrans is Toxocara canis. It occurs most commonly in children and presents as fever, hepatosplenomegaly, and pneumonitis.

Larva migrans is generally seen when human infection occurs with nonhuman species of nematodes. It can be classified into cutaneous or visceral types, depending on whether the larval migration takes place in the skin or in deeper tissues.

Visceral larva migrans is caused by the migration of nematode larvae that are acquired via the oral route. Cutaneous larva migrans is caused by the migration of nematode larvae (most commonly Ancylostoma braziliense) that are acquired via skin penetration.

Patients having visceral larva migrans may develop neurological disturbances (neural larva migrans) and endophthalmitis (ophthalmic larva migrans).

Differences between cutaneous larva migrans and visceral larva migrans:

	Cutaneous larva migrans	Visceral larva migrans
Tissue involved	Skin	Various organs of the body li ke the liver, lungs, and eyes
Infecting Organis m	Mostly by non-human nemat odes	Mostly by non-human nemat odes (mainly by dog and cat)
Portal of Entry	Penetration of skin	Ingestion of infected eggs
Eosinophilia	Mild	Persistent high
Serodiagnosis	Not developed	Well developed
Treatment	Thiabendazole	Diethylcarbamazine and pred nisolone

Sold by @itachibot

Solution to Question 11:

The given clinical scenario of iron deficiency anemia and the stool microscopy showing the egg is suggestive of infection by Ancylostoma duodenale.

Eggs of both Ancylostoma duodenale (old world hookworm) and Necator americanus (new world hookworm) are identical.

The hookworm eggs are oval or elliptical in shape, colorless, and non-bile stained. It has a thin transparent hyaline shell membrane. The ovum is segmented with 4 to 8 blastomeres. There is a clear space between the segmented ovum and eggshell. Eggs float in the saturated salt solution.

Humans are infected when filariform larvae in moist soil penetrate the skin or subcutaneous tissue. The larvae are then carried in the venous circulation to the heart and lungs, where they may cause mild transient pneumonitis or bronchitis. They migrate up the respiratory tract to the pharynx, are swallowed, and reach the small intestine producing epigastric pain, dyspepsia, nausea, vomiting, and diarrhea.

A. duodenale or N. americanus ingest blood, secrete anticoagulants, and interfere with the absorption of iron, B12, and folic acid. This leads to iron deficiency anemia and protein-energy malnutrition resulting from blood loss.

The average blood loss of the host per worm per day is 0.2 mL with Ancylostoma duodenale and 0.03 mL with Necator americanus.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed

Solution to Question 12:

Loeffler's syndrome is an eosinophilic inflammation of the lung caused by Ascaris lumbricoides, Strongyloides stercoralis and Necator americanus. Enterobius vermicularis does not cause Loeffler's syndrome.

It is associated with pneumonia characterized by dry cough, low-grade fever, wheezing, urticaria, and eosinophilia.

Parasites causing Loeffler's syndrome are:

- Migrating larva of:
- Ascaris lumbricoides
- Strongyloides stercoralis
- Ancylostoma duodenale
- Necator americanus
- Echinococcus granulosus
- Cryptosporidium parvum
- Eggs of Paragonimus westermani
- Entamoeba histolytica
- Trichomonas tenax

Sold by @itachibot

Solution to Question 13:

The given clinical scenario and the image showing serpiginous skin lesions are suggestive of cutaneous larva migrans or creeping eruption or ground itch. This is most commonly caused by Ancylostoma braziliense.

Cutaneous larva migrans is caused by non-human hookworms like Ancylostoma braziliense and Ancylostoma caninum. It is acquired via direct skin contact with soil contaminated by dog or cat feces. The larvae remain confined to the outer layers of skin, as they are unable to penetrate the basement membrane of the human skin.

It presents as erythematous, pruritic papules at the site of entry and serpiginous tracks in the epidermis. The symptoms begin to appear within the first two weeks of the patient's return from a tropical or endemic country. The larval track progresses at approximately 1 cm/h with cutaneous lesions only progressing a few millimeters to a few centimeters daily.

Diagnosis is based mainly on clinical features.

The treatment includes:

• Thiabendazole topical application

• Albendazole 400mg/day X 3 days or

• Ivermectin 200mcg/kg single dose stat

The differences between cutaneous larva migrans and visceral larva migrans are:

Note: A rapidly moving lesion (10cm/hr) produced by Strongyloides stercoralis is known as larva currens (running larva). It occurs as an otherwise asymptomatic lesion in immunocompetent individuals. In immunocompromised, hyper-infestation syndrome occurs with paralytic ileus and secondary Gram-negative sepsis.

	Cutaneous larva migrans	Visceral larva migrans
Tissue involved	Skin	Various organs of the body li ke the liver, lungs, and eyes
Infecting Organis m	Mostly by non-human nemat odes	Mostly by non-human nemat odes (mainly by dog and cat)
Portal of Entry	Penetration of skin	Ingestion of infected eggs
Eosinophilia	Mild	Persistent high
Serodiagnosis	Not developed	Well developed
Treatment	Thiabendazole	Diethylcarbamazine and pred nisolone

Solution to Question 14:

Mansonella ozzardi is generally non-pathogenic to humans.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot

Option A: Onchocerca volvulus infection may manifest as skin nodules or onchocercoma which are formed as a result of fibroblastic reaction around adult worms deep in the subcutaneous tissues. Microfilariae cause lesions in the skin and eyes.

Options B and C: In lymphatic filariasis (Wuchereria and Brugia), the typical manifestations are caused by the adult worms blocking lymph nodes and vessels. Microfilaria (larval form) are responsible for allergic manifestations seen in occult filariasis.

Solution to Question 15:

The given clinical picture is suggestive of lymphatic filariasis (elephantiasis). The best next step would be to test the night blood for worms since the filarial species is nocturnally periodic (between 10 pm and 4 am) in India.

Lymphatic filariasis is caused by Wuchereria bancrofti and Brugia malayi.

Elephantiasis is a delayed sequel to repeated lymphangitis, obstruction, and lymphedema. Repeated leakage of lymph into tissues first results in lymphedema, then to elephantiasis.

Diethylcarbamazine is the drug of choice.

Option A: PCR is negative in chronic filariasis.

Options B & D: Lymphangiography and lymphoscintigraphy may be used to demonstrate abnormal lymphatic urinary fistula leading to chyluria in chronic filariasis.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 16:

The given clinical scenario and the peripheral smear showing sheathed microfilaria with two nuclei at the tail tip is suggestive of infection with Brugia malayi.

Acute infection may manifest as acute adenolymphangitis which is characterized by high fever, lymphatic inflammation (lymphangitis and lymphadenitis) and transient local edema. The most commonly affected lymph nodes are inguinal nodes followed by axillary nodes which become enlarged, painful and tender.

The following are the characteristic features of microfilaria of Brugia malayi:

• Smaller in size compared to microfilaria of W. bancrofti.

• Shows kinks and secondary curves.

• Cephalic space is longer.

• Carries double stylets at the anterior end.

• Nuclear column appears blurred in Giemsa-stained films.

• Tail tip carries 2 distinct nuclei, one terminal and the other subterminal.

The below image shows the microfilaria of Brugia malayi.

Sold by @itachibot



Sold by @itachi... If you purchased this from some... else, you may have been scammed.

Solution to Question 17:

The given clinical scenario with a peripheral smear showing sheathed microfilaria with a pointed tail free of nuclei is suggestive of Wuchereria bancrofti. The third stage filariform larva is the infective form.

Wuchereria bancrofti infection is usually asymptomatic. The symptoms are due to obstruction of lymphatics by the filarial stage of adult worms.

The lymphatic vessels of the male genitalia are most commonly affected in bancroftian filariasis, producing episodic funiculitis, epididymitis, and orchitis. Adenolymphangitis of the extremities is less common.

Hydrocele is the most common sign of chronic bancroftian filariasis, followed by lymphedema, elephantiasis, and chyluria. The swelling involves the whole leg, the whole arm, the scrotum, the vulva, or the breast.

DEC provocation test is used to demonstrate microfilaria during the day.

The image below shows the life cycle of Wuchereria bancrofti.

Sold by @itachibot



The image below shows the difference between microfilaria of various filarial nematodes.

Solution to Question 18:

The clinical scenario and image showing sclerosing keratitis are suggestive of onchocerchiasis (river blindness). The drug of choice is ivermectin.

It is caused by Onchocerca volvulus. It is also known as convoluted filaria or blinding filaria.

The vector is blackfly (Simulium species) which breed in fast-flowing rivers, and therefore the disease is most common along the course of rivers. Hence, it is also called river blindness.

The usual presentation is formation of a subcutaneous nodule or onchocercoma. It is a firm non-tender nodule arising from bony sites like scalp, elbow, ribs, and iliac crest. It is associated with pruritus and rashes.

Sold by @itachibot

The microfilaria in the eyes cause photophobia and gradual loss of vision. The most severe manifestation is sclerosing keratitis and chorioretinitis which causes blindness. It is the 2nd major infective cause of blindness in the world after trachoma.

Suramin and diethylcarbamazine (DEC) were used earlier but are not used nowadays. DEC destroys microfilariae but usually causes Mazzotti reaction consisting of pruritus, rash, lymphadenopathy, fever, hypotension, and occasionally, eye damage.

Note: Onchocerciasis is not endemic in India but sporadic cases have been reported from North-Eastern India.

Solution to Question 19:

The given clinical scenario and the image showing Calabar swelling is suggestive of loiasis caused by Loa loa.

Calabar swellings are temporary foci of inflammation that appear as Loa loa wanders through the subcutaneous tissues. They are up to 3 cm in size and are also called fugitive swellings because they disappear in a few days only to reappear elsewhere.

Loa loa is transmitted by the bite of Chrysops species (deerflies). The infective form is the third stage microfilaria. After entry into human skin, it grows into an adult worm.

The ocular manifestations occur when the worm reaches the subconjunctival tissue during its wanderings. They include:

• Granuloma in bulbar conjunctiva

• Painless edema of the eyelids

• Proptosis.

The image below shows Calabar swelling.

The image below shows a live, adult Loa loa in the anterior chamber of the eye.

If you purchased this from someone else, it may have been scammed. Sold by @itachibot

Sold by @itachibot



If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 20:

Tropical pulmonary eosinophilia is also known as Weingarten syndrome.

Tropical eosinophilia occurs as a result of hypersensitivity reaction to microfilarial antigens and not directly due to lymphatic involvement. It is a manifestation of occult filariasis (Meyer's-Kouwenaar syndrome).

Clinical features of occult filariasis include massive eosinophilia and hepatosplenomegaly.

Synonyms of tropical pulmonary eosinophilia include -

- Weingarten syndrome
- Pseudo-tuberculosis of the lung
- Eosinophilic pseudo-leukemia
- Tropical eosinophilic asthma
- Frimodt-Moller and Barton's syndrome

Solution to Question 21:

Dracunculus medinensis is known as serpent worm.

It is a viviparous nematode.

This has been eradicated from India and all of the southeast Asian region in the year 2000.

Solution to Question 22:

Sold by @itachibot

Parasite	Definitive host	First intermediate host	Second intermediate host
Diphyllobothrium	Humans	Cyclops	Cyclops and fish
Paragonimus	Humans	Snail	Crayfish and crab
Fasciola	Sheep and humans	Snail	Aquatic vegetations
Brugia	Humans	Mansonia mosquito	-

Sold by @Itachibot
If you purchased this from someone else,
you may have been scammed.

Sold by @itachibot

Applied Microbiology

Question 1:

While assessing the quality of drinking water in a village, you detect the presence of coliform bacilli by the multiple tube technique. Following this, you decide to perform the Eijkman test. What is this test also known as?

a) Presumptive coliform count

b) Differential coliform test

c) Colony plate count

d) Methylene blue reduction test

Question 2:

Faecal contamination of a water source is suspected by the mayor of a municipal corporation. He asks you to test the water for evidence of recent contamination. The presence of which of the following organisms would suggest this?

a) Shigella

b) Enterococcus

c) Clostridium

d) Staphylococcus

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Question 3:

Which of the following tests is not used for bacteriological examination of milk?

a) Methylene blue reduction test

b) Phosphatase test

c) Resazurin test

d) Pasteurisation

Question 4:

The following image depicts:

Sold by @itachibot



a) Production of monoclonal antibodies

b) Cell lines for viral culture

c) Vaccine synthesis

d) Process of genetic engineering

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Question 5:

Which of the following statements about hybridoma is false?

a) Specific antibody-producing cells are integrated with myeloma cells.

b) Myeloma cells using salvage pathway grow well in HAT medium.

c) Aminopterin, a folate antagonist, inhibits de novo pathway.

d) HGPRTase and thymidylate synthetase are required for salvage pathway.

Question 6:

Which of the following is both a diffusion and a dilution method of testing antibiotic sensitivity?

a) Kirby–Bauer method

b) Stokes method

c) Epsilometer method

d) Tube dilution method

Sold by @itachibot

Question 7:

In which of the following is the disc diffusion method for antibiotic susceptibility testing applied?

a) Agar dilution method

b) Kirby-Bauer method

c) Broth dilution method

d) PCR based assay

Question 8:

Which of the following methods of antibiotic susceptibility testing cannot be used to calculate the minimum inhibitory concentration of an antimicrobial drug?

a) Kirby–Bauer method

b) Tube dilution method

c) Agar dilution method

d) Epsilometer method

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Question 9:

A microbiologist is about to run a polymerase chain reaction assay on the serum of a patient suspected to have Q fever. He wishes to quantify the PCR by monitoring the amplification process while the PCR is ongoing. Which of the following types of PCR should he perform?

a) Multiplex PCR

b) Nested PCR

c) Reverse transcriptase PCR

d) Real time PCR

Question 10:

Which of the following investigations does not use antibody marking?

a) ELISA

b) Radioimmunoassay

c) Immunofluorescence

Sold by @itachibot

d) Hemagglutination Test

Question 11:

Which of the following is not used as an immediate test in the screening of urinary tract infections?

a) Pour plate culture

b) Griess Nitrite test

c) Leukocyte esterase test

d) Wet mount examination

Question 12:

The CLED medium is preferred over MacConkey agar for the culture of organisms in case of UTI because:

a) It inhibits proteus swarming

b) It promotes the growth of candida, streptococcus and staphylococcus

c) It differentiates between lactose-fermenting and non-lactose fermenting bacteria

d) It promotes the growth of pseudomonas

Question 13:

A patient who had been receiving ceftazidime and amikacin via central venous line for a week suddenly spiked a fever. Blood culture is positive for a vancomycin-resistant, catalase-negative organism whose Gram-stained image is given below. Identify the organism.

Sold by @itachibot
If you purchased this from someone else, you may have been scammed.

Sold by @itachibot



a) Staphylococcus aureus

b) Viridans streptococci

c) Enterococcus faecalis

d) Coagulase-negative staphylococcus

Sold by @itachi... else, If you purchased this from some... you may have been scammed.

Question 14:

Which of the following is a bioterrorism category A agent?

a) Nipah virus

b) Anthrax

c) Coxiella

d) Brucella

Question 15:

The test kit shown in the image below works on the principle of:

Sold by @itachibot

©MARROW

a) ELISA

b) Immunochromatography

c) Immunofluroscence

d) Chemiluminiscence

Sold by @itachi... purchased this from some... else, If you have been scammed.

Answer Key

Question No.	Correct Option
1	b
2	b
3	d
4	a
5	b
6	c
7	b
8	a
9	d
10	d
11	a
12	b
13	c
14	b

Sold by @itachibot

15	b

Detailed Explanations

Solution to Question 1:

Eijkman test is also known as differential coliform test.

Drinking water should be free of any pathogenic microorganisms. The presence of coliform bacteria in the water may indicate faecal pollution, while the detection of E.coli confirms it.

The presumptive coliform test is done by the multiple tube technique. Following this, the Eijkman test is done to confirm the presence of E. coli. They can grow at 44°C (faecal thermotolerant bacilli), ferment lactose and cause indole and gas production.

Option A: Presumptive coliform count detects the probable number of coliform bacilli in water. It can be positive due to the presence of organisms other than E.coli also.

Option C: Colony counts provide an estimate of the general bacterial purity of water.

Option D: Methylene blue reduction test is used to assess the bacterial contamination of milk.

Solution to Question 2:

Recent fecal contamination of water is suggested by the presence of enterococcus (fecal streptococci).

Clostridium perfringens may be present for a long duration. If it is present in water, it suggests that contamination is not recent.

If Escherichia coli is detected, it is definitive proof of fecal contamination. However, it is non-specific for suggesting the duration of the contamination.

Solution to Question 3:

Pasteurization is not used for the bacteriological examination of milk.

Pasteurization is the process of applying heat, to milk, for a specified period for the purpose of killing pathogenic bacteria. There are two methods: holder method and flash method.

Option A: Methylene blue reduction test depends on the reduction of methylene blue by bacteria in milk when incubated at 37 °C in complete darkness.

Option B: Resazurin test is similar to methylene blue reduction test, but the dye resazurin on reduction passes through a series of colour changes from blue to pink to colourless. The shade of colour after incubation with milk for a particular period of time depends on the degree of contamination.

Sold by @itachibot
If you purchased this from someone else
you may have been scammed.

Sold by @itachibot

Option C: Phosphatase test is used to check whether milk has been pasteurised. Positive test after pasteurisation of milk shows contamination after pasteurisation.

Solution to Question 4:

The image depicts the production of monoclonal antibodies by hybridoma technology.

The following are the steps involved in hybridoma technology:

• Mice are immunized with an antigen. After several weeks of immunization, blood samples are obtained from mice for measurement of serum antibodies.

• When the antibody titer is high enough, mice are commonly boosted by injecting antigen without adjuvant intraperitoneally or intravenously. Then the mice are euthanized and their spleens removed for in vitro hybridoma cell production.

• Fusing antibody-producing spleen cells, which have a limited life span with myeloma cells (derived from an immortal tumor of lymphocytes) results in a hybridoma that is capable of unlimited growth.

• The cells are cultured in a medium containing hypoxanthine, aminopterin, and thymidine (HAT medium) which allows only the fused cells to survive in culture.

• Monoclonal antibodies are then harvested.



Solution to Question 5:

Aminopterin is a folate antagonist present in the HAT medium that inhibits the de novo pathway of nucleotide synthesis. Since myeloma cells cannot use the salvage pathway for nucleotide synthesis due to the lack of HGPRTase enzyme, they cannot survive in the HAT medium.

Sold by @itachibot

Hybridoma technology is a method for producing large numbers of identical antibodies (monoclonal antibodies).

Hybridomas are somatic cell hybrids produced by fusing the antibody-forming spleen cells with myeloma cells. The resultant hybrid thus has both the antibody-producing ability of the B-cell and the ability to multiply indefinitely of the myeloma.

The following are the steps involved in hybridoma technology:

• Mice are immunized with an antigen. After several weeks of immunization, blood samples are obtained from mice for measurement of serum antibodies.

• When the antibody titer is high enough, mice are commonly boosted by injecting antigen without adjuvant intraperitoneally or intravenously. Then the mice are euthanized and their spleens removed for in vitro hybridoma cell production.

• Fusing antibody-producing spleen cells, which have a limited life span with myeloma cells (derived from an immortal tumor of lymphocytes) results in a hybridoma that is capable of unlimited growth.

• The cells are cultured in a medium containing hypoxanthine, aminopterin, and thymidine (HAT medium) which allows only the fused cells to survive in culture.

• Monoclonal antibodies are then harvested.



Solution to Question 6:

Epsilometer (E-test) is both a diffusion and a dilution method of testing antibiotic sensitivity.

It is a quantitative technique for determining the antimicrobial susceptibility (AST) and minimum inhibitory concentration (MIC) in µg/mL of gram-negative and gram-positive aerobic bacteria.

Sold by @itachibot



The antibiotic sensitivity tests are classified as follows:

• Diffusion methods: Kirby–Bauer method, Stokes method

• Dilution methods: Tube dilution method, Agar dilution method

• Diffusion and dilution method: E–test (Epsilometer)

If you purchased this from someone else, you may have been scammed. Sold by @itachibot

Solution to Question 7:

The disc diffusion method for antibiotic susceptibility testing is applied in the Kirby-Bauer method.

To perform Kirby-Bauer's disc diffusion method, a suitable standard dilution of a broth culture of the test bacterium is inoculated on the surface of a solid medium (Cation-adjusted Mueller-Hinton agar) as a lawn culture. After drying, various antibiotic discs are placed on its surface. Zones of inhibition around the disc are recorded and interpreted according to the zone diameters available in the tables of the Clinical and Laboratory Standards Institute (CLSI) guidelines. The results obtained are qualitative (resistant, sensitive, or partially sensitive).

Laboratory testing for antibiotic sensitivity can be done by:

• Diffusion method- Kirby-Bauer disc diffusion method, Stokes disc diffusion method

• Dilution method- broth dilution method, agar dilution method

• Combination of diffusion and dilution method- E–test (Epsilometer)

The image below shows Kirby-Bauer disc diffusion method.

Sold by @itachibot

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed.

Solution to Question 8:

Minimum inhibitory concentration (MIC) cannot be calculated using Kirby-Bauer's disk diffusion method and Stokes disk diffusion method as they are semi-quantitative methods.

MIC is the lowest concentration of an antimicrobial agent that will inhibit the visible growth of a microorganism.

Methods to determine MIC include:

- Tube dilution method
- Agar dilution method
- E-test method

Solution to Question 9:

Real-time polymerase chain reaction (PCR) can be used to quantify the PCR by monitoring the amplification process while the PCR is ongoing.

Types of PCR:

- Reverse transcriptase PCR- The enzyme reverse transcriptase (RT) is used to synthesize a complementary copy of DNA (cDNA) from RNA and this cDNA is then used as a template for PCR.
- Real-time PCR- This helps quantify the PCR and gives an estimate of the pathogen DNA load in the specimen. It uses fluorescent dyes or specific DNA probes. It is also known as qPCR (Quantitative PCR)
- Multiplex PCR- Primers to more than one pathogen are used simultaneously and hence multiple products are obtained.

Sold by @itachibot

• Nested PCR- It involves the sequential use of two primer sets. The first primer set is used for amplification and the amplicon obtained is then used as a target sequence for second amplification. This ensures specificity.

Solution to Question 10:

The hemagglutination test is not based on the principle of antibody marking.

When antibodies bind to antigens on the surface of red blood cells, the resultant clumping reaction is referred to as hemagglutination. It is routinely performed to type red blood cells in ABO blood-typing.

The image below shows hemagglutination.



In ELISA, radioimmunoassay and immunofluorescence, the antigens or antibodies are labeled or marked with enzymes, fluorescent dyes, etc.

Note: If no hemagglutination occurs it means the patient's serum has anti-haemagglutinin antibodies that bind to hemagglutinin and prevent the clumping reaction. This hemagglutination inhibition reaction is a method of detection of viruses and antiviral-antibodies and is used to make a diagnosis of influenza.

Solution to Question 11:

Pour plate culture (quantitative test) is not used as an immediate test for screening of UTI. However, it is considered the best diagnostic test for confirmation of UTI due it's high sensitivity and specificity.

The screening tests for UTI are as follows:

• Wet mount examination- It is done to demonstrate pus cells in urine.

Sold by @itachibot

• Leukocyte esterase test- It is a rapid and cheaper method and detects leukocyte esterases secreted by pus cells present in urine.

• Nitrate reduction test (Griess test)- Nitrate reducing bacteria like E. coli gives a positive result.

• Catalase test- E. coli and other bacteria that produce the enzyme catalase give a positive result.

Quantitative Culture:

• Standardized loop technique

• Pour plate method

Solution to Question 12:

CLED (Cysteine lactose electrolyte-deficient) medium is preferred over MacConkey agar for the culture of organisms involved in urinary tract infections because it is less inhibitory and promotes the growth of gram-positive bacteria (except β hemolytic streptococcus) and candida.

It is used as an alternative to the combination of blood agar and MacConkey agar, for the processing of urine specimens.

CLED medium is not similar to blood agar in the fact that it can prevent swarming of proteus. It is similar to MacConkey agar in that it differentiates between lactose fermenters and non-fermenters.



Solution to Question 13:

From the given case scenario and image we can deduce that the most likely organism involved is Enterococcus faecalis.

The image shows gram-positive cocci in chains, that are described as catalase negative. This helps rule out Staphylococcoccus aureus and coagulase-negative staphylococcus.

Sold by @itachibot

Streptococcus viridans is usually sensitive to penicillin and vancomycin hence it is ruled out.

The mechanism of vancomycin resistance seen in E. faecalis is the alteration of the D-alanyl–D-alanine chain in the cell wall.

Solution to Question 14:

Bacillus anthracis which causes anthrax is categorized under category A bioterrorism agent.

Solution to Question 15:

The above image shows a HbsAg card test kit which works on the principle of immunochromatography.

The immunochromatographic membrane is used for antigen detection, while a nitrocellulose membrane is used to absorb the antigen from a specimen. A coloured reaction appears directly on the membrane with the sequential addition of conjugate followed by the substrate. In some cases, the antigen is captured by a bound antibody directed against the antigen. These assays have the advantage of being rapid and also frequently include a built-in internal control.

Sold by @itachibot
If you purchased this from someone else,
you may have been scammed

Sold by @itachibot